Dysregulation of M segment gene expression contributes to influenza A virus host

restriction

Brenda M. Calderon[1], Shamika Danzy[1], Gabrielle K. Delima[1], Nathan T. Jacobs[1], Ketaki Ganti[1],

Megan R. Hockman[1], Graeme L. Conn[2], Anice C. Lowen[1*], and John Steel[1#*]

Departments of Microbiology and Immunology[1] and Biochemistry[2], Emory University School of

Medicine, Atlanta, GA, USA

*Address correspondence to John Steel at john.steel@emory.edu or Anice Lowen at

anice.lowen@emory.edu

[#]Current address: Influenza Division, Centers for Disease Control and Prevention, Atlanta, GA,

USA

Abstract

The M segment of the 2009 pandemic influenza A virus (IAV) has been implicated in its emergence into human populations. To elucidate the genetic contributions of the M segment to host adaptation, and the underlying mechanisms, we examined a panel of isogenic viruses that carry avian- or human-derived M segments. Avian, but not human, M segments restricted viral growth and transmission in mammalian model systems, and the restricted growth correlated with increased expression of M2 relative to M1. M2 overexpression was associated with intracellular accumulation of autophagosomes, which was alleviated by interference of the viral proton channel activity by amantadine treatment. As M1 and M2 are expressed from the M mRNA through alternative splicing, we separated synonymous and non-synonymous changes that differentiate human and avian M segments and found that dysregulation of gene expression leading to M2 overexpression diminished replication, irrespective of amino acid composition of M1 or M2. Moreover, in spite of efficient replication, virus possessing a human M segment that expressed avian M2 protein at low level did not transmit efficiently. We conclude that (i) determinants of transmission reside in the IAV M2 protein, and that (ii) control of M segment gene expression is a critical aspect of IAV host adaptation needed to prevent M2-mediated dysregulation of vesicular homeostasis.

Author summary

Influenza A virus (IAV) pandemics arise when a virus adapted to a non-human host overcomes species barriers to successfully infect humans and sustain human-to-human transmission. To gauge the adaptive potential and therefore pandemic risk posed by a particular IAV, it is critical to understand the mechanisms underlying viral adaptation to human hosts. Here, we focused on the role of one of IAV's eight gene segments, the M segment, in host adaptation. Comparing the growth of IAVs with avian- and human-derived M segments in avian and mammalian systems revealed that the avian M segment restricts viral growth specifically in mammalian cells. We

show that the mechanism underlying this host range phenotype is a dysregulation of viral gene

expression when the avian IAV M segment is transcribed in mammalian cells. In particular,

excess production of the M2 protein results in viral interference with cellular functions on which

the virus relies. Our results therefore reveal that the use of cellular machinery to control viral

gene expression leaves the virus vulnerable to over- or under-production of critical viral gene

products in a new host species.

Introduction

Influenza A virus (IAV) epidemics and pandemics result in widespread and often severe disease

[1-3], as well as considerable societal economic costs [4]. IAV lineages endemic in humans are

ultimately derived from those circulating in wild waterfowl. Sporadic transmission of IAV from

avian reservoirs to humans occurs mainly through domesticated intermediate hosts such as

chickens and pigs [5]. For example, infection of humans with H7N9 and H5N1 subtype IAVs

occurs through direct exposure to infected poultry. Although they often cause severe disease,

these zoonotic cases have not led to sustained onward transmission to date and have therefore

not caused a pandemic. Despite abundant circulation of IAVs at the animal-human interface,

pandemics occur only rarely owing to the host specificity of IAV infection. As with all viruses,

IAV must exploit host cell functions and overcome antiviral barriers to execute its life cycle. Viral

replication in a new host species therefore requires adaptation to re-establish virus-host

interactions broken by species-specific features of host cellular processes. IAV can undergo

such adaptation through reassortment of intact gene segments between avian IAVs and those

already adapted to mammals or through more incremental changes brought about by

polymerase error [6]. With the goal of anticipating and mitigating IAV emergence in humans, it is

essential to understand the species-specific barriers to infection and the mechanisms by which

IAV evolution allows these impediments to be overcome.

Changes to viral receptor specificity and polymerase function necessary to overcome host

species barriers have been well documented [7-11], but it is clear that additional features of

avian IAVs restrict their growth in mammalian systems [12-14]. Seminal studies, performed in

the 1970s, indicated a potential role for the M segment. Avian IAVs were shown to express low

levels of M1 protein during abortive infection of mammalian cells, but not avian cells [15-17],

and the limited quantity of viral particles formed during such abortive infections correlated with

the amount of M1 protein expressed. Significantly, these studies were conducted prior to the

identification of a second gene product expressed from a spliced form of the M segment mRNA:

the M2 protein [18,19]. Subsequent studies demonstrated that low M1 expression during

abortive avian IAV infection in mammalian cells was accompanied by increased M2 mRNA and

protein expression, relative to productive infection in avian cells, leading to new hypotheses

regarding the mechanisms leading to abortive outcomes [20-22].

Additional evidence for a role of the M segment in host range comes from the *in silico*

identification of positive selection acting on the M segment of the Eurasian avian-like swine

lineage, following transfer of this IAV from birds to swine [23]. Moreover, others and we have

shown that the M segment of the 2009 pandemic H1N1 (pH1N1) strain carries determinants of

transmissibility, suggesting that adaptation of this segment to humans contributed to the

emergence of the pH1N1 lineage [24-28]. The mechanistic basis for this contribution; the

specific roles, if any, of M1 and M2 proteins to host range; and whether the M segment plays a

broader role in IAV host adaptation, remain unclear.

It is now recognized that the M segment of IAV encodes the M1 matrix and the M2 proton

channel proteins via alternative splicing (**Suppl. Figure 1A)** [29,30]. M1 forms a structural layer

under the viral envelope and is a major determinant of virion morphology [24,26,31,32]. Upon

endocytosis of virus particles into cells, low pH and high [K+] trigger a conformational change in

M1 required for release of viral ribonucleoprotein (vRNP) complexes into the cytoplasm [33]. M1

is also critical for the transport of newly replicated vRNPs from the nucleus to the cytoplasm [34-

37]. At late stages of infection, cytoplasmic M1 is thought to recruit vRNP complexes to the

budding site [38-40].

M2 is a transmembrane protein that tetramerizes to form a proton channel. Within the envelope

of incoming virions, this channel allows diffusion of protons into the virion core, thereby

triggering the release of vRNPs from M1, as well as facilitating the intracellular transport of HA

protein [41-44]. M2 also plays an important role in the final stages of virion morphogenesis,

mediating ESCRT-independent membrane scission [45]. Importantly for the findings reported

herein, multiple groups have reported that M2 blocks the fusion of autophagosomes with

lysosomes [46-48]. Under normal conditions, this fusion event allows degradation and recycling

of organelles and protein aggregates contained within autophagosomes [49], and is thus

important for cellular homeostasis. M2-mediated interference with autolysosome formation

therefore stalls autophagic flux and leads to the accumulation of autophagosomes within the cell

[46-48].

Many elegant molecular virology studies have revealed that viral strategies for regulating gene

expression are central to orchestration of the viral life cycle [50-53]. Here, we report that

regulation of M2 expression of IAV is an important determinant of host range. In particular, our

data show that the expression of M1 and M2 from alternatively spliced M segment mRNAs is

dysregulated when an avian IAV replicates in mammalian cells, and that the resultant

overexpression of the M2 proton channel attenuates viral growth by disrupting the cellular

autophagy pathway. The observed block in autophagy is consistent with that reported previously

for lab-adapted strains of IAV [46-48]. Importantly, however, we see that lab-adapted IAVs

express M2 at high levels, comparable to those seen for avian IAV M segments in mammalian

cells. In contrast, IAVs carrying M segments well adapted to mammalian systems express

relatively little M2 and do not trigger the accumulation of autophagosomes to the same extent as

avian-origin and lab-adapted strains. These findings suggest that IAV-induced stalling of

autophagic turnover is a result of aberrantly high M2 expression upon infection of mammalian

cells with avian IAVs, and which is overcome in human-adapted influenza A lineages. Critically,

the mechanism leading to this host-restricted phenotype occurs at the level of gene expression

and is driven by host-specific differences in viral nucleic acid sequence, not amino acid changes.

This mechanism constitutes a novel paradigm in RNA virus host adaptation, and reveals a new

species barrier for IAV, which may be highly relevant for the emergence of avian IAVs into

humans.

Results

Generation of isogenic IAVs differing only in the M segment

To assess the contribution of the M segment to host adaptation, we generated a panel of IAVs

possessing human- or avian-derived M segments in the A/Puerto Rico/8/34 (H1N1) [PR8]

background (**Table 1; Figure 1A; Suppl. Figure 1B, C, D**). The M segments used were

selected with the aims of representing circulating human IAVs and capturing the breadth of

diversity of M segment sequences present in wild waterfowl. In addition to three wild-type avian

M segments, we included in our panel a mutant of the dk/Alb/76 M segment that carries a

uniquely identified mutation in M2 (S89). Reversion of this unique amino acid residue to glycine

yields an M2 protein matching the consensus amino acid sequence of IAVs in wild bird

reservoirs. The human-adapted M segment analyzed most extensively herein is that of the

pandemic A/NL/602/09 (H1N1) strain [NL09].

Table 1. Origin of IAV M segments used in study

Strain	Abbreviation	% Nucleotide identity to avian H1N1 subtype consensus	% M1 a.a. identity to avian consensus (no. of differences)	% M2 a.a. acid identity to avian consensus (no. of differences)
A/NL/602/2009 (H1N1)	NL09	91.7	96.4 (9)	92.8 (7)
A/Panama/2007/99 (H3N2)	Pan99	91.0	95.6 (11)	85.6 (14)
A/Bethesda/55/2015 (H3N2)	Beth15	90.5	95.6 (11)	83.5 (16)
A/mallard/MN/19906/99 (H3N8)	MallMN99	98.9	100	97.9 (2)

A/rhea/North Carolina/39482/93 (H7N1)	RhNC93	98.7	100	100
A/duck/Alberta/35/76 (H1N1)	DkAlb76	96.6	100	99 (1)
A/duck/Alberta/35/76 (H1N1) M2 89G	DkAlb76 89G	96.7	100	100

140   a.a.: Amino acid residue.

IAV M segments of avian origin confer a host-restricted phenotype

The relative fitness of PR8-based viruses carrying M segments from the human-adapted NL09

virus or from avian IAVs was evaluated in three avian substrates: embryonated chicken eggs

(ECE), chicken origin DF-1 cells, and quail origin QT-6 cells. When grown from low multiplicities

of infection (MOI) in either ECE or DF-1 cell substrates, all viruses exhibited similar kinetics and

reached comparable titers (**Figure 1B and 1C**). Also, under high MOI growth conditions in DF-1

and QT-6 cells, the growth phenotypes of the PR8 NL09 M and PR8 avian M viruses were

similar, although the NL09 M-containing virus trended towards increased growth relative to the

avian M-containing viruses in DF-1 cells (**Figure 1D and 1E**). Nonetheless, at no time did the

NL09 M-encoding virus grow to significantly higher titers than the avian M-encoding viruses in

any avian host substrate. In contrast, in mammalian cells, human-adapted IAV M segments

were found to support improved growth relative to avian-adapted M segments. Viral growth was

monitored from low and high MOI in human A549 cells and canine MDCK cells (**Figure 2**), and

additionally from high MOI in human 293T cells (**Suppl. Figure 2A**). In each case, PR8-based

viruses carrying the NL09 M segment grew with more rapid kinetics and to higher titers than

isogenic viruses carrying avian M segments, although the differences in growth from low MOI

did not reach statistical significance. To extend our findings to an independent human lineage of

IAV, we evaluated the growth from high MOI in A549 cells of PR8 Pan99 M and PR8 Beth15 M

viruses, each of which have an M segment derived from the human seasonal H3N2 lineage

(**Suppl. Figure 1C, D**). We found that both of these human-adapted M segments supported

significantly faster kinetics and higher magnitude of growth than the dkAlb76 89G M segment

(**Suppl. Figure 2B**). Thus, results from multiple different avian and mammalian culture systems

reveal defects in the kinetics and magnitude of viral growth associated with avian IAV M

segments specifically in mammalian host cells and provide evidence of the acquisition of

adaptive changes in the M segment of pH1N1 and H3N2 human influenza A lineages.

Avian IAV M segments confer lower viral infectivity, growth and transmission than NL09

M segment in a guinea pig model

To evaluate the impact of M segment host adaptation on viral phenotypes in an intact host, we

evaluated the PR8 NL09 M virus and the four PR8-based viruses carrying avian M segments in

a guinea pig model [54,55]. A low inoculation dose of 10 PFU per animal was used to maximize

the sensitivity of the system. Each inoculated animal was co-caged with a naïve guinea pig at

24 h post-inoculation and viral infectivity, magnitude and kinetics of growth, and transmission

were monitored by collecting nasal lavage samples every other day. Among those animals that

were productively infected, total virus sampled in nasal washes over the course of infection was

evaluated by calculating the area under the shedding curves shown in **Figure 3A**. Area under

the curve values did not differ significantly among the four viruses with avian M segments, but

were significantly higher for PR8 NL09 M virus (**Figure 3A**). As shown in **Figure 3B**, fewer

guinea pigs were productively infected with 10 PFU when the inoculating virus carried an avian

M segment. Differences in the kinetics of viral growth were determined by assessing changes in

growth over time using repeated measures ANOVA (**Suppl. Figure 3**). Each virus encoding an

avian host-derived M segment replicated with significantly slower kinetics than the PR8 NL09 M

virus.

The efficiency of transmission was assessed by calculating the proportion of contact animals

that contracted infection, considering only cages in which donor guinea pigs were productively

infected through intranasal inoculation. While transmission occurred in 7/9 instances for PR8

NL09 M virus, a significantly lower proportion of animals (1/20; $p<0.0001$) were infected with a

virus carrying an avian M segment transmitted from her cage mate (**Figure 3B**). Thus, data from

a mammalian animal model reveal defects in infectivity, growth and transmission associated

with avian IAV M segments. Our results suggest the acquisition of adaptive changes in the M

segment of pH1N1 human IAV since its introduction from the avian reservoir into mammals.

The M2 proton channel is aberrantly overexpressed from avian IAV M segments in

mammalian cells

To investigate the mechanism underlying the host-restricted phenotypes conferred by avian-

adapted M segments, the steady state levels of the M1 and M2 proteins were evaluated in

human A549 and avian DF-1 cells infected at high MOI. Western immunoblotting with the

monoclonal antibody E10, which recognizes the shared N-terminus of M1 and M2, was used to

monitor levels of both proteins concurrently.

In the avian cell line, the PR8 NL09 M virus expressed lower levels of M2 than M1 (**Figure 4A**).

When M1 and M2 levels were normalized to those of vinculin and the relative levels of M1 and

M2 calculated, the NL09 M2 protein comprised <10% of the total M segment-derived protein

expression. Very similar results were seen for four PR8-based viruses carrying avian M

segments grown in DF-1 cells. In each case, M2 comprised <20% of the total M segment-

derived protein, and in the case of three avian M segment-containing viruses (PR8 dkAlb76 M;

PR8 dkAlb76 M 89G; and PR8 rheaNC93 M), the levels of M1 and M2 proteins were not

significantly different from those expressed by the PR8 NL09 M virus (**Figure 4B, 4C, 4D**).

In A549 cells, the PR8 NL09 M virus expressed M2 at ~20% of the total M protein, similar to the

expression levels observed in infected DF-1 cells (**Figure 5A**). In contrast, avian M segments

yielded markedly higher levels of the M2 protein than were observed for the same viruses grown

in avian cells. Here, M2 levels were approximately equal to those of M1 (**Figure 5B, 5C, 5D**).

Moreover, when compared to the NL09 M segment in human cells, levels of M2 protein were

significantly higher for avian M segments (P<0.0001) (**Figure 5C, 5D**). Thus, compared to

matched virus-host pairings in both human and avian systems, M2 was markedly

overexpressed when avian-adapted M segments were introduced into human cells.

Overexpression of M2 protein in mammalian cells corresponds to increased levels of the

M2 mRNA (mRNA 10)

The M segment vRNA is transcribed by the viral polymerase to give rise to a co-linear mRNA

($mRNA_7$), which can be spliced by cellular splicing factors to yield $mRNA_{10}$ or $mRNA_{11}$ [18,56].

The M1 protein is translated from the unspliced $mRNA_7$, while M2 is expressed from the spliced

$mRNA_{10}$. To date, no polypeptide corresponding to the short $mRNA_{11}$ ORF has been identified.

Aberrantly high levels of IAV $mRNA_{10}$ in mammalian cells have been previously reported for the

avian IAV, fowl plague virus [20,21]. We therefore hypothesized that the high expression levels

of M2 protein, observed for avian-adapted M segments in human cells, was due to increased

splicing of $mRNA_7$ to yield excessive $mRNA_{10}$. We used a RT primer extension assay to quantify

the levels of each M segment-derived mRNA species in DF-1 and A549 cells infected with either

PR8 NL09 M virus or one of four viruses carrying an avian-adapted M segment.

As expected, based on the levels of the encoded proteins, $mRNA_{10}$ was of low abundance

compared to $mRNA_7$ for all viruses in DF-1 cells (**Figure 6A**). Quantification of band intensities

revealed that, in DF-1 cells infected with PR8 NL09 M virus, $mRNA_7$ comprised ~75%, while

$mRNA_{10}$ comprised ~5% of the total M segment-derived mRNA (**Figure 6B**). For three avian M

viruses (PR8 dkAlb76 M; PR8 dkAlb76 M 89G; and PR8 rheaNC93 M), relative levels of $mRNA_7$

were ~70-75% and were not significantly different from those of PR8 NL09 M virus (**Figure 6B**).

For all viruses, $mRNA_{10}$ was expressed at low levels (10-40% of total segment 7 mRNA), and in

the case of two avian M segment-containing viruses (PR8 dkAlb76 M; PR8 dkAlb76 M 89G), the

levels of $mRNA_{10}$ were not significantly different from those expressed by the PR8 NL09 M virus

(**Figure 6C**). $mRNA_{11}$ was expressed in each infection at approximately 20% of the total M

segment-derived mRNA, and the relative level did not differ significantly with any virus (**Figure**

6D). Overall, the relative levels of each of the three mRNA species were similar when

synthesized from avian- or human-origin encoded M segments in DF-1 cells, correlating with

Western immunoblot data on M segment protein expression in these cells.

In A549 cells, the PR8 NL09 M virus showed relatively high abundance of $mRNA_7$ and low

abundance of $mRNA_{10}$, as was seen in DF-1 cells (**Figure 7A**). By contrast, for viruses carrying

avian-adapted M segments, $mRNA_{10}$ levels exceeded those of $mRNA_7$ in A549 cells (**Figure 7B,**

7C). Both decreases in $mRNA_{10}$ levels and increases in $mRNA_7$ levels contributed to the altered

relative abundance. Again, no significant differences in relative $mRNA_{11}$ levels among the

viruses were noted (**Figure 7D**) suggesting that changes in splicing to produce $mRNA_{11}$ do not

contribute to changes in M segment mRNA and protein expression observed in A549 cells.

Overall, our data point to increased splicing to produce $mRNA_{10}$ as the underlying mechanism

leading to heightened M2 expression from avian M segments in A549 cells, and suggest that

cis-acting signals present on the M segment are responsible for driving the disparate M2

expression levels.

M2 expressed from avian M segments localizes in perinuclear puncta

To gain initial insight into the phenotypic consequences of M2 overexpression, we examined the

subcellular localization of M2 in A549 cells infected with PR8 NL09 M virus or one of four PR8-

based viruses with avian M segments. At 8 hours-post-infection (hpi) in PR8 NL09 M virus-

infected cells, M2 showed mainly dispersed cytoplasmic staining, along with weak cytoplasmic

membrane localization and some perinuclear accumulation. By contrast, in cells infected with

viruses that encoded any of the four avian origin M segments, we noted more abundant M2

protein staining localized at the plasma membrane, and in large puncta near the cell nucleus

(**Figure 8A**). Similar data were obtained from virus-infected 293T cells at 8 hpi (**Figure 8B**) and

from both human cell lines at 12 hpi (**Suppl. Figure 6**). This staining pattern, along with data in

previously published studies [46-48], prompted us to investigate whether highly expressed M2

protein was interacting with the cellular autophagy pathway.

Human cells exhibit increased LC3B II levels when infected with IAV carrying avian M segments

An interaction between the IAV M2 protein and the cellular autophagy pathway has been reported previously [46-48]. To test whether M2 expression from avian vs. human M segments has differential effects on autophagy, we first evaluated the level of LC3B lipidation in PR8 NL09 M- and PR8 avian M-infected cells. Upon activation of the autophagy pathway, LC3B is modified through the addition of phosphotidylethanolamine [57,58]. This lipidated form, called LC3B II, remains associated with the membranes of maturing and mature autophagosomes. Thus, increased levels of LC3B II in the cell are indicative of autophagosome formation; however, under normal conditions LC3B I and II are cycled through an autophagic flux which involves autophagosome maturation and degradation via fusion with lysosomes [59]. A block in this fusion event can therefore lead to accumulation of autophagosomes in the cytoplasm and, consequently, accumulation of LC3B II in the cell.

Western immunoblot analysis of LC3B revealed decreased LC3B I and increased LC3B II protein levels in virus-infected cells compared to mock, providing evidence of autophagy activation and/or block during IAV infection of either avian or human cells (**Figure 4A, 4E, 4F; 5A, 5E, 5F**). In avian DF-1 cells, levels of LC3B I were slightly decreased and levels of LC3B II were slightly increased over mock, but LC3B I and LC3B II levels were generally comparable among human M- and avian M-containing viruses (**Figure 4E, 4F**). Importantly, human A549 cells infected with PR8 avian M-containing viruses consistently showed lower levels of LC3B I, and higher levels of LC3B II than A549 cells infected with PR8 NL09 M as well as other human M-containing viruses (**Figure 5E, 5F).** Additionally, the same phenotypes were obtained from the panel of viruses upon infection of human origin 293T cells (**Suppl. Figure 4A-4F**), or canine origin MDCK cells (**Suppl. Figure 4G-4L**).

Taken together, these data reveal a correlation between levels of M2 expression and LC3B

lipidation (accumulation of LC3B II), suggesting that high levels of M2 present in PR8 avian M-

infected cells may be inducing an over-activation of autophagy, or a block in the turnover of

autophagosomes.

Infection of human cells with IAV carrying avian M segments triggers accumulation of

autophagosomes

To confirm that M2 protein was co-localizing with LC3-positive autophagic vesicles in

mammalian cells, we monitored sub-cellular localization of an overexpressed LC3B-GFP fusion

protein in infected cells. Human 293T cells were primarily used for this purpose as they

transduce well with GFP-LC3. Under normal conditions, LC3B-GFP can be seen throughout the

cell with a diffuse distribution (**Figure 9, row 1: mock infected cells**). Treatment with

chloroquine, an inhibitor of autophagy that decreases autophagosome-lysosome fusion, results

in the concentration of LC3B-GFP into punctate cytoplasmic structures (**Figure 9, row 7: CQ-**

treated cells). This relocalization is indicative of the accumulation of autophagosomes [60]. In

LC3B-GFP expressing 293T cells infected with PR8 NL09 M virus, we observed little to no

increase in GFP-positive cytoplasmic puncta (**Figure 9, row 2: NL09 M infected cells**). In cells

infected with PR8 avian M-encoding viruses, by contrast, extensive accumulation was apparent,

with both the size and number of intracellular vesicles increased relative to PR8 NL09 M

infection (**Figure 9, rows 3-6: Avian M infected cells**). In addition to these cytoplasmic puncta,

viruses with avian M segments were found to redirect a small proportion of LC3B-GFP to the

plasma membrane. Notably, the M2 protein co-localized with LC3B-GFP in PR8 avian M-

infected cells, both in cytoplasmic puncta and at the plasma membrane. Similar phenotypes

were observed in virus-infected A549 cells, at 8 hpi **(Suppl. Figure 7)**. Overall, these results

suggest that expression of avian M2 protein at high levels interferes with the turnover of

autophagosomes and with the localization of LC3B-GFP. In addition, the results reveal a

reduction in the tight perinuclear localization of M2 upon LC3B overexpression.

As an additional test designed to differentiate autophagy induction from autophagy block in cells

infected with IAV carrying avian M segments, we used chloroquine treatment followed by

Western immunoblotting for LC3B I and LC3B II. We hypothesized that, if autophagic flux is

blocked in cells infected with PR8 avian M viruses, then treatment with chloroquine would not

lead to a further increase in LC3B II levels. Conversely, if virus infection of these cells induces

autophagy, then blocking their turnover with chloroquine would result in heightened LC3B II over

mock-infected, chloroquine-treated cells. As shown in **Suppl. Figure 8A and 8C**, we observed a

significant difference in the activation of LC3B II following infection with human- or avian M-

encoding viruses. This difference in LC3B II activation is lost in the presence of chloroquine.

Additionally, as we observed no change in LC3B II levels upon treatment of virus-infected cells

with chloroquine, relative to chloroquine-treated, mock-infected cells, we conclude that the high

levels of IAV M2 protein expressed from avian M segments, but not human M segments,

precipitate a block in the autophagy pathway such that chloroquine treatment cannot bring

about further accumulation of autophagosomes.

Ion channel activity of IAV M2 contributes to LC3B II accumulation in A549 cells

To evaluate the contribution of IAV M2 ion channel activity to the observed stalling of autophagy

in PR8 avian M virus-infected cells, we inhibited channel activity with amantadine [61].

Amantadine was added at 1 hpi and the level of LC3B lipidation was assessed at 8 hpi in PR8

NL09 M and PR8 avian M virus-infected A549 cells. The timing of amantadine addition was

designed to avoid disruption of the early function of M2 in delivery of the viral genome to the

cytoplasm, but allow inhibition of M2 activity in autophagosomes.

As noted above (**Figure 5; Suppl. Figure 4**), Western immunoblot analysis of LC3B revealed

decreased LC3B I and increased LC3B II upon infection of A549, 293T or MDCK cells, with

avian or human M-encoding IAV, with the effect being stronger for viruses possessing avian M

segments. Interestingly, addition of amantadine reversed the intracellular accumulation of LC3B

II in both human and avian M-infected A549 cells (**Figure 10**). Quantitation of M segment-

encoded proteins by Western immunoblot analysis showed no difference in the expression

levels of M1 or M2 protein upon treatment with amantadine (**Figure 10 A, 10B, 10C, 10D**), but

revealed a statistically significant increase in LC3B I (**Figure 10E**) and decrease in LC3B II

(**Figure 10F**) to levels comparable to mock-infected cells.

To ensure that the amantadine was working to prevent LC3B II accumulation through inhibition

of M2 ion channel activity, we assessed the impact of amantadine treatment on the inhibition of

phagosome-lysosome fusion by chloroquine. Amantadine had no impact on LC3B II

accumulation in the absence of M2 protein (**Figure 10G-I**), indicating that the mechanism of

amantadine block of autophagosome accumulation was specifically mediated through

interference with M2 function.

As a complementary approach to monitor the effects of amantadine, we again used

overexpressed LC3B-GFP fusion protein in infected 293T cells at 12 hpi. Using confocal

microscopy, we tracked LC3B-GFP localization in the presence and absence of the ion channel

blocker. In line with the Western immunoblot analyses obtained in A549 cells, we saw that

amantadine reduced the accumulation of LC3B-GFP within cytoplasmic puncta and at the

plasma membrane of infected cells (**Figure 11**). Of note, amantadine treatment did not alter

levels or localization of M2 protein and, moreover, had no detectable effect on LC3B-GFP

appearance in chloroquine-treated cells (**Figure 11**).

Overall, our results indicate that the intracellular ion channel activity of IAV M2 protein is

necessary to mediate the observed block in autophagy, and that addition of amantadine can

reverse the autophagic phenotypes induced by both human and avian M-encoding IAVs.

Improved viral growth in the presence of amantadine implicates autophagy block in viral growth restriction

To test for a causal link between the observed block in autophagy and the restriction of growth seen for PR8 avian M-encoding viruses in mammalian cells, we monitored the effect on viral growth of relieving the autophagy block with amantadine treatment. Specifically, we evaluated single-cycle growth in A549 cells using PR8 NL09 M and PR8 dkAlb76 M 89G viruses. Amantadine was again added at 1 hpi to avoid disruption of the earliest steps of the viral life cycle. While amantadine had no effect on growth of the PR8 NL09 M virus (P=0.271), a modest but consistent increase in the titer of PR8 dkAlb76 M 89G virus was noted throughout the time-course (P=0.0007) (**Figure 12**). This result indicates that M2 ion channel-induced disruption of vesicular homeostasis contributes to the host restriction conferred by avian IAV M segments.

Viral growth restriction and autophagy block are linked to M2 expression levels, not amino acid sequence

To determine whether the observed effects of avian-origin M segments on the viral life cycle and the autophagy pathway are attributable to avian-adapted amino acid sequences or rather to the overexpression of the viral proton channel, we generated a series of chimeric M segments in which the coding changes found in the NL09 M segment were transferred to an avian-adapted RNA background, and vice versa. We introduced all 16 coding changes into M1 and M2 simultaneously, or introduced only the nine or seven changes into M1 or M2, respectively. The genotypes of this panel of six chimeric viruses are outlined in **Table 2** and **Suppl. Figure 9**.

Table 2. Genotypes of avian-human chimeric M segments

	Nucleic acid background	M1 amino acid sequence	M2 amino acid sequence
NL M 16mut	NL09 M	Avian consensus M1	Avian consensus M2
NL M 9mut	NL09 M	Avian consensus M1	NL09 M2
NL M 7mut	NL09 M	NL09 M1	Avian consensus M2
Av M 16mut	Dk Alb 76 M 89G	NL09 M1	NL09 M2

| Av M 9mut | Dk Alb 76 M 89G | NL09 M1 | Avian consensus M2 |
| Av M 7mut | Dk Alb 76 M 89G | Avian consensus M1 | NL09 M2 |

Western immunoblot analysis of M1 and M2 proteins expressed from these six chimeric M

segments in infected A549 cells is shown in **Figure 13A**. Among the viruses with the NL09 M

nucleic acid background, the PR8 NL M 16mut and PR8 NL M 9mut viruses both showed

increased levels of M2 relative to the PR8 NL09 M virus control, indicating that all or some of

the coding changes introduced into the M1 ORF impact regulation of M2 gene expression. The

PR8 NL M 7mut virus showed similar M1 and M2 expression levels as the PR8 NL09 M control,

however. This virus therefore expresses the avian consensus M2 protein at low levels. All

viruses with the nucleic acid background of the avian M segment showed relatively high levels

of M2, similar to those seen for the PR8 dkAlb76 M 89G virus control. Thus, in the avian nucleic

acid background, we were able to fully separate M2 expression level from M2 and M1 amino

acid sequence.

Single-cycle growth analysis of the chimeric viruses in A549 cells revealed a strong correlation

between M2 expression level and viral yields (**Figure 14A, B and C**). Notably, all viruses with

the dkAlb76 M 89G-derived nucleic acid background, which express high levels of M2 protein,

grew to approximately 10-fold lower titers than the PR8 NL09 M virus, despite expression of

NL09 M1 and/or M2 proteins. In addition, the PR8 NL M 7mut virus, which expresses the human

M1 and avian M2 proteins at levels typical of human-adapted strains, showed equivalent growth

to that seen with the PR8 NL09 M virus, indicating that amino acid changes in M2 are not

required for efficient replication in human cells. Notably, the PR8 Av M 9mut virus, which

encodes identical M1 and M2 proteins as PR8 NL M 7mut virus (**Suppl. Figure 9**), exhibits

approximately 10-fold reduced viral replication relative to PR8 NL M 7mut virus (P=0.0003)

(**Figure 14B and C**). This result confirms that overexpression of M2 protein is a dominant

negative trait that is deleterious to virus growth.

Examination of LC3B lipidation by Western immunoblot analysis also revealed a clear

correlation between M2 expression levels and those of LC3B II within infected cells (**Figure**

13A). Importantly, overexpression of either human or avian M2 during infection boosted LC3B II

levels (compare lanes 2, 3, 5, 6, 7 and 8 in **Figure 13A**), while expression of avian M2 at levels

comparable to those of PR8 NL09 M virus gave low LC3B II levels (compare lanes 1 and 4 in

Figure 13A). Confocal images of 293T cells transduced with LC3B-GFP and infected with the

chimeric viruses were consistent with these findings. As seen for wild-type avian M segments,

high M2 expression again led to the accumulation of LC3B-GFP in perinuclear puncta and at the

plasma membrane (**Suppl. Figure 10**). Thus, as with viral growth, the accumulation of

autophagosomes within infected cells appears to be triggered by high levels of M2, regardless

of amino acid sequence.

To verify that the levels of M1 and M2 protein exhibited by the chimeric M segment-encoding

viruses are modulated at the level of mRNA expression, we conducted RT primer extension

assays. These data confirmed that differences in the levels of segment 7 mRNA expression

(**Suppl. Figure 11**) mirrored the observed differences in M1 and M2 protein expression levels.

Overall, these data support the premise that splicing signals are at least partly encoded in non-

synonymous nucleotide changes residing in the M1 ORF.

Chimeric M segment expressing NL09-like levels of NL09 M1 and avian M2 supports

robust growth *in vivo*

To evaluate whether correcting expression of the avian M2 protein to levels seen for viruses

with human-adapted M segments yielded a well-adapted phenotype in a mammalian animal

model, we evaluated growth and transmission of PR8 NL09 M 7mut virus in guinea pigs.

Titration of nasal wash samples revealed that the PR8 NL09 M 7mut virus (which expresses

avian M2 at low levels) had a similar growth phenotype to that of PR8 NL09 M virus: growth

kinetics and area under the curve values were not significantly different (**Figure 15; Suppl.**

Figure 12). In contrast, analysis of nasal wash samples collected from contact animals

indicated that only one of eight inoculated guinea pigs transmitted the PR8 NL09 M 7mut virus

to her cage mate. This level of transmission is significantly decreased relative to the PR8 NL09

M virus (P=0.0339) (**Figure 3B**). Taken together, these results indicate that, when expressed at

levels typical for a human-adapted M2 protein, an avian M2 supports efficient replication but

inhibits transmission in a mammalian host.

Lab-adapted IAV, but not recent human isolates, express abundant M2 and block

autophagy in human cells

The M2-induced block in autophagic flux reported previously was based on multiple IAVs that

were originally isolated from human hosts but passaged extensively in laboratory substrates

[46-48]. To assess whether these prior data were consistent with our own, we compared M2

expression levels exhibited by one of the lab-adapted strains used, PR8. Our results show that

the relative expression levels of M1 and M2 in PR8-infected 293T cells are comparable to those

seen in cells infected with avian M-encoding viruses (**Suppl. Figure 13**). These results suggest

that prior reports of M2-induced block in autophagy reflect a feature of mal-adaptation to the

host cell.

Discussion

Our data indicate that dysregulation of gene expression from the IAV M segment contributes to

the limited replicative capacity of avian IAVs in mammalian hosts. When transcribed within

mammalian cells, avian IAV M segments gave rise to abundant mRNA$_{10}$ and correspondingly

high levels of the encoded M2 proton channel. Comparison to matched virus-host systems

indicated that this M2 expression profile was an aberrant feature of avian IAV replication in

mammalian cells. When present at high levels, M2 had marked effects on the infected cell,

blocking the turnover of autophagic vesicles and redirecting LC3B II, the activated form of a

critical autophagy mediator, to the plasma membrane. These effects were found to rely on the

proton channel activity of M2 and occurred when either the avian or human M2 protein was

overexpressed in infected cells. Importantly, the reduction in viral growth conferred by avian M

segments in mammalian systems could be attributed at least in part to disruption of vesicular

homeostasis and was fully attributed to the altered expression levels of M1 and M2. Thus, our

data identify the regulation of viral gene expression as a novel host-dependent feature of the

IAV lifecycle that contributes to the host range restriction of this virus (**Figure 16**).

Quantification of IAV encoded mRNAs within infected cells indicated that overexpression of M2

from avian IAV M segments stems from over-production of the corresponding mRNA. M2 is

encoded by the $mRNA_{10}$ transcript, which is generated through splicing of the M1 message,

$mRNA_7$ [18,56]. Thus, regulation of M2 gene expression is intimately linked to the cellular

splicing machinery. Our data indicate that this is a finely balanced interaction between IAV and

its host cell, which is susceptible to disruption upon transfer to new species. In this regard, it is

notable that the M segment splice donor and acceptor sites are identical between the human

and avian IAVs examined. M segment sequences outside of these canonical splice sites have,

however, been shown to modulate splicing efficiency [62-65]. Interestingly, $mRNA_7$ association

with mammalian SF2/ASF [64], and with mammalian hnRNPK and NS1-binding protein [65],

have been shown to alter the efficiency of splicing. Nucleotide differences in these binding

regions exist between human- and avian-adapted M segments. An interesting area of further

investigation will be to test whether there are differences in the binding of human and avian

$mRNA_7$ molecules to mammalian SF2/ASF, hnRNPK, and NS1-binding proteins.

The gene segments of IAVs circulating in humans are ultimately derived from the avian IAV

gene pool. Thus, the contrasting phenotypes conferred by avian- and human-adapted M

segments in mammalian cells suggest that a resetting of M segment gene expression to reduce

M2 levels has been positively selected following emergence of avian IAVs into mammalian

populations. Indeed, our data reveal that this adaptation has occurred in at least two

independent incidences: i) within the Eurasian avian-like swine lineage M, that emerged from

birds in the late 1970s and contributed the M segment to the pH1N1 virus prior to 2009; and ii)

within the human seasonal lineage derived from the 1918 pandemic, which contributed its M

segment to the H3N2 lineage represented by Pan99 virus. The timing of the adaptive changes

is unclear in each case and cannot be inferred from our data. Importantly, M2 expression in

human IAV mimics that of avian IAV in avian cells. Thus, restriction of splicing efficiency to

achieve low M2 levels appears to be an important feature of IAV host adaptation. Our data

furthermore suggest that the negative impact of an autophagy block on viral growth constitutes

an important pressure driving this selection.

To date, most publications related to autophagy in IAV-infected cells have used the PR8 or

WSN lab-adapted strains, the X31 variant of PR8, or avian IAVs [46,47,66,67]. We found that

PR8 virus expresses levels of M2 in mammalian cells that are comparable with avian IAVs. This

phenotype may be a result of adaptation to growth in chicken eggs. Owing to the use of viruses

carrying lab-adapted M segments, the current literature suggests that IAV infection routinely

induces a block in autophagy [46,47,66,67]. We also see evidence of such a block when using

viruses that express high levels of M2, but we propose that this phenotype is a symptom of poor

adaptation to the mammalian host cell. Analysis of viruses carrying human M segments in

mammalian cells suggests that low levels of M2 and activation, but not blocking, of autophagy

are the norm in IAV-infected cells.

While the previously reported block in autophagy induced by IAV has generally been interpreted

as a viral defense against an antiviral mechanism, a pro-viral role for autophagy is not without

precedent. In Dengue virus infection, autophagy has been reported to support viral replication.

Specifically, autophagic turnover of lipid droplets in infected cells is thought to provide energy

needed for viral growth and to supply phospholipids important during virion assembly [68-70].

For IAV, the availability of activated LC3B protein was reported to be important for viral

morphogenesis [46]. Specifically, M2 was implicated in recruitment of LC3B II to the plasma

membrane, and disruption of this recruitment was found to affect budding and reduce stability of

released virions [46]. We also noted strong redistribution of LC3B-GFP to the plasma

membrane during infection with viruses that express high levels of M2. This relocalization

occurred to a lesser extent in a well-matched IAV-host system (NL09 M in human cells), and

may play a functionally important role in this context. A role for LC3B II in morphogenesis is

consistent with the activation of autophagy having a pro-viral effect. Conversely, our data

indicate that a block in autophagic turnover reduces viral growth, potentially owing to disruption

of vesicular trafficking that is normally exploited by the virus [71-73], or due to disruption of

cellular metabolic functions on which the virus relies.

By inhibiting the proton channel activity of the avian M2 with amantadine, we found that a

functional channel is needed to bring about the observed effects on autophagy. Indeed, M2 has

been shown previously to block the fusion of autophagosomes with lysosomes in a proton

channel-dependent fashion [66]. Surprisingly, however, amantadine treatment was also found to

reduce LC3B II accumulation in PR8 NL09 M-infected cells. The pH1N1 M2 protein carries the

S31N mutation, which confers resistance to amantadine. Our data suggest that this resistance

is not effective at a relatively high concentration of 200 μM amantadine, at least in the context of

intracellular viral infection.

The mutagenesis of the M segment that we carried out with the goal of separating the effects of

M2 expression level from M2 amino acid sequence revealed an important constraint on M1

coding capacity. Namely, the human / avian amino acid polymorphisms introduced into the M1

ORF were found to impact regulation of M2 gene expression. Some or all of these non-

synonymous changes to M1 likely modulate the efficiency of $mRNA_7$ splicing. With this new

insight in mind, a fresh look at viral phenotypes previously attributed to M1 amino acid variations

is warranted. Effects of M1 amino acid sequence on virion morphology [25,31,32,74,75], for

example, may in fact be directly or indirectly mediated through changes in intracellular M2

expression level.

Notably, while an avian M2 protein was sufficient to support robust viral replication in

mammalian systems when expressed at low levels, this sufficiency did not extend to

transmission. Virus expressing the avian M2 from a human M segment transmitted poorly,

indicating that the IAV M2 carries viral determinants of transmission and that these determinants

are not well conserved between avian and human strains. This novel observation also supports

the notion that high replicative capacity of a virus in a given host species is not necessarily

sufficient for transmission between individuals. In this context, it will be interesting to evaluate

the impact of avian M2 amino acid signatures on virion stability in the environment.

In summary, our data suggest that adaptive change in the M segment is needed to maintain low

expression of M2 in mammalian cells and that, in the absence of such adaptive changes,

excess M2 limits viral growth by blocking the autophagy pathway. These findings establish

regulation of viral gene expression as a feature of the virus-host interface critical for IAV host

range determination and therefore emergence into novel host populations.

Materials and Methods

<u>Ethics Statement</u>

The Institutional Animal Care and Use Committee (IACUC) of Emory University approved the

study protocol under approval number PROTO201700595.

<u>Cells</u>

Madin-Darby Canine Kidney (MDCK) cells (a kind gift of Peter Palese, Icahn School of Medicine

at Mount Sinai) were maintained in minimal essential medium (Gibco) supplemented with 10%

fetal bovine serum (FBS) and penicillin-streptomycin. A549 and 293T cells were obtained from

the ATCC and maintained in Dulbecco's minimal essential medium (DMEM; Gibco)

supplemented with 10% FBS and penicillin-streptomycin. DF-1 cells were obtained from ATCC

and maintained in DMEM plus 5% FBS and penicillin-streptomycin. QT-6 cells were obtained

from ATCC and maintained in F-12K medium plus 5% FBS, 10% tryptose phosphate and

penicillin-streptomycin. All cells were incubated at 37^{o}C with 5% CO_2.

Viruses

All viruses used in this work were generated using reverse genetics techniques [76]. The rescue

system for PR8 was a gift of Peter Palese (Icahn School of Medicine at Mount Sinai). The

rescue system for A/Panama/2007/99 (H3N2) [Pan/99] virus was initially described in [7]; and

that for A/Netherlands/602/2009 (H1N1) [NL09] virus was a gift of Ron Fouchier (Erasmus

Medical Center) [77]. The NL09 virus was propagated in MDCK cells for two passages. All other

virus stocks were generated in 10-11 day old embryonated chicken eggs.

Generation of chimeric M segments

Non-synonymous nucleotide changes differentiating NL09 and avian consensus M segments

were introduced in a reciprocal fashion into i) M1 and M2 to generate 16 mut viruses, ii) M1-only

to generate 9 mut viruses, and iii) M2-only to generate 7 mut viruses. The modified viral cDNAs

were synthesized by Genewiz and subcloned into the pDZ reverse genetic vector [78].

Sequence alignments and generation of consensus sequences

Alignment of the FASTA sequences downloaded from Genbank (or GISAID) databases was

carried out using DNASTAR Lasergene 13 MegAlign software running the Clustal W

phylogenetic alignment algorithm.

To obtain a consensus nucleotide sequence and to assess the degree of variability in the M

segment of H1N1 subtype avian IAVs, we searched the Genbank nucleotide database for full-

length M segment sequences derived from avian H1N1 subtype IAV collected between 1976

and 2008 from any temperate Northern hemisphere location. We obtained 262 sequences,

which included sequences with partial 3' and 5' UTRs, and manually curated the sequences to

(i) remove swine and human lineage derived H1N1 M segments, retaining 247 sequences, and

(ii) trim UTR sequences from the retained segments. Alignment of the sequences by Clustal W

provided a working consensus sequence and revealed up to 7% variability at the nucleotide

level among the selected avian host-derived sequences.

To assess the degree of amino acid variability in the M segment of avian IAV of any subtype, we

searched Genbank for full-length M1 or M2 protein sequences derived from avian H1Nx to

H16Nx subtype IAV, collected between 1970 and 2000, from any geographical location. These

dates were chosen to avoid selecting sequences representing isolates originating from the

repeated H5 subtype HPAI incursions into poultry populations which dominate the Genbank

database from the early 2000s onwards, and which would likely bias the subsequent analysis.

For each hemagglutinin subtype (H1Nx to H16Nx), we aligned the available sequences to

generate individual consensus M1 and M2 amino acid sequences (**Supplemental Table 1**), and

aligned those consensus sequences to the avian H1N1 subtype consensus M1 and M2

sequences, in order to assess amino acid diversity. The matrix proteins of each subtype, with

the exception of H9Nx, were 100% conserved. The H9Nx subtype matrix protein shared 97.6%

identity to the avian H1N1 sequence. The M2 proteins of each subtype were 100% conserved,

excepting H9Nx, H13Nx and H16Nx subtypes, which shared 97.2%, 94.8% and 93.8% identity

respectively, to the avian H1N1 consensus sequence. Thus, the amino acid composition of M1

and M2 proteins are highly conserved within wild waterfowl, particularly within isolates that

circulate among dabbling ducks and geese. In these host species, both M1 and M2 proteins are

at, or close to, 100% identity at the consensus amino acid level.

To identify a consensus nucleotide sequence for the 2009 pH1N1 IAV, and compare it to the

avian consensus sequence, we searched Genbank for full-length M segments derived from

pH1N1 subtype IAV isolated in North America, collected between the 7[th] June and the 20[th] June

2009, which was shortly after emergence of the pH1N1 lineage into the human population. 189

individual nucleotide sequences were obtained, curated, and aligned, to provide a working

consensus sequence for the pH1N1 M segment. The pH1N1 M segments possessed >2%

variability at the nucleotide level. The A/NL/602/09 (H1N1) (NL09) M segment is a 100% match

to the consensus sequence of the 2009 pH1N1 lineage at the nucleotide level, and the M

segment from this strain was adopted for use throughout the current study. Interestingly,

comparison of the NL09 M segment to the avian consensus sequence revealed 8.3% difference

at the nucleotide level, and 9 and 7 amino acid changes in M1 and M2 proteins, respectively

(**Table 2**). These data suggest that pH1N1 M segment possesses a nucleotide sequence that is

highly divergent from M segments obtained from viruses circulating within the wildfowl reservoir,

and which may encode host adaptive changes.

<u>Evaluation of viral growth in cell culture</u>

To assess growth, subconfluent cell monolayers in 6-well dishes were inoculated at an MOI of 5

PFU/cell (for single-cycle growth assays) or 0.01 PFU/cell (for multi-cycle growth assays) in a

200 µl volume. Following incubation for 1 h at 37^{o}C, inocula were removed, cells washed 3x

with PBS and 2 ml viral growth medium (MEM (for MDCK cells), DMEM (for 293T, and DF1

cells), or F-12K (for A549, and QT-6 cells) medium plus 3% bovine serum albumin, penicillin

streptomycin, and TPCK-treated trypsin) was added. Infected cells were incubated at 37^{o}C for

the remainder of the time-course. At the indicated time points, 120 µl medium was sampled from

each dish and 120 µl fresh medium was added to maintain a 2 ml volume. Samples were stored

at -80^{o}C and later titered by plaque assay on MDCK cells. Each infection was performed in

triplicate and at least three biological replicate infections were carried out on different days. Data

are plotted as mean with SD and analyzed by repeated measures ANOVA.

<u>Evaluation of viral growth in guinea pigs</u>

Female, Hartley strain, guinea pigs weighing 300-350 g were obtained from Charles River

Laboratories. Prior to inoculation or nasal lavage, animals were sedated with a mixture of

ketamine (30 mg/kg) and xylazine (4 mg/kg). Virus used for inoculation was diluted in PBS to

allow intranasal inoculation of guinea pigs with 1×10^1 to 1×10^3 PFU in a 300 µl volume. Nasal

wash samples were collected as described previously [54], with PBS as the collection fluid.

Animals were housed in a Caron 6040 environmental chamber set to $10^{\circ}C$ and 20% RH

throughout the seven-day period and lids were left off of the cages during this time to ensure

environmental control within the cages [79].

<u>Immunoblotting</u>

A549 or DF-1 cells in 6-well plates were infected with the indicated IAVs and cells were lysed

with 2X Laemmli sample buffer (Bio-rad) plus 2% beta-mercaptoethanol at 8 hpi. Samples were

boiled at 95 ºC for 10 min and resolved on 4-20% gradient SDS-PAGE gels, followed by transfer

to nitrocellulose membranes (Bio-rad), and incubation of membranes with 5% non-fat dry

milk/TBST blocking buffer. Antibodies used for immunoblotting were: anti-vinculin monoclonal

antibody (catalog no. V9131; Sigma-Aldrich) at 1:5000 dilution, IAV nucleoprotein monoclonal

antibody (HT-103; catalog no. EMS010, Kerafast) at 1:1000 dilution, IAV matrix protein

monoclonal antibody (E10; catalog no. EMS009, Kerafast) at 1:1000 dilution, and LC3B

polyclonal antibody (Thermo Fisher Scientific) at 1:1000 dilution. Bands were quantified using

ImageLab software (Bio-rad) after normalization to vinculin. At least three biological replicate

viral infections were carried out, with two technical replicates of the immunoblotting for each

infection. Data shown with error bars representing SD and statistical analysis was carried out

using one-way ANOVA with Tukey's multiple comparisons in GraphPad Prism software.

<u>RT primer extension assay</u>

A549 or DF-1 cells in 6-well plates were infected with the specified IAVs and total RNA was

extracted 8 hpi using a RNeasy Kit (Qiagen). Total RNA was eluted in RNase-free water and

stored at -80 ºC until needed.

Radiolabeling of primers: 1 µmol of oligo DNA primer was incubated with 10 µCuries of [γ-

^{32}P]ATP and 10 U of T4 PNK (New England Biolabs) in 70 mM Tris-HCl (pH 7.6) containing 10

mM $MgCl_2$ and 5 mM DTT. Radiolabeling reactions were carried out at 37 ºC for 1 h and diluted

to 30 µL with RNase-free water after termination of the reaction. Radiolabeled primers were

stored at –20 ºC until needed. 500 ng of total RNA was incubated with 0.45 µL unlabeled 5S

rRNA primer, 0.05 µL radiolabeled 5S rRNA primer, 0.25 µL radiolabeled IAV segment 7 vRNA

primer, and 0.25 µL radiolabeled IAV segment 7 mRNA primer in a final volume of 5 µL. Primers

were annealed to RNA by sequential incubations at 95 ºC for 3 min and ice for 3 min. Reactions

were pre-warmed at 50 ºC for 5 min. Transcription was initiated by the addition of 50 mM Tris-

HCl (pH 8.3) buffer containing 75 mM KCl, 3 mM MgCL2, 10 mM DTT, 2mM dNTP, and 50 units

of SuperScript III Reverse Transcriptase (Thermo Fisher Scientific). Reactions were incubated

at 50 ºC for 1 h and terminated by the addition of Gel Loading Buffer II (Thermo Fisher

Scientific) and incubation at 95 ºC for 10 min. Reactions were resolved on denaturing

sequencing polyacrylamide gels (7M urea, 6% acrylamide). Gels were subsequently dried and

exposed to a phosphor storage screen overnight. The intensity of bands was analyzed using a

Typhoon Trio Imager (GE Healthcare) and quantified with ImageQuant software (GE

Healthcare) after normalization to 5S rRNA. At least three biological replicate viral infections

were carried out, with two technical replicates of the RT primer extension assay for each

infection. Data are shown with error bars representing SD and statistical analysis was carried

out using one-way ANOVA with Tukey's multiple comparisons in GraphPad Prism software.

<u>Fluorescence microscopy</u>

To allow tracking of LC3B II and viral protein, cells grown on collagen-coated coverslips were

transduced with lentivirus expressing LC3B-GFP (BacMamm 2.0; catalog no. P36235,

ThermoFisher), as follows.

18h post-transduction, 293T or A549 cells were infected with the indicated IAVs at an MOI of 5

PFU/cell and incubated at 37^{o}C. At 8h or 12 h post-IAV infection, cells were washed once with

PBS and fixed with 4% paraformaldehyde. Following permeabilization with 0.1% Triton X-100,

cells were incubated overnight at 4^{o}C with anti-M2 monoclonal antibody (E10; catalog no.

EMS009, Kerafast). Three washes were performed prior to addition of goat anti-mouse Alexa

Fluor-647 secondary antibody (catalog no. A21241, Life Technologies) and incubation for 2 h at

room temperature. Antibody dilutions and washes were performed with PBS plus 1% Tween-20.

Cells were treated with DAPI, and coverslips were mounted on slides using Vectashield (Vector

Labs) mounting medium. Images were obtained with on a Nikon FV1000 confocal microscope at

the Emory Integrated Cellular Imaging core facility.

Drug treatments

Where indicated, chloroquine (Invitrogen) was added to cell culture medium at a final

concentration of 60 or 120 µM. Amantadine HCl (catalog no. 1018505, Sigma) was added to cell

culture medium at a final concentration of 200 µM. Both drugs were added to infected cells 1 h

after IAV inoculation.

Statistical analyses

Statistical analyses were performed in GraphPad Prism.

Multi-cycle and single-cycle growth in cell culture was assessed in three independent

experiments, with three technical sample replicates per experiment. Overall, statistical

significance was determined using repeated measures, two-way, multiple comparisons ANOVA

on log-transformed mean values, with Bonferroni correction applied to account for comparison

of a limited number of means, as well as by assessing pairwise comparisons of viruses at each

time point, except for single-cycle growth in cell culture in the presence of amantadine, which

was assessed in four independent experiments, with three technical sample replicates per

experiment. Overall statistical significance was determined using repeated measures, two-way,

multiple comparisons ANOVA on log-transformed values, with Sidak's correction applied, as

well as by assessing pairwise comparisons of viruses at each time point.

Growth of each virus in the guinea pig model was assessed in three independent experiments

with four guinea pigs per experiment. For transmission analysis, each guinea pig was treated as

an independent biological sample. Statistical significance in the magnitude of replication was

determined by comparing AUC of growth curves using two-tailed Student's t-test. Statistical

significance in transmission efficiency was determined using unpaired two-tailed Student's t-test.

Significance of differences in the kinetics of replication was determined by assessing the

interaction of time and virus using repeated measures, two-way ANOVA on mean log-

transformed values, with Bonferroni correction applied to account for comparison of a limited

number of means.

For experiments presented throughout the manuscript, unless specific values are provided, P

values are represented as follows: *< 0.05; **<0.01; ***< 0.001; ****< 0.0001.

Acknowledgements

We thank Paul Digard for his excellent advice relating to RT Primer Extension assays, and

Emory Integrated Cellular Imaging Core Facility for assistance with confocal imaging techniques.

Figure Legends

Figure 1. pH1N1 M segment does not confer improved growth relative to avian M

segment in avian cells at 37°C. (A) A panel of PR8-based viruses was rescued by reverse

genetics. The viruses differ only in the identity of the M segment, which were derived from

A/mallard/MN/99 (H3N8), A/rhea/NC/93 (H7N1), A/duck/Alberta/76 (H1N1), or A/NL/602/09

(H1N1) viruses. **(B)** 11-day-old embryonated chicken eggs (ECE) or (**C**) chicken DF-1 cells

were infected at low MOI with the indicated viruses. **(D)** Quail QT-6 cells or **(E)** DF-1 cells were

infected at a high MOI of 5 PFU/cell with the indicated viruses. In all avian systems assessed,

the pH1N1 M segment did not confer an increase in magnitude or kinetics of growth, relative to

any avian-origin M segment. Multi-cycle and single-cycle growth were assessed in three

independent experiments, with three technical replicates per experiment. Graphs show the

means with SD for the three experiments. Statistical significance was determined using

repeated measures, two-way, multiple ANOVA on log-transformed values, with Bonferroni

correction applied to account for comparison of a limited number of means.

Figure 2. pH1N1 M segment confers higher growth relative to avian M segment in

mammalian cells at 37^{o}C. (A) Canine MDCK cells or **(B)** human A549 cells were inoculated

with the indicated viruses at a low MOI of 0.01 PFU/cell. **(C)** MDCK cells or **(D)** A549 cells were

inoculated with the indicated viruses at a high MOI of 5 PFU/cell. In mammalian cells, the

pH1N1 M segment conferred more rapid kinetics and higher peak titers than any avian-origin M

segment. Multi-cycle and single-cycle growth were assessed in three independent experiments,

with three technical sample replicates per experiment. Graphs show the means with SD for the

three experiments. Statistical significance was determined using repeated measures, two-way,

multiple ANOVA on log-transformed values, with Bonferroni correction applied to account for

comparison of a limited number of means. At each time point, pairwise comparisons between

viruses were performed, and asterisks corresponding to the most significant difference observed

is shown.

Figure 3. pH1N1 influenza A virus M segment confers efficient replication and
transmission of PR8-based virus among guinea pigs. Groups of four guinea pigs were
inoculated with 10 PFU of each avian M-encoding virus or NL09 M-encoding virus, as indicated.
(A) Virus replication in nasal wash was measured by plaque titration at days 2, 4, 6, and 8 post-
infection and the area under the curve was calculated. Graphs show mean AUC with SD for
three experiments. All avian M-encoding viruses exhibited lower levels of growth *in vivo* than the
isogenic virus possessing the NL09 M segment. The replication differences between avian M
and NL09 M-based viruses were statistically significant, whereas those among avian M-based
viruses were not. Statistical significance was evaluated using an unpaired, two-tailed Student's
t-test. **(B)** To assess transmission, each inoculated guinea pig was treated as an independent
biological sample. The pH1N1 M segment conferred efficient transmission to naïve animals that
were contact exposed (7/9 transmissions; 78%). In contrast, significantly poorer (12%), or no
transmission was observed from guinea pigs infected with PR8 avian M viruses to naïve cage
mates. Statistical significance of transmission differences were evaluated using an unpaired,
two-tailed Student's t-test.

Figure 4. Ratio of M1 to M2 protein expression is high in DF-1 Cells irrespective of viral M
segment host origin. DF-1 cells were inoculated at a MOI of 5 PFU/cell, with PR8 viruses
encoding avian or human-derived M segments, and incubated at 37^{o}C for 8 h, then lysed.
Western immunoblot analysis of virus-infected DF-1 cells: **(A)** Vinculin expression was
measured to allow normalization of viral protein levels. NP expression was measured to assess
viral replication. Levels of M1 and M2 protein expression were assessed using an antibody
(Mab E10) to a common epitope at the amino terminus of M1 and M2 proteins, allowing relative
expression to be assessed. Levels of LC3B I and II were assessed using an antibody that
detects both the precursor and activated forms of LC3B protein. **(B)** M1 protein and **(C)** M2
protein were normalized to vinculin, quantitated and displayed as a percentage of total protein

expressed from the M gene. **(D)** The ratio of M1:M2 protein expression**. (E)** LC3B I protein and

(F) LC3B II protein were normalized, quantitated and displayed as a percentage of total LC3B

protein. Graphs in **B-F** show the means with SD from three independent experiments. For each

experiment, two replicate Western immunoblots were performed and quantitated. Statistical

significance was assessed using ordinary one-way ANOVA.

Figure 5. High expression ratio of M1 to M2 protein in A549 cells is dependent upon viral

M segment host origin. A549 cells were inoculated at a MOI of 5 PFU/cell with PR8 viruses

encoding avian or human-derived M segments and incubated at 37°C for 8 h, then lysed.

Western immunoblot analysis of virus-infected A549 cells: **(A)** Vinculin expression was

measured to allow normalization of viral protein levels. NP expression was measured to assess

viral replication. Levels of M1 and M2 protein expression were assessed using an antibody

(Mab E10) to a common epitope at the amino terminus of M1 and M2 proteins, allowing relative

expression to be assessed. Levels of LC3B I and II were assessed using an antibody that

detects both the precursor and activated forms of LC3B protein. **(B)** M1 protein and **(C)** M2

protein were normalized to vinculin, quantitated and displayed as a percentage of total protein

expressed from the M gene. **(D)** The ratio of M1:M2 protein expression. **(E)** LC3B I protein and

(F) LC3B II protein were normalized, quantitated and displayed as a percentage of total LC3B

protein. Graphs in **B-F** show the means with SD from three independent experiments. For each

experiment, two replicate Western immunoblots were performed and quantitated. Statistical

significance was assessed using ordinary one-way ANOVA.

Figure 6. Ratio of mRNA$_7$ (encoding M1) to mRNA$_{10}$ (encoding M2) expression is high in

DF-1 cells irrespective of viral M Segment host origin.

DF-1 cells were inoculated at a MOI of 5 PFU/cell with PR8 viruses encoding avian or human-

derived M segments and incubated at 37°C for 8 h prior to RNA extraction. RT primer extension

radiogram of virus-infected DF-1 cells: **(A)** 5S rRNA levels were measured to allow

normalization of viral RNA. Segment 7 vRNA expression was measured to assess viral

replication. Levels of $mRNA_7$, $mRNA_{10}$, and $mRNA_{11}$ mRNA expression were assessed using

radiolabeled probes. **(B)** $mRNA_7$, **(C)** $mRNA_{10}$, and **(D)** $mRNA_{11}$ was quantitated and displayed

as a percentage of total M gene-expressed mRNA. Graphs in **B-D** show the means with SD

from three independent experiments. For each experiment, two replicate radiograms were

quantitated. Statistical significance was assessed using ordinary one-way ANOVA.

Figure 7. Ratio of $mRNA_7$ (encoding M1) to $mRNA_{10}$ (encoding M2) expression in A549

cells is dependent upon viral M segment host origin. A549 cells were inoculated at a MOI of

5 PFU/cell with PR8 viruses encoding avian or human-derived M segments and incubated at

$37^{o}C$ for 8 h prior to RNA extraction. RT primer extension radiogram of virus-infected A549 cells:

(A) 5S rRNA levels were measured to allow normalization of viral RNA. Segment 7 vRNA

expression was measured to assess viral replication. Levels of $mRNA_7$, $mRNA_{10}$, and $mRNA_{11}$

mRNA expression were assessed using radiolabeled probes. **(B)** $mRNA_7$, **(C)** $mRNA_{10}$, and **(D)**

$mRNA_{11}$ was quantitated and displayed as a percentage of total M gene expressed mRNA.

Graphs in **B-D** show the means with SD from three independent experiments. For each

experiment, two replicate radiograms were quantitated. Statistical significance was assessed

using ordinary one-way ANOVA.

Figure 8. In human cells, avian M2 was expressed at higher levels and localized more

strongly to perinuclear bodies than human M2. A549 and 293T cells were inoculated with

the indicated viruses, encoding avian or human M segments, at a MOI of 5 PFU/cell. At 8 hpi

cells were stained with anti-M2 (Mab E10; red) and DAPI (blue) and imaged by confocal

microscopy. Examples of optical sections are shown, either as merged 2-color images or the

red and blue channels alone (shown in grey scale). **(A)** A549 cells with 63x optical and 3x digital

magnification (189x total magnification). **(B)** 293T cells with 63x optical and 3x digital

magnification (189x total magnification). Brightness was adjusted for optimal clarity, with all

images treated equally.

Figure 9. Visualization of LC3 and M2 co-localization.

293T cells were transduced with GFP-LC3 protein and inoculated 24 h later with the indicated

IAVs, encoding avian or human-derived M segments, at a MOI of 5 PFU/cell. Cells were fixed 8

h later and stained with anti-M2 (Mab E10; red) and DAPI (blue) followed by imaging with

confocal microscopy. CQ: chloroquine. Examples of optical sections are shown, either as

merged 3-color images, or the red, green, and blue channels alone (in grey scale). Images are

at 63x optical and 3x digital magnification (189x total magnification). Brightness was adjusted for

optimal clarity, with all images treated equally.

Figure 10. Inhibition of M2 ion channel results in loss of activation of LC3B.

A549 cells were inoculated at an MOI of 5 PFU/cell with PR8 viruses encoding avian or human-

derived M segments and incubated in the presence or absence of 200 µM amantadine from 1

hpi. Cells were lysed at 8 hpi. **(A)** Representative Western immunoblot. Vinculin expression was

measured to allow normalization of viral protein levels. NP expression was measured to assess

viral replication. Levels of M1 and M2 protein were assessed using an antibody (Mab E10) to a

common epitope of M1 and M2 proteins, allowing relative expression to be assessed. Levels of

LC3B I and II were assessed using an antibody that detects both the precursor and activated

forms of LC3B protein. **(B)** M1 protein and **(C)** M2 protein were normalized to vinculin and

displayed as a percentage of total M gene-expressed protein. **(D)** The ratio of M1:M2 protein

expression. **(E)** LC3B I and **(F)** LC3B II were normalized to vinculin and displayed as a

percentage of total LC3B protein. **(G)** As a control to ensure the specificity of amantadine, we

incubated chloroquine-treated A549 cells in the presence or absence of 200 µM amantadine for

8 h. **(H)** LC3B I and **(I)** LC3B II were normalized to vinculin and displayed as a percentage of

total LC3B protein. Graphs in **B-F, H-I** show the means with SD from three independent

experiments. For each experiment, two replicate Western immunoblots were performed and

quantitated. Statistical significance was assessed using ordinary one-way ANOVA.

Figure 11. Inhibition of proton channel activity relieves M2-induced accumulation of

LC3B positive vesicles. 293T cells transduced with LC3-GFP were inoculated at a MOI of 5

PFU/cell with PR8 viruses possessing M segments from a human or an avian strain and

incubated in the presence or absence of 200 µM amantadine from 1 hpi. Cells were fixed at 12

hpi and stained with anti-M2 (Mab E10; red) and DAPI (blue) followed by imaging with confocal

microscopy. Examples of optical sections are shown, either as merged 3-color images or the

red, green, and blue channels alone (in grey scale). 3x zoom of a 63x magnification is shown.

Brightness was adjusted for optimal clarity, with all images treated equally. CQ: chloroquine.

Figure 12. Amantadine improves growth of avian M segment-encoding virus in A549 cells.

PR8-based NL09 M or avian M-encoding viruses were inoculated at an MOI of 5 PFU/cell onto

A549 cells. At 1hpi, cells were treated with 200µM amantadine, and incubated at 37°C for 24 h

in the continuous presence of amantadine. Virus released into supernatant was collected at the

indicated time points, and virus growth was measured by plaque titration. The pH1N1 M

segment conferred more rapid kinetics and higher peak titers of growth than the avian-origin M

segment, however amantadine had no impact on the growth of the pH1N1 M encoding virus. In

contrast, growth of the avian M-encoding virus was improved in the presence of drug (P=

0.0007). Single-cycle growth was assessed in four independent experiments, with three

technical sample replicates per experiment. Graphs show the means with SD for the four

experiments. Statistical significance was determined using repeated measures, two-way,

multiple comparisons ANOVA on log-transformed values, with Sidak's correction applied.

Figure 13. Activation of LC3B II correlates with high expression of M2 protein in A549 cells regardless of M2 amino acid composition. PR8 NL09 M virus and PR8 avian M virus, along with six chimeric variant PR8-based viruses, were inoculated at a MOI of 5 PFU/cell onto A549 cells. Cells were incubated for 8 h, then lysed. **(A)** Western immunoblot analysis of virus-infected A549 cells. Vinculin was measured to allow normalization of viral protein levels. NP was measured to assess viral replication. Levels of M1 and M2 protein were assessed using an antibody (Mab E10) to a common epitope at the amino terminus of M1 and M2 proteins, allowing their direct comparison. Levels of LC3B I and II were assessed using an antibody that detects both the precursor and activated forms of LC3B protein. **(B)** M1 protein and **(C)** M2 protein were normalized to vinculin and displayed as a percentage of total protein expressed from the M gene. **(D)** The ratio of M1:M2 protein expression. **(E)** LC3B I protein and **(F)** LC3B II protein were normalized to vinculin and displayed as a percentage of total LC3B protein. Graphs in **B-F** show the means with SD from three independent experiments. For each experiment, two replicate Western immunoblots were performed and quantitated. Statistical significance was assessed using ordinary one-way ANOVA.

Figure 14. pH1N1 M amino acid sequences are not sufficient to confer high growth phenotype in A549 cells. PR8 NL09 M virus and PR8 avian M virus, along with six PR8-based viruses carrying avian-human chimeric M segments, were inoculated at a MOI of 5 PFU/cell onto A549 cells. Virus growth was measured by plaque titration. The effects on viral growth of amino acid changes in **(A)** both M1 and M2, **(B)** M1 only and **(C)** M2 only are shown. Single-cycle growth was assessed in at least three independent experiments, with three technical sample replicates per experiment. Graphed data show the means with SD for at least three experiments. Statistical significance was determined using repeated measures, two-way, multiple ANOVA on log-transformed values, with Bonferroni correction applied to account for

comparison of a limited no of means. Experiments involving PR8 NL09 9mut or 7mut viruses as

well as PR8 Av M 9mut or 7mut viruses were performed at the same time, but are displayed on

two separate graphs with the same PR8 NL09 M and PR8 avian M control virus data, to

improve clarity of presentation.

Figure 15. Reduced expression of avian M2 improves replication efficiency *in vivo* but is

not sufficient for efficient transmission in guinea pigs.

Groups of twelve guinea pigs were inoculated with 10 PFU of PR8-based viruses carrying avian

M, NL09 M, chimeric NL09 M 7mut M, or chimeric avian M 9mut segments, as indicated. **(A)**

Virus replication in nasal washings was measured by plaque titration at days 2, 4, 6, and 8 post-

infection and the total area under the curve was calculated. The replication differences between

viruses with NL09 M and NL09 M 7mutAv segments were not significant (P=0.720). Differences

between viruses encoding the NL09 M 7mut segment and the avian M 9mut segment trended

towards, but were not statistically significant (P=0.115). (**B**) Transmission to naïve cage mates

that were exposed at 24 hpi. Neither chimeric M segment conferred efficient transmission (7/9

transmissions; 78%). Statistical significance of replication differences was determined using a

two-tailed Student's t-test. Statistical significance of transmission differences was determined

using an unpaired two-tailed Student's t-test. NL M and avian M data are identical to data

presented in Figure 3 and are reproduced here for clarity.

Figure 16. Regulation of M segment gene expression contributes to host range

restriction of IAV. When an avian IAV M segment is transcribed within mammalian cells,

excessive splicing of mRNA$_7$ results in aberrant overexpression of the M2 proton channel. When

highly expressed, this channel accumulates in autophagic vesicles. Abundant proton channel

activity in the membranes of autophagosomes blocks their turnover by preventing fusion with

lysosomes. This stalling of a critical cellular housekeeping function in turn reduces the efficiency

of viral replication, contributing to the species barrier that limits the zoonotic potential of avian

IAVs.

References

1. Patterson KD, Pyle GF (1991) The geography and mortality of the 1918 influenza pandemic.
Bull Hist Med 65: 4-21.
2. Viboud C, Simonsen L, Fuentes R, Flores J, Miller MA, et al. (2016) Global Mortality Impact of
the 1957-1959 Influenza Pandemic. J Infect Dis 213: 738-745.
3. Viboud C, Grais RF, Lafont BA, Miller MA, Simonsen L, et al. (2005) Multinational impact of
the 1968 Hong Kong influenza pandemic: evidence for a smoldering pandemic. J Infect
Dis 192: 233-248.
4. Molinari NA, Ortega-Sanchez IR, Messonnier ML, Thompson WW, Wortley PM, et al. (2007)
The annual impact of seasonal influenza in the US: measuring disease burden and costs.
Vaccine 25: 5086-5096.
5. Reperant LA, Kuiken T, Osterhaus AD (2012) Adaptive pathways of zoonotic influenza viruses:
from exposure to establishment in humans. Vaccine 30: 4419-4434.
6. Lowen AC (2017) Constraints, Drivers, and Implications of Influenza A Virus Reassortment.
Annu Rev Virol.
7. Steel J, Lowen AC, Mubareka S, Palese P (2009) Transmission of influenza virus in a
mammalian host is increased by PB2 amino acids 627K or 627E/701N. PLoS Pathog 5:
e1000252.
8. Subbarao EK, London W, Murphy BR (1993) A single amino acid in the PB2 gene of influenza
A virus is a determinant of host range. J Virol 67: 1761-1764.
9. Linster M, van Boheemen S, de Graaf M, Schrauwen EJ, Lexmond P, et al. (2014)
Identification, characterization, and natural selection of mutations driving airborne
transmission of A/H5N1 virus. Cell 157: 329-339.
10. Long JS, Giotis ES, Moncorge O, Frise R, Mistry B, et al. (2016) Species difference in ANP32A
underlies influenza A virus polymerase host restriction. Nature 529: 101-104.
11. Rogers GN, Paulson JC (1983) Receptor determinants of human and animal influenza virus
isolates: differences in receptor specificity of the H3 hemagglutinin based on species of
origin. Virology 127: 361-373.
12. Tanner WD, Toth DJ, Gundlapalli AV (2015) The pandemic potential of avian influenza
A(H7N9) virus: a review. Epidemiol Infect 143: 3359-3374.
13. Capua I, Munoz O (2013) Emergence of influenza viruses with zoonotic potential: open
issues which need to be addressed. A review. Vet Microbiol 165: 7-12.
14. Cauldwell AV, Long JS, Moncorge O, Barclay WS (2014) Viral determinants of influenza A
host range. J Gen Virol.
15. Bosch FX, Hay AJ, Skehel JJ (1978) RNA and protein synthesis in a permissive and an abortive
influenza virus infection. In: Mahy BWJ, Barry RD, editors. Negative Strand Viruses and
the Host Cell. London, New York, San Francisco: Academic Press. pp. 465-474.
16. Valcavi PP, Conti G, Schito GC (1978) Macromolecular syntheses during abortive infection of
KB cells by influenza virus. In: Mahy BWJ, Barry RD, editors. Negative Strand Viruses and
the Host Cell. London, New York, San Francisco: Academic Press. pp. 475-482.
17. Lohmeyer J, Talens LT, Klenk HD (1979) Biosynthesis of the influenza virus envelope in
abortive infection. J Gen Virol 42: 73-88.

18. Lamb RA, Lai CJ, Choppin PW (1981) Sequences of mRNAs derived from genome RNA
1003   segment 7 of influenza virus: colinear and interrupted mRNAs code for overlapping
1004   proteins. Proc Natl Acad Sci U S A 78: 4170-4174.
1005   19. Lamb RA, Choppin PW (1981) Identification of a second protein (M2) encoded by RNA
1006   segment 7 of influenza virus. Virology 112: 729-737.
1007   20. Inglis SC, Brown CM (1984) Differences in the control of virus mRNA splicing during
1008   permissive or abortive infection with influenza A (fowl plague) virus. J Gen Virol 65 (Pt
1009   1): 153-164.
1010   21. Lau SC, Scholtissek C (1995) Abortive infection of Vero cells by an influenza A virus (FPV).
1011   Virology 212: 225-231.
1012   22. Bogdanow B, Eichelbaum K, Sadewasser A, Wang X, Husic I, et al. (2018) The dynamic
1013   proteome of influenza A virus infection identifies M segment splicing as a host range
1014   determinant. BioRxiv: https://doi.org/10.1101/438176.
1015   23. Joseph U, Vijaykrishna D, Smith GJD, Su YCF (2018) Adaptive evolution during the
1016   establishment of European avian-like H1N1 influenza A virus in swine. Evol Appl 11: 534-
1017   546.
1018   24. Campbell PJ, Danzy S, Kyriakis CS, Deymier MJ, Lowen AC, et al. (2014) The M segment of
1019   the 2009 pandemic influenza virus confers increased NA activity, filamentous
1020   morphology and efficient contact transmissibility to A/Puerto Rico/8/1934-based
1021   reassortant viruses. J Virol.
1022   25. Campbell PJ, Kyriakis CS, Marshall N, Suppiah S, Seladi-Schulman J, et al. (2014) Residue 41
1023   of the Eurasian avian-like swine influenza a virus matrix protein modulates virion
1024   filament length and efficiency of contact transmission. J Virol 88: 7569-7577.
1025   26. Seladi-Schulman J, Steel J, Lowen AC (2013) Spherical Influenza Viruses Have a Fitness
1026   Advantage in Embryonated Eggs, while Filament-Producing Strains Are Selected In Vivo.
1027   J Virol 87: 13343-13353.
1028   27. Chou YY, Albrecht RA, Pica N, Lowen AC, Richt JA, et al. (2011) The M segment of the 2009
1029   new pandemic H1N1 influenza virus is critical for its high transmission efficiency in the
1030   guinea pig model. J Virol 85: 11235-11241.
1031   28. Lakdawala SS, Lamirande EW, Suguitan AL, Jr., Wang W, Santos CP, et al. (2011) Eurasian-
1032   origin gene segments contribute to the transmissibility, aerosol release, and
1033   morphology of the 2009 pandemic H1N1 influenza virus. PLoS Pathog 7: e1002443.
1034   29. Rossman JS, Lamb RA (2011) Influenza virus assembly and budding. Virology 411: 229-236.
1035   30. Pielak RM, Chou JJ (2010) Flu channel drug resistance: a tale of two sites. Protein Cell 1:
1036   246-258.
1037   31. Bourmakina SV, Garcia-Sastre A (2003) Reverse genetics studies on the filamentous
1038   morphology of influenza A virus. J Gen Virol 84: 517-527.
1039   32. Elleman CJ, Barclay WS (2004) The M1 matrix protein controls the filamentous phenotype of
1040   influenza A virus. Virology 321: 144-153.
1041   33. Stauffer S, Feng Y, Nebioglu F, Heilig R, Picotti P, et al. (2014) Stepwise priming by acidic pH
1042   and a high K+ concentration is required for efficient uncoating of influenza A virus cores
1043   after penetration. J Virol 88: 13029-13046.
1044   34. O'Neill RE, Talon J, Palese P (1998) The influenza virus NEP (NS2 protein) mediates the
1045   nuclear export of viral ribonucleoproteins. EMBO J 17: 288-296.

35. Akarsu H, Burmeister WP, Petosa C, Petit I, Muller CW, et al. (2003) Crystal structure of the
M1 protein-binding domain of the influenza A virus nuclear export protein (NEP/NS2).
EMBO J 22: 4646-4655.
36. Brunotte L, Flies J, Bolte H, Reuther P, Vreede F, et al. (2014) The nuclear export protein of
H5N1 influenza A viruses recruits Matrix 1 (M1) protein to the viral ribonucleoprotein to
mediate nuclear export. J Biol Chem 289: 20067-20077.
37. Watanabe K, Handa H, Mizumoto K, Nagata K (1996) Mechanism for inhibition of influenza
virus RNA polymerase activity by matrix protein. J Virol 70: 241-247.
38. Zhang J, Pekosz A, Lamb RA (2000) Influenza virus assembly and lipid raft microdomains: a
role for the cytoplasmic tails of the spike glycoproteins. J Virol 74: 4634-4644.
39. Ali A, Avalos RT, Ponimaskin E, Nayak DP (2000) Influenza virus assembly: effect of influenza
virus glycoproteins on the membrane association of M1 protein. J Virol 74: 8709-8719.
40. Barman S, Adhikary L, Chakrabarti AK, Bernas C, Kawaoka Y, et al. (2004) Role of
transmembrane domain and cytoplasmic tail amino acid sequences of influenza a virus
neuraminidase in raft association and virus budding. J Virol 78: 5258-5269.
41. Shimbo K, Brassard DL, Lamb RA, Pinto LH (1996) Ion selectivity and activation of the M2 ion
channel of influenza virus. Biophys J 70: 1335-1346.
42. Grambas S, Hay AJ (1992) Maturation of influenza A virus hemagglutinin--estimates of the
pH encountered during transport and its regulation by the M2 protein. Virology 190: 11-
18.
43. Ciampor F, Thompson CA, Grambas S, Hay AJ (1992) Regulation of pH by the M2 protein of
influenza A viruses. Virus Res 22: 247-258.
44. Takeuchi K, Lamb RA (1994) Influenza virus M2 protein ion channel activity stabilizes the
native form of fowl plague virus hemagglutinin during intracellular transport. J Virol 68:
911-919.
45. Rossman JS, Jing X, Leser GP, Lamb RA (2010) Influenza virus M2 protein mediates ESCRT-
independent membrane scission. Cell 142: 902-913.
46. Beale R, Wise H, Stuart A, Ravenhill BJ, Digard P, et al. (2014) A LC3-interacting motif in the
influenza A virus M2 protein is required to subvert autophagy and maintain virion
stability. Cell Host Microbe 15: 239-247.
47. Gannage M, Dormann D, Albrecht R, Dengjel J, Torossi T, et al. (2009) Matrix protein 2 of
influenza A virus blocks autophagosome fusion with lysosomes. Cell Host Microbe 6:
367-380.
48. Ren Y, Li C, Feng L, Pan W, Li L, et al. (2016) Proton Channel Activity of Influenza A Virus
Matrix Protein 2 Contributes to Autophagy Arrest. J Virol 90: 591-598.
49. Kovacs AL, Reith A, Seglen PO (1982) Accumulation of autophagosomes after inhibition of
hepatocytic protein degradation by vinblastine, leupeptin or a lysosomotropic amine.
Exp Cell Res 137: 191-201.
50. Barr JN, Whelan SP, Wertz GW (1997) cis-Acting signals involved in termination of vesicular
stomatitis virus mRNA synthesis include the conserved AUAC and the U7 signal for
polyadenylation. J Virol 71: 8718-8725.
51. Barr JN, Whelan SP, Wertz GW (1997) Role of the intergenic dinucleotide in vesicular
stomatitis virus RNA transcription. J Virol 71: 1794-1801.

52. Pagan I, Holmes EC, Simon-Loriere E (2012) Level of gene expression is a major determinant
of protein evolution in the viral order Mononegavirales. J Virol 86: 5253-5263.
53. Abraham G, Banerjee AK (1976) Sequential transcription of the genes of vesicular stomatitis
virus. Proc Natl Acad Sci U S A 73: 1504-1508.
54. Lowen AC, Mubareka S, Tumpey TM, Garcia-Sastre A, Palese P (2006) The guinea pig as a
transmission model for human influenza viruses. Proc Natl Acad Sci U S A 103: 9988-
9992.
55. Lowen AC, Bouvier NM, Steel J (2014) Transmission in the guinea pig model. Curr Top
Microbiol Immunol 385: 157-183.
56. Inglis SC, Brown CM (1981) Spliced and unspliced RNAs encoded by virion RNA segment 7 of
influenza virus. Nucleic Acids Res 9: 2727-2740.
57. Fujita N, Itoh T, Omori H, Fukuda M, Noda T, et al. (2008) The Atg16L complex specifies the
site of LC3 lipidation for membrane biogenesis in autophagy. Mol Biol Cell 19: 2092-
2100.
58. Hanada T, Noda NN, Satomi Y, Ichimura Y, Fujioka Y, et al. (2007) The Atg12-Atg5 conjugate
has a novel E3-like activity for protein lipidation in autophagy. J Biol Chem 282: 37298-
37302.
59. Barth S, Glick D, Macleod KF (2010) Autophagy: assays and artifacts. J Pathol 221: 117-124.
60. Mauthe M, Orhon I, Rocchi C, Zhou X, Luhr M, et al. (2018) Chloroquine inhibits autophagic
flux by decreasing autophagosome-lysosome fusion. Autophagy 14: 1435-1455.
61. Skehel JJ, Hay AJ, Armstrong JA (1978) On the mechanism of inhibition of influenza virus
replication by amantadine hydrochloride. J Gen Virol 38: 97-110.
62. van Wielink R, Harmsen MM, Martens DE, Peeters BP, Wijffels RH, et al. (2012) Mutations in
the M-gene segment can substantially increase replication efficiency of NS1 deletion
influenza A virus in MDCK cells. J Virol 86: 12341-12350.
63. Spronken MI, van de Sandt CE, de Jongh EP, Vuong O, van der Vliet S, et al. (2017) A
compensatory mutagenesis study of a conserved hairpin in the M gene segment of
influenza A virus shows its role in virus replication. RNA Biol 14: 1606-1616.
64. Shih SR, Krug RM (1996) Novel exploitation of a nuclear function by influenza virus: the
cellular SF2/ASF splicing factor controls the amount of the essential viral M2 ion channel
protein in infected cells. EMBO J 15: 5415-5427.
65. Thompson MG, Munoz-Moreno R, Bhat P, Roytenberg R, Lindberg J, et al. (2018) Co-
regulatory activity of hnRNP K and NS1-BP in influenza and human mRNA splicing. Nat
Commun 9: 2407.
66. Ren Y, Li C, Feng L, Pan W, Li L, et al. (2015) Proton Channel Activity of Influenza A Virus
Matrix Protein 2 Contributes to Autophagy Arrest. J Virol 90: 591-598.
67. Zhirnov OP, Klenk HD (2013) Influenza A virus proteins NS1 and hemagglutinin along with
M2 are involved in stimulation of autophagy in infected cells. J Virol 87: 13107-13114.
68. Heaton NS, Randall G (2011) Dengue virus and autophagy. Viruses 3: 1332-1341.
69. Heaton NS, Randall G (2010) Dengue virus-induced autophagy regulates lipid metabolism.
Cell Host Microbe 8: 422-432.
70. Zhang J, Lan Y, Li MY, Lamers MM, Fusade-Boyer M, et al. (2018) Flaviviruses Exploit the
Lipid Droplet Protein AUP1 to Trigger Lipophagy and Drive Virus Production. Cell Host
Microbe 23: 819-831 e815.

71. Henkel JR, Weisz OA (1998) Influenza virus M2 protein slows traffic along the secretory
pathway. pH perturbation of acidified compartments affects early Golgi transport steps.
J Biol Chem 273: 6518-6524.
72. Henkel JR, Apodaca G, Altschuler Y, Hardy S, Weisz OA (1998) Selective perturbation of
apical membrane traffic by expression of influenza M2, an acid-activated ion channel, in
polarized madin-darby canine kidney cells. Mol Biol Cell 9: 2477-2490.
73. Henkel JR, Popovich JL, Gibson GA, Watkins SC, Weisz OA (1999) Selective perturbation of
early endosome and/or trans-Golgi network pH but not lysosome pH by dose-
dependent expression of influenza M2 protein. J Biol Chem 274: 9854-9860.
74. Burleigh LM, Calder LJ, Skehel JJ, Steinhauer DA (2005) Influenza a viruses with mutations in
the m1 helix six domain display a wide variety of morphological phenotypes. J Virol 79:
1262-1270.
75. Roberts PC, Lamb RA, Compans RW (1998) The M1 and M2 proteins of influenza A virus are
important determinants in filamentous particle formation. Virology 240: 127-137.
76. Fodor E, Devenish L, Engelhardt OG, Palese P, Brownlee GG, et al. (1999) Rescue of
influenza A virus from recombinant DNA. J Virol 73: 9679-9682.
77. Chutinimitkul S, Herfst S, Steel J, Lowen AC, Ye J, et al. (2010) Virulence-Associated
Substitution D222G in the Hemagglutinin of 2009 Pandemic Influenza A(H1N1) Virus
Affects Receptor Binding. Journal of Virology 84: 11802-11813.
78. Quinlivan M, Zamarin D, Garcia-Sastre A, Cullinane A, Chambers T, et al. (2005) Attenuation
of equine influenza viruses through truncations of the NS1 protein. J Virol 79: 8431-8439.
79. Lowen AC, Mubareka S, Steel J, Palese P (2007) Influenza virus transmission is dependent on
relative humidity and temperature. PLoS Pathog 3: 1470-1476.

1156
1157
1158

1159 **Supporting Information Legends**

1160

1161 **Supplementary Figure 1. Schematic of M segment mRNAs and gene products.**

1162

(A) The M segment of influenza virus is template for synthesis of $mRNA_7$ (encoding M1), mRNA

M_{10} (encoding M2), and $mRNA_{11}$ (which encodes a putative but unconfirmed 10 amino acid

peptide from a short open reading frame). (**B**) Pandemic H1N1 influenza virus M1 and M2

proteins differ from the avian consensus sequences by 9 residues in M1 and 7 residues in M2.

The M segments differ by 8.3% at the nucleotide level. (**C**) Seasonal H3N2 influenza virus strain

A/Panama/2007/99 M1 and M2 proteins differ from the avian consensus sequences by 11

residues in M1 and 14 residues in M2. The M segments differ by 9% at the nucleotide level. (**D**)

Seasonal H3N2 influenza virus strain A/Bethesda/55/15 M1 and M2 proteins differ from the

avian consensus sequences by 11 residues in M1 and 16 residues in M2. The M segments

differ by 9.5% at the nucleotide level.

1173

1174   **Supplementary Figure 2. Human host-derived M segments confer higher growth to PR8-**

1175   **based viruses than avian host-derived M segments in mammalian cells at 37^{o}C.**

1176

PR8-based viruses were inoculated at a MOI of 5 PFU/cell onto human-derived 293T cells (**A**),

or A549 cells (**B**). Cells were incubated at 37^{o}C for up to 24 h. Virus released into supernatant

was collected at the indicated time points, and virus growth was measured by plaque titration.

Data obtained from viruses possessing human M segments are represented with blue lines:

A/NL/602/09 M (**A,B**), A/Panama/2007/99 M (**B**), and A/Bethesda/15 M (**B**), while data from

viruses encoding avian M segments are represented with red lines. In each cell type, the human

M segments conferred more rapid kinetics and higher peak titers of growth than any avian-origin

M segment. Single-cycle growth was assessed in three independent experiments, with three

technical sample replicates per experiment. Graphs show the means with SD for the three

experiments. Statistical significance was determined using repeated measures, two-way,

multiple ANOVA on log-transformed data, with Bonferroni correction applied as there were a

limited no of means to compare.

1189

1190   **Supplementary Figure 3. pH1N1 influenza virus M segment increases kinetics of**

1191   **replication of PR8-based viruses among guinea pigs.**

1192

Groups of four guinea pigs were inoculated with 10 PFU of each avian M-encoding virus, or

NL09 M-encoding virus, as indicated. Graphs show individual titers obtained from animals used

in three independent experiments. **(A-E)** Virus replication in nasal wash of inoculated animals

was measured by plaque titration at days 2, 4, 6, and 8 post-infection and the titers at each time

point were plotted (dotted lines). The differences between PR8 NL09 M and each avian M-

encoding virus were considered significant. Statistical significance in kinetics of growth was

determined by assessing the interaction of time and virus using repeated measures, two-way,

multiple comparisons ANOVA on mean values, with Bonferroni correction applied to account for

comparison of a limited no of means.

1202

1203 **Supplementary Figure 4. High expression ratio of M1 to M2 protein in human cells is**

1204 **dependent upon viral M segment host origin.**

1205

293T and MDCK cells were inoculated at a MOI of 5 PFU/cell with PR8 viruses encoding avian

or human-derived M segments and incubated at 37°C for 8 h, then cells were lysed. Western

immunoblot analysis of virus-infected 293T cells **(A)** and MDCK cells **(G)**. Vinculin expression

was measured to allow normalization of viral protein levels. NP expression was measured to

assess viral replication. Levels of M1 and M2 protein expression were assessed using an

antibody (Mab E10) to a common epitope at the amino terminus of M1 and M2 proteins,

allowing relative expression to be assessed. Levels of LC3B I and II were assessed using an

antibody that detects both the precursor and activated forms of LC3B protein. **(B, H)** M1 protein

and **(C, I)** M2 protein were normalized to vinculin, quantitated and displayed as a percentage of

total protein expressed from the M gene. **(D, J)** The ratio of M1:M2 protein expression. **(E, K)**

LC3B I protein and **(F, L)** LC3BII protein were normalized, quantitated and displayed as a

percentage of total LC3B protein. Graphs in **B-F**, and **H-K** show the means with SD from three

independent experiments. For each experiment, two replicate Western immunoblots were

performed and quantitated. Statistical significance was assessed using ordinary one-way

ANOVA.

1221

1222

Supplementary Figure 5. High expression ratio of M1 to M2 protein in human cells is dependent upon viral M segment host origin.

1225

A549 cells were inoculated at a MOI of 5 PFU/cell with PR8 viruses encoding avian or human-derived M segments and incubated at 37^{o}C for 8 h, then cells were lysed. Human M segments were derived from the following viruses: A/NL/602/09 (H1N1) (NL09); A/Panama/2007/99 (H3N2) (Pan99); and A/Bethesda/15 (H3N2) (Beth15). Vinculin expression was measured to allow normalization of viral protein levels. NP expression was measured to assess viral replication. Levels of M1 and M2 protein expression were assessed using an antibody (Mab E10) to a common epitope at the amino terminus of M1 and M2 proteins, allowing relative expression to be assessed. Levels of LC3B I and II were assessed using an antibody that detects both the precursor and the activated forms of LC3B protein. Data presented are representative of Western immunoblots from three independent experiments.

1236

Supplementary Figure 6. Visualization of M2 localization by immunofluorescence microscopy at 12 h post-infection in A549 and 293T cells.

1239

A549 **(A, B)**, or 293T **(C, D)** cells were inoculated with the indicated viruses, encoding avian or human M segments, at a MOI of 5 PFU/cell. Cells were fixed at 12 hpi, permeabilised, and stained with anti-M2 (Mab E10; red) and DAPI (blue) followed by imaging with confocal microscopy. Examples of optical sections are shown, either as merged 2-color images or the red and blue channels alone (in grey scale). **(A)** A549 cells with 63x magnification. **(B)** 3x magnification of the same images shown in **A. (C)** 293T cells with 63x magnification. **(D)** 3x magnification of the same images shown in **C.** Brightness was adjusted for optimal clarity, with

1247   all images treated equally.

1248

1249   **Supplementary Figure 7. Visualization of LC3 and M2 co-localization by**

1250   **immunofluorescence microscopy.**

1251

A549 cells were transduced with GFP-LC3 protein and inoculated 24 h later with the indicated

IAVs, encoding avian- or human-derived M segments, at a MOI of 5 PFU/cell. Cells were fixed 8

h later and stained with anti-M2 (Mab E10; red) and DAPI (blue) followed by imaging with

confocal microscopy. Examples of optical sections are shown, either as merged 3-color images

or the red, green, and blue channels alone (in grey scale). **(A)** 63x magnification. **(B)** 3x

magnification of the same images shown in **A.** Brightness was adjusted for optimal clarity, with

all images treated equally. CQ: chloroquine.

1259

1260   **Supplementary Figure 8. Chloroquine treatment results in loss of activation of LC3B.**

1261

A549 cells were inoculated at a MOI of 5 PFU/cell with PR8 viruses encoding avian or human-

derived M segments and incubated in the presence or absence of 60 µM chloroquine from 1 hpi.

Cells were lysed with whole cell lysis buffer following 8 h incubation at 37^{o}C. Western

immunoblot of virus-infected A549 cells: **(A)** Vinculin expression was measured to allow

normalization of viral protein levels. NP expression was measured to assess viral replication.

Levels of M1 and M2 protein expression were assessed using an antibody (Mab E10) to a

common epitope at the amino terminus of M1 and M2 proteins, allowing relative expression to

be assessed. Levels of LC3B I and II were assessed using an antibody that detects both the

precursor and the activated forms of LC3B protein. Data presented are representative Western

immunoblots from three independent experiments. LC3B I protein and LC3B II protein **(B, C)**

were normalized, quantitated and displayed as a percentage of total LC3B protein in the

absence **(B)** or presence **(C)** of 60 µM chloroquine. Data presented in **B-C**, show the means

with SD from three independent experiments. For each experiment, two replicate radiograms

were quantitated. Statistical significance was assessed using ordinary two-way ANOVA. CQ:

chloroquine.

1277

1278   **Supplementary Figure 9. Schematic depicting chimeric M Segment RNAs**

1279

(A) Pandemic H1N1 influenza virus M1 and M2 proteins differ from the avian consensus

sequences by 9 residues in M1 and 7 residues in M2. The pH1N1 amino acid identities are

indicated in blue at their approximate positions within linear representations of M1 and M2

proteins. The pH1N1 M segment differs from the avian consensus by 8.3% at the nucleotide

level. Here, pH1N1 RNA sequence is indicated by blue coloring. **(B-D)** Chimeric human-avian M

segments were constructed in which only the non-synonymous changes in the avian consensus

were introduced into the A/NL/602/09 (H1N1) M segment. **(E)** Avian consensus M1 and M2

proteins differ from the NL09 sequences by 9 residues in M1 and 7 residues in M2. The avian

amino acid identities are indicated in red and avian RNA sequence is indicated by red coloring.

(F-H) A second set of chimeric M segments was constructed in which the NL09 amino acid

identities were introduced into the M segment of A/duck/Alberta/76 (H1N1) virus, yielding

segments that encode avian consensus protein(s) but retain much of the nucleotide sequence

of the NL09 M segment.

1293

1294   **Supplementary Figure 10. Visualization of LC3 and M2 co-localization in 293T cells**

1295   **infected with viruses carrying avian-human chimeric M segments.**

1296

293T cells were transduced with GFP-LC3 protein and inoculated 24 h later with the indicated

IAVs, encoding avian, human or chimeric M segments, at a MOI of 5 PFU/cell. Cells were fixed

12 h later and stained with anti-M2 (Mab E10; red) and DAPI (blue) followed by imaging with

confocal microscopy. Examples of optical sections are shown, either as merged 3-color images

or the red, green, and blue channels alone (in grey scale). 3x magnification of 63x images are

shown. Brightness was adjusted for optimal clarity, with all images treated equally.

1303

1304 **Supplementary Figure 11. Levels of mRNA$_7$ (encoding M1) and mRNA$_{10}$ (encoding M2)**

1305 **transcripts in A549 cells infected with viruses carrying human, avian or chimeric M**

1306 **segments.**

1307

A549 cells were inoculated at a MOI of 5 PFU/cell with PR8 NL09 M virus and PR8 Av M virus,

along with six PR8-based viruses with chimeric M segments. Cells were incubated at 37^{o}C for 8

h and then lysed with a RNeasy Kit. **(A)** RT primer extension radiogram of virus-infected A549

cells. 5S rRNA levels were measured to allow normalization of viral RNA. Segment 7 vRNA

expression was measured to assess viral replication. **(B)** Levels of mRNA$_7$, **(C)** mRNA$_{10}$, and

(D) mRNA$_{11}$ were quantitated and displayed as a percentage of total M gene expressed mRNA.

Graphs in **B-D** show the means with SD from three independent experiments. For each

experiment, two replicate radiograms were quantitated. Statistical significance was assessed

using ordinary one-way ANOVA.

1317

1318 **Supplementary Figure 12. High expression of M2 protein reduces kinetics of replication**

1319 **of PR8-based viruses in guinea pigs.**

1320

Groups of four guinea pigs were inoculated with 10 PFU of each chimeric M-encoding virus, as

indicated. Graphs show individual titers obtained from animals used in three independent

experiments. Virus replication in nasal wash of inoculated animals was measured by plaque

titration at days 2, 4, 6, and 8 post-infection and the titers at each time point were plotted (dotted

lines). The differences between PR8 NL09 M 7 mut Av and avian M 9 mut NL encoding viruses

were considered significant. Statistical significance in kinetics of growth was determined by

assessing the interaction of time and virus using repeated measures, two-way, multiple

comparisons ANOVA on mean values, with Bonferroni correction applied to account for

comparison of a limited no of means.

1330

1331 **Supplementary Figure 13. PR8 M2 protein is overexpressed and partly localized in**

1332 **perinuclear vesicles in 293T cells.**

1333

293T cells were inoculated with the indicated IAVs, encoding avian-, human- or PR8-derived M

segments, at a MOI of 5 PFU/cell, or treated with 60 µM chloroquine. Cells were fixed 8 h later

and stained with anti-M2 (Mab E10; red) and DAPI (blue) followed by imaging with confocal

microscopy. Examples of optical sections are shown, either as merged 2-color images or the

red and blue channels alone (in grey scale). 63x magnification with 3x optical zoom. Brightness

was adjusted for optimal clarity, with all images treated equally. CQ: chloroquine.

Figure 1

Figure 8

Figure 12

A

B

M mRNA

Excess splicing

M2 mRNA

Excess M2

High proton channel activity in autophagic vesicles

Failure to fuse with lysosomes leads to accumulation of autophagosomes

Restriction of avian IAV in mammalian cells