Spatial density estimates of Eurasian lynx (*Lynx lynx*) in the French Jura and Vosges Mountains

Olivier Gimenez[1], Sylvain Gatti[2], Christophe Duchamp[3], Estelle Germain[4], Alain Laurent[2], Fridolin Zimmermann[5], Eric Marboutin[2]

[1]CEFE, CNRS, Univ Montpellier, Univ Paul Valéry Montpellier 3, EPHE, IRD, Montpellier, France

[2]Office National de la Chasse et de la Faune Sauvage, ZI Mayencin, Gières, France.

[3]Office National de la Chasse et de la Faune Sauvage, Parc Micropolis, Gap, France.

[4]Centre de Recherche et d'Observation sur les Carnivores (CROC), 4 rue de la banie, 57590, Lucy, France

[5]KORA, Thunstrasse 31, 3074, Muri, Switzerland

Abstract

Reliable density estimates are the cornerstone of effective conservation and management planning. For large carnivores however, estimating density is difficult because these species are elusive and wide-ranging. Here, we focus on the Eurasian lynx (*Lynx lynx*) in the French Jura and Vosges Mountains (Mts) and provide the first density estimates. We set up a total of 413 camera trapping sites (with 2 cameras per site) between January 2011 and April 2016 in seven study areas across seven counties of the French Jura and Vosges Mts. We obtained 592 lynx detections over 19,035 trap days in the Jura Mts and 0 detection over 6,804 trap days in the Vosges Mts. Based on coat patterns, we identified a total number of 92 unique individuals from photographs, including 16 females, 13 males and 63 individuals of unknown sex. Using

spatial capture-recapture (SCR) models, we estimated abundance in the study areas between 5 (SE = 0.1) and 29 (0.2) lynx and density between 0.24 (SE = 0.02) and 0.91 (SE = 0.03) lynx per 100 km^2. Our study is yet another example of the advantage of combining SCR methods and non-invasive sampling techniques to estimate density for elusive and wide-ranging species like large carnivores. While the estimated densities in the French Jura Mts are comparable to other lynx populations in Europe, the situation in the Vosges Mts is alarming. Connectivity should be encouraged between the French Jura Mts and the Palatinate Forest in Germany where a reintroduction program is currently ongoing. Our density estimates will help in setting a baseline conservation status for the lynx population in France.

Introduction

Obtaining animal density is a key step in providing sound conservation and management strategies for wildlife [1]. For large carnivores however, estimating density is difficult because these species are elusive and wide-ranging, resulting in low detection rates [2]. To deal with these issues, non-invasive techniques, such as camera trapping and DNA sampling, are increasingly used [3]. These non-invasive techniques generate data that can be analyzed with capture-recapture methods to estimate densities [4].

Standard capture-recapture models for closed populations [5] have long been used to estimate animal abundance and density, including for large carnivores [6,7]. However, when converting abundance into density, density estimates are highly sensitive to the size of the sampling area [8]. Besides, heterogeneity in the detection due to spatial variation in the distance of individuals to the sampling devices may lead to biased density estimates [5]. Spatial capture-recapture (SCR) models deal with these issues by explicitly incorporating

spatial locations of detections [9–12]. SCR models are increasingly used to estimate large carnivores densities [13–18].

Here, we focus on the Eurasian lynx (*Lynx lynx*) in the French Jura and Vosges Mountains (Mts). Like in many regions of western Europe [19], lynx went extinct in France between the 17[th] and 20[th] centuries due to habitat degradation, human persecution and decrease in prey availability [20]. The species was reintroduced in Switzerland in the 1970s [21], then re-colonized France through the French Jura Mts in the 1980s [20]. Reintroductions also occurred in the French Vosges Mts between 1983 and 1993 contributing to the population establishment [22]. The species is listed as endangered under the 2017 IUCN Red list and is of conservation concern in France due to habitat fragmentation, poaching and collisions with vehicles. Currently, the French population of lynx is restricted to three mountain ranges, the Vosges, in northeastern France, the Jura and the Alps, with little connectivity between the first two due to human-made linear infrastructures. While the Northern Alps are slowly being recolonized with lynx mostly coming from the Jura [23], the Jura holds the bulk of the French lynx population. In contrast, the lynx presence in the Vosges Mts remained stable following the reintroductions but and has continuously been decreasing since 2005.

However, little information on abundance and density of lynx in France exist. In this study, we used SCR and standard capture-recapture models to provide the first estimate of lynx abundance and density using camera-trap surveys implemented in the French Jura and Vosges Mts from 2011 to 2016. Based on these results, we discuss research and management priorities for the effective conservation of lynx in France.

Methods

Ethics statement

We used non-invasive methods for data collection, which did not involve manipulation or handling of any living organism. Therefore, approval from an animal ethics committee was not required. Cameras were set on public or private forests with the permission of local authorities or local owners respectively. We advertised the study and the presence of camera traps to the local stakeholders and the public visiting the areas. In agreement with French legislation, photographs permitting to identify goods or people were deleted.

Study area and sampling design

The study area encompassed three counties of the French Jura Mts, namely Ain, Doubs and Jura and four counties of the Vosges Mts, namely Vosges, Haut-Rhin, Bas-Rhin and Moselle (Figure 1). Elevation ranged from 163 and 1,718 m above sea level in the Jura Mts and from 104 to 1,422 m in the Vosges Mts. The human population reached an average density of 88 inhabitants per km^2 in the Jura Mts and 170 inhabitants per km^2 in the Vosges Mts. Forests cover 39% on average of the Jura Mts and 38% of the Vosges Mts. Sampling occurred over 6 years, from 2011 to 2016. We considered two study areas in 2011, 2014 and 2015, three study areas in 2013 and one study area in 2012 and 2016 through camera trapping (Figure 1).

[Figure 1 about here]

We divided each study area into a grid of 2.7 x 2.7 km cells applying a systematic design where one out of two cells was sampled [24], hence ensuring that at least one camera trap was set in each potential lynx home range. To maximize detectability, we set camera traps at optimal locations (on forest roads, hiking trails and to a lesser extent on game paths) based on previous signs of lynx presence and on local knowledge [25]. At each trapping

location, we set two Xenon white flash camera traps (models: Capture, Ambush and Attack; Cuddeback, WI, USA) with passive infrared trigger mechanisms to photograph both flanks of an animal. We checked camera traps weekly to change memory cards, batteries and to remove fresh snow after heavy snowfall. Based on unique coat patterns, we identified the lynx individually on photographs [26] using Extractcompare© software (http://conservationresearch.org.uk/Home/ExtractCompare/lynx.html). We recorded date, time, and location of each photographic capture of a lynx. During the time of year our study took place; juvenile lynx (< 1 year old) can still be with their mother [27]. In our analysis, we retained only independent lynx, i.e. adult lynx or emancipated individuals based on physical characteristics or previous knowledge of their age or status (from photographic evidence). We defined a capture occasion as 5 successive trap nights [25], dissociating trapping events from individual photo to avoid pseudo-replications.

Spatial capture-recapture analyses

We used spatial capture-recapture (SCR) models to estimate lynx densities [4]. In contrast with standard (non-spatial) capture-recapture models, SCR models use the spatial locations of captures to infer activity center (or home range) of each individual. We assumed that individual encounters are Bernoulli random variables with individual- and trap-specific detection probabilities. More precisely, the detection probability p_{ij} of an individual i at trap j is assumed to decrease as the distance d_{ij} from its activity center increases according to a detection function. We used the half-normal detection function, $p_{ij} = p_0 \exp(- d_{ij}^2/(2\sigma^2))$, where p_0 is the probability of detecting an individual when the trap is located exactly at its center of activity and σ is the spatial scale (or movement) parameter that controls the shape of the detection function. For one of the two study areas in the French Jura Mts in years 2011 and 2013, we detected only few individuals. To increase the sample size, we shared

information between the two sampling areas by estimating density separately for each study area but using common detection and spatial parameters for both areas (e.g., [28]). We defined a state-space, i.e. the area encompassing all potential activity centers of the observed individuals, by building a grid that buffered outermost camera trap locations by 15 km (corresponding to at least 2σ [4]) with a resolution of 1.5 km (or pixels of area 2.25 km^2). We fitted SCR models in the maximum likelihood framework using the R package oSCR [29].

For comparison, we also estimated abundance using standard (non-spatial) capture-recapture models [5]. We dropped the spatial information and considered only the detections and non-detections for each individual. We considered two models, M0 in which the detection probability is the same for all individuals, and Mh in which the detection probability varies among individuals. We fitted standard models in the maximum likelihood framework using the R package Rcapture [30]. We estimated density as the ratio of estimated abundance over an effective trapping area (ETA). ETA was estimated by adding a buffer to the trapping area equal to the mean maximum distance moved (MMDM) or half of it (HMMDM). We calculated the MMDM by averaging the maximum distances between capture locations for all individuals detected at more than one site.

Results

We collected data between January 2011 and April 2016 from 413 camera trapping sites (2 camera traps were set per site) resulting in 25,839 trap days (Table 1). In total, we identified 92 lynx over 532 detection events in the Jura Mts, including 16 females, 13 males and 63 individuals of unknown sex. The number of detections per individual was 2.6 on average and varied from 1 up to 11. In contrast, we identified no lynx in the Vosges Mts, therefore we did not proceed with SCR analyses in this area.

[Table 1 about here]

For the Jura Mts, abundance estimates were similar whether we used spatial or non-spatial models, although always slightly higher for the former. Estimated abundance varied between 5 (SE = 0.1) and 29 (0.2) lynx in the spatial analyses, between 4 (0.7) and 23 (0.7) with model M0, and between 5 (1.7) and 28 (3.6) with model Mh. Estimated density varied between 0.24 (0.02) and 0.91 (0.03) lynx per 100 km^2 in the spatial analyses (Table 2). In the non-spatial analyses, the density varied between 0.31 (0.05) and 0.78 (0.02) lynx per 100 km^2 under model M0 and between 0.34 (0.06) and 0.95 (0.12) under model Mh when the MMDM was used. When we used HMMDM, the density varied between 0.57 (0.10) and 1.46 (0.16) lynx per 100 km^2 under model M0 and between 0.67 (0.12) and 1.43 (0.16) under model Mh.

[Table 2 about here]

From the spatial analyses, we used the model estimates to produce density surfaces within the state-space (Figure 2). The density per pixel of area 2.25 km^2 ranged from 0 to 0.20 individuals in the Jura Mts.

[Figure 2 about here]

Discussion

By using camera-trap sampling and SCR models, we provided the first multi-site density estimates for lynx that will help in setting a baseline conservation status for the French lynx

population. Our study is yet another example of the potential of combining SCR methods and non-invasive sampling techniques to estimate abundance and density for elusive and wide-ranging species like large carnivores [13–18].

Our density estimates in the French Jura Mts were of similar magnitude to other lynx populations in Europe: 1.47 and 1.38 lynx / 100 km^2 in the Northwestern Swiss Alps [13], 0.58 (Štiavnica Mts) and 0.81 individuals / 100 km^2 (Velká Fatra National Park) in Slovakia [31] and 0.9 individuals / 100 km^2 in the Bavarian Forest National Park in Germany [32].

While [13] and [31] used SCR models, [32] used standard capture-recapture models with HMMDM to estimate densities, which makes them difficult to compare. Indeed, in other carnivores studies, the use of HMMDM did produce similar density estimates to SCR models [13], while in others, including ours, the SCR estimates were closer to the MMDM estimates [2] or intermediate between the MMDM and HMMDM estimates [33]. When looking at reference values for densities across the distribution range of the species, it may be biologically meaningful to use the MMDM density estimate as a reference as it covers the whole potential of animal movements.

On the other hand, because SCR models make space explicit whereas standard model-based densities are sensitive to the definition of the effective sampling area, we recommend the use of SCR models to estimate lynx densities.

When examining densities across study areas in the Jura Mts, we found spatial variation between the three counties, with Doubs area having the lowest densities, Ain the highest densities, and Jura intermediate densities.

Our lynx density estimates might suffer from potential sources of bias that need to be discussed. First, the period of sampling is important to account for when setting up camera trap surveys [34]. We conducted our survey outside the dispersal period, during the lynx mating season (February-March mostly). We did so to avoid capturing transient individuals

and to increase detectability because of high lynx activity and relatively reduced human activities [26]. However, some individuals might have gone in and out the study area, especially males who cover greater distances during the mating season. Whereas the presence of non-resident individuals can affect the calculation of (H)MMDM, and in turn density estimated with standard capture-recapture models, SCR density estimates were found to be robust to the presence of transient individuals [35]. Second, males have larger home ranges than females [13], which leads to heterogeneity in the SCR model parameter estimates. Because there were too few males and females identified and lots of individuals with unknown sex, sex-specific SCR analyses [36] produced unreliable abundance and density estimates (results not shown). If detection heterogeneity is ignored in capture-recapture models, abundance is underestimated [37], therefore our density estimates are probably biased low and should be considered as a conservative metric. Last, we did not detect any individuals in the Vosges Mts while the sampling effort in the Vosges Mts was similar to that implemented in the Jura Mts (Table 1). This finding is likely to be representative of the current critical situation in the Vosges Mts despite the reintroduction program conducted 30 years ago.

We envision several perspectives to our work. First, while density estimates are of primary interest for conservation, understanding the mechanisms underlying trends in abundance is required to make sound conservation decisions [1]. SCR models have been extended to open populations [38] and can be used to estimate demographic parameters (survival, reproduction) of large carnivores [39]. Unfortunately, because of logistic constraints, we could not sample the same areas over several years, which precludes a standard application of these models. A solution may lie in the combination of the data we collected through systematic camera-trap surveys with additional data in the SCR framework, such as occupancy data [40] or opportunistic camera-trap data [41]. Second, in addition to

traffic-induced mortality and conflicts with human activities, the expansion of lynx populations is limited by habitat fragmentation [42], hence the need to assess connectivity with other populations [43]. SCR models can be used to quantify landscape connectivity by replacing the Euclidean distance between camera traps and home range centers by the least-cost path [44,45]. For lynx, this will require setting up traps across a gradient of habitat types, not only forested habitats, and also to assess the effects of wildlife crossings, so that resistance to movement can be estimated.

In conclusion, our lynx density estimates for the French Jura Mts complement nicely the estimates recently provided for the Northwestern Swiss Alps [13]. The use of camera-trapping coupled with SCR models in both France and Switzerland was the result of a cooperation between the two countries with the perspective of a transboundary monitoring [46,47]. This approach would prove useful to accurately estimate densities in other areas where habitats and prey availability might differ, and overall lynx detectability varies.

The case can be made for monitoring the return of the lynx in the French Alps. Indeed, small-scale camera-trapping surveys and opportunistic observations are currently active and producing signs of lynx presence. However, the lack of a coordinated and intensive sampling effort prevents from estimating abundance and density and inferring trends.

In contrast, the situation in the Vosges Mts is alarming with no individuals detected over the study period. The Vosges Mts being located between the French Jura Mts and the Palatinate Forest in Germany where a reintroduction program is ongoing (program LIFE13 NAT/DE/000755), the lynx colonization in the Vosges Mts remains possible both by the north and the south. Incidentally, two cases of lynx dispersal in the Vosges Mts from neighboring mountains have been recently observed ([48]; program LIFE13 NAT/DE/000755). In this context, obtaining accurate and comparable lynx densities will be crucial to closely monitor population trends at the national scale.

Acknowledgments

We thank the staff from the French National Game and Wildlife Agency (ONCFS), the CROC, the Forest National Agency, the "Directions Départementales des Territoires", the "Fédérations Départementales des Chasseurs" the Regional Natural Parks, the environmental protection associations and all the volunteers from the "Réseau Loup Lynx" who collected the photographs during the camera-trapping session. OG was funded by CNRS and the "Mission pour l'Interdisciplinarité" through the "Osez l'Interdisciplinarité" initiative. CD, EM and SG were funded by ONCFS. CEFE and CROC were funded by CIL&B, MTES (ITTECOP) and FRB through the research program ERC-Lynx. CROC was funded by the European Union within the framework of the Operational Program FEDER-FSE "Lorraine et Massif des Vosges 2014–2020", the "Commissariat à l'Aménagement du Massif des Vosges" for the FNADT ("Fonds National d'Aménagement et de Développement du Territoire"), the DREAL Grand Est ("Direction Régionale pour l'Environnement, l'Aménagement et le Logement"), the "Région Grand Est", the "Zoo d'Amnéville", the "Fondation d'entreprise UEM", the "Fondation Nature & Découvertes", the "Fondation Le Pal Nature", and the *Chasseur d'images* magazine. This study could not have been conducted without authorizations from and agreements with municipalities, environmental managers and owners.

Author contributions

OG wrote the paper and all co-authors commented on the manuscript. OG and SG analyzed the data. AL, CD, EG, EM and SG coordinated the study designs, the data collection and interpretation, with help from FZ for setting the experimental design in the Jura Mts.

References

1. Williams BK, Nichols JD, Conroy MJ. Analysis and management of animal populations. San Diego: Academic Press; 2002.

2. Obbard ME, Howe EJ, Kyle CJ. Empirical comparison of density estimators for large carnivores. Journal of Applied Ecology. 2010;47: 76–84. doi:10.1111/j.1365-2664.2009.01758.x

3. Kelly M, Betsch J, Wultsch C, Mesa B, Mills L. Noninvasive Sampling for Carnivores. Carnivore Ecology and Conservation: A Handbook of Techniques. Oxford University Press. 2012. pp. 47–69.

4. Royle JA, Chandler RB, Sollmann R, Gardner B. Spatial Capture-Recapture [Internet]. New York, USA: Academic Press, New York; 2014. doi:10.1016/C2012-0-01222-7

5. Otis DL, Burnham KP, White GC, Anderson DR. Statistical Inference from Capture Data on Closed Animal Populations. Wildlife Monographs. The Wildlife Society; 1978. pp. 3–135. doi:10.2307/2287873

6. Mumma MA, Zieminski C, Fuller TK, Mahoney SP, Waits LP. Evaluating noninvasive genetic sampling techniques to estimate large carnivore abundance. Molecular Ecology Resources. 2015;15: 1133–1144. doi:10.1111/1755-0998.12390

7. Gerber BD, Ivan JS, Burnham KP. Estimating the abundance of rare and elusive carnivores from photographic-sampling data when the population size is very small. Population Ecology. 2014;56: 463–470. doi:10.1007/s10144-014-0431-8

8. White GC, Anderson DR, Burnham KP, Otis DL. Capture-recapture and removal methods for sampling closed populations [Internet]. Los Alamos, NM: Los Alamos National Laboratory; 1982 p. 14. Available: http://pubs.er.usgs.gov/publication/70120390

9. Royle JA, Young KV. A hierarchical model for spatial capture-recapture data. Ecology. 2008;89: 2281–2289. doi:10.1890/07-0601.1

10.   Borchers DL, Efford MG. Spatially Explicit Maximum Likelihood Methods for Capture–Recapture Studies. Biometrics. 2008;64: 377–385. doi:10.1111/j.1541-0420.2007.00927.x

11.   Efford M. Density estimation in live-trapping studies. Oikos. 2004;106: 598–610. doi:10.1111/j.0030-1299.2004.13043.x

12.   Borchers D. A non-technical overview of spatially explicit capture-recapture models. Journal of Ornithology. 2012;152: 435–444.

13.   Pesenti E, Zimmermann F. Density estimations of the Eurasian lynx (*Lynx lynx*) in the Swiss Alps. Journal of Mammalogy. 2013;94: 73–81. doi:10.1644/11-MAMM-A-322.1

14.   Stetz JB, Mitchell MS, Kendall KC. Using spatially-explicit capture-recapture models to explain variation in seasonal density patterns of sympatric ursids. Ecography. 2018; doi:10.1111/ecog.03556

15.   López-Bao JV, Godinho R, Pacheco C, Lema FJ, García E, Llaneza L, et al. Toward reliable population estimates of wolves by combining spatial capture-recapture models and non-invasive DNA monitoring. Scientific Reports. 2018;8. doi:10.1038/s41598-018-20675-9

16.   Goldberg JF, Tempa T, Norbu N, Hebblewhite M, Mills LS, Wangchuk TR, et al. Examining Temporal Sample Scale and Model Choice with Spatial Capture-Recapture Models in the Common Leopard Panthera pardus. Aegerter CM, editor. PLOS ONE. 2015;10: e0140757. doi:10.1371/journal.pone.0140757

17.   Broekhuis F, Gopalaswamy AM. Counting Cats: Spatially Explicit Population Estimates of Cheetah (Acinonyx jubatus) Using Unstructured Sampling Data. Festa-Bianchet M, editor. PLOS ONE. 2016;11: e0153875. doi:10.1371/journal.pone.0153875

18.   Alexander JS, Gopalaswamy AM, Shi K, Riordan P. Face Value: Towards Robust Estimates of Snow Leopard Densities. Margalida A, editor. PLOS ONE. 2015;10: e0134815. doi:10.1371/journal.pone.0134815

19.   Breitenmoser U. Large predators in the Alps: The fall and rise of man's competitors. Biological Conservation. 1998;83: 279–289. doi:10.1016/S0006-3207(97)00084-0

20.   Vandel J-M, Stahl P. Distribution trend of the Eurasian lynx Lynx lynx populations in France. Mammalia. 2005;69. doi:10.1515/mamm.2005.013

21.   Breitenmoser U, Breitenmoser-Würsten C, Capt S. Re-introduction and present status of the lynx (Lynx Lynx) in Switzerland. Hystrix It J Mamm. 1998;10. doi:10.4404/hystrix-10.1-4118

22.   Vandel J-M, Stahl P, Herrenschmidt V, Marboutin E. Reintroduction of the lynx into the Vosges mountain massif: From animal survival and movements to population development. Biological Conservation. 2006;131: 370–385. doi:10.1016/j.biocon.2006.02.012

23.   Marboutin E, Duchamp C, Moris P, Briaudet P-E, Léonard Y, Catusse M. Survey of the Lynx distribution in the French Alps□: 2005-2009 update. Acta biologica Slovenica. 2012: 29–34.

24.   Zimmermann F, Breitenmoser-Würsten C, Molinari-Jobin A, Breitenmoser U. Optimizing the size of the area surveyed for monitoring a Eurasian lynx (Lynx lynx) population in the Swiss Alps by means of photographic capture-recapture. Integr Zool. 2013;8: 232–243. doi:10.1111/1749-4877.12017

25.   Blanc L, Marboutin E, Gatti S, Gimenez O. Abundance of rare and elusive species: Empirical investigation of closed versus spatially explicit capture-recapture models with lynx as a case study. The Journal of Wildlife Management. 2013;77: 372–378. doi:10.1002/jwmg.453

26.   Zimmermann F, Foresti D. Capture-recapture methods for density estimation. Camera Trapping for Wildlife Research. Pelagic Publishing. Exeter, UK; 2016. pp. 95–141.

27.   Zimmermann F, Breitenmoser-Würsten C, Breitenmoser U. Natal dispersal of

Eurasian lynx (Lynx lynx) in Switzerland. Journal of Zoology. 2005;267: 381–395. doi:10.1017/S0952836905007545

28. Rocha DG da, Sollmann R, Ramalho EE, Ilha R, Tan CKW. Ocelot (Leopardus pardalis) Density in Central Amazonia. Hagen CA, editor. PLOS ONE. 2016;11: e0154624. doi:10.1371/journal.pone.0154624

29. Sutherland C, Muñoz DJ, Miller DAW, Grant EHC. Spatial Capture–Recapture: A Promising Method for Analyzing Data Collected Using Artificial Cover Objects. Herpetologica. 2016;72: 6–12. doi:10.1655/HERPETOLOGICA-D-15-00027

30. Baillargeon S, Rivest L-P. Rcapture: Loglinear Models for Capture-Recapture in R. Journal Of Statistical Software. Citeseer; 2007;19: 1–31.

31. Kubala J, Smolko P, Zimmermann F, Rigg R, Tám B, Iľko T, et al. Robust monitoring of the Eurasian lynx Lynx lynx in the Slovak Carpathians reveals lower numbers than officially reported. Oryx. 2017; 1–9. doi:10.1017/S003060531700076X

32. Weingarth K, Heibl C, Knauer F, Zimmermann F, Bufka L, Heurich M. First estimation of Eurasian lynx (Lynx lynx) abundance and density using digital cameras and capture–recapture techniques in a German national park. Animal Biodiversity and Conservation. 2012; 11.

33. Reppucci J, Gardner B, Lucherini M. Estimating detection and density of the Andean cat in the high Andes. Journal of Mammalogy. 2011;92: 140–147. doi:10.1644/10-MAMM-A-053.1

34. Weingarth K, Zeppenfeld T, Heibl C, Heurich M, Bufka L, Daniszová K, et al. Hide and seek: extended camera-trap session lengths and autumn provide best parameters for estimating lynx densities in mountainous areas. Biodiversity and Conservation. 2015;24: 2935–2952. doi:10.1007/s10531-015-0986-5

35. Royle JA, Fuller AK, Sutherland C. Spatial capture–recapture models allowing

Markovian transience or dispersal. Population Ecology. 2016;58: 53–62. doi:10.1007/s10144-015-0524-z

36.   Sollmann R, Furtado MM, Gardner B, Hofer H, Jácomo ATA, Tôrres NM, et al. Improving density estimates for elusive carnivores: Accounting for sex-specific detection and movements using spatial capture–recapture models for jaguars in central Brazil. Biological Conservation. 2011;144: 1017–1024. doi:10.1016/j.biocon.2010.12.011

37.   Cubaynes S, Pradel R, Choquet R, Duchamp C, Gaillard J-M, Lebreton J-D, et al. Importance of accounting for detection heterogeneity when estimating abundance: the case of French wolves. Conservation Biology. 2010;24: 621–6. doi:10.1111/j.1523-1739.2009.01431.x

38.   Gardner B, Reppucci J, Lucherini M, Royle JA. Spatially explicit inference for open populations: Estimating demographic parameters from camera-trap studies. Ecology. 2010;91: 3376–3383.

39.   Whittington J, Sawaya MA. A Comparison of Grizzly Bear Demographic Parameters Estimated from Non-Spatial and Spatial Open Population Capture-Recapture Models. Boyce MS, editor. PLOS ONE. 2015;10: e0134446. doi:10.1371/journal.pone.0134446

40.   Blanc L, Marboutin E, Gatti S, Zimmermann F, Gimenez O. Improving abundance estimation by combining capture-recapture and occupancy data: Example with a large carnivore. Journal of Applied Ecology. Blackwell Publishing Ltd; 2014;51: 1733–1739.

41.   Tenan S, Pedrini P, Bragalanti N, Groff C, Sutherland C. Data integration for inference about spatial processes: A model-based approach to test and account for data inconsistency. Rocchini D, editor. PLOS ONE. 2017;12: e0185588. doi:10.1371/journal.pone.0185588

42.   Kramer-Schadt S, Revilla E, Wiegand T, Breitenmoser U. Fragmented landscapes, road mortality and patch connectivity: Modelling influences on the dispersal of Eurasian lynx.

Journal of Applied Ecology. 2004;41: 711–723. doi:10.1111/j.0021-8901.2004.00933.x

43.   Zimmermann F, Breitenmoser U. Potential distribution and population size of the Eurasian lynx Lynx lynx in the jura Mountains and possible corridors to adjacent ranges. Wildlife Biology. 2007;13: 406–416. doi:10.2981/0909-6396(2007)13[406:PDAPSO]2.0.CO;2

44.   Royle JA, Chandler RB, Gazenski KD, Graves TA. Spatial capture–recapture models for jointly estimating population density and landscape connectivity. Ecology. 2013;94: 287–294. doi:10.1890/12-0413.1

45.   Sutherland C, Fuller AK, Royle JA. Modelling non-Euclidean movement and landscape connectivity in highly structured ecological networks. Methods in Ecology and Evolution. 2015;6: 169–177. doi:10.1111/2041-210X.12316

46.   Gervasi V, Brøseth H, Gimenez O, Nilsen EB, Odden J, Flagstad ., et al. Sharing data improves monitoring of trans-boundary populations: The case of wolverines in central Scandinavia. Wildlife Biology. 2016;22: 95–106. doi:10.2981/wlb.00142

47.   Vitkalova AV, Feng L, Rybin AN, Gerber BD, Miquelle DG, Wang T, et al. Transboundary cooperation improves endangered species monitoring and conservation actions: A case study of the global population of Amur leopards. Conservation Letters. 2018; e12574. doi:10.1111/conl.12574

48.   Hurstel A, Laurent A. Première preuve de dispersion du Lynx d'Eurasie (Lynx lynx) du Jura vers les Vosges. Ciconia. 2016: 1–6.

Figures

Figure 1: Map of the study area in the French Jura and Vosges Mts. The study area encompassed seven counties (Ain, Jura and Doubs in the Jura Mts and Vosges, Haut-Rhin, Bas-Rhin and Moselle in the Vosges Mts) that were monitored through 418 camera trapping sites (298 in the Jura Mts and 115 in the Vosges Mts; two camera traps were set per site), each within a 2.7 x 2.7 km cell.

Figure 2: Lynx density maps in the French Jura Mts. The density scale is in lynx per 2.25 km^2 (pixel resolution is 1500m x 1500m). We obtain the estimated abundance in each map by summing up the densities in each pixel altogether. Yellow is for low densities, green for medium densities and blue for high densities. Note that density scales are specific to each map.

Table 1: Main characteristics and results of the lynx camera-trap survey carried out in a) the French Jura Mts and b) the French Vosges Mts.

a. Year/County	2011/Doubs	2011/Jura	2012/Jura & Doubs	2013/Doubs	2013/Ain & Jura	2014/Ain	2015/Ain
Period of trap activity	Jan-Apr	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr	Feb-Apr	Feb-May
Number of active camera traps	48	66	148	44	142	118	30
Number of trapping days (average/area)	63	59	69	63	58	59	99
Number of capture occasions	15	15	17	14	13	13	21
Number of detections	22	42	130	25	117	158	38
Number of detected individuals	4	9	21	6	19	23	10
Number of females, unknown, males	1, 1, 2	1, 7, 1	2, 14, 5	1, 4, 1	2, 13, 4	4, 16, 3	2, 8, 0
Number of detections / ind: mean, min, max	3, 2, 4	2.8, 1, 6	2.5, 1, 10	2.7, 1, 6	3.6, 1, 11	3.3, 1, 9	2.2, 1, 5

b. Year/County	2013/Haut-Rhin & Vosges	2014/Bas-Rhin & Moselle	2015/Bas-Rhin & Moselle	2016/Bas & Haut-Rhin
Period of trap activity	Dec-Jan	Feb-Apr	Feb-Apr	Feb-Apr
Number of active traps	60	50	60	60
Number of trapping days (average/area)	52	59	57	59
Number of detections	0	0	0	0

Table 2: Lynx abundance and density estimates obtained from spatial and non-spatial capture-recapture analyses of camera-trapping data collected in the French Jura Mts. Densities are provided in number of lynx per 100 km^2. ETA is for Effective Trapping Area, MMDM for Mean Maximum Distance Moved, HMMDM for Half Mean Maximum Distance Moved, SCR for spatial capture-recapture, M0 for the (non-spatial) capture-recapture model with homogeneous detection probability, Mh for the (non-spatial) capture-recapture model with heterogeneous detection probability and SE for standard error. Note: For 2011 and 2013, parameters of the spatial capture-recapture model (p_0 and σ) are common to both areas in each year.

Year/County	2011/Doubs	2011/Jura	2012/Jura-Doubs	2013/Doubs	2013/Ain-Jura	2014-Ain	2015-Ain
SCR abundance (SE)	5 (0.1)	12 (0.1)	29 (0.2)	7 (0.1)	21 (0.1)	29 (0.2)	12 (0.1)
SCR density (SE)	0.24 (0.02)	0.44 (0.02)	0.67 (0.02)	0.36 (0.02)	0.54 (0.02)	0.91 (0.03)	0.64 (0.03)
p_0 logit scale (SE)	-2.94 (0.24)		-2.01 (0.20)	-2.57 (0.20)		-2.34 (0.19)	-3.01 (0.42)
σ log scale (SE)	8.89 (0.14)		8.54 (0.08)	8.95 (0.06)		8.80 (0.07)	8.97 (0.19)
M0 abundance (SE)	4 (0.7)	9 (0.7)	21 (0.6)	6 (0.3)	19 (0.8)	23 (0.7)	11 (1.2)
Mh abundance (SE)	5 (1.7)	10 (1.8)	25 (2.8)	7 (1.2)	25 (4.1)	28 (3.6)	11 (1.2)
MMDM (km)	9.1	16.2	8.9	9.1	18.2	13.6	12.1
ETA with MMDM (km^2)	1991	2930	3089	1171	4954	2936	1549

M0 density MMDM (SE)	0.31 (0.05)	0.31 (0.02)	0.68 (0.02)	0.51 (0.02)	0.38 (0.02)	0.78 (0.02)	0.71 (0.08)
Mh density MMDM (SE)	0.39 (0.13)	0.34 (0.06)	0.81 (0.09)	0.60 (0.10)	0.50 (0.08)	0.95 (0.12)	0.70 (0.08)
ETA with HMMDM (km^2)	697	1491	2111	659	2673	1668	753
M0 density HMMDM (SE)	0.57 (0.10)	0.60 (0.05)	0.99 (0.03)	0.91 (0.05)	0.71 (0.03)	1.38 (0.04)	1.46 (0.16)
Mh density HMMDM (SE)	0.72 (0.24)	0.67 (0.12)	1.18 (0.13)	1.06 (0.18)	0.93 (0.15)	1.68 (0.21)	1.43 (0.16)

Figure 1

Figure 2