Peptidoglycan layer and disruption processes in *Bacillus subtilis* cells visualized using quick-freeze, deep-etch electron microscopy

Isil Tulum[a,b,#], Yuhei O Tahara[a,b,#], Makoto Miyata[a,b,*]

[a] Graduate School of Science, Osaka City University, Osaka, 558-8585, Japan.

Tel +81 6 6605 3157, Fax +81 6 6605 3158.

[b] The OCU Advanced Research Institute for Natural Science and Technology (OCARINA), Osaka City University, Osaka, 558-8585, Japan.

Running title: Peptidoglycan layer of *Bacillus subtilis*

*To whom correspondence should be addressed. E-mail: miyata@sci.osaka-cu.ac.jp. ORCID ID: https://orcid.org/0000-0002-7478-7390

[#]I.T. and Y.O.T. contributed equally to this work.

Keywords: Lysozyme, Fluorescence, Penicillin, Piecrust, L-form, Turgor

The total number of pages: 20 pages

The total number of figures: 5

Y.O.T ORCID ID: https://orcid.org/0000-0002-2736-9233

ABSTRACT

Peptidoglycan, which is the main component of the bacterial cell wall, is a heterogeneous polymer of glycan strands crosslinked with short peptides and is synthesized in cooperation with the cell division cycle. Although it plays a critical role in bacterial survival, its architecture is not well understood. Herein, we visualized the architecture of the peptidoglycan surface in *Bacillus subtilis* at the nanometer resolution, using quick-freeze, deep-etch electron microscopy. Filamentous structures were observed on the entire surface of the cell, where filaments about 11-nm wide formed concentric circles on cell poles, filaments about 13-nm wide formed a circumferential mesh-like structure on the cylindrical part, and a "piecrust" structure was observed at the boundary. When growing cells were treated with lysozyme, the entire cell mass migrated to one side and came out from the cell envelope. Fluorescence labeling showed that lysozyme preferentially bound to a cell pole and cell division site, where the peptidoglycan synthesis was not complete. Ruffling of surface structures was observed during electron microscopy. When cells were treated with penicillin, the cell mass came out from a cleft around the cell division site. Outward curvature of the protoplast at the cleft seen using electron microscopy suggested that turgor pressure was applied as the peptidoglycan was not damaged at other positions. When muropeptides were depleted, surface filaments were lost while the rod shape of the cell was maintained. These changes can be explained on the basis of the working points of the chemical structure of peptidoglycan.

INTRODUCTION

Bacteria, the major inhabitants of the Earth, are in a constant battle to outlast

their competitors in the environment and the immune system of host organisms.

Most bacterial cells are surrounded by a rigid shield called the "peptidoglycan

layer," which protects them from chemical agents, including lytic enzymes and

antibiotics, that are produced by their competitors [1-4].

Peptidoglycan is an essential component of the bacterial cell wall that is found

on the outside of the cytoplasmic membrane of almost all bacterial cells except

the class *Mollicutes*. This polymer provides strength, rigidity, and shape stability

by maintaining turgor pressure. In peptidoglycan, the glycan strands comprise

alternating β-1,4-linked *N*-acetylglucosamine (GlcNAc) and *N*-acetylmuramic

acid (MurNAc), and the peptide stems are covalently linked to the glycan

strands with an amide bond to the carboxyl carbon of the MurNAc. The

peptidoglycan layer is also the site of action for antimicrobial agents. Lysozyme,

an antimicrobial enzyme critical in animal host defense, is one of the most

abundant proteins present on the mucosal surfaces and in body secretions,

such as saliva and tears [5, 6]. The epithelial cells secrete lysozyme to protect

the host's mucosal surfaces from infectious bacteria. Lysozyme is also present

in white blood cells, especially in granules of phagocytes, where it helps in the

elimination of infectious bacteria within phagolysosomes. The underlying

mechanism of action of lysozyme involves breaking the bond between GlcNAc

and MurNAc (muramidase activity) leading to the degradation of peptidoglycan.

On the other hand, fungi and bacteria produce secondary metabolites in

defense against predators and competitors [7, 8]. A well-known group of

secondary metabolites is β-lactams, a broad class of antibiotics that include

penicillin derivatives, cephalosporins, monobactams, and carbapenems. β-

lactams inhibit peptidoglycan synthesis by covalently binding to the active site of

transpeptidases, known as penicillin-binding proteins (PBPs), cause changes in

bacterial cell shape and lead to cell lysis.

This property makes peptidoglycan a vitally important target of β-lactam

antibiotics. Therefore, peptidoglycan architecture and the processes that are

used to disrupt it are valuable to understand the survival strategies of bacteria

and to control pathogenic bacteria. However, the architecture of peptidoglycan

is not well understood, because the structure is featured with low density, high

flexibility, and is multilayered, which are characteristics that make it unsuitable

to be observed using transmission electron microscopy (EM) [2, 9-13].

Quick-freeze, deep-etch replica EM was introduced in order to visualize

synaptic transmission processes in 1979, and it has emerged as a useful tool

that can be applied for the visualization of many other biological phenomena

[14, 15]. It is an advanced technology that is used to visualize biological

specimens in an active state as a shot image, with spatial resolution of the order

of nanometers and time resolution of sub milliseconds. In this method, the

specimen is frozen in less than a millisecond by pressing it against a metal

block chilled with liquid helium or liquid nitrogen, resulting in much faster fixation

than with chemical methods [15]. Then, the frozen specimen is exposed by

fracturing and etching, and shadowed by platinum. The metal shadowing gives

a high image contrast, allowing visualization of shot images without image

averaging, unlike cryo EM [16, 17]. Moreover, metal contact freezing can be

applied to various conditions and specimens, with much more flexibility than

with the freezing process for cryo EM. Therefore, quick-freeze, deep-etch

replica EM has great advantages, especially when used to visualize low density

and flexible structures, in comparison to other methods of transmission EM.

Bacillus subtilis is a rod-shaped, Gram-positive, non-pathogenic bacterium

that belongs to the phylum *Firmicutes* [1]. The genus *Bacillus* also includes

human pathogens such as *Bacillus anthracis* and *Bacillus cereus* [18] and is

related to the genus *Clostridium*. Therefore, *B. subtilis* can be an attractive

model for the clarification of the architecture and the roles of the cell wall. In this

study, the detailed structures of the peptidoglycan layer and its disruption

processes in *B. subtilis* were analyzed using the quick-freeze, deep-etch EM

and optical microscopy.

METHODS

Bacterial strains and media. *B. subtilis* 168 CA and LR2, a strain inducible for

L-form derived from *B. subtilis* 168 CA [5, 19] were used. Nutrient agar (NA,

Oxoid) was used for routine selection and maintenance of *B. subtilis* 168 CA.

Luria-Bertani (LB), nutrient broth (NB, Oxoid), and SMM-defined minimal

medium (Spizizen) containing 0.5% xylose or 1 mM isopropyl β-D-1-

thiogalactopyranoside (IPTG) were used when required.

Treatment of cells. Cultured cells around optical density (OD) at 0.2 in 600 nm

wavelength, a middle exponential stage, were collected by centrifugation at

8,000 × *g*, 23 °C for 3 minutes. For lysozyme treatment, the cells were

suspended in phosphate-buffered saline consisting of 75 mM sodium phosphate

(pH 7.3) and 68 mM NaCl to be the original cell density. Lysozyme was added

to the cell suspension at final concentration of 0.1 mg/mL and kept at 37 °C

without shaking. For Penicillin G treatment, cultured cells were collected and

suspended in a fresh medium at the original cell density. PenG was added at

the final concentration of 0.1 mg/mL and kept at 37°C with shaking. To induce

the L-form transition, 2 × MSM osmoprotective medium (40 mM $MgCl_2$, 1 M

sucrose and 40 mM maleic acid, pH 7.0) was mixed with the same volume of 2

× NB or 2 × NA. L-form liquid cultivation was done in NB/MSM at 30°C without

shaking [19]. Lysozyme hydrochloride, from egg white (Wako Pure Chemical

Industries, Osaka, Japan) was labeled with DyLight 488 NHS Ester (Thermo

Fisher Scientific, Rockford, IL), according to the instruction.

Optical microscopy. The cells were inserted into a tunnel chamber with a 5-

mm interior width, a 22-mm length, and an 86-µm wall thickness [20]. The

tunnel chamber was constructed with a coverslip and a glass slide and

assembled with double-sided tape. Fluorescence microscopy was performed

with a BX50 fluorescence microscope equipped with a YFP filter unit (U-

MYFPHQ) and a phase-contrast setup (Olympus, Tokyo, Japan). Images were

captured with a WAT-120NRC charge-coupled-device (CCD) camera (Watec,

Yamagata, Japan) and analyzed using ImageJ 1.52.

Quick-freeze, deep-etch EM. The original cells were collected by centrifugation

at 8,000 × g, at 23 °C for 5 min and suspended in water (original cells) or a

buffer consisting of 10 mM HEPES (pH 7.6), 150 mM NaCl, 1 mM $MgCl_2$, and

0.1 mg/mL DNase I (treated cells) to achieve a 20-fold higher cell density.

DNase I and $MgCl_2$ were included to reduce the viscosity due to bacterial DNA

from cells damaged by lysozyme, PenG or *murE* repression. This process was

repeated and the cell suspension was mixed with a slurry that included mica

flakes, placed on a rabbit lung slab, and frozen by a CryoPress (Valiant

Instruments, St. Louis, MO) cooled by liquid helium [21]. The slurry was used to

retain an appropriate amount of water before freezing. The specimens were

fractured and etched for 15 minutes at -104 °C, in a JFDV freeze-etching device

(JEOL Ltd, Akishima, Japan) [22]. The exposed cells were rotary-shadowed by

platinum at an angle of 20 degrees to be 2 nm in thickness and backed with

carbon. Replicas were floated off on full-strength hydrofluoric acid, rinsed in

water, cleaned with a commercial bleach, rinsed in water, and picked up onto

copper grids as described [23, 24]. Replica specimens were observed by a

JEM-1010 transmission electron microscope (JEOL, Tokyo, Japan) at 80 kV

equipped with a FastScan-F214 (T) CCD camera (TVIPS, Gauting, Germany).

For tomography, cell replicas were observed by Talos F200C G2 (Thermo

Fisher Scientific, Waltham, MA) at 200 kV, and image sets were acquired every

degree of angle for 96 steps, by a complementary metal-oxide-semiconductor

(CMOS) camera (Ceta camera, FEI) [25]. The images were analyzed by

ImageJ.

RESULTS

Surface structure of *Bacillus subtilis*

To visualize the structure of the peptidoglycan layer, *B. subtilis* was observed for

the first time by quick-freeze, deep-etch EM. Cells in the exponential growth

phase were collected by centrifugation and placed on glass, frozen, fractured,

deeply etched, shadowed with platinum, and then the platinum replicas were

recovered and observed. The shape and dimensions of the cells were

consistent with the images of living cells obtained by optical microscopy (Fig.

1A, B). Filamentous structures were clearly observed on the cell surface and

can be distinguished between the cell pole and the cylindrical part of the rod-

shaped cells (Fig. 1C, D and Movie S1). Thick filaments were aligned in a

partial circumferential manner on the cylindrical part and thin filaments were

concentrically arranged around both poles. The filament widths were measured

based on the image profile, assuming that the mid density of the image profile

positions at the boundary of the filament being focused on (Fig. S1). As the

structures were coated by platinum to be 2 nm in thickness, the original filament

width should be 4 nm thinner than the value obtained. Thus, we concluded that

the widths of thick and thin filaments were 8.7 ± 0.4 and 6.9 ± 0.3 nm,

respectively, and the thin filaments were aligned with a pitch of 11.3 ± 0.4 nm

(Fig. 1E, F).

On elongated cells prior to cell division, invagination was observed at the

central position of the axis, and an obvious boundary was found between the

cylindrical part and the surface of the invagination (Fig. 1G). Obvious

filamentous structures were not found on the surface of the invagination. A wall

24.8 ± 1.6 nm wide was observed between the surface of the invagination and

the cylindrical part (Fig. 1G). This small wall was also found in cells after

division (Fig. 1H), but not in isolated single cells (Fig. 1D).

Cells aligned vertically were frequently fractured by a knife, and a fractured

cytosol, membrane surface, and cross section of the peptidoglycan layer were

visualized (Fig. 1I, J). The height of the peptidoglycan layer was estimated as

30.5 ± 4.1 nm (n = 26).

Peptidoglycan disruption by lysozyme

Next, we examined how the peptidoglycan layer is disrupted by lysozyme. We

added lysozyme from the egg white to growing cells, and observed the changes

occurring on the cell structures by phase-contrast optical microscopy (Fig. 2A).

The effects of lysozyme on *B. subtilis* cells were consistent with previous

reports, although we traced the changes in more detail [5, 26, 27]. In the

present observation, the cell mass started to detach from the envelope structure

around 30 minutes after the addition of lysozyme and moved to one side over

time. Finally, all the mass was localized as a sphere at one side of the cell

envelope. After a 60 minute treatment, 100% of the cells (n = 72) were

converted to protoplasts.

To locate the sites where lysozyme works on the cell in this process, we

labeled lysozyme fluorescently with DyLight 488 through amino groups of

protein and using fluorescence microscopy, traced its attachment point (Fig.

2B). After 10 minutes, the labeled lysozyme bound to the whole cell surface but

preferentially bound to the division site and a cell pole. The changes in

lysozyme localization over time were traced through live imaging (Fig. 2C). The

signal found at a pole decreased with time and disappeared at 40 minutes. The

signal on parts other than the cell poles remained for 40 minutes, and then it

moved to one side. The disappearance of lysozyme signal should suggest the

dissociation of lysozyme from envelope, caused by the complete digestion of

the target sites on the peptidoglycan layer.

Next, we observed the structural changes on the cell surface at high

resolution using the quick-freeze, deep-etch EM (Fig. 2D, E). The images

featured filamentous structures covering the entire field. They are likely to be

filaments of eutectic mixtures, resulting from the cytosol released from damaged

cells, because generally, some ions and polymers form filaments of eutectic

mixtures in the quick-freeze, deep-etch EM [28]. This assumption is consistent

with the fact that the covering of filaments increased with the time of treatment

with lysozyme, and PenG and *murE* repression (see below). Eutectic filaments

can be confused with the real surface in images; however, they can be

distinguished because eutectics are continuous from a position distant from the

cell. At 20 minutes after the addition of lysozyme, ruffling of the surface

structure was observed around a cell pole, which is probably corresponding to

the area to which lysozyme bound on the cell. The peptidoglycan layer should

detach from the cell partly by the degradation of the sugar chain. At 60 minutes,

the ruffling parts became restrictive, which is consistent with the observation

that the area of lysozyme binding became restricted in the fluorescence

microscopy. Instead of ruffling, we found coarse patterns on the cylindrical part

of cells and spherical structures with a smooth surface, suggesting that

lysozyme moved and digested the peptidoglycan at the cylindrical parts.

Peptidoglycan disruption by penicillin

The process of peptidoglycan disruption by the effect of penicillin was examined

(Fig. 3). Beginning 30 minutes after the addition of 100 μg/mL Penicillin G

(PenG) to the growing culture, a less dense, flexible structure was observed to

project from a side position on a cell under optical microscopy (Fig. 3A). At 120

minutes, 89% of the cells (n = 172) were released from the peripheral structure

probably as a protoplast.

Next, the cells damaged by PenG treatment for 120 minutes were observed

by the quick-freeze, deep-etch EM (Fig. 3B, C). The cell images showed various

stages of the process by which a protoplast comes out from the cell envelope.

In many cases, the cytoplasm came out from the central position of the cell

where the cell division should be scheduled (Fig. 3C). Unlike the effect of

lysozyme, the cytoplasm in the early stage of cell disruption seemed to be

subjected to strong turgor, because the protoplasts coming through the cleft had

some outward curvature.

Peptidoglycan disruption by *murE* operon repression

The repression of the *murE* operon ceases supplying muropeptides, induces

the gentle degradation of the peptidoglycan layer, and then results in the

transfer of *B. subtilis* cells into L-form [29, 30]. This process was visualized by

phase-contrast optical microscopy (Fig. 4A). When the *murE* operon was

repressed for 80 minutes, the cell morphology was disturbed mostly around cell

poles, featured by a tapered pole, and the detachment of cell mass from the

pole. Time lapse imaging of cells showed that these two features occurred in a

sequential manner. In those cells, finally the cell became spherical, which may

proliferate as L-form. These processes agree with those previously reported

[29, 30]. Next, we visualized the cells by the quick-freeze, deep-etch EM.

Irregular cell shapes were obvious under this method, including asymmetric cell

division, tapered cell pole (Fig. 4B), and spherical extension at a cell pole (Fig.

4C, left). The filaments on the cell surfaces were unclear compared to the

original cells, or totally lacking (Fig. 4C, left and middle). On the spherical cells,

no filaments were found on the surface, supporting our assumption that the

filaments observed in the quick-freeze, deep-etch EM are derived from the

peptidoglycan layer (Fig. 4C, right).

DISCUSSION

Visualization of peptidoglycan layer by quick-freeze, deep-etch EM

The images obtained by the quick-freeze, deep-etch EM may appear like those from Scanning Electron Microscopy (SEM), but they are featured much higher spatial resolution and sub-millisecond time resolution [14, 15]. This method should be very efficient for research of microorganisms whose interaction with the environment on their surface is critical for many activities [15, 31, 32]. In the present study, the quick-freeze, deep-etch EM was applied to the visualization of the peptidoglycan and its disruption process, for the first time. The results showed the details of filamentous structures at nanometer resolution and their change, as expected, with time resolution. In previous studies, peptidoglycan layers have been visualized by other methods including negative-staining EM, cryo electron tomography, sectioning EM, SEM, AFM, and freeze fracture EM. However, the quick-freeze, deep-etch EM has advantages over other methods. As negative-staining EM and cryo electron tomography need to isolate the peptidoglycan layer, the alignment of the structure on the cell cannot be observed [10, 33]. Sectioning EM does not show the images three dimensionally [34, 35]. SEM can provide three dimensional images but chemical fixation may modify the original structure and the resolution of SEM is much lower than that of the quick-freeze, deep-etch EM [14, 15]. Atomic force microscopy (AFM) visualizes the surface three dimensionally but cross-section images are not available and the resolution is lower than that of the quick-freeze, deep-etch EM, especially when the target structure is vertically aligned against the surface [12, 36]. Although the image obtained with freeze fracture EM is similar to that of the quick-freeze, deep-etch EM, the chemical fixation

involved in the former may modify the original structure [37].

In normally-growing cells, the alignments of the peptidoglycan filaments

were clearly distinguished between the poles and the cylindrical part (Figs. 1

and 5A left). This was consistent with the knowledge from molecular biology that

peptidoglycan synthase is localized by MreB when cells are elongating and the

synthase is localized at the division site by FtsZ during cell division [38, 39]. The

concentric pattern in the pole is considered to be common to the structure

previously observed by a related method, the freeze fracture of fixed cells of

Staphylococcus [37]. Atomic Force Microscopy (AFM) of the peptidoglycan

showed circular pattern for *B. subtilis* [12] and also for *Lactococcus lactis* [11].

The circular pattern observed in the present study may be general in *Firmicute*

bacterial species, although the appearance depends on the visualizing

methods. A small wall structure was seen at the boundary between the surfaces

of the invaginating part and the cylindrical part (Fig. 1G, H). This structure

should correspond to a structure named "wall band" observed in sectioned EM

images [34, 35]. A similar structure is known as "piecrust" in SEM observation of

Staphylococcus aureus [37], although it has not been observed in SEM images

of *B. subtilis* [2]. In previous observations of *B. subtilis* using SEM, the low

resolution and chemical fixation processes interfered with the visualization of

the piecrusts. The filament pattern observed on the cylindrical part is a

circumferential mesh-like structure similar to the pattern observed for the

isolated peptidoglycan layer of *Escherichia coli* using AFM, although the

filament widths are different [9]. Perhaps, the filaments in the cylindrical part

may be aligned roughly in a circumferential way, generally in rod shaped

bacteria [10, 12, 13].

Based on the observation here, we can suggest a scheme for cell surface

structures in cell division cycles (Fig. 5A left).

Lysozyme starts from new pole and division site

In the present study, the disruption process of the peptidoglycan layer was

visualized for each of the three factors with different working points (Fig. 5A

right, B). In the process, lysozyme preferentially bound to a cell pole and a cell

division site (Fig. 2B). Probably, the newly synthesized peptidoglycan has many

gaps, which lysozyme molecules can access easily. The tracking of lysozyme

showed that it detached from the initially bound position earlier than it did the

other parts, suggesting that the peptidoglycan digestion was completed at the

cell pole (Fig. 2C). Even when pathogenic or parasitic bacteria invade the host,

the division site and the new pole is preferentially attacked by lysozyme. We

focused on *B. subtilis* inhabiting the environment in the present study, but the

process by which lysozyme attacks pathogenic or parasitic bacteria should also

be very interesting.

Quick-freeze, deep-etch EM showed that the action of lysozyme loosens

the peptidoglycan. As lysozyme cleaves carbohydrate chains that mainly form

the filaments of the peptidoglycan layer (Fig. 5B) [5, 6], the peptidoglycan layer

separates from the cell surface (Fig. 2E). This damage on the entire

peptidoglycan layer is as effective as the protection system that is involved in

the resistance against invasion by pathogenic or parasitic bacteria.

Turgor by penicillin treatment

In the early stages of cell disruption by PenG, large turgor appeared to be

applied to the cytoplasm at the cell division site from the cell inside, because the

cell membrane at a cleft of the cell envelope showed curvature to outside (Figs.

3C and 5A right). This observation is reasonable if we consider the working

mechanism of PenG, in which the transpeptidase activity is inhibited [7, 8].

In the process involving lysozyme, such turgor was not observed. This

difference can be explained by the disruption mechanisms (Fig. 5B)[5, 6, 40]. As

PenG inhibits only *de novo* crosslinking of peptidoglycan, the matured parts of

the peptidoglycan layer apply turgor to the cell. However, the damage by

lysozyme disrupts large parts of the cell envelope.

Change in surface pattern after *murE* operon repression

When the supply of muropeptide was stopped, the cell shape was disturbed

around the cell pole (Figs. 4 and 5A right). At that time, the features of the

pattern on the surface were also lost. Since the peptidoglycan layer of *B. subtilis*

is 20–40 nm thick, insertion of newly synthesized filaments into the

peptidoglycan layer through muropeptide supply is thought to occur on the side

close to the membrane [41, 42]. If it is true, the influence on the surface pattern

of the muropeptide depletion should occur in the final stage of morphological

change. In fact, a noticeable change in pattern was observed when the cells

were still maintaining the rod shape (Fig. 4C). This may suggest that there is

fluidity in the existing peptidoglycan filaments [42, 43].

CONCLUDING REMARKS

In this study, we showed that quick-freeze deep-etch EM is useful to visualize

the peptidoglycan layer and its disruption process of *B. subtilis* in high spatial

and time resolutions. Information on the structure of peptidoglycan is critical to

understanding the survival strategy and evolution of bacteria in general. The

application of quick-freeze deep-etch EM to peptidoglycan study should

contribute to our understanding of bacteria.

Funding

This work was supported by a Grant-in-Aid for Scientific Research on the

Innovative Area "Harmonized Supramolecular Motility Machinery and Its

Diversity" (MEXT KAKENHI Grant Number 24117002), by Grant-in-Aids for

Scientific Research (B) and (A) (MEXT KAKENHI Grant Numbers 24390107

and 17H01544) and by the Osaka City University (OCU) Strategic Research

Grant 2019 for top priority researches to MM.

Acknowledgements

We appreciate the helpful input from Eisaku Katayama at Osaka City University

and Daisuke Shiomi at Rikkyo University, gift of *B. subtilis* strains from

Yoshikazu Kawai at Newcastle University, and technical support from Junko

Shiomi at Osaka City University. The application of the quick-freeze, deep-etch

EM technique to microbiology was developed as the general supporting project

for the Grant-in-Aid for Scientific Research on Innovative Areas "Harmonized

Supramolecular Motility Machinery and Its Diversity" (25117501) directed by

MM.

Author contributions: All authors contributed to the research design, analyzing

data and writing the paper; I.T. and Y.O.T. performed the experiments. The

authors declare no conflict of interest.

References

1. Errington J. (2013) L-form bacteria, cell walls and the origins of life. *Open Biol* **3:**120143.

2. Turner RD, Vollmer W, Foster SJ. (2014) Different walls for rods and balls: the diversity of peptidoglycan. *Mol Microbiol* **91:**862-74.

3. Erickson HP. (2017) How bacterial cell division might cheat turgor pressure - a unified mechanism of septal division in Gram-positive and Gram-negative bacteria. *Bioessays* **39:**1700045.

4. Osawa M, Erickson HP. (2018) Turgor pressure and possible constriction mechanisms in bacterial division. *Front Microbiol* **9:**111.

5. Kawai Y, Mickiewicz K, Errington J. (2018) Lysozyme counteracts beta-lactam antibiotics by promoting the emergence of L-form bacteria. *Cell* **172:**1038-1049 e10.

6. Ragland SA, Criss AK. (2017) From bacterial killing to immune modulation: Recent insights into the functions of lysozyme. *PLoS Pathog* **13:**e1006512.

7. Stulberg ER, Lozano GL, Morin JB, Park H, Baraban EG, Mlot C, Heffelfinger C, Phillips GM, Rush JS, Phillips AJ, Broderick NA, Thomas MG, Stabb EV, Handelsman J. (2016) Genomic and secondary metabolite analyses of *Streptomyces* sp. 2AW provide insight into the evolution of the cycloheximide pathway. *Front Microbiol* **7:**573.

8. Kunzler M. (2018) How fungi defend themselves against microbial competitors and animal predators. *PLoS Pathog* **14:**e1007184.

9. Turner RD, Mesnage S, Hobbs JK, Foster SJ. (2018) Molecular imaging of glycan chains couples cell-wall polysaccharide architecture to bacterial cell morphology. *Nat Commun* **9:**1263.

10. Beeby M, Gumbart JC, Roux B, Jensen GJ. (2013) Architecture and assembly of the Gram-positive cell wall. *Mol Microbiol* **88:**664-72.

11. Andre G, Kulakauskas S, Chapot-Chartier MP, Navet B, Deghorain M, Bernard E, Hols P, Dufrene YF. (2010) Imaging the nanoscale organization of peptidoglycan in living *Lactococcus lactis cells*. *Nat Commun* **1:**27.

12. Hayhurst EJ, Kailas L, Hobbs JK, Foster SJ. (2008) Cell wall peptidoglycan architecture in *Bacillus subtilis*. *Proc Natl Acad Sci U S A* **105:**14603-8.

13. Gan L, Chen S, Jensen GJ. (2008) Molecular organization of Gram-negative peptidoglycan. *Proc Natl Acad Sci U S A* **105:**18953-7.

14. Heuser JE, Reese TS, Dennis MJ, Jan Y, Jan L, Evans L. (1979) Synaptic vesicle exocytosis captured by quick freezing and correlated with quantal transmitter release. *J Cell Biol* **81:**275-300.

15. Heuser JE. (2011) The origins and evolution of freeze-etch electron microscopy. *J Electron Microsc (Tokyo)* **60 Suppl 1:**S3-29.

16. Pilhofer M, Ladinsky MS, McDowall AW, Jensen GJ. (2010) Bacterial TEM: new insights

from cryo-microscopy. *Methods Cell Biol* **96:**21-45.

17. Zhao X, Norris SJ, Liu J. (2014) Molecular architecture of the bacterial flagellar motor in

cells. *Biochemistry* **53:**4323-33.

18. Swick MC, Koehler TM, Driks A. (2016) Surviving between hosts: sporulation and

transmission. *Microbiol Spectr* **4**.

19. Mercier R, Kawai Y, Errington J. (2013) Excess membrane synthesis drives a primitive

mode of cell proliferation. *Cell* **152:**997-1007.

20. Kasai T, Hamaguchi T, Miyata M. (2015) Gliding motility of *Mycoplasma mobile* on uniform

oligosaccharides. *J Bacteriol* **197:**2952-7.

21. Heuser JE. (1983) Procedure for freeze-drying molecules adsorbed to mica flakes. *J Mol*

Biol **169:**155-95.

22. Numata T, Murakami T, Kawashima F, Morone N, Heuser JE, Takano Y, Ohkubo K,

Fukuzumi S, Mori Y, Imahori H. (2012) Utilization of photoinduced charge-separated state

of donor-acceptor-linked molecules for regulation of cell membrane potential and ion

transport. *J Am Chem Soc* **134:**6092-5.

23. Miyata M, Petersen JD. (2004) Spike structure at the interface between gliding

Mycoplasma mobile cells and glass surfaces visualized by rapid-freeze-and-fracture

electron microscopy. *J Bacteriol* **186:**4382-6.

24. Kunoh T, Suzuki T, Shiraishi T, Kunoh H, Takada J. (2015) Treatment of leptothrix cells with

ultrapure water poses a threat to their viability. *Biology (Basel)* **4:**50-66.

25. Morone N, Fujiwara T, Murase K, Kasai RS, Ike H, Yuasa S, Usukura J, Kusumi A. (2006)

Three-dimensional reconstruction of the membrane skeleton at the plasma membrane

interface by electron tomography. *J Cell Biol* **174:**851-62.

26. Weibull C. (1953) The isolation of protoplasts from *Bacillus megaterium* by controlled

treatment with lysozyme. *J Bacteriol* **66:**688-95.

27. Muchova K, Wilkinson AJ, Barak I. (2011) Changes of lipid domains in *Bacillus subtilis* cells

with disrupted cell wall peptidoglycan. *FEMS Microbiol Lett* **325:**92-8.

28. Miller KR, Prescott CS, Jacobs TL, Lassignal NL. (1983) Artifacts associated with quick-

freezing and freeze-drying. *J Ultrastruct Res* **82:**123-33.

29. Dominguez-Cuevas P, Mercier R, Leaver M, Kawai Y, Errington J. (2012) The rod to L-form

transition of *Bacillus subtilis* is limited by a requirement for the protoplast to escape from

the cell wall sacculus. *Mol Microbiol* **83:**52-66.

30. Leaver M, Dominguez-Cuevas P, Coxhead JM, Daniel RA, Errington J. (2009) Life without

a wall or division machine in *Bacillus subtilis*. *Nature* **457:**849-53.

31. Hung C, Zhou Y, Pinkner JS, Dodson KW, Crowley JR, Heuser J, Chapman MR,

Hadjifrangiskou M, Henderson JP, Hultgren SJ. (2013) *Escherichia coli* biofilms have an

organized and complex extracellular matrix structure. *MBio* **4:**e00645-13.

32. Yoshida S, Handa Y, Suzuki T, Ogawa M, Suzuki M, Tamai A, Abe A, Katayama E,

Sasakawa C. (2006) Microtubule-severing activity of Shigella is pivotal for intercellular

spreading. *Science* **314:**985-9.

33. Verwer RW, Nanninga N. (1976) Electron microscopy of isolated cell walls of *Bacillus*

subtilis var. niger. *Arch Microbiol* **109:**195-7.

34. Burdett ID, Higgins ML. (1978) Study of pole assembly in *Bacillus subtilis* by computer

reconstruction of septal growth zones seen in central, longitudinal thin sections of cells. *J*

Bacteriol **133:**959-71.

35. Burdett ID. (1979) Electron microscope study of the rod-to-coccus shape change in a

temperature-sensitive rod- mutant of *Bacillus subtilis*. *J Bacteriol* **137:**1395-405.

36. Dufrene YF. (2014) Atomic force microscopy in microbiology: new structural and functional

insights into the microbial cell surface. *MBio* **5:**e01363-14.

37. Amako K, Umeda A, Murata K. (1982) Arrangement of peptidoglycan in the cell wall of

Staphylococcus spp. *J Bacteriol* **150:**844-50.

38. Cabeen MT, Jacobs-Wagner C. (2007) Skin and bones: the bacterial cytoskeleton, cell

wall, and cell morphogenesis. *J Cell Biol* **179:**381-7.

39. Typas A, Banzhaf M, Gross CA, Vollmer W. (2011) From the regulation of peptidoglycan

synthesis to bacterial growth and morphology. *Nat Rev Microbiol* **10:**123-36.

40. Monahan LG, Turnbull L, Osvath SR, Birch D, Charles IG, Whitchurch CB. (2014) Rapid

conversion of *Pseudomonas aeruginosa* to a spherical cell morphotype facilitates tolerance

to carbapenems and penicillins but increases susceptibility to antimicrobial peptides.

Antimicrob Agents Chemother **58:**1956-62.

41. Kim SJ, Chang J, Singh M. (2015) Peptidoglycan architecture of Gram-positive bacteria by

solid-state NMR. *Biochim Biophys Acta* **1848:**350-62.

42. Zhao H, Patel V, Helmann JD, Dorr T. (2017) Don't let sleeping dogmas lie: new views of

peptidoglycan synthesis and its regulation. *Mol Microbiol* **106:**847-860.

43. Shi H, Bratton BP, Gitai Z, Huang KC. (2018) How to build a bacterial cell: MreB as the

foreman of *E. coli* construction. *Cell* **172:**1294-1305.

LEGENDS

Fig. 1. Surface structures of *Bacillus subtilis* 168 CA cells visualized by quick-

freeze, deep-etch EM. **A)** Phase-contrast optical microscopy of cells. **B)** Field

image of cells. **C-H)** Magnified cell images. **C**) Side view. A flagellum is marked

by a blue triangle. A boxed area is magnified in (D). **D)** Magnified cell surface

image. Filaments on the cylindrical part are marked by red triangles. The

boundary between the cylindrical and pole parts is marked by a yellow triangle.

The area positioning at the yellow triangle is shown in an inset and colored for

the filamentous structures at cell pole (blue), piecrust (yellow), and cylindrical

part. **E**) Pole view. A boxed area is magnified in (F). **F)** Magnified pole surface

image. Concentric circles are marked by yellow triangles. **G)** Dividing cell. The

invagination is marked by a green triangle. **H)** Daughter cells, probably just after

cell division. Piecrust structures are marked by red triangles. **I)** A fractured cell

showing three different surfaces. The cytosol, cell membrane, and a cross

section of the peptidoglycan layer are marked by yellow, orange, and red

arrows, respectively. **J)** Magnified image of fractured part in (I). A cross section

of the peptidoglycan layer is marked by an arrow.

Fig. 2. Damages on cell structure by lysozyme. **A)** Phase-contrast microscopic

images after addition of lysozyme taken with 10 minutes intervals. **B)**

Localization of fluorescently labeled lysozyme on cell. Fluorescence of labeled

lysozyme was visualized at 10 minutes after the addition. Phase-contrast and

fluorescence images are shown as the left and the right panels, respectively, in

each of three image sets. A boxed cell in the middle-paired panels is magnified

in the right panel set. The signals were found preferentially at a pole and the

division site as marked by yellow triangles. **C)** Phase-contrast and fluorescence

images of a single cell treated by labeled lysozyme taken at 5-minute intervals.

D) Field image of cells treated with lysozyme for 20 and 60 minutes taken by

the quick-freeze, deep-etch EM. The mica surface is covered by eutectics

appearing as thin filaments. A typical eutectic is shown in the inset with an

arrow. **E)** Quick-freeze, deep-etch EM images of cells treated with lysozyme for

20 and 60 minutes, as shown in the bottom. The focused regions boxed in the

upper panels are magnified in the lower panels. Ruffling, coarse pattern and

smooth surfaces are marked by yellow, orange and red triangles, respectively.

Fig. 3. Damages on cell structure by PenG. **A)** Phase-contrast microscopic

images after addition of PenG taken at 30-minute intervals. **B, C)** Quick-freeze,

deep-etch EM images of cells treated with PenG for 120 minutes. **B**) Field

image. **C**) Magnified cell images. The boxed regions in the upper panels are

magnified more in the lower panels.

Fig. 4. Effects on cell structure by *murE* operon repression. **A)** Phase-contrast

microscopic images taken at 5-minute intervals commencing at 60 minutes after

repression. The cell marked by a yellow triangle changed its structure

drastically. A tapered polar, a bulge, and swollen cells can be observed

sequentially with time on the cell. **B, C)** Quick-freeze, deep-etch EM images of

cells repressed for *murE* for 85 minutes. **C**) Cell features resulted by *murE*

repression. The area boxed in the upper panels are magnified in the lower

panels.

Fig. 5. Schematic presentations of peptidoglycan architecture and working

points of inhibitory factors. **A)** Structural features on surfaces of cells in the

division cycle (left) and peptidoglycan disruption (right). **B)** Chemical processes.

The peptidoglycan layer consists of strands of repeating GlcNAc and *N*-

acetylmuramic acid (MurNAc) subunits. Transpeptidase (TP) forms a crosslink

between two sugar chains, forming peptidoglycan network. Glycosyltransferase

forms linkages between GlcNAc and MurNAc residues. Peptidoglycan synthesis

starts in the cytoplasm, where the nucleotide precursors are synthesized by the

Mur enzymes (MurA, MurB, MurC, MurD, MurE and MurF). MurNAc-

pentapeptide is coupled to a membrane carrier, undecaprenyl pyrophosphate,

by MraY. Then, lipid II is formed through coupling with GlcNAc by MurG, and

transferred to the outside of the cytoplasmic membrane.

The antibiotic β-lactams represented by PenG bind to and inhibit the activity of

the transpeptidase by forming a highly stable penicilloyl-enzyme intermediate.

Lysozyme cleaves the bond between MurNAc and the fourth carbon atom of

GlcNAc. The depletion of MurE blocks the supply of muropeptides.

Movie S1. Tomogram of isolated cell for 96 degrees. The cylindrical and cell

pole parts are colored differently in a still image at zero angle.

Fig. 1 (Double column)

Fig. 2 (Double column)

Fig. 3 (Double column)

Fig. 4 (Double column)

Fig. 5 (Double column)

Fig. S1.

Measurement of filament width. a) Cell surface image by quick-freeze, deep-etch EM. b, c) Magnified image of areas marked by colored box in panel (a). d, e) Image profiles of red arrow area in panels (b) and (c). The positions at the mid density was defined as the boundary of focusing filament.