ERASE: Extended Randomization for assessment of annotation enrichment in ASE datasets

Karishma D'Sa,[1,2] Regina H. Reynolds,[1] Sebastian Guelfi,[1] David Zhang,[1] Sonia Garcia Ruiz,[1] International Parkinson's Disease Genomics Consortium (IPDGC), System Genomics of Parkinson's Disease (SGPD), John Hardy,[1,3] Sarah A. Gagliano Taliun,[4] Kerrin S. Small,[5] Mina Ryten,[1†]* Juan A. Botía,[1,6†]

[1] Department of Neurodegenerative Disease, University College London (UCL) Institute of Neurology, London, WC1N 3BG, UK

[2] Department of Medical & Molecular Genetics, King's College London, London, SE1 1UL, UK

[3] UK Dementia Research Institute at University College London, London, WC1E 6BT, UK

[4] Center for Statistical Genetics, University of Michigan, Ann Arbor, Michigan, MI 48109, USA

[5] Department of Twin Research and Genetic Epidemiology, King's College London, London, SE1 7EH, UK

[6] Departamento de Ingeniería de la Información y las Comunicaciones. Universidad de Murcia, Murcia, 30100, Spain

[†]These authors contributed equally.

*Corresponding author

Correspondence- mina.ryten@ucl.ac.uk

Full consortia membership is available in the supplemental data file (Supplemental Text 1)

Abstract

Genome-wide association studies (GWAS) have identified thousands of genetic variants associated with various human phenotypes and many of these loci are thought to act at a molecular level by regulating gene expression. Detection of allele specific expression (ASE), namely preferential usage of an allele at a transcribed locus, is an increasingly important means of studying the genetic regulation of gene expression. However, there are currently a paucity of tools available to link ASE sites with GWAS risk loci. Existing integration methods first use ASE sites to infer cis-acting expression quantitative trait loci (eQTL) and then apply eQTL-based approaches. ERASE is a method that assesses the enrichment of risk loci amongst ASE sites directly. Furthermore, ERASE enables additional biological insights to be made through the addition of other SNP level annotations. ERASE is based on a randomization approach and controls for read depth, a significant confounder in ASE analyses. In this paper, we demonstrate that ERASE can efficiently detect the enrichment of eQTLs and risk loci within ASE data and that it remains sensitive even when used with underpowered GWAS datasets. Finally, using ERASE in combination with GWAS data for Parkinson's disease and data on the splicing potential of individual SNPs, we provide evidence to suggest that risk loci for Parkinson's disease are enriched amongst ASEs likely to affect splicing. Thus, we show that ERASE is an important new tool for the integration of ASE and GWAS data, capable of providing novel insights into the pathophysiology of complex diseases.

Introduction

Over the last decade our understanding of complex genetic disorders has been transformed by the use of genome-wide association studies (GWAS) with 71,673 SNP-trait associations reported in the latest release of the NHGRI-EBI GWAS catalogue[1]. Complex neurological and neuropsychiatric disorders are no exception. The most recent GWAS for Parkinson's disease (PD) reported 78 risk loci[2], while that for schizophrenia[3] reported 145. While there is a lag in terms of understanding the underlying biological processes involved in contributing to disease risk, it is thought that risk loci largely act by regulating gene expression as opposed to changing protein function[4]. This observation has led to an increasing interest in the genetic regulation of gene expression and the identification of expression quantitative trait loci (eQTLs, genetic variants that are associated with variation in gene expression) across the genome. eQTL studies have helped in the interpretation of GWAS findings across a range of disorders[5; 6]. However, one limitation of eQTL analyses is the requirement for large sample sizes. Since this can be difficult to achieve for some human tissues, such as specific brain regions, alternative approaches more robust to small sample sets are required.

Allele specific expression (ASE) analysis is another means of studying the genetics of gene expression, which is less dependent on sample size. This form of analysis quantifies expression variation between the two haplotypes of a diploid individual distinguised by heterozygous sites. While most heterozygous sites will have equal expression, this is not always true and differential expression at a transcribed position (termed an ASE site) can occur due to imprinting[7-10], protein truncating variants triggering nonsense mediated decay[11; 12] and cis-regulatory effects[13; 14]. ASE analysis being a within individual comparison, can be effective even with limited sample sizes as

it is not confounded by factors such as age, sex or co-morbidity. Furthermore, since most ASE

sites (particularly common SNPs) are driven by cis-regulatory effects[15-17], ASE analyses could be

integrated with GWAS to provide novel disease insights.

Existing tools for ASE-GWAS integration have been based on inferring cis-eQTLs from ASE data

and then examining their association to disease[18; 19]. This approach requires an aggregation of

information across the gene and can restrict the potential for inferring within-gene events relevant

to the phenotype, namely splicing. Given that there is increasing evidence that common disease

risk can be mediated through splicing[20-23], this is a significant limitation. While there are other

methods, which could be adapted for integration of ASE datasets with GWAS including

SNPSnap[24; 25] and stratified LD score regression[26], they have been designed primarily with

alternative forms of variant annotation in mind. For example, SNPSnap aims to provide control

SNPs subsets for any set of SNPs. This implies accounting for a number of factors that become

irrelevant when the source SNPs are ASEs (namely minor allele frequency and distance to the

nearest gene, as ASEs are predominantly within exons). In stratified LD score regression[26], ASE

SNPs would be used as a form of annotation in heritability enrichment analyses. However, often

ASE calling datasets are small, and thus would be underpowered in such an analysis. The intuitive

approach would be to use ASE data directly with the GWAS using an ASE-specific method.

However, there are intrinsic biases in ASE detection, which make such an approach challenging.

ASEs are discovered only in transcribed regions, which means they are primarily located within

exons, and are inevitably easier to detect confidently in a highly expressed genomic region. Given

that GWAS loci are also enriched within regions as defined by open chromatin[27-29] or tissue-

specific expression features[3; 26; 30], overlap between ASEs and GWAS hits have a high likelihood

of occurring by chance. This makes it hard to know whether ASEs are truly providing useful

disease insights.

In this paper we propose Extended Randomization for assessment of annotation enrichment in

ASE datasets (ERASE), a tool that integrates ASE results with any SNP-based annotation dataset

associated with a continuous score, such as eQTL or GWAS summary statistics. We test ERASE

using multiple large publicly available ASE datasets. These datasets have been generated

independently with different methods allowing us to test ERASE in the context of variable ASE

calling pipelines. When we integrate the ASEs with an annotation dataset such as GWAS summary

statistics, we take the shared set of SNPs, namely the SNPs both analysed for their ASE potential

and their effect on the phenotype of interest. We then ask if ASE sites are enriched for loci linked

to disease more than we would expect by chance. We use a randomization-based approach to assess

enrichment. This approach has proven useful in the context of eQTLs[24]. ERASE has an added

advantage in that it does not require individual-level genotype data to test whether detected ASEs

are enriched for loci with higher scores in the annotation datasets and to determine the contribution

of each annotation dataset to the enrichment. When ERASE compares the observed ASE calling

set with a randomly chosen set of non-ASE SNPs it matches for the read depth. This is an important

step because it ensures that any enrichment seen is due to the presence of an ASE and does not

simply reflect high expression of a given gene. Moreover, we further extend this method to

incorporate multiple annotation datasets in our enrichment analyses in order to better understand

the specific properties of the ASEs. We do so by studying the relative contribution of each

annotation dataset to the enrichment.

Material and Methods

ERASE

In this section, we describe the specifics of the method. We also provide a publicly available R package, available from GitHub (see Web Resources), implementing it.

ERASE - single annotation enrichment (SAE)

We wish to test whether significant ASEs in a study are also enriched for notable results in a SNP-based annotation dataset and apply a version of parametric bootstrapping to address this problem (Figure 1). Let the values in the annotation dataset be transformed such that large positive values represent more significant results (for example, a minus-$\log_{10}$ transformation would achieve this for p-values). Let A be the set of SNPs tested for ASE, let G be the set of SNPs in the SNP annotation dataset and let C be the common SNPs present in both A and G. Let $Ase(+)$ be the set of SNPs in C with a significant ASE signal, and let $Ase(-)$ be the set of SNPs in C with a non-significant ASE signal. We use $Ase(-)$ to model the null distribution of SNPs across the genome. To control for the confounding effect of read depth, SNPs are randomly selected with replacement from $Ase(-)$ to create a SNP set with the same size and read distribution as that of $Ase(+)$. This is done by placing the SNPs in $Ase(-)$ into bins based on their average read depth across samples (using read depth information from the study that generated the ASE data). By default, a bin corresponds to an interval of average reads with width 2, such that SNPs with average read depth of 3 and 4 will be in one bin while those with a read depth of 5 and 6 will be in the next bin and so on. All SNPs with an average read depth greater than 200 are placed in a single final bin. For each SNP in the $Ase(+)$ set, a 'null' counterpart is selected at random from the corresponding bin that matches its read depth, until a full set of 'null' SNPs is obtained. By default, the random selection

of 'null' SNP sets is repeated 10^4 times, and the mean annotation SNP score (for example, mean(-

$\log_{10}$ (p-value)) if a GWAS summary statistic dataset is used) is calculated for each set.

Let r_i be the mean annotation SNP score for the i-th iteration. The mean ($\bar{x}$) and standard deviation

$\sigma(x)$ of the random r_i values are used to calculate the z-score for the observed mean value from

our set of significant ASE SNPs, $z^{Ase} = \frac{y - \bar{x}}{\sigma(x)}$, where y is the mean SNP score for the $Ase(+)$ SNP

set. The final p-value for the enrichment of SNPs for the annotation amongst ASEs versus non-

ASEs is obtained from a one right-tailed z-test on z^{Ase}.

ERASE - multiple annotation enrichment (MAE)

The multiple annotation version of ERASE (Figure S1) tests simultaneously for enrichment of

ASEs in more than one SNP annotation dataset. Simultaneous enrichment in two or more SNP

annotation datasets means that we effectively test whether the same ASE SNPs tagging SNPs in

one source also tag SNPs in the other, by using only SNPs common to the ASE calling and SNP

annotation datasets. Starting from this basic concept, we also examine the relative contribution of

each annotation dataset to the enrichment. To illustrate the approach in the case of two annotation

datasets, we will consider one set of SNPs, G, with GWAS summary association statistics, and

another set of SNPs, S, with transcript specificity values (as obtained, for example, from the

SPIDEX[31] database). As with ERASE-SAE, we assume the annotation values are expressed in a

way that large positive values represent more notable results.

To perform the ERASE-MAE test, we will consider the set of SNPs, C, which are common to A,

G and S. The randomization procedure proceeds in a similar way as for ERASE-SAE, but is now

run twice, once for each of the two annotation types. Let $Ase(+)$ be the set of SNPs in C with a

significant ASE signal, and let *Ase(-)* be the set of SNPs in *C* with a non-significant ASE signal.

For each annotation type, subsets of SNPs are randomly selected from the *Ase(-)* set using the same

matching procedure as in ERASE-SAE, and the mean value of the relevant annotation scores are

stored. By default, the randomization step is repeated 10^4 times. In order to provide a consistent

basis for comparison between the two annotation types, the stored values are then standardized by

subtracting their respective means and dividing by their respective standard deviations, to create

two sets of transformed values, T_G^{ran} and T_S^{ran}. Similarly, the observed mean annotation values in

the *Ase(+)* set are standardized in the same way to create two further transformed values, T_G^{Ase}

and T_S^{Ase}.

The scores for the two annotations sources are then integrated as follows. Consider, $\alpha \in$ *[0,1]* to

be a calibration parameter representing the relative weight assigned to each SNP annotation

dataset, i.e. α being the weight for *G* and $(1 - \alpha)$ the weight for *S*. We define the observed value

of significant ASE sites in the integration of *G* and *S* as

167
$$I^{Ase} = \alpha\, T_G^{Ase} + (1 - \alpha)\, T_S^{Ase}$$

168   and the integrated value for the *i*th random set is then

169
$$I^{ran(i)} = \alpha\, T_G^{ran(i)} + (1 - \alpha)\, T_S^{ran(i)}, 1 \le i \le 10^4.$$

The p-value for the enrichment of *G* and *S* SNPs amongst ASEs versus non-ASEs is obtained with

a one right-tailed z-test using the z-score for *Ase,* calculated using the mean and standard deviation

of the $I^{ran(i)}, 1 \le i \le 10^4$ values and the I^{Ase} for a given value of α. We assume that the

population values of $T_G^{ran(i)}$ and $T_S^{ran(i)}$ are both normal (i.e. z-scores measure the deviation

from the mean, in standard deviation units), thus $I^{ran(i)}$ should also be normal. Note that the

value of I^{Ase} depends upon the relative contribution of each SNP annotation dataset.

Therefore, we would aim to find the value of α that optimizes I^{Ase} (i.e. the one that yields the

highest enrichment). The extension of the method for the N annotation datasets case follows

naturally when we define a series of weights w_i for N, such that $\sum_{i=1}^{N} w_i = 1$. Finding the value

of the weights when N=2 is reduced to vary one of the weights (i.e. alpha) in [0,1] such that

$$\alpha = argmin_{0 \leq \alpha \leq 1} P[I^{Ase}(\alpha) > \widehat{I^{ran}}].$$

When N ≥ 3, the brute force approach we suggest using (i.e. trying a representative set of value

combinations for all w_i's) is less applicable, due to computational complexity. Approaches to deal

with this problem in a computationally efficient manner exist. For example, using dynamic

programming we can decompose finding the N weights problem into two subproblems: finding

the values for N-1 weights corresponding to dimensions 1…N-1 and the value for w_N in the N-th

dimension. And again, we can decompose finding the N-1 weights into finding N-2 weights and

weight w_{N-1}. Once we reach directly solvable problems, we can reconstruct composite solutions

backtracking towards the initial problem. We are currently exploring this approach.

Stratified LD score regression (LDSC)

Stratified LDSC[26] partitions common SNP-heritability (estimated from GWAS summary

statistics) across customizable sets of SNPs; in this case we use ASE annotations. All annotations

were constructed in a binary format (i.e. using 1 if the SNP was present within the annotation and

0 otherwise), using only SNPs with a minor allele frequency > 5%. Annotations were added

individually to the baseline model of 53 annotations provided by Finucane et al. (version 1.1, see

Web Resources), comprising genome-wide annotations reflecting genetic architecture, such as

coding regions and histone marks. HapMap Project Phase 3 (HapMap3)[32] SNPs and 1000

Genomes Project[33] Phase 3 European population SNPs were used for the regression and LD

reference panels, respectively. The MHC region was excluded from all analyses due to the complex

and long-range LD patterns in this region. For all stratified LDSC analyses, in addition to the

enrichment, enrichment p-value and regression coefficient, we report a one-tailed p-value

(coefficient p-value) based on the coefficient z-score outputted by stratified LDSC. A one-tailed

test was used as we were only interested in annotation categories with a significantly positive

contribution to trait heritability, conditional upon the baseline model.

Results

Datasets

ERASE´s ability to handle relatively low sample numbers emerges from the fact that ASE analyses

work well with small sample sizes. Thus, the best suited dataset for ASE analyses is one based on

specific brain regions, which can be hard to obtain in large numbers. With this in mind, we tested

ERASE on ASE calling datasets generated from human brain and lymphoblastoid cell lines. The

former was generated from 49 putamen and 35 substantia nigra tissue samples, of the UK Brain

Expression Consortium dataset[34]. These samples were obtained from neuropathologically normal

individuals of European descent. All sites having > 5 reads in a sample were tested for ASE. In

some analyses, data from putamen and substantia nigra were considered together, to distinguish

between lack of power and no real enrichment, and the minimum FDR was taken across both

tissues. The second ASE calling dataset, comprised of lymphoblastoid cell lines (LCLs)[15] from

462 individuals of the 1000 Genomes Project where sites with at least 30 reads and both alleles

seen were tested for ASE. From both the ASE calling datasets, SNPs with an FDR < 5% were

considered ASE sites.

We used 1000 Genomes SNP data[35] (phase 3 genotype data – see Web Resources) to simulate a

set of genome-wide SNPs representing null ASEs. The 1000 Genomes data was selected as it has

formed the basis of analyses in many studies[26; 36; 37]. SNPs were selected from 503 individuals of

European ancestry from the following 5 populations: Utah Residents (CEPH) with Northern and

Western European Ancestry (CEU), Toscani in Italia (TSI), Finnish in Finland (FIN), British in

England and Scotland (GBR), Iberian Population in Spain (IBS). The 3 types of SNP annotation

datasets we used to test ERASE are eQTLs, GWAS full summary statistic datasets and the

alternative splicing prediction SPIDEX database. We used the gene-level eQTL data generated

from 111 putamen and 69 substantia nigra tissue samples of the UK Brain Expression Consortium

dataset[34] to test enrichment of eQTLs amongst ASEs. Depending on the ASE dataset analyzed,

different GWAS summary datasets were used when testing for enrichment of disease risk loci. For

UKBEC-putamen dataset, Schizophrenia[3; 38; 39] , PD (excluding 23andMe data)[40] (access obtained,

with permissions from IPDGC and SGPD) , waist-to-hip ratio adjusted for body mass index

(BMI)[41] were used, with the former two used as positive controls and the latter as a negative

control. In the case of the LCL dataset, rheumatoid arthritis[42] and epilepsy[43] were used as positive

and negative controls respectively. GWAS summary data files for individuals of European

ancestry were downloaded (see Web Resources) for consistency with the origin of individuals in

the ASE calling dataset. Waist-to-hip ratio adjusted for BMI summary statistics were converted

from hg18 to hg19 using the liftOver tool[44]. The other summary statistics already used hg19 co-

ordinates. SPIDEX[31] database (see Web Resources) serves as data source for transcript specificity

while examining the specific properties of ASEs. It is a database of pre-computed list splicing

variants with the percent inclusion ratio of the exon where the variant is present ($\Delta\Psi$). It reports

the maximum $\Delta\Psi$ value across 16 tissues.

ERASE detects known enrichment of eQTLs amongst ASEs

Besides imprinting and nonsense mediated decay, it is known that cis-regulation can also generate

ASE signals[13]. In fact, this process has been previously shown to account for most of the ASE

sites[15-17]. We tested whether ERASE detects the known enrichment of eQTLs amongst ASEs using

ASEs and gene-level eQTLs identified in human post-mortem putamen[34]. We applied ERASE-

SAE to the 154,214 SNPs, which had been analysed for both evidence of allelic imbalance and

eQTL activity. ASEs were placed into bins, based on the average read depth overlying the SNP of

interest. We did the same for SNPs negative for the ASE test (termed non-ASEs). They were placed

into bins for random selection such that the average read depth of the randomly selected SNPs

showed a similar distribution as that of the ASEs. As expected, we found that the ASEs were

enriched for gene-level eQTLs in putamen (1.96×10^{-150}, Figure S2).

We repeated this analysis varying bin width (from 2 to 10) to model the impact of less accurately

matching the read distribution of non-ASEs to ASEs and also calculated enrichment when no

attempt was made to control for differences in read depth. We observed that with increasing bin

sizes the p-value for eQTL enrichment amongst ASEs is stable across the bins but when read depth

was totally ignored as a confounding factor, a highly inflated p-value was observed (Figure S2,

File S1). Enrichment examined with a negative control, waist-to-hip ratio adjusted for BMI[41]

GWAS summary dataset, resulted in a significant p-value (2.23×10^{-2}) being observed only when

read depth was not accounted for. This suggests that controlling for read depth is effective and can

help prevent a false positive result. Moreover, in this analysis we note the following trade-off:

while the smallest bin width of 2 may provide us with the most accurate estimate of enrichment,

since the average read distribution of the randomly selected SNPs closely matches that of the ASEs

it is also true that the proportion of unique SNPs randomly selected on each iteration is lower than

when larger bin sizes are used (Figure S3, File S2). This is simply because the pool of SNPs which

are non-ASEs shrinks with the bin size. Thus, the choice of average read intervals needs to be

balanced against the ability to generate independent randomizations. The higher the number of

non-ASE SNPs with respect to ASEs falling within each bin, the more independent randomizations

we will obtain. Having said this, we note that enrichment p-values were reasonably stable across

a range of average read depth intervals in our test case (Figure S2).

ERASE yields a controlled Type I error rate

We conducted simulation studies to estimate the Type I (false positives) error rates incurred by

ERASE using the example of eQTL enrichment amongst ASEs detected in putamen[34]. This

analysis was performed by generating null ASE datasets by selecting subsets of SNPs from

amongst those identified within the 1000 Genomes dataset. To ensure consistency with the origins

of individuals from the UKBEC dataset, only SNPs from individuals of European ancestry in the

1000 Genomes dataset were considered (see Datasets section for details). The generation of the

null ASE datasets was performed accounting for the main biases affecting ASE discovery. SNPs

were further selected on the basis of biallelic status, exonic location and presence within a gene

known to be expressed in human putamen[34]. This led to the identification of 595,348 SNPs that

could be considered as false ASEs, and these were used to construct null ASE calling datasets.

Each time we tested enrichment of gene-level eQTLs against the null, we randomly selected 8,527

SNPs (matching the number of true ASEs that overlapped with the putamen gene-level eQTL

summary SNPs) and ran ERASE with the gene-level eQTL dataset. In order to get enough null

experiments, we repeated the process 1000 times and the Type I error rate was estimated using the

proportion of ERASE tests yielding enrichment (ERASE p-values < 0.05). We found an acceptable

ratio of Type I errors of 4.3%. (Figure 2, File S3).

ERASE detects known enrichment of GWAS risk loci for rheumatoid arthritis amongst ASEs

identified in lymphoblastoid cell lines

Given that ERASE was developed to enable the integration of publicly available ASE data with

disease-relevant annotation datasets, such as GWAS data we wanted to test our method in this

context. With this in mind, we investigated the enrichment of risk loci for rheumatoid arthritis and

epilepsy, selected to serve as a positive and negative controls, amongst ASEs identified in

lymphoblastoid cell lines (LCLs)[15], a commonly used model of B- lymphocytes[45] with regulatory

sites already implicated in a range of autoimmune diseases. ERASE-SAE was run separately on

the 51,403 and 35,199 sites that were examined for ASE and present in the rheumatoid arthritis[42]

and epilepsy[43] GWAS summary datasets respectively. A bin size of 10 was used to enable a better

sampling of the non-ASEs such that they matched the average read distribution of the ASEs in the

LCL dataset. In line with previous studies, we found that the ASEs in LCLs were enriched for rick

loci linked to rheumatoid arthritis (p-value $= 1.40 \times 10^{-79}$) and not for epilepsy (p-value $= 0.19$).

Thus, we demonstrate the utility of ERASE by successfully running it on two independent, public

ASE datasets.

ERASE detects significant enrichment of risk loci amongst ASEs in underpowered GWAS's

Studies have shown that GWAS's require large sample sizes to achieve adequate statistical power across a full range of risk allele frequencies and effect sizes[46; 47]. As a consequence, for many diseases, including schizophrenia, GWAS analyses have been repeated using increasing numbers of cases and controls[3; 38; 39; 48] with the aim of finding more risk loci each time. We tested ERASE at varying levels of GWAS size in order to calibrate its behavior in potentially underpowered conditions. This analysis was performed by running ERASE on three schizophrenia GWAS summary datasets with varying sample sizes (including cases and controls) ranging from 33,100 (Ripke et al., 2013[38]) to 105,318 (Pardiñas et al.,2018[3]) (Table 1). We examined if ERASE could detect enrichment of risk loci for schizophrenia in putamen ASE signals when sample sizes were small, noting that medium spiny neurons of the putamen (amongst other cell types including pyramidal cells and interneurons) have recently been implicated in the pathogenesis of schizophrenia[49]. As ASEs generated tend to be in highly expressed genes and these tend to be neuronal in the case of schizophrenia, not accounting for read depth as a confounder can result in high expression being considered as the reason for any enrichment observed. Controlling for read depth ensures that the enrichment seen is due to the ASEs. We tested for enrichment of loci linked to waist-to-hip ratio adjusted for BMI[41] as a negative control. As a comparison to our method, we used stratified LD score regression[10] (LDSC) (see Material and Methods), a recently developed and popular tool. As expected, ERASE detected an enrichment of schizophrenia risk loci amongst ASEs identified in putamen using the largest available schizophrenia dataset[3] (40,675 cases, 64,643 controls) (Table 1) with no significant enrichment for waist-to-hip ratio adjusted for BMI (p-value = 0.13). Similarly, stratified LDSC demonstrated a significant enrichment of schizophrenia heritability amongst ASEs (coefficient p-value = 1.82×10^{-2}) with no significant

enrichment for waist-to-hip ratio (coefficient p-value = 0.07). However, while ERASE was able

to detect the enrichment of risk loci amongst ASEs even when schizophrenia GWAS datasets were

smaller and consequently less well-powered, the enrichment of heritability amongst ASEs was

non-significant when stratified LDSC was used with smaller schizophrenia GWAS datasets due to

the larger error margins. Thus, we demonstrate that ERASE performs favorably in terms of

detecting enrichment of risk loci in GWAS with small sample sizes.

340
ERASE detects existing enrichment when ASEs are under-called

Next, we tested the sensitivity of ERASE in detecting the enrichment of risk loci amongst small

ASE calling datasets (i.e. varying levels of ASE test stringency). We approached this by down

sampling true ASEs. This analysis was performed by calculating the enrichment of risk loci for

schizophrenia as determined by Pardinas and colleagues[3] amongst decreasing percentages of ASEs

identified in human putamen and substantia nigra[34]. In order to ensure this process was robust, for

each subset of ASEs generated (including 5% and 10% increments- 5%, 10%, ..., 30%, 40%...,

90%) the random subsampling was repeated twenty times for each tested set, varying the random

seed. We applied both ERASE-SAE and stratified LDSC to these ASE datasets. As expected, both

ERASE and stratified LDSC generated a positive correlation between ASE dataset size and the

significance of enrichment observed. However, using ERASE a significant enrichment of

schizophrenia risk loci amongst ASEs was observed (median p-value = 4.01×10^{-11}; Bonferroni cut-

off = 2.07×10^{-4}), even when the ASE dataset was down sampled to 30%, equating to 4,543 SNPs

(Figure 3(a), File S4). On the other hand, stratified LDSC did not identify a significant enrichment

of heritability for schizophrenia even when 80% of the original ASE annotation was included

(median coefficient p-value at 80% = 5.27×10^{-4}; Bonferroni cut-off = 4.95×10^{-4}) (Figure 3(b),

File S5). It only manages to reach a significant enrichment with all the ASEs (coefficient p-value

= 4.19×10^{-4}). We noted that in the case of stratified LDSC, enrichment in heritability within the

ASE annotation was always observed but as the ASE annotation increased in size the confidence

in these estimates increased, enabling identification of a significant coefficient p-value (Figure

S4). Thus, we see that ERASE identifies a significant enrichment of risk loci within smaller subsets

of ASEs compared to stratified LDSC.

ERASE demonstrates ASEs predicted to have stronger effects on splicing, drive enrichment of risk

loci

ERASE works well when testing the enrichment of ASE in single annotation datasets, as evidenced

by its ability to detect enrichments of schizophrenia and PD GWAS loci in ASEs called in putamen

and substantia nigra[34]. ERASE also allows the integration of more than one annotation datasets,

thus providing further insight into the nature of these ASE signals. To illustrate this, we used

ERASE-MAE to examine if ASEs derived from putamen and substantia nigra, which were

predicted by Guelfi et al. to have stronger splicing effects[34] also drive the enrichment of risk loci

linked to schizophrenia[3] and PD[40]. We used SPIDEX[31], a database of splicing variants, generated

using machine learning, with the percent inclusion ratio of the exon where the variant is present

($\Delta\Psi$), for transcript specificity.

ERASE-MAE was applied to the set of 30,892 SNPs that were common to the ASE calling dataset

and two annotation datasets, namely schizophrenia GWAS summary data and the SPIDEX variant

dataset. Similarly, we applied this approach to the 35,922 SNPs common to the ASE calling

dataset, PD GWAS summary data and the SPIDEX variant dataset. In each case, enrichment was

examined by varying the relative importance of the two annotation datasets (the GWAS and

SPIDEX) when calculating enrichment amongst ASEs. This was done by assigning the calibration

parameter, α, which indicates the relative weight of the GWAS, with values ranging from 0 to 1

with 0.1 increments, such that an α value of 1 indicates an enrichment p-value obtained by using

the GWAS annotation dataset alone, while an α value of 0 refers to an enrichment p-value based

on the SPIDEX annotation dataset alone. We observed a very significant enrichment of risk loci

for schizophrenia and PD amongst ASEs, as predicted by the previously observed enrichment of

schizophrenia and PD risk loci amongst ASEs derived from putamen and substantia nigra (Figure

4, File S6).

In addition, using ERASE-MAE and integrating the SPIDEX annotation, further increased the

enrichment of risk loci amongst ASEs for both diseases. For schizophrenia, an enrichment peak

was observed when the relative weight of GWAS to SPIDEX was 0.7:0.3 (p-value = 1.80×10^{-27})

(Figure 4(a)), while for PD an enrichment peak was observed when the relative weight of GWAS

to SPIDEX was 0.5:0.5 (p-value = 1.11×10^{-9}) (Figure 4(b)). Thus, factoring in the splicing

potential of ASEs improved the enrichment of risk loci for schizophrenia and PD. This suggests

that the ASEs in these tissues are likely to be disease-associated splice sites.

Discussion

ASE is a potentially powerful tool due to its ability to provide regulatory information even in the

context of small numbers of RNA sequencing samples. However, the effectiveness of ASE

analyses in understanding the biology of complex genetic diseases has been limited by the

challenges of robustly integrating ASE data with other forms of annotation. This is analogous to

existing methods used to calculate GWAS enrichment amongst eQTLs. In this study, we address

this issue by presenting a method, ERASE, to calculate enrichment of a ranked annotation dataset

amongst a set of ASEs. Using ASE data from two different public datasets and integrating this

with cis-eQTL and GWAS data, we show that ERASE can reliably enable this form of analysis

with a low Type I error rate (4.3%). We also demonstrate that read depth is a confounder when

examining enrichment in ASEs and not accounting for it results in a highly inflated p-value. It

should be noted that a key component of ERASE is its ability to control for read depth, an

important bias in ASE detection. This is achieved by randomly sampling non-ASE sites within

bins, such that they have a read depth distribution similar to that of ASEs. While the default bin

size (set at 2) provides a conservative enrichment p-value, the proportion of unique SNPs randomly

selected may be low when the numbers of non-ASE sites at the matching read depth are limited.

This means that the bin size selection needs to be balanced against the proportion of unique SNPs.

However, we also show that the enrichment p-value is not inconsistent or highly inflated at varying

bin sizes, the key point being read depth is accounted for.

ERASE is easy to use requiring only the SNP set examined for ASE, with their corresponding read

depths and whether an ASE was observed or not, and a ranked SNP annotation dataset. By directly

using ASEs rather than leveraging ASE data across the gene to infer cis-acting eQTLs, ERASE

makes use of all available ASE data and in particular retains ASEs, which are generated through

splicing QTLs. This is important given the growing recognition that risk loci for complex diseases

can operate by affecting transcript usage[21; 31; 50; 51] and may explain the sensitivity of this method

in the context of smaller GWAS or ASE analyses. ERASE detected the enrichment of

schizophrenia risk loci amongst ASEs identified in human putamen across all schizophrenia

GWAS analyses tested including where the smallest GWAS was used. While this enrichment was

validated using an independent approach, namely stratified LDSC, this was only possible using

the larger schizophrenia GWAS. Similarly, we found that even when the ASE sites identified in

basal ganglia were down-sampled to only 4,543 SNPs (30% of all ASE sites) ERASE was still

able to detect a significant enrichment of schizophrenia risk loci amongst ASEs. In the case of

stratified LDSC, while enrichment in heritability was still apparent amongst this subset of ASE

sites, the enrichment was not significant. Given the growing interest in using iPSC-derived cell

types to identify cell and state-specific regulatory sites[52-54], the sensitivity of ERASE is particularly

relevant as generating, differentiating and maintaining the large iPSC-derived sample sets required

for eQTL analysis remains time-consuming and expensive, while applying ASE analyses to small

iPSC-derived sample sets is more feasible.

Importantly, we show how ERASE, in particular ERASE-MAE, can be used to provide insights

into the pathogenesis of complex diseases. From a methodological standpoint, ERASE-MAE is

useful to test ASE enrichment in more than one source simultaneously, namely whether the same

subset of ASE SNPs enriches in all annotation sources. However, it can also help unravel unequal

contributions from different annotation sources to the global enrichment. We illustrate this utility

by showing that the enrichment of schizophrenia and PD risk loci is driven in part by the splicing-

related properties of ASE sites. Our results provide additional evidence for the role of splicing in

schizophrenia and PD, where rick loci have already been found to be enriched for splicing QTLs

identified in human frontal cortex[22; 55; 56].In summary, ERASE, a sensitive and robust tool, has

been designed to use ASE data directly, assessing if it is enriched for SNPs in single or multiple

annotation datasets. ERASE-MAE is the arm of the tool that evaluates the enrichment in more than

one SNP annotation dataset and the relative contribution of each annotation to the overall

enrichment. Notably, ERASE uses all ASEs rather than summarizing information across a gene,

thus retaining ASEs generated due to splicing QTLs. Given the growing popularity of ASE

analyses [57-60] and the increased interest in splicing, ERASE's ability to detect enrichment in small

ASE datasets and multiple annotations has the potential to add significantly to biological

knowledge.

Supplemental Data

Supplemental Data include four figures and 6 tables (as excel files).

Acknowledgments

Mina Ryten, Karishma D'Sa, David Zhang and Sonia Garcia Ruiz were supported by the UK Medical Research Council (MRC) through the award of Tenure-track Clinician Scientist Fellowship to Mina Ryten (MR/N008324/1). Regina Reynolds was supported through the award of a Leonard Wolfson Doctoral Training Fellowship in Neurodegeneration. Sebastian Guelfi was supported by Alzheimer's Research UK through the award of a PhD Fellowship (ARUK-PhD2014-16). This work was supported by the UK Dementia Research Institute. Full consortia acknowledgements are available in the supplemental data file (Supplemental Text 2)

Declaration of Interests

The authors declare no competing interests.

Web Resources

1000 Genomes genotype data, Phase 3, was downloaded on 18/4/18 from ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/release/20130502/; Baseline LDSC annotations, https://data.broadinstitute.org/alkesgroup/LDSCORE/; LDSC, https://github.com/bulik/ldsc ; SPIDEX, was downloaded on 27/2/17 from https://www.deepgenomics.com/spidex-noncommercial-download ; UK Brain Expression Consortium data http://braineacv2.inf.um.es/;

GWAS summary datasets downloaded were Schizophrenia [3; 38; 39] from

https://www.med.unc.edu/pgc/results-and-downloads , Waist-to-hip ratio adjusted for body mass

index (BMI)[41] from the GIANT consortium website

https://www.broadinstitute.org/collaboration/giant/index.php/GIANT_consortium_data_files ,

Rheumatoid arthritis[42] from http://plaza.umin.ac.jp/~yokada/datasource/software.htm, Epilepsy[43]

from http://www.epigad.org/gwas_ilae2014/ ; ERASE https://github.com/karishdsa/ERASE

References

1. Buniello, A., MacArthur, J.A.L., Cerezo, M., Harris, L.W., Hayhurst, J., Malangone, C., McMahon, A.,
Morales, J., Mountjoy, E., Sollis, E., et al. The NHGRI-EBI GWAS Catalog of published genome-wide
association studies, targeted arrays and summary statistics 2019.
2. Nalls, M.A., Blauwendraat, C., Vallerga, C.L., Heilbron, K., Bandres-Ciga, S., Chang, D., Tan, M., Kia, D.A.,
Noyce, A.J., Xue, A., et al. (2019). Expanding Parkinson's disease genetics: novel risk loci, genomic
context, causal insights and heritable risk. bioRxiv, 388165.
3. Pardiñas, A.F., Holmans, P., Pocklington, A.J., Escott-Price, V., Ripke, S., Carrera, N., Legge, S.E., Bishop,
S., Cameron, D., Hamshere, M.L., et al. (2018). Common schizophrenia alleles are enriched in
mutation-intolerant genes and in regions under strong background selection. Nature genetics 50,
381-389.
4. Price, A.L., Spencer, C.C., and Donnelly, P. (2015). Progress and promise in understanding the genetic
basis of common diseases. Proceedings Biological sciences 282, 20151684.
5. Nica, A.C., and Dermitzakis, E.T. (2013). Expression quantitative trait loci: present and future.
Philosophical transactions of the Royal Society of London Series B, Biological sciences 368,
20120362.
6. Gamazon, E.R., Segrè, A.V., van de Bunt, M., Wen, X., Xi, H.S., Hormozdiari, F., Ongen, H., Konkashbaev,
A., Derks, E.M., Aguet, F., et al. (2018). Using an atlas of gene regulation across 44 human tissues
to inform complex disease- and trait-associated variation. Nature genetics 50, 956-967.
7. Yan, H., Yuan, W., Velculescu, V.E., Vogelstein, B., and Kinzler, K.W. (2002). Allelic Variation in Human
Gene Expression. Science (New York, NY) 297, 1143.
8. Santoni, F.A., Stamoulis, G., Garieri, M., Falconnet, E., Ribaux, P., Borel, C., and Antonarakis, S.E. (2017).
Detection of Imprinted Genes by Single-Cell Allele-Specific Gene Expression. American journal of
human genetics 100, 444-453.
9. Babak, T., DeVeale, B., Tsang, E.K., Zhou, Y., Li, X., Smith, K.S., Kukurba, K.R., Zhang, R., Li, J.B., van der
Kooy, D., et al. (2015). Genetic conflict reflected in tissue-specific maps of genomic imprinting in
human and mouse. Nat Genet 47, 544-549.
10. Baran, Y., Subramaniam, M., Biton, A., Tukiainen, T., Tsang, E.K., Rivas, M.A., Pirinen, M., Gutierrez-
Arcelus, M., Smith, K.S., Kukurba, K.R., et al. (2015). The landscape of genomic imprinting across
diverse adult human tissues. Genome Research 25, 927-936.
11. Montgomery, S.B., Lappalainen, T., Gutierrez-Arcelus, M., and Dermitzakis, E.T. (2011). Rare and
Common Regulatory Variation in Population-Scale Sequenced Human Genomes. PLoS genetics 7,
e1002144.
12. Rivas, M.A., Pirinen, M., Conrad, D.F., Lek, M., Tsang, E.K., Karczewski, K.J., Maller, J.B., Kukurba, K.R.,
DeLuca, D.S., Fromer, M., et al. (2015). Human genomics. Effect of predicted protein-truncating
genetic variants on the human transcriptome. Science (New York, NY) 348, 666-669.
13. Pickrell, J.K., Marioni, J.C., Pai, A.A., and Degner, J.F. (2010). Understanding mechanisms underlying
human gene expression variation with RNA sequencing. Nature 464, 768-772.
14. (2015). Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene
regulation in humans. Science (New York, NY) 348, 648-660.
15. Lappalainen, T., Sammeth, M., Friedlander, M.R., t Hoen, P.A.C., Monlong, J., Rivas, M.A., Gonzalez-
Porta, M., Kurbatova, N., Griebel, T., Ferreira, P.G., et al. (2013). Transcriptome and genome
sequencing uncovers functional variation in humans. Nature 501, 506-511.
16. Castel, S.E., Levy-Moonshine, A., Mohammadi, P., Banks, E., and Lappalainen, T. (2015). Tools and best
practices for data processing in allelic expression analysis. Genome Biology 16, 1-12.

17. Consortium, G.T., Aguet, F., Brown, A.A., Castel, S.E., Davis, J.R., He, Y., Jo, B., Mohammadi, P., Park,
Y., Parsana, P., et al. (2017). Genetic effects on gene expression across human tissues. Nature 550,
204.
18. van de Geijn, B., McVicker, G., Gilad, Y., and Pritchard, J.K. (2015). WASP: allele-specific software for
robust molecular quantitative trait locus discovery. Nat Meth 12, 1061-1063.
19. Kumasaka, N., Knights, A.J., and Gaffney, D.J. (2016). Fine-mapping cellular QTLs with RASQUAL and
ATAC-seq. Nat Genet 48, 206-213.
20. Battle, A., Khan, Z., Wang, S.H., Mitrano, A., Ford, M.J., Pritchard, J.K., and Gilad, Y. Genomic variation.
Impact of regulatory variation from RNA to protein.
21. Gandal, M.J., Zhang, P., Hadjimichael, E., Walker, R.L., Chen, C., Liu, S., Won, H., van Bakel, H.,
Varghese, M., Wang, Y., et al. (2018). Transcriptome-wide isoform-level dysregulation in ASD,
schizophrenia, and bipolar disorder. Science (New York, NY) 362, eaat8127.
22. Takata, A., Matsumoto, N., and Kato, T. (2017). Genome-wide identification of splicing QTLs in the
human brain and their enrichment among schizophrenia-associated loci. Nat Commun 8, 14519.
23. Odhams, C.A., Cortini, A., Chen, L., Roberts, A.L., Vinuela, A., Buil, A., Small, K.S., Dermitzakis, E.T.,
Morris, D.L., Vyse, T.J., et al. (2017). Mapping eQTLs with RNA-seq reveals novel susceptibility
genes, non-coding RNAs and alternative-splicing events in systemic lupus erythematosus. Hum
Mol Genet 26, 1003-1017.
24. Nicolae, D.L., Gamazon, E., Zhang, W., Duan, S., Dolan, M.E., and Cox, N.J. (2010). Trait-Associated
SNPs Are More Likely to Be eQTLs: Annotation to Enhance Discovery from GWAS. PLoS genetics
6, e1000888.
25. Pers, T.H., Timshel, P., and Hirschhorn, J.N. (2015). SNPsnap: a Web-based tool for identification and
annotation of matched SNPs. Bioinformatics 31, 418-420.
26. Finucane, H.K., Bulik-Sullivan, B., Gusev, A., Trynka, G., Reshef, Y., Loh, P.-R., Anttila, V., Xu, H., Zang,
C., Farh, K., et al. (2015). Partitioning heritability by functional annotation using genome-wide
association summary statistics. Nature genetics 47, 1228-1235.
27. Ernst, J., Kheradpour, P., Mikkelsen, T.S., Shoresh, N., Ward, L.D., Epstein, C.B., Zhang, X., Wang, L.,
Issner, R., Coyne, M., et al. (2011). Mapping and analysis of chromatin state dynamics in nine
human cell types. Nature 473, 43-49.
28. The, E.P.C., Dunham, I., Kundaje, A., Aldred, S.F., Collins, P.J., Davis, C.A., Doyle, F., Epstein, C.B.,
Frietze, S., Harrow, J., et al. (2012). An integrated encyclopedia of DNA elements in the human
genome. Nature 489, 57.
29. Maurano, M.T., Humbert, R., Rynes, E., Thurman, R.E., Haugen, E., Wang, H., Reynolds, A.P.,
Sandstrom, R., Qu, H., Brody, J., et al. (2012). Systematic localization of common disease-
associated variation in regulatory DNA. Science (New York, NY) 337, 1190-1195.
30. Reynolds, R.H., Botia, J.A., Nalls, M.A., Hardy, J., Gagliano, S.A., and Ryten, M. (2018). Moving beyond
neurons: the role of cell type-specific gene regulation in Parkinson's disease heritability. bioRxiv,
442152.
31. Xiong, H.Y., Alipanahi, B., Lee, L.J., Bretschneider, H., Merico, D., Yuen, R.K.C., Hua, Y., Gueroussov, S.,
Najafabadi, H.S., Hughes, T.R., et al. (2015). The human splicing code reveals new insights into the
genetic determinants of disease. Science (New York, NY) 347, 1254806-1254806.
32. The International HapMap, C., Altshuler, D.M., Gibbs, R.A., Peltonen, L., Altshuler, D.M., Gibbs, R.A.,
Peltonen, L., Dermitzakis, E., Schaffner, S.F., Yu, F., et al. (2010). Integrating common and rare
genetic variation in diverse human populations. Nature 467, 52.
33. The Genomes Project, C., McVean, G.A., Altshuler, D.M., Durbin, R.M., Abecasis, G.R., Bentley, D.R.,
Chakravarti, A., Clark, A.G., Donnelly, P., Eichler, E.E., et al. (2012). An integrated map of genetic
variation from 1,092 human genomes. Nature 491, 56.

34. Guelfi, S., D'Sa, K., Botia, J.A., Vandrovcova, J., Reynolds, R.H., Zhang, D., Trabzuni, D., Collado-Torres,
L., Thomason, A., Quijada Leyton, P., et al. (2019). Regulatory sites for known and novel splicing
in human basal ganglia are enriched for disease-relevant information. bioRxiv, 591156.
35. The Genomes Project, C., Auton, A., Abecasis, G.R., Altshuler, D.M., Durbin, R.M., Abecasis, G.R.,
Bentley, D.R., Chakravarti, A., Clark, A.G., Donnelly, P., et al. (2015). A global reference for human
genetic variation. Nature 526, 68.
36. Pickrell, Joseph K. (2014). Joint Analysis of Functional Genomic Data and Genome-wide Association
Studies of 18 Human Traits. American journal of human genetics 94, 559-573.
37. Kuruppumullage Don, P., Ananda, G., Chiaromonte, F., and Makova, K.D. (2013). Segmenting the
human genome based on states of neutral genetic divergence. Proc Natl Acad Sci U S A 110,
14699-14704.
38. Ripke, S., O'Dushlaine, C., Chambert, K., Moran, J.L., Kähler, A.K., Akterin, S., Bergen, S., Collins, A.L.,
Crowley, J.J., Fromer, M., et al. (2013). Genome-wide Association Analysis Identifies 14 New Risk
Loci for Schizophrenia. Nature genetics 45, 1150-1159.
39. Schizophrenia Working Group of the Psychiatric Genomics, C., Ripke, S., Neale, B.M., Corvin, A.,
Walters, J.T.R., Farh, K.-H., Holmans, P.A., Lee, P., Bulik-Sullivan, B., Collier, D.A., et al. (2014).
Biological Insights From 108 Schizophrenia-Associated Genetic Loci. Nature 511, 421-427.
40. Nalls, M.A., Blauwendraat, C., Vallerga, C.L., Heilbron, K., Bandres-Ciga, S., Chang, D., Tan, M., Kia, D.A.,
Noyce, A.J., Xue, A., et al. (2018). Parkinson's disease genetics: identifying novel risk loci,
providing causal insights and improving estimates of heritable risk. bioRxiv.
41. Shungin, D., Winkler, T.W., Croteau-Chonka, D.C., Ferreira, T., Locke, A.E., Magi, R., Strawbridge, R.J.,
Pers, T.H., Fischer, K., Justice, A.E., et al. (2015). New genetic loci link adipose and insulin biology
to body fat distribution. Nature 518, 187-196.
42. Okada, Y., Wu, D., Trynka, G., Raj, T., Terao, C., Ikari, K., Kochi, Y., Ohmura, K., Suzuki, A., Yoshida, S.,
et al. (2014). Genetics of rheumatoid arthritis contributes to biology and drug discovery. Nature
506, 376-381.
43. (2014). Genetic determinants of common epilepsies: a meta-analysis of genome-wide association
studies. The Lancet Neurology 13, 893-903.
44. Hinrichs, A.S., Karolchik D Fau - Baertsch, R., Baertsch R Fau - Barber, G.P., Barber Gp Fau - Bejerano,
G., Bejerano G Fau - Clawson, H., Clawson H Fau - Diekhans, M., Diekhans M Fau - Furey, T.S.,
Furey Ts Fau - Harte, R.A., Harte Ra Fau - Hsu, F., Hsu F Fau - Hillman-Jackson, J., et al. The UCSC
Genome Browser Database: update 2006.
45. Neitzel, H. (1986). A routine method for the establishment of permanent growing lymphoblastoid cell
lines. Human Genetics 73, 320-326.
46. Park, J.-H., Wacholder, S., Gail, M.H., Peters, U., Jacobs, K.B., Chanock, S.J., and Chatterjee, N. (2010).
Estimation of effect size distribution from genome-wide association studies and implications for
future discoveries. Nature genetics 42, 570-575.
47. Klein, R.J. (2007). Power analysis for genome-wide association studies. BMC Genetics 8, 58-58.
48. Manolio, T.A. (2017). A decade of shared genomic associations. Nature 546, 360.
49. Skene, N.G., Bryois, J., Bakken, T.E., Breen, G., Crowley, J.J., Gaspar, H.A., Giusti-Rodriguez, P., Hodge,
R.D., Miller, J.A., Muñoz-Manchado, A.B., et al. (2018). Genetic identification of brain cell types
underlying schizophrenia. Nature Genetics 50, 825-833.
50. Tazi, J., Bakkour, N., and Stamm, S. (2009). Alternative splicing and disease. Biochimica et biophysica
acta 1792, 14-26.
51. Li, Y.I., van de Geijn, B., Raj, A., Knowles, D.A., Petti, A.A., Golan, D., Gilad, Y., and Pritchard, J.K. (2016).
RNA splicing is a primary link between genetic variation and disease. Science (New York, NY) 352,
600-604.

52. Alasoo, K., Rodrigues, J., Mukhopadhyay, S., Knights, A.J., Mann, A.L., Kundu, K., Hale, C., Dougan, G.,
Gaffney, D.J., and Consortium, H. (2018). Shared genetic effects on chromatin and gene
expression indicate a role for enhancer priming in immune response. Nature Genetics 50, 424-
431.
53. Schwartzentruber, J., Foskolou, S., Kilpinen, H., Rodrigues, J., Alasoo, K., Knights, A.J., Patel, M.,
Goncalves, A., Ferreira, R., Benn, C.L., et al. (2018). Molecular and functional variation in iPSC-
derived sensory neurons. Nature Genetics 50, 54-61.
54. Banovich, N.E., Li, Y.I., Raj, A., Ward, M.C., Greenside, P., Calderon, D., Tung, P.Y., Burnett, J.E., Myrthil,
M., Thomas, S.M., et al. (2018). Impact of regulatory variation across human iPSCs and
differentiated cells. Genome Res 28, 122-131.
55. Gusev, A., Mancuso, N., Won, H., Kousi, M., Finucane, H.K., Reshef, Y., Song, L., Safi, A., McCarroll, S.,
Neale, B.M., et al. (2018). Transcriptome-wide association study of schizophrenia and chromatin
activity yields mechanistic disease insights. Nat Genet 50, 538-548.
56. Li, Y.I., Wong, G., Humphrey, J., and Raj, T. (2019). Prioritizing Parkinson's disease genes using
population-scale transcriptomic data. Nature Communications 10, 994.
57. Rao, X., Thapa, K.S., Chen, A.B., Lin, H., Gao, H., Reiter, J.L., Hargreaves, K.A., Ipe, J., Lai, D., Xuei, X., et
al. (2019). Allele-Specific Expression and High-Throughput Reporter Assay Reveal Functional
Variants in Human Brains with Alcohol Use Disorders. bioRxiv, 514992.
58. Liu, Z., Dong, X., and Li, Y. (2018). A Genome-Wide Study of Allele-Specific Expression in Colorectal
Cancer.
59. Llavona, P., Pinelli, M., Mutarelli, M., Marwah, V.S., Schimpf-Linzenbold, S., Thaler, S., Yoeruek, E.,
Vetter, J., Kohl, S., and Wissinger, B. (2017). Allelic Expression Imbalance in the Human Retinal
Transcriptome and Potential Impact on Inherited Retinal Diseases. Genes 8, 283.
60. Rhee, J.-K., Lee, S., Park, W.-Y., Kim, Y.-H., and Kim, T.-M. (2017). Allelic imbalance of somatic
mutations in cancer genomes and transcriptomes. Scientific Reports 7, 1653.

Figures

Figure 1

Overview of ERASE-single annotation enrichment (SAE) method

The figure represents a graphical overview of the general approach used in ERASE. The ERASE-

SAE version of the tool just needs as input the set of SNPs examined for ASE and the SNP

annotation dataset. It starts with finding the SNPs common (C) to both datasets. To construct a

null distribution of random samples, the non-ASE SNPs from C are randomly selected 10^4 times

such that their average read distribution matches that of the ASE SNPs in C. This accounts for

biases in ASE discovery due to high read depth and exonic structure. The mean and the standard

distribution obtained from the randomization are used to calculate the z-score for the observed set

of ASEs. The p-value is generated by a one right-tailed z-test.

Figure 2

QQ-plot of simulations performed to assess Type I error

The estimate for the Type I error in ERASE we report is obtained by using the tool 1000 times with random sets of SNPs as ASEs. In each simulation, we generate null ASE datasets from the 1000 Genomes SNPs and test them for enrichment with ERASE. The tested enrichment is on gene level eQTLs from human post-mortem putamen samples. The figure shows a QQ-plot representing the null population of p-values we obtain. The grey area region marks the not significant p-values region. The pool of SNPs selected from the 1000 Genomes for the simulations were from individuals of European ancestry, of biallelic status, exonic in location and present within a gene known to be expressed in human putamen. The Type I error rate estimate (i.e. proportion of p-values greater than 0.05 out of the 1000 null p-values) obtained in the simulation was 4.3%.

Assessment of sensitivity in ERASE and stratified LDSC with ASE SNP datasets of various

sizes, when testing for observed enrichment of schizophrenia risk loci (Pardiñas et al.,2018).

ASEs derived from putamen and substantia nigra were randomly subsampled to form sample sets

of various sizes and 20 sample sets per size varying the random seed. These sample sets were

tested with ERASE and stratified LDSC.

Each box plot corresponds to the p-values or coefficient p-values, obtained with (a) ERASE or (b)

stratified LDSC respectively, tested on the 20 sample sets. The x-axis shows the different sampling

sizes of ASEs, i.e. the percentage of ASEs sampled from the total, followed by the number of

ASEs that it equates to, within brackets. The y-axis shows the $-\log_{10}$ scale for the p-values or

coefficient p-values. The red dashed lines indicate the corresponding Bonferroni cut-off applied to

the results obtained from each of the tools. (a) ERASE identifies significant enrichment of

schizophrenia risk loci amongst the ASEs (total of 15,144 signals) even when the number of ASEs

was lowered to 30% (4,543 signals) of the total ASEs. (Bonferroni cut-off = 2.07×10^{-4}). (b)

Stratified LDSC identifies significant enrichment of heritability for schizophrenia only when the

entire ASE annotation (15,144 signals) was included (Bonferroni cut-off = 4.95×10^{-4}). Notably, an

enrichment in heritability within the ASE annotation was always observed with stratified LDSC

and as the ASE annotation increased in size the confidence in these estimates increased, enabling

a significant coefficient p-value to be identified (Figure S4).

Figure 4

Results of ERASE-MAE, with varying the relative contributions of the two SNP annotation

datasets to the final enrichment.

ERASE-MAE was applied to ASEs derived from human putamen and substantia nigra using SNP

annotation datasets SPIDEX and (a) Schizophrenia (Pardiñas et al.2018), (b) Parkinson's disease

(Nalls et al.2018). The relative contribution of each SNP annotation dataset to the final enrichment

was varied by assigning α, a calibration parameter, a range of values from 0 to 1 incremented by

0.1, such that an α value of 1 indicates the enrichment contribution is from the GWAS dataset

alone and value of 0 from SPIDEX. The x-axis shows the relative weights assigned to GWAS and

SPIDEX. The y-axis shows the $-\log_{10}$ scale for the p-values. The red dashed line indicates the

Bonferroni cut-off of 4.55×10^{-3}. In plot (a) we see an enrichment peak for schizophrenia risk loci

and transcript specificity at 07:03. This reflects clearly that both contribute to the enrichment. It

also suggests that the contribution is unbalanced, i.e. the contribution of schizophrenia risk loci is

more relevant. In plot (b) the case is different. We see a balanced case, 0.5:0.5, in which both PD

risk loci and SPIDEX equally contribute.

Tables

Table 1

Enrichment of risk loci for schizophrenia in putamen ASEs observed with ERASE and

LDSC, using schizophrenia GWAS's of varying power. * indicates p-values and co-efficient

p-values < 0.05.

Schizophrenia GWAS	Number of cases	Number of controls	ERASE p-value	Stratified LDSC coefficient p-value
Ripke et al.[38]	14,395	18,705	*6.01×10^{-12}	1.23×10^{-1}
Schizophrenia Working Group of the Psychiatric Genomics Consortium[39]	33,640	43,456	*2.90×10^{-21}	6.55×10^{-2}
Pardiñas et al.[3]	40,675	64,643	*1.52×10^{-20}	*1.83×10^{-2}