Lethal mutagenesis of Rift Valley fever virus induced by favipiravir

Belén Borrego[a], Ana I. de Ávila[b] , Esteban Domingo[b,c] and Alejandro Brun[a]#

[a] Centro de Investigación en Sanidad Animal (CISA-INIA), Valdeolmos, Madrid, Spain

[b] Centro de Biología Molecular "Severo Ochoa" (CSIC-UAM), Madrid, Spain

[c] Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas

(CIBERehd) del Instituto de Salud Carlos III, 28029, Madrid, Spain,

Running Head: Mutagenic effect of favipiravir on RVFV

#Address correspondence to brun@inia.es (A. Brun)

15
16

ABSTRACT

Rift Valley fever virus (RVFV) is an emerging, mosquito-borne, zoonotic pathogen

with recurrent outbreaks paying a considerable toll of human deaths in many African

countries, for which no effective treatment is available. In cell culture studies and with

laboratory animal models, the nucleoside analogue favipiravir (T-705) has demonstrated

great potential for the treatment of several seasonal, chronic and emerging RNA virus

infections of humans, suggesting applicability to control some viral outbreaks.

Treatment with favipiravir was shown to reduce the infectivity of Rift Valley fever virus

both in cell cultures and in experimental animal models, but the mechanism of this

protective effect is not understood. In this work we show that favipiravir at

concentrations well below the toxicity threshold estimated for cells is able to extinguish

RVFV from infected cell cultures. Nucleotide sequence analysis has documented RVFV

mutagenesis associated with virus extinction, with a significant increase in G to A and C

to U transition frequencies, and a decrease of specific infectivity, hallmarks of lethal

mutagenesis.

INTRODUCTION

Rift Valley fever virus (RVFV) is a mosquito-borne bunyavirus belonging to the recently reclassified phlebovirus genus (Family *Phenuiviridae*, Order Bunyavirales). RVFV causes an important disease in ruminants often transmitted to humans after the occurrence of epizootic outbreaks. Although the disease has been reported only in African countries with some incursions in the Middle East and Indian Ocean islands, there are concerns for its potential spread to other locations, including Europe (1, 2). Currently, there is no available treatment or licensed RVF vaccine in Europe, therefore the development of effective control strategies is an essential field of research. The use of antiviral agents for treatment in livestock is generally not affordable, although it is often being proposed for viral veterinary diseases as a strategy to fill the gap between the time of the infection and the effective development of the host immune response (3, 4). In contrast, in the case of humans infected or at risk of infection with RVFV, antiviral treatment is fully warranted. Antiviral base and nucleoside analogues effective against RVFV are available (5-7). Among the nucleoside analogs exerting anti-RVFV activity, ribavirin [1-β-D-ribofuranosyl-1-H-1,2,4-triazole-3-carboxamide], favipiravir [6-fluoro-3-hydroxy-2-pyrazinecarboxamide] and more recently BCX4430 [(2S,3S,4R,5R)-2-(4-amino-5H-pyrrolo[3,2-d]pyrimidin-7-yl)-5-(hydroxymethyl)pyrrolidine-3,4-diol] (Galidesivir) have been described (6-8). Particularly, favipiravir displays potent antiviral activity against different RNA viruses (9-16). Upon cell intake, favipiravir is converted by cellular enzymes into its active form (favipiravir-4-ribofuranosyl-5-triphosphate), which functions as a purine nucleotide analogue. Favipiravir can act as an RNA chain terminator (17, 18), and as lethal mutagen for several RNA viruses, including hepatitis C virus (19), foot-and-mouth disease virus (20), West Nile fever virus (21), norovirus (22), and influenza virus

(23), among others. Here we show that the mechanism of action of favipiravir against

RVFV in cell culture is due, at least in part, to the accumulation of mutations in the viral

genome that leads to a progressive decrease in viable viral progeny. This conclusion is

based on a significant increase of mutation frequency measured in a 1060 bp fragment

from the Gc (glycoprotein C)-coding region, and a concomitant decrease of specific

infectivity, associated with virus extinction, supporting the use of favipiravir for the

control of emerging RVFV infections in humans.

RESULTS

Effect of favipiravir (T-705) on RVFV 56/74 progeny virus yield.

The aim of this work was to determine whether favipiravir induces lethal mutagenesis

upon RVFV infection in cell culture. We first determined the effect of different

concentrations of favipiravir on the yield of RVFV isolate 56/74 in Vero cells. The half

inhibitory concentration (IC_{50}, concentration required for a 50% decrease in the

production of infectious progeny), was calculated to be 6.22 ± 1.5 µM (average of 3

determinations) **(Figure 1).** Since the cytotoxic concentration 50 (CC_{50}) of T-705 for

Vero cells is >10000 µM (21), the therapeutic index for RVFV (TI = CC_{50} / IC_{50}) is >

1600.

The RVFV 56/74 isolate was subjected to serial passages in Vero cells in the absence or

presence of different concentrations of favipiravir (**Figure 2a**). The titration of

supernatants indicated loss of infectivity in the passages carried out in the presence of

80 µM favipiravir, as well as for higher doses, such as 160, 200 and 500 µM, but not at

the lowest concentrations tested. A concentration of 20 µM produced a slight reduction

in viral yield during the consecutive passages, whilst at 40 µM the progeny production

declined progressively, and was not detected after the fifth, sixth and seventh passage.

Interestingly, virus yields were recovered at passages eighth and ninth. These data

suggested the selection of a resistant mutant under certain suboptimal inhibitory

concentrations of the drug. To test this hypothesis, viral yields in the presence of

different concentrations of favipiravir were analyzed for this virus. As expected from

the previous results (**Figure 1**), favipiravir concentrations over 5 μM significantly

reduced the yield of either the parental or the serially passaged virus RVFV 56/74 not

treated with the drug, whilst for the virus selected after 8 passages in the presence of 40

μM favipiravir a 50% reduction in viral yield was achieved at a higher concentration,

rendering an IC_{50} of 19.2 μM (**Figure 2b**). These results clearly indicated a 3-fold

change in susceptibility to the drug when compared to the parental strain, although it

remained susceptible to higher concentrations of the drug. The characterization of this

resistant variant is currently under investigation.

To confirm virus extinction, cell culture supernatants from passages carried out in the

presence of favipiravir where no cytopathic effect (cpe) had been observed were

subjected to up to five additional passages in the absence of the drug (**Figure 2c**). No

infectivity was rescued from samples that were collected after at least two serial

passages in the presence of 80 μM favipiravir or higher. Viral infectivity could be

recovered from supernatants of cells previously treated with 40 μM favipiravir. These

data confirmed the previous observation, indicating that the deleterious effect on virus

propagation at this concentration was not complete.

Analysis of RVFV passaged in absence or presence of favipiravir

Decrease of specific infectivity, broadening of the mutant spectrum and invariance of

the consensus sequence of the viral population are diagnostic of lethal mutagenesis (19-

21, 23-28). To explore these parameters, RVFV populations at passages 1 to 4 in

absence or presence of 40 μM favipiravir were examined. Extracellular viral titers and

viral RNA decreased significantly in the presence of favipiravir (**Figure 3**). The

changes in infectivity and viral RNA production resulted in a gradual decrease of

specific infectivity in the populations passaged in the presence of favipiravir (up to 8-

fold in four passages) but not in its absence (p<0.0001 for passages 1 to 4; t test). Thus,

the first diagnosis criterium of lethal mutagenesis is fulfilled for favipiravir acting on

RVFV.

To examine whether the decrease in progeny production was associated with increased

mutagenesis of the viral RNA, total RNA from the cell culture supernatant of passage 4

in the absence or presence of 40 μM favipiravir was extracted and a 1.1kb DNA

fragment encoding a glycoprotein Gc region comprising nucleotides 2091 to 3223

(numbering according to GenBank sequence of RVFV strain SA-75, accession number

DQ380189) was amplified by RT-PCR and subjected to molecular cloning and

nucleotide sequencing. For each condition (with/without drug) a total of 29 clones were

analyzed, corresponding to 31,610 nucleotides sequenced. The results (**Table 1**)

indicated a significant 2.5-fold increase of minimum mutation frequency associated with

replication in the presence of favipiravir (p< 0.0001; χ2 test). Of the total 142 different

mutations identified in the populations analyzed, 89 were detected only in the

population passaged in the presence of favipiravir, 4 were detected only in the

population passaged in the absence of favipiravir, and 49 were common in the two

populations (indicated in **Table S1**). Once the common mutations are excluded, the

minimum mutation frequencies are 1.3×10^{-4} and 2.8×10^{-3} in absence and presence of

favipiravir, respectively, and the difference is statistically significant (P<0.0001; χ2

test). The distribution of the number of Gc sequences was analyzed as a function of the

number of mutations per clone (**Figure 4**). While the viral population that was passaged

in absence of drug yielded 7 clones without mutations and 7 clones with only one

mutation (24.1% in both cases), only 1 clone without mutations and 2 with one mutation

were retrieved from the population that was passaged in the presence of favipiravir

(1/29, 3.4%; 2/29, 6.9%, respectively). Finally, the consensus genomic nucleotide

sequence (analyzed at passage 4) did not change as a result of replication in the

presence of favipiravir, an observation already found during lethal mutagenesis of other

RNA viruses (19, 27-30).

Analysis of mutation types

In order to analyze whether passage of RVFV in the presence of favipiravir resulted in

any deviation of mutation types, we compared the mutation matrices in the mutant

spectrum of the populations passaged in absence or presence of the drug. The results

indicate a predominance of G to A and C to U transitions associated with replication in

the presence of favipiravir: without drug the number of changes G→A + C→U was 24

(10+14), while with favipiravir it was 98 (52+46), thus resulting in 3-fold increase of

the ratio (G→A) + (C→U) / (A→G) + (U→C) (**Figure 5**). The difference between the

type of mutations was statistically significant ($p < 0.0001$; $\chi 2$ test), and reveals that

favipiravir produces a bias in favor of GA and CU transitions; the transition to

transversion ratio was comparable: 12.2 and 18.7 upon passage in absence and presence

of favipiravir, respectively.

These results indicate that favipiravir is a mutagenic agent for RVFV and suggest

that its anti-RVFV activity may be exerted at least in part through lethal mutagenesis.

DISCUSSION

The broad spectrum antiviral activity of favipiravir has been established with

representatives of several families of RNA viruses, so it stands as a promising drug to

be used against viral pathogens. In the case of Rift Valley fever, for which no

therapeutic or prophylactic treatments are available for humans, favipiravir could be a

powerful therapeutic tool to provide immediate treatment to farmers, veterinary and

research personnel at risk of infection after being in contact with the virus or virus-

infected animals. Favipiravir was initially considered an inhibitor of viral RNA

synthesis, exerted mainly through RNA chain termination (17, 18). However, model

studies with several viruses in cell culture and *in vivo* suggest that the antiviral activity

of favipiravir may include lethal mutagenesis.

Previous studies have shown that favipiravir treatment of laboratory animals exposed to

RVFV increased their survival, and counteracted acute or neurological disease

syndromes (6, 7). Furthermore, treatment efficacy was enhanced when favipiravir was

combined with ribavirin treatment (7). Knowledge of the mechanism of action of this

promising drug is important to improve antiviral designs. Our data clearly show at least

a 2.5-fold increased RVFV mutation frequencies when the virus was propagated in the

presence of 40 µM favipiravir, a dose which is 10- to 25-fold lower than the dose

proven effective for other RNA viruses tested in a similar way (19-21). The limitation

of non-synonymous mutations introduced during the four passages in the absence of

favipiravir (9.4% and 5.6% of the total 53 and 375 for different and total mutations,

respectively) is diminished in the passages in the presence of favipiravir (38.4% and

18.4%, respectively), consistent with random introduction of mutations that decrease

viral fitness.

One of the most advantageous features of favipiravir is that it shows a high barrier for

the generation of resistance mutants (9). Since its activity against human influenza virus

was first described (11), only one resistant mutant has been reported (31). Attempts to

isolate resistant variants of polio- and influenza viruses seem to have been unsuccessful

(32-34). In addition, influenza virus pairs isolated from patients before and after

favipiravir administration showed no changes in the susceptibility to the drug (35).

Similar results were reported in norovirus-infected mice after favipiravir treatment (22).

Moreover, previous studies reporting favipiravir mediated viral extinction by lethal

mutagenesis did not describe selection of specific mutations or resistant variants (19-21,

23). In our system, 40 µM favipiravir seemed not to lead to the extinction of the virus,

since after a lag phase of three consecutive blind passages the presence of cpe was

detected again in cultures. The comparison of the IC_{50} value of the virus recovered in

supernatant after the lag phase with that of the original samples confirmed the selection

of variants with increased favipiravir resistance, suggesting the selection of mutant

viruses with increased resistance to the drug. Work is currently underway in order to

characterize this novel resistance variant both at genomic and phenotypic level *in vitro*

and *in vivo*.

The appearance of variants showing decreased susceptibility to the drug has been shown

earlier (Delang et al., 2014) and it highlights the importance of deciphering the exact

mechanisms exerted by new promising antiviral drugs, such as favipiravir. In this sense,

drug concentration could be proposed as a crucial determinant on the antiviral effect.

Thus, between high concentrations -leading to a total abolishment of infectivity- and

low concentrations -with none or low effects on viral replication- there might be a

critical intermediate concentration that, even though forcing the virus into lethal

mutagenesis, promotes the selection for drug resistant variants before the virus is

extinguished. Nonetheless, it is worth noting that other studies found that drug

concentrations close to the IC_{50} led to viral extinction of influenza virus without the

emergence of variants (23) perhaps indicating that other unknown factors play a role the

inhibitory mechanism of T-705. Still, the antiviral activity mechanism of favipiravir is

not fully understood. Most nucleoside analogues used as antiviral agents display several

mechanisms of activity (36-38). This is also the case for favipiravir with a dual chain

terminating and mutagenic activity by ambiguous base pairing, as well as a possible role

in modulation of innate immunity (10).

In summary, what our results suggest is that favipiravir can act as a lethal mutagen for

RVFV but they do not exclude that other mechanisms, alone or together with lethal

mutagenesis, are required for virus extinction.

MATERIALS AND METHODS

Cells, viruses and infections

Vero cells (ATCC Cat.No. CCL-81) were grown in Dulbecco's modified Eagle's

medium supplemented with 5% –10% fetal calf serum (FCS), and L-glutamine (2 mM),

penicillin (100 U/ml) and streptomycin (100 μg/ml), in a humid atmosphere of 5% CO_2

at 37°C.

The virus used in this study originates from a sheep experimentally infected in our

laboratory (39) with RVFV isolate 56/74 (40, 41). The virus was re-isolated from

infected sheep plasma by culturing in a c6/36 mosquito cell line (ATCC Cat.No. CRL

1660). The infection of cells in the absence of drugs and the assays for titration of

infectivity in liquid medium to determine tissue culture infective doses (TCID) were

performed as previously described (42). Titers were determined according to (43).

Assays to quantify plaque-forming units (pfu) were done in semisolid medium including

1% Carboxymethylcellulose (CMC, Sigma). Monolayers were fixed and stained 3-5

days post infection.

Infection in the presence of favipiravir

Favipiravir T-705 (Atomax chemicals Co. Ltd., China) was dissolved in DMSO at a

concentration of 100mM, aliquoted, and stored at -70ºC. For use in the experiments, the

solution was thawed and diluted in DMEM to reach the desired concentration indicated

in each experiment. Cells were pretreated with favipiravir 14 h prior to infection, the

virus was adsorbed for 1 h to cells in absence of drug, and then infection continued in

the presence of the same drug concentration.

Serial passages in the presence of favipiravir were performed by infecting Vero cell

monolayers seeded on M6 plates, in 2 ml of cell culture medium; 0.5 ml of the

supernatant was collected at 72 h post-infection, and used for the next infection. The

first passage was carried out at a multiplicity of infection (MOI) of 0.1 PFU/cell (4 $\times 10^5$

cells infected with 4 $\times 10^4$ PFU of virus). Mock-infected cells were maintained in

parallel to control for the absence of contaminating virus. When no cytopathic effect

(cpe) was observed, after several passages in the presence of favipiravir, the cell culture

supernatant was used to perform up to 5 blind passages in the absence of drug, to

ascertain that no infectious virus could be recovered. Lack of detectable cpe before and

after blind passages was considered as criterion of RVFV extinction. The IC_{50} of T-705

was calculated as described previously (19).

RNA extraction, RT-PCR, and nucleotide sequencing

RNA was extracted from the supernatants of infected cells using the Speedtools RNA

virus extraction kit (Biotools B&M Labs, S.A., Madrid, Spain) according to the

manufacturer's instructions. RT-PCR was performed using SuperScript IV reverse

transcriptase and Phusion High-Fidelity DNA polymerase (Thermo Scientific). cDNA

was obtained by retrotranscription from RVFV M-segment RNA using primer

MRV1ag (5'-CAAATGACTACCAGTCAGC-3'; positions 772-790; antigenomic

orientation) as described (44). The cDNA was then amplified by PCR using

oligonucleotides startGc forward (5'-ACTCGCATTGTCGACGTTTT) (positions 2091-

2109; genomic orientation) and sm3 reverse (5'-

GGATTAAGGAAGCGGGAAAAGCCC) (positions 3200-3223; antigenomic

orientation). The amplified cDNA is an 1100 nt fragment that corresponds to the amino-

terminal half of glycoprotein Gc, and it was chosen because it is highly conserved in

RVFV (45). Amplifications in the absence of RNA were carried out in parallel to

ascertain the absence of contaminating templates. For the analysis of mutant spectra,

amplifications were carried out with template preparations diluted 1:10, 1:100 and

1:1000. When amplification of the 1:100 diluted template yielded a detectable DNA

fragment, cloning was performed with the amplified fragment obtained with the

undiluted template, as described previously (46). This procedure ensures an excess of

template viral RNA molecules thus avoiding redundant amplifications of the same

initial RNA molecules. Amplified DNA fragments were then ligated to pJET2.1

plasmid (Thermo Scientific), and the products transformed into *E.coli* DH5α. The

pJET2.1 vector contains the lethal gene eco47IR enabling positive selection of

recombinant plasmids. Plasmid DNA from individual bacterial colonies was purified

and subjected to nucleotide sequencing, as previously described (47).

Quantification of RVFV RNA

For the calculation of specific infectivity (the ratio of infectivity to viral RNA in

a virus sample) a real time quantitative one-step PCR (RT-qPCR) of a fragment within

L-segment RNA was carried out using RNA HighScript Master mix (Biotools B&M),

according to manufacturer's instructions. Primers used were L-forward (5'-

TTCTTTGCTTCTGATACCCTCTG-3') [5' position 2872; sense orientation] and L-

reverse (5'-GTTCCACTTCCTTGCATCATCTG) [5' position 3006; antisense

orientation] that yield a 135bp fragment within the RdRp-coding region (41). Detection

was achieved by a L-segment specific hydrolysis probe (5'-[6-

FAM]TTGCACAAGTCCACACAGGCCCCT[BHQ1]-3'). Quantification of RNA was

relative to a standard curve obtained with known amounts of RVFV RNA, synthesized

by *in vitro* transcription of a NdeI-linearized pGEM-T plasmid containing the 135bp

cDNA target fragment using the T7 RiboMAX in vitro transcription kit (Promega).

Negative controls (without template RNA and RNA from mock-infected cells) were run

in parallel with each amplification reaction, to ascertain absence of contamination with

undesired templates.

Statistical analysis

Data analysis was performed using GraphPad Prism software version 6.

Acknowledgements

We thank Nuria de la Losa for excellent technical assistance. This work was supported

by grants S2013/ABI-2906 (PLATESA), P2018/BAA-4370 (PLATESA2) from

Comunidad de Madrid/FEDER. AGL2014-57430-R and AGL2017-83326-R from

Ministerio de Economía y Competitividad. SAF2014-52400-R and SAF2017-87846-R

from Ministerio de Ciencia, Innovación y Universidades. CIBERehd (Centro de

Investigación en Red de Enfermedades Hepáticas y Digestivas) is funded by Instituto de

Salud Carlos III. Institutional grants from the Fundación Ramón Areces and Banco

Santander to the CBMSO are also acknowledged.

References

1. Chevalier V. 2013. Relevance of Rift Valley fever to public health in the European Union. Clin Microbiol Infect 19:705-8.

2. Rolin AI, Berrang-Ford L, Kulkarni MA. 2013. The risk of Rift Valley fever virus introduction and establishment in the United States and European Union. Emerg Microbes Infect 2:e81.

3. Dal Pozzo F, Thiry E. 2014. Antiviral chemotherapy in veterinary medicine: current applications and perspectives. Rev Sci Tech 33:791-801.

4. De Vleeschauwer AR, Lefrebvre DJ, De Clercq K. 2017. Antiviral therapies for foot-and-mouth disease, p 357-384. *In* Sobrino F, Domingo E (ed), Foot-and-mouth Disease Virus Current Research and Emerging Trends. Caister Academic Press, Norfolk, UK.

5. Gowen BB, Wong MH, Jung KH, Sanders AB, Mendenhall M, Bailey KW, Furuta Y, Sidwell RW. 2007. In vitro and in vivo activities of T-705 against arenavirus and bunyavirus infections. Antimicrob Agents Chemother 51:3168-3176.

6. Caroline AL, Powell DS, Bethel LM, Oury TD, Reed DS, Hartman AL. 2014. Broad spectrum antiviral activity of favipiravir (T-705): protection from highly lethal inhalational Rift Valley Fever. PLoS Negl Trop Dis 8:e2790.

7. Scharton D, Bailey KW, Vest Z, Westover JB, Kumaki Y, Van Wettere A, Furuta Y, Gowen BB. 2014. Favipiravir (T-705) protects against peracute Rift Valley fever virus infection and reduces delayed-onset neurologic disease observed with ribavirin treatment. Antiviral Res 104:84-92.

8. Westover JB, Mathis A, Taylor R, Wandersee L, Bailey KW, Sefing EJ, Hickerson BT, Jung KH, Sheridan WP, Gowen BB. 2018. Galidesivir limits Rift

Valley fever virus infection and disease in Syrian golden hamsters. Antiviral Res

156:38-45.

9.   Delang L, Abdelnabi R, Neyts J. 2018. Favipiravir as a potential countermeasure

against neglected and emerging RNA viruses. Antiviral Res 153:85-94.

10.   de la Torre JC. 2018. Extending the Antiviral Value of Favipiravir. J Infect Dis

218:509-511.

11.   Furuta Y, Takahashi K, Fukuda Y, Kuno M, Kamiyama T, Kozaki K, Nomura

N, Egawa H, Minami S, Watanabe Y, Narita H, Shiraki K. 2002. In vitro and in

vivo activities of anti-influenza virus compound T-705. Antimicrob Agents

Chemother 46:977-81.

12.   Furuta Y, Komeno T, Nakamura T. 2017. Favipiravir (T-705), a broad spectrum

inhibitor of viral RNA polymerase. Proc Jpn Acad Ser B Phys Biol Sci 93:449-

463.

13.   Qiu L, Patterson SE, Bonnac LF, Geraghty RJ. 2018. Nucleobases and

corresponding nucleosides display potent antiviral activities against dengue

virus possibly through viral lethal mutagenesis. PLoS Negl Trop Dis

12:e0006421.

14.   Guedj J, Piorkowski G, Jacquot F, Madelain V, Nguyen THT, Rodallec A,

Gunther S, Carbonnelle C, Mentre F, Raoul H, de Lamballerie X. 2018.

Antiviral efficacy of favipiravir against Ebola virus: A translational study in

cynomolgus macaques. PLoS Med 15:e1002535.

15.   Bassi MR, Sempere RN, Meyn P, Polacek C, Arias A. 2018. Extinction of Zika

Virus and Usutu Virus by Lethal Mutagenesis Reveals Different Patterns of

Sensitivity to Three Mutagenic Drugs. Antimicrob Agents Chemother 62.

16. Goldhill DH, Langat P, Xie H, Galiano M, Miah S, Kellam P, Zambon M,
Lackenby A, Barclay WS. 2019. Determining the Mutation Bias of Favipiravir
in Influenza Virus Using Next-Generation Sequencing. J Virol 93.

17. Jin Z, Smith LK, Rajwanshi VK, Kim B, Deval J. 2013. The ambiguous base-
pairing and high substrate efficiency of T-705 (Favipiravir) Ribofuranosyl 5'-
triphosphate towards influenza A virus polymerase. PLoS One 8:e68347.

18. Sangawa H, Komeno T, Nishikawa H, Yoshida A, Takahashi K, Nomura N,
Furuta Y. 2013. Mechanism of action of T-705 ribosyl triphosphate against
influenza virus RNA polymerase. Antimicrob Agents Chemother 57:5202-8.

19. de Avila AI, Gallego I, Soria ME, Gregori J, Quer J, Esteban JI, Rice CM,
Domingo E, Perales C. 2016. Lethal Mutagenesis of Hepatitis C Virus Induced
by Favipiravir. PLoS One 11:e0164691.

20. de Avila AI, Moreno E, Perales C, Domingo E. 2017. Favipiravir can evoke
lethal mutagenesis and extinction of foot-and-mouth disease virus. Virus Res
233:105-112.

21. Escribano-Romero E, Jimenez de Oya N, Domingo E, Saiz JC. 2017. Extinction
of West Nile Virus by Favipiravir through Lethal Mutagenesis. Antimicrob
Agents Chemother 61:e01400-17.

22. Arias A, Thorne L, Goodfellow I. 2014. Favipiravir elicits antiviral mutagenesis
during virus replication in vivo. Elife 3:e03679.

23. Baranovich T, Wong SS, Armstrong J, Marjuki H, Webby RJ, Webster RG,
Govorkova EA. 2013. T-705 (Favipiravir) Induces Lethal Mutagenesis in
Influenza A H1N1 Viruses In Vitro. J Virol 87:3741-3751.

24. Loeb LA, Essigmann JM, Kazazi F, Zhang J, Rose KD, Mullins JI. 1999. Lethal

mutagenesis of HIV with mutagenic nucleoside analogs. Proc Natl Acad Sci U S

A 96:1492-7.

25. Sierra S, Davila M, Lowenstein PR, Domingo E. 2000. Response of foot-and-

mouth disease virus to increased mutagenesis: influence of viral load and fitness

in loss of infectivity. J Virol 74:8316-23.

26. Ortega-Prieto AM, Sheldon J, Grande-Perez A, Tejero H, Gregori J, Quer J,

Esteban JI, Domingo E, Perales C. 2013. Extinction of hepatitis C virus by

ribavirin in hepatoma cells involves lethal mutagenesis. PLoS ONE 8:e71039.

27. Sanchez-Campos S, Dominguez-Huerta G, Diaz-Martinez L, Tomas DM,

Navas-Castillo J, Moriones E, Grande-Perez A. 2018. Differential Shape of

Geminivirus Mutant Spectra Across Cultivated and Wild Hosts With Invariant

Viral Consensus Sequences. Front Plant Sci 9:932.

28. Grande-Pérez A, Gómez-Mariano G, Lowenstein PR, Domingo E. 2005.

Mutagenesis-induced, large fitness variations with an invariant arenavirus

consensus genomic nucleotide sequence. J Virol 79:10451-9.

29. Domingo E. 2016. Virus as Populations. Academic Press, Elsevier, Amsterdam.

30. González-López C, Gómez-Mariano G, Escarmís C, Domingo E. 2005.

Invariant aphthovirus consensus nucleotide sequence in the transition to error

catastrophe. Infection Genetics and Evolution 5:366-374.

31. Goldhill DH, Te Velthuis AJW, Fletcher RA, Langat P, Zambon M, Lackenby

A, Barclay WS. 2018. The mechanism of resistance to favipiravir in influenza.

Proc Natl Acad Sci U S A 115:11613-11618.

32. Daikoku T, Yoshida Y, Okuda T, Shiraki K. 2014. Characterization of
susceptibility variants of influenza virus grown in the presence of T-705. J
Pharmacol Sci 126:281-4.

33. Daikoku T, Mizuguchi M, Obita T, Yokoyama T, Yoshida Y, Takemoto M,
Shiraki K. 2018. Characterization of susceptibility variants of poliovirus grown
in the presence of favipiravir. J Microbiol Immunol Infect 51:581-586.

34. Vanderlinden E, Vrancken B, Van Houdt J, Rajwanshi VK, Gillemot S, Andrei
G, Lemey P, Naesens L. 2016. Distinct Effects of T-705 (Favipiravir) and
Ribavirin on Influenza Virus Replication and Viral RNA Synthesis. Antimicrob
Agents Chemother 60:6679-6691.

35. Takashita E, Fujisaki S, Shirakura M, Nakamura K, Kishida N, Kuwahara T,
Shimazu Y, Shimomura T, Watanabe S, Odagiri T. 2016. Influenza
A(H1N1)pdm09 virus exhibiting enhanced cross-resistance to oseltamivir and
peramivir due to a dual H275Y/G147R substitution, Japan, March 2016. Euro
Surveill 21.

36. Beaucourt S, Vignuzzi M. 2014. Ribavirin: a drug active against many viruses
with multiple effects on virus replication and propagation. Molecular basis of
ribavirin resistance. Curr Opin Virol 8:10-5.

37. Parker WB. 2005. Metabolism and antiviral activity of ribavirin. Virus Res
107:165-71.

38. Snell NJ. 2001. Ribavirin-current status of a broad spectrum antiviral agent.
Expert Opin Pharmacother 2:1317-24.

39. Lorenzo G, Lopez-Gil E, Ortego J, Brun A. 2018. Efficacy of different DNA
and MVA prime-boost vaccination regimens against a Rift Valley fever virus
(RVFV) challenge in sheep 12 weeks following vaccination. Vet Res 49:21.

40. Barnard BJ, Botha MJ. 1977. An inactivated rift valley fever vaccine. J S Afr
Vet Assoc 48:45-8.

41. Busquets N, Xavier F, Martin-Folgar R, Lorenzo G, Galindo-Cardiel I, del Val
BP, Rivas R, Iglesias J, Rodriguez F, Solanes D, Domingo M, Brun A. 2010.
Experimental infection of young adult European breed sheep with Rift Valley
fever virus field isolates. Vector Borne Zoonotic Dis 10:689-96.

42. Lorenzo G, Martin-Folgar R, Hevia E, Boshra H, Brun A. 2010. Protection
against lethal Rift Valley fever virus (RVFV) infection in transgenic IFNAR(-/-)
mice induced by different DNA vaccination regimens. Vaccine 28:2937-44.

43. Lindenbach BD. 2009. Measuring HCV infectivity produced in cell culture and
in vivo. Methods Mol Biol 510:329-36.

44. Takehara K, Min MK, Battles JK, Sugiyama K, Emery VC, Dalrymple JM,
Bishop DH. 1989. Identification of mutations in the M RNA of a candidate
vaccine strain of Rift Valley fever virus. Virology 169:452-7.

45. Bird BH, Khristova ML, Rollin PE, Ksiazek TG, Nichol ST. 2007. Complete
genome analysis of 33 ecologically and biologically diverse Rift Valley fever
virus strains reveals widespread virus movement and low genetic diversity due
to recent common ancestry. J Virol 81:2805-16.

46. Airaksinen A, Pariente N, Menendez-Arias L, Domingo E. 2003. Curing of foot-
and-mouth disease virus from persistently infected cells by ribavirin involves
enhanced mutagenesis. Virology 311:339-49.

47. Agudo R, Ferrer-Orta C, Arias A, de la Higuera I, Perales C, Perez-Luque R,
Verdaguer N, Domingo E. 2010. A multi-step process of viral adaptation to a
mutagenic nucleoside analogue by modulation of transition types leads to
extinction-escape. PLoS Pathog 6:e1001072.

Legend to Figures

Figure 1. **Effect of favipiravir treatment on viral yields in cell culture.** Vero cells

were pretreated overnight with the indicated concentrations of drug and then infected

with RVFV 56/74 at a moi of 0.1 PFU/cell. Infection proceeded for 72 hpi in the

presence of the same concentration of favipiravir. Results are shown either as

$TCID_{50}$/ml (continuous line; left Y axis) or as percentages of RVFV titer with respect to

the one obtained in untreated cells (dashed line; right Y axis). Values under the limit of

sensitivity of the assay (estimated to be 10, indicated with a grid line) were arbitrarily

represented as 2. Red line indicates 50%. Error bars denote SD.

Figure 2. **Serial passages of RVFV in Vero cells in the absence or presence of**

favipiravir. A) Viral titers of RVFV after each passage in the absence or presence of

different concentrations of favipiravir (T-705), as indicated. For concentrations higher

than 80 µM, only 3 passages were performed. Viral titers in the supernatant of infected

cells were determined at 72 hpi. Error bars denote SD. B) Comparison of the viral titers

obtained from virus recovered after passage 8^{th} (previously propagated or no with 40

µM favipiravir) and parental RVFV56/74 in the presence of a 5-80 µM range of

favipiravir concentrations. Titers are represented as percentages with respect to those

obtained without drug. C) Lack of detectable infectivity after blind passages as a

criterion for RVFV extinction. Cell culture supernatants from passages carried out in the

presence of favipiravir where no cytopathic effect (cpe) was observed (Figure 2A,

samples whose titres were below the limit of sensitivity of the assay) were subjected to

additional passages (up to five) in the absence of drug. Original samples are named

according to their passage number in the presence of the indicated favipiravir

concentration (in µM). Flattened columns indicate no cpe detection; lifted columns

indicate evident cpe.

Figure 3. **Effect of favipiravir in the specific activity of RVFV.** A) Viral titers of

RVFV after each passage (n=4) in the absence (white columns) or presence (black

columns) of 40 µM favipiravir (redrawn from data shown in Figure 2A). B)

Quantification by qRT-PCR (in triplicate) of RNA extracted from the supernatants of

untreated (white columns) or 40 µM favipiravir-treated (black columns) Vero cells after

each passage (n=4). C) Specific infectivity upon passage of RVFV in the absence (open

symbols) or presence (closed symbols) of 40 µM favipiravir. Values correspond to the

ratio between infectivity (panel A) and the amount of viral RNA (panel B). Error bars

denote SD. Statistically significant differences are indicated by asterisks (*p≤0.05,

**p≤0.01, ***p≤0.005, ****p≤0.0001; Student's t-test).

Figure 4. Distribution of mutations among individual clones. Plot of the distribution

of the number of mutations per molecular clone in the RVFV populations passaged in

the absence or presence of 40 µM favipiravir.

Figure 5. Mutational spectrum induced by favipiravir on RVFV. Matrices of

mutation types found in the glycoprotein Gc coding region of RVFV passaged 4 times

in Vero cells in the absence or presence of 40 µM favipiravir. The boxes below each

matrix quantify the mutational bias, according to the ratio shown on the left.

Table 1. Mutant spectrum analysis by molecular cloning and Sanger sequencing of glycoprotein Gc coding region of RVFV passaged 4
times in Vero cells

RVFV population[a]	Number of nucleotides analysed (clones/haplotypes)[b]	Different (total) mutations[c]	Minimum mutation frequency[d]	Maximum mutation frequency[e]	Different synonymous (non-synonymous) mutations	Total synonymous (non-synonymous) mutations
56/74 p4, no drug	31,610 (29/21)	53 (375)	1.7×10^{-3}	1.2×10^{-2}	48 (5)	354 (21)
56/74 p4 + favipiravir	31,610 (29/29)	138 (348)	4.4×10^{-3}	1.1×10^{-2}	85 (53)	284 (64)

[a]The origin and passage history of the viral populations analysed is described in Fig. 2 and Materials and Methods.

[b]The genomic region analysed spans residues 2110-3199 of glycoprotein Gc coding region; the residue numbering is that of the SA-75 genome.
The values in parenthesis indicate the number of clones analysed followed by the number of haplotypes (number of different RNA sequences).

[c]The number of different and total mutations were counted relative to the consensus sequence of the corresponding population. Different and
total mutations were used to calculate the minimum and maximum mutation frequency, respectively.

[d]Data represent the average number of different mutations per nucleotide in the mutant spectrum relative to the consensus sequence of the
corresponding population. The results indicate a significant 2.5-fold increase of minimum mutation frequency associated with replication in the
presence of favipiravir ($p<0.0001$; χ^2 test)

[e]Data represent the average number of total mutations per nucleotide in the mutant spectrum relative to the consensus sequence of the
corresponding population.

Figure 1

Figure 2

A

B

C

Figure 3

Figure 4

Figure 5

	No drug						Favipiravir 40 μM			
↗	U	C	G	A		↗	U	C	G	A
U		15	0	1		U		21	0	1
C	14		0	0		C	46		0	1
G	0	1		10		G	0	1		52
A	1	1	10			A	1	3	12	

$\dfrac{[(G{\rightarrow}A)+(C{\rightarrow}U)]}{[(A{\rightarrow}G)+(U{\rightarrow}C)]}$	0.96	2.97