Lethal mutagenesis of Rift Valley fever virus induced by favipiravir

Belén Borrego[a], Ana I. de Ávila[b] , Esteban Domingo[b,c] and Alejandro Brun[a]#

[a] Centro de Investigación en Sanidad Animal (CISA-INIA), Valdeolmos, Madrid, Spain

[b] Centro de Biología Molecular "Severo Ochoa" (CSIC-UAM), Madrid, Spain

[c]Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas

(CIBERehd) del Instituto de Salud Carlos III, 28029, Madrid, Spain,

Running Head: Mutagenic effect of favipiravir on RVFV

#Address correspondence to brun@inia.es (A. Brun)

15
16

ABSTRACT

Rift Valley fever virus (RVFV) is an emerging, mosquito-borne, zoonotic pathogen with

recurrent outbreaks paying an unacceptable toll of human beings in many African

countries, for which no effective treatment is available. In cell culture studies and with

laboratory animal models, the nucleoside analogue favipiravir (T-705) has demonstrated

great potential for the treatment of several seasonal, chronic and emerging RNA virus

infections of humans, suggesting applicability to control some viral outbreaks. Treatment

with favipiravir was shown to reduce the infectivity of Rift Valley fever virus both in cell

cultures and in experimental animal models, but the mechanism of this protective effect

is not understood. In this work we show that favipiravir at concentrations within the doses

proven effective *in vivo* is able to extinguish RVFV from infected cell cultures.

Nucleotide sequence analysis has documented RVFV mutagenesis associated with virus

extinction, with a significant increase in G to A and C to U transition frequencies, and a

decrease of specific infectivity, hallmarks of lethal mutagenesis.

INTRODUCTION

Rift Valley fever virus (RVFV) is a mosquito-borne bunyavirus belonging to the recently

reclassified phlebovirus genus (Family *Phenuiviridae*, Order Bunyavirales). RVFV

causes an important disease in ruminants often transmitted to humans after the occurrence

of epizootic outbreaks. Although the disease has been reported only in African countries

with some incursions in the Middle East and Indian Ocean islands, there are concerns for

its potential spread to other locations, including Europe (Chevalier, 2013; Rolin, Berrang-

Ford, and Kulkarni, 2013). Currently, there is no available treatment or licensed RVF

vaccine in Europe, therefore the development of effective control strategies is an essential

field of research. The use of antiviral agents for treatment in livestock is generally not

affordable, although it is often being proposed for viral veterinary diseases as an strategy

to fill the gap between the time of the infection and the effective development of the host

immune response (Dal Pozzo and Thiry, 2014; De Vleeschauwer, Lefrebvre, and De

Clercq, 2017). In contrast, in the case of humans infected or at risk of infection with

RVFV, antiviral treatment is fully warranted. Antiviral base and nucleoside analogues

effective against RVFV are available (Caroline et al., 2014; Gowen et al., 2007; Scharton

et al., 2014). Among the nucleoside analogs exerting anti-RVFV activity, ribavirin [1-β-

D-ribofuranosyl-1-H-1,2,4-triazole-3-carboxamide], favipiravir [6-fluoro-3-hydroxy-2-

pyrazinecarboxamide] and more recently BCX4430 [(2S,3S,4R,5R)-2-(4-amino-5H-

pyrrolo[3,2-d]pyrimidin-7-yl)-5-(hydroxymethyl)pyrrolidine-3,4-diol] (Galidesivir)

have been described (Caroline et al., 2014; Scharton et al., 2014; Westover et al., 2018).

Particularly, favipiravir displays potent antiviral activity against different RNA viruses

(Bassi et al., 2018; de la Torre, 2018; Delang, Abdelnabi, and Neyts, 2018; Furuta,

Komeno, and Nakamura, 2017; Furuta et al., 2002; Goldhill et al., 2019; Guedj et al.,

2018; Qiu et al., 2018). Upon cell intake, favipiravir is converted by cellular enzymes into

its active form (favipiravir-4-ribofuranosyl-5-triphosphate), which functions as a purine

nucleotide analogue. Favipiravir can act as an RNA chain terminator (Goldhill et al.,

2019; Jin et al., 2013; Sangawa et al., 2013), and as lethal mutagen for several RNA

viruses, including hepatitis C virus (de Avila et al., 2016), foot-and-mouth disease virus

(de Avila et al., 2017), West Nile fever virus (Escribano-Romero et al., 2017), norovirus

(Arias, Thorne, and Goodfellow, 2014), and influenza virus (Baranovich et al., 2013),

among others. Here we show that the mechanism of action of favipiravir against RVFV

in cell culture is due, at least in part, to the accumulation of mutations in the viral genome

that leads to a progressive decrease in viable viral progeny. This conclusion is based on a

significant increase of mutation frequency measured in a 1060 bp fragment from the Gc

(glycoprotein C)-coding region, and a concomitant decrease of specific infectivity,

associated with virus extinction, supporting the use of favipiravir for the control of

emerging RVFV infections in humans.

RESULTS

Effect of favipiravir (T-705) on RVFV 56/74 progeny virus yield.

The aim of this work was to determine whether favipiravir induces lethal mutagenesis

upon RVFV infection in cell culture. We first determined the effect of different

concentrations of favipiravir on the yield of RVFV isolate 56/74 in Vero cells. The half

inhibitory concentration (IC_{50}, concentration required for a 50% decrease in the

production of infectious progeny), was calculated to be 6.22 ± 1.5 μM (average of 3

determinations) **(Figure 1).** Since the cytotoxic concentration 50 (CC_{50}) of T-705 for

Vero cells is >10000 μM (Escribano-Romero et al., 2017), the therapeutic index for

RVFV (TI = CC_{50} / IC_{50}) is > 1600.

The RVFV 56/74 isolate was subjected to serial passages in Vero cells in the absence or

presence of different concentrations of favipiravir (**Figure 2a**). The titration of

supernatants indicated loss of infectivity in the passages carried out in the presence of 80

μM favipiravir, as well as for higher doses, such as 160, 200 and 500 μM, but not at the

lowest concentrations tested. A concentration of 20 μM produced a slight reduction in

viral yield during the consecutive passages, whilst at 40 μM the progeny production

declined progressively, and was not detected after the fifth, sixth and seventh passage.

Interestingly, virus yields were recovered at passages eighth and ninth.

To confirm virus extinction, cell culture supernatants from passages carried out in the

presence of favipiravir where no cytopathic effect (cpe) had been observed were subjected

to up to five additional passages in the absence of the drug (**Figure 2b**). No infectivity

was rescued from samples that were collected after serial passages in the presence of 80

μM favipiravir or higher. Viral infectivity could be recovered from supernatants of cells

previously treated with 40 μM favipiravir. These data confirmed the previous

observation, indicating that the deleterious effect on virus propagation at this

concentration was not complete.

Analysis of RVFV passaged in absence or presence of favipiravir

Decrease of specific infectivity, broadening of the mutant spectrum and invariance of the

consensus sequence of the viral population are diagnostic of lethal mutagenesis

(Baranovich et al., 2013; de Avila et al., 2016; de Avila et al., 2017; Escribano-Romero

et al., 2017; Grande-Pérez et al., 2005; Loeb et al., 1999; Ortega-Prieto et al., 2013;

Sanchez-Campos et al., 2018; Sierra et al., 2000). To explore these parameters, RVFV

populations at passages 1 to 4 in absence or presence of 40 μM favipiravir were examined.

Extracellular viral titers and viral RNA decreased significantly in the presence of

favipiravir (**Figure 3**). The changes in infectivity and viral RNA production resulted in a

gradual decrease of specific infectivity of the populations passaged in the presence of

favipiravir (up to 8-fold in four passages) but not in its absence (p<0.0001 for passages 1

to 4; t test). Thus, the first diagnosis criterium of lethal mutagenesis is fulfilled for

favipiravir acting on RVFV.

To examine whether the decrease in progeny production was associated with increased

mutagenesis of the viral RNA, total RNA from the cell culture supernatant of passage 4

in the absence or presence of 40 μM favipiravir was extracted and a 1.1kb DNA fragment

encoding a glycoprotein Gc region comprising nucleotides 2091 to 3223 (numbering

according to GenBank sequence of RVFV strain SA-75, accession number DQ380189)

was amplified by RT-PCR and subjected to molecular cloning and nucleotide sequencing.

For each condition (with/without drug) a total of 29 clones were analyzed, corresponding

to 31,610 nucleotides sequenced. The results (**Table 1**) indicated a significant 2.5-fold

increase of minimum mutation frequency associated with replication in the presence of

favipiravir (p< 0.0001; $\chi2$ test). The distribution of the number of Gc sequences was

analyzed as a function of the number of mutations per clone (**Figure 4**). While the viral

population that was passaged in absence of drug yielded 7 clones without mutations and

7 clones with only one mutation (24.1% in both cases), only 1 clone without mutations

and 2 with one mutation were retrieved from the population that was passaged in the

presence of favipiravir (1/29, 3.4%; 2/29, 6.9%, respectively). Finally, the consensus

genomic nucleotide sequence (analyzed at passage 4) did not change as a result of

replication in the presence of favipiravir, an observation already found during lethal

mutagenesis of other RNA viruses (de Avila et al., 2016; Domingo, 2016; González-

López et al., 2005; Grande-Pérez et al., 2005; Sanchez-Campos et al., 2018).

Analysis of mutation types

In order to analyze whether passage of RVFV in the presence of favipiravir resulted in

any deviation of mutation types, we compared the mutation matrices in the mutant

spectrum of the populations passaged in absence or presence of the drug. The results

indicate a predominance of G to A and C to U transitions associated with replication in

the presence of favipiravir: without drug the number of changes G→A + C→U was 24

(10+14), while with favipiravir it was 98 (52+46), thus resulting in 3-fold increase of the

ratio (G→A) + (C→U) / (A→G) + (U→C) (**Figure 5**). The difference between the type

of mutations was statistically significant (p< 0.0001; χ2 test), and reveals that favipiravir

produces a bias in favor of GA and CU transitions; the transition to transversion ratio was

comparable: 12.2 and 18.7 upon passage in absence and presence of favipiravir,

respectively.

These results indicate that favipiravir is a mutagenic agent for RVFV and suggest that

its anti-RVFV activity may be exerted at least in part through lethal mutagenesis.

DISCUSSION

The broad spectrum antiviral activity of favipiravir has been established with

representatives of several families of RNA viruses, so it stands as a promising drug to be

used against viral pathogens. In the case of Rift Valley fever, for which no therapeutic or

prophylactic treatments are available for humans, favipiravir could be a powerful

therapeutic tool to provide immediate treatment to farmers, veterinary and research

personnel at risk of infection after being in contact with the virus or virus-infected

animals. Favipiravir was initially considered an inhibitor of viral RNA synthesis, exerted

mainly through RNA chain termination (Goldhill et al., 2019; Jin et al., 2013; Sangawa

et al., 2013). However, model studies with several viruses in cell culture and *in vivo*

suggest that the antiviral activity of favipiravir may include also lethal mutagenesis.

Previous studies have shown that favipiravir treatment of laboratory animals exposed to

RVFV increased their survival, and counteracted acute or neurological disease syndromes

(Caroline et al., 2014; Scharton et al., 2014). Furthermore, treatment efficacy was

enhanced when favipiravir was combined with ribavirin treatment (Scharton et al., 2014).

Knowledge of the mechanism of action of this promising drug is important to improve

antiviral designs. Our data clearly show at least a 2.5-fold increased RVFV mutation

frequencies when the virus was propagated in the presence of 40 μM favipiravir, a dose

which is 10- to 25-fold lower than the dose proven effective for other RNA viruses (de

Avila et al., 2016; de Avila et al., 2017; Escribano-Romero et al., 2017). The comparison

of TI values favors the therapeutic use of this drug in humans or animals exposed to the

virus.

One of the most advantageous features of favipiravir is that it shows a high barrier for the

generation of resistance mutants (Delang, Abdelnabi, and Neyts, 2018). Since its activity

against human influenza virus was first described (Furuta et al., 2002), only one resistant

mutant has been reported (Goldhill et al., 2018). Attempts to isolate resistant variants of

polio- and influenza viruses seem to have been unsuccessful (Daikoku et al., 2018;

Daikoku et al., 2014; Vanderlinden et al., 2016). In addition, influenza virus pairs isolated

from patients before and after favipiravir administration showed no changes in the

susceptibility to the drug (Takashita et al., 2016). Similar results were reported in

norovirus-infected mice after favipiravir treatment (Arias, Thorne, and Goodfellow,

2014). Moreover, previous studies reporting favipiravir mediated viral extinction by

lethal mutagenesis did not describe selection of specific mutations or resistant variants

(Baranovich et al., 2013; de Avila et al., 2016; de Avila et al., 2017; Escribano-Romero

et al., 2017). In our system, 40 μM favipiravir seemed not to lead to the extinction of the

virus, since after a lag phase of three consecutive blind passages the presence of cpe was

detected again in cultures. This observation could be due to the selection of mutant viruses

with increased resistance to the drug. Work is currently underway in order to test this

hypothesis.

Nonetheless, the appearance of variants showing decreased susceptibility to the drug

(Delang et al., 2014) highlights the importance of deciphering the exact mechanisms

exerted by new promising antiviral drugs, such as favipiravir. In this sense, it is worth

noting that other studies found that drug concentrations close to the IC_{50} led to viral

extinction of influenza virus without the emergence of variants (Baranovich et al., 2013)

perhaps indicating that other unknown factors play a role the inhibitory mechanism of T-

705. Still, the antiviral activity mechanism of favipiravir is not fully understood. Most

nucleoside analogues used as antiviral agents display several mechanisms of activity

(Beaucourt and Vignuzzi, 2014; Parker, 2005; Snell, 2001). This is also the case for

favipiravir with a dual chain terminating and mutagenic activity by ambiguous base

pairing, as well as a possible role in modulation of innate immunity (de la Torre, 2018).

In summary, what our results suggest is that favipiravir can act as a lethal mutagen for

RVFV but they do not exclude that other mechanisms, alone or together with lethal

mutagenesis, are required for virus extinction.

MATERIALS AND METHODS

Cells, viruses and infections

Vero cells (ATCC Cat.No. CCL-81) were grown in Dulbecco's modified Eagle's medium

supplemented with 5% −10% fetal calf serum (FCS), and L-glutamine (2 mM), penicillin

(100 U/ml) and streptomycin (100 μg/ml), in a humid atmosphere of 5% CO_2 at 37°C.

The virus used in this study originates from a sheep experimentally infected in our

laboratory (Lorenzo et al., 2018) with RVFV isolate 56/74 (Barnard and Botha, 1977;

Busquets et al., 2010). The virus was re-isolated from infected sheep plasma by culturing

in a c6/36 mosquito cell line (ATCC Cat.No. CRL 1660). The infection of cells in the

absence of drugs and the assays for titration of infectivity in liquid medium to determine

tissue culture infective doses (TCID) were performed as previously described(Lorenzo et

al., 2010). Titers were determined according to Lindenbach (2009). Assays to quantify

plaque-forming units (pfu) were done in semisolid medium including 1%

Carboxymethylcellulose (CMC, Sigma). Monolayers were fixed and stained 3-5 days

post infection.

Infection in the presence of favipiravir

Favipiravir T-705 (Atomax chemicals Co. Ltd., China) was dissolved in DMSO at a

concentration of 100mM, aliquoted, and stored at -70ºC. For use in the experiments, the

solution was thawed and diluted in DMEM to reach the desired concentration indicated

in each experiment. Cells were pretreated with favipiravir 14 h prior to infection, the virus

was adsorbed for 1 h to cells in absence of drug, and then infection continued in the

presence of the same drug concentration.

Serial passages in the presence of favipiravir were performed by infecting Vero cell

monolayers seeded on M6 plates, in 2 ml of cell culture medium; 0.5 ml of the supernatant

was collected at 72 h post-infection, and used for the next infection. The first passage was

carried out at a multiplicity of infection (MOI) of 0.1 PFU/cell (4 x10^5 cells infected with

4 x10^4 PFU of virus). Mock-infected cells were maintained in parallel to control for the

absence of contaminating virus. When no cytopathic effect (cpe) was observed, after

several passages in the presence of favipiravir, the cell culture supernatant was used to

perform up to 5 blind passages in the absence of drug, to ascertain that no infectious virus

could be recovered. Lack of detectable cpe before and after blind passages was considered

as criterion of RVFV extinction. The IC_{50} of T-705 was calculated as described previously

(de Avila et al., 2016).

RNA extraction, RT-PCR, and nucleotide sequencing

RNA was extracted from the supernatants of infected cells using the Speedtools RNA

virus extraction kit (Biotools B&M Labs, S.A., Madrid, Spain) according to the

manufacturer's instructions. RT-PCR was performed using SuperScript IV reverse

transcriptase and Phusion High-Fidelity DNA polymerase (Thermo Scientific). cDNA

was obtained by retrotranscription from RVFV M-segment RNA using primer MRV1ag

(5'-CAAATGACTACCAGTCAGC-3'; positions 772-790; antigenomic orientation) as

described (Takehara et al., 1989). The cDNA was then amplified by PCR using

oligonucleotides startGc forward (5'-ACTCGCATTGTCGACGTTTT) (positions 2091-

2109;   genomic   orientation)   and   sm3   reverse   (5'-

GGATTAAGGAAGCGGGAAAAGCCC)   (positions   3200-3223;   antigenomic

orientation). The amplified cDNA is an 1100 nt fragment that corresponds to the amino-

terminal half of glycoprotein Gc, and it was chosen because it is highly conserved in

RVFV (Bird et al., 2007). Amplifications in the absence of RNA were carried out in

parallel to ascertain the absence of contaminating templates. For the analysis of mutant

spectra, amplifications were carried out with template preparations diluted 1:10, 1:100

and 1:1000. When amplification of the 1:100 diluted template yielded a detectable DNA

fragment, cloning was performed with the amplified fragment obtained with the undiluted

template, as described previously (Airaksinen et al., 2003). This procedure ensures an

excess of template viral RNA molecules thus avoiding redundant amplifications of the

same initial RNA molecules. Amplified DNA fragments were then ligated to pJET2.1

plasmid (Thermo Scientific), and the products transformed into *E.coli* DH5α. The

pJET2.1 vector contains the lethal gene eco47IR enabling positive selection of

recombinant plasmids. Plasmid DNA from individual bacterial colonies was purified and

subjected to nucleotide sequencing, as previously described (Agudo et al., 2010).

Quantification of RVFV RNA

For the calculation of specific infectivity (the ratio of infectivity to viral RNA in

a virus sample) a real time quantitative one-step PCR (RT-qPCR) of a fragment within

L-segment RNA was carried out using RNA HighScript Master mix (Biotools B&M),

according to manufacturer's instructions. Primers used were L-forward (5'-

TTCTTTGCTTCTGATACCCTCTG-3') [5' position 2872; sense orientation] and L-

reverse (5'-GTTCCACTTCCTTGCATCATCTG) [5' position 3006; antisense

orientation] that yield a 135bp fragment within the RdRp-coding region (Busquets et al.,

2010). Detection was achieved by a L-segment specific hydrolysis probe (5'-[6-

[FAM]TTGCACAAGTCCACACAGGCCCCT[BHQ1]-3'). Quantification of RNA was

relative to a standard curve obtained with known amounts of RVFV RNA, synthesized

by *in vitro* transcription of a NdeI-linearized pGEM-T plasmid containing the 135bp

cDNA target fragment using the T7 RiboMAX in vitro transcription kit (Promega).

Negative controls (without template RNA and RNA from mock-infected cells) were run

in parallel with each amplification reaction, to ascertain absence of contamination with

undesired templates.

Statistical analysis

Data analysis was performed using GraphPad Prism software version 6.

Acknowledgements

We thank Nuria de la Losa for excellent technical assistance. This work was supported by grants S2013/ABI-2906 (PLATESA), P2018/BAA-4370 (PLATESA2) from Comunidad de Madrid/FEDER. AGL2014-57430-R and AGL2017-83326-R from Ministerio de Economía y Competitividad. SAF2014-52400-R and SAF2017-87846-R from Ministerio de Ciencia, Innovación y Universidades. CIBERehd (Centro de Investigación en Red de Enfermedades Hepáticas y Digestivas) is funded by Instituto de Salud Carlos III. Institutional grants from the Fundación Ramón Areces and Banco Santander to the CBMSO are also acknowledged.

References

Agudo, R., Ferrer-Orta, C., Arias, A., de la Higuera, I., Perales, C., Perez-Luque, R.,

Verdaguer, N., and Domingo, E. (2010). A multi-step process of viral adaptation

to a mutagenic nucleoside analogue by modulation of transition types leads to

extinction-escape. *PLoS Pathog* 6(8), e1001072.

Airaksinen, A., Pariente, N., Menendez-Arias, L., and Domingo, E. (2003). Curing of

foot-and-mouth disease virus from persistently infected cells by ribavirin involves

enhanced mutagenesis. *Virology* 311(2), 339-49.

Arias, A., Thorne, L., and Goodfellow, I. (2014). Favipiravir elicits antiviral mutagenesis

during virus replication in vivo. *Elife* 3, e03679.

Baranovich, T., Wong, S. S., Armstrong, J., Marjuki, H., Webby, R. J., Webster, R. G.,

and Govorkova, E. A. (2013). T-705 (Favipiravir) Induces Lethal Mutagenesis in

Influenza A H1N1 Viruses In Vitro. *J Virol* 87(7), 3741-3751.

Barnard, B. J., and Botha, M. J. (1977). An inactivated rift valley fever vaccine. *J S Afr*

Vet Assoc 48(1), 45-8.

Bassi, M. R., Sempere, R. N., Meyn, P., Polacek, C., and Arias, A. (2018). Extinction of

Zika Virus and Usutu Virus by Lethal Mutagenesis Reveals Different Patterns of

Sensitivity to Three Mutagenic Drugs. *Antimicrob Agents Chemother* 62(9).

Beaucourt, S., and Vignuzzi, M. (2014). Ribavirin: a drug active against many viruses

with multiple effects on virus replication and propagation. Molecular basis of

ribavirin resistance. *Curr Opin Virol* 8, 10-5.

Bird, B. H., Khristova, M. L., Rollin, P. E., Ksiazek, T. G., and Nichol, S. T. (2007).

Complete genome analysis of 33 ecologically and biologically diverse Rift Valley

fever virus strains reveals widespread virus movement and low genetic diversity

due to recent common ancestry. *J Virol* 81(6), 2805-16.

Busquets, N., Xavier, F., Martin-Folgar, R., Lorenzo, G., Galindo-Cardiel, I., del Val, B.

P., Rivas, R., Iglesias, J., Rodriguez, F., Solanes, D., Domingo, M., and Brun, A.

(2010). Experimental infection of young adult European breed sheep with Rift

Valley fever virus field isolates. *Vector Borne Zoonotic Dis* 10(7), 689-96.

Caroline, A. L., Powell, D. S., Bethel, L. M., Oury, T. D., Reed, D. S., and Hartman, A.

L. (2014). Broad spectrum antiviral activity of favipiravir (T-705): protection

from highly lethal inhalational Rift Valley Fever. *PLoS Negl Trop Dis* 8(4),

e2790.

Chevalier, V. (2013). Relevance of Rift Valley fever to public health in the European

Union. *Clin Microbiol Infect* 19(8), 705-8.

Daikoku, T., Mizuguchi, M., Obita, T., Yokoyama, T., Yoshida, Y., Takemoto, M., and

Shiraki, K. (2018). Characterization of susceptibility variants of poliovirus grown

in the presence of favipiravir. *J Microbiol Immunol Infect* 51(5), 581-586.

Daikoku, T., Yoshida, Y., Okuda, T., and Shiraki, K. (2014). Characterization of

susceptibility variants of influenza virus grown in the presence of T-705. *J

Pharmacol Sci* 126(3), 281-4.

Dal Pozzo, F., and Thiry, E. (2014). Antiviral chemotherapy in veterinary medicine:

current applications and perspectives. *Rev Sci Tech* 33(3), 791-801.

de Avila, A. I., Gallego, I., Soria, M. E., Gregori, J., Quer, J., Esteban, J. I., Rice, C. M.,

Domingo, E., and Perales, C. (2016). Lethal Mutagenesis of Hepatitis C Virus

Induced by Favipiravir. *PLoS One* 11(10), e0164691.

de Avila, A. I., Moreno, E., Perales, C., and Domingo, E. (2017). Favipiravir can evoke

lethal mutagenesis and extinction of foot-and-mouth disease virus. *Virus Res* 233,

105-112.

de la Torre, J. C. (2018). Extending the Antiviral Value of Favipiravir. *J Infect Dis* 218(4),

509-511.

De Vleeschauwer, A. R., Lefrebvre, D. J., and De Clercq, K. (2017). Antiviral therapies

for foot-and-mouth disease. *In* "Foot-and-mouth Disease Virus. Current Research

and Emerging Trends" (F. Sobrino, and E. Domingo, Eds.), pp. 357-384. Caister

Academic Press, Norfolk, UK.

Delang, L., Abdelnabi, R., and Neyts, J. (2018). Favipiravir as a potential countermeasure

against neglected and emerging RNA viruses. *Antiviral Res* 153, 85-94.

Domingo, E. (2016). "Virus as Populations." Academic Press, Elsevier, Amsterdam.

Escribano-Romero, E., Jimenez de Oya, N., Domingo, E., and Saiz, J. C. (2017).

Extinction of West Nile Virus by Favipiravir through Lethal Mutagenesis.

Antimicrob Agents Chemother 61:e01400-17(11).

Furuta, Y., Komeno, T., and Nakamura, T. (2017). Favipiravir (T-705), a broad spectrum

inhibitor of viral RNA polymerase. *Proc Jpn Acad Ser B Phys Biol Sci* 93(7), 449-

463.

Furuta, Y., Takahashi, K., Fukuda, Y., Kuno, M., Kamiyama, T., Kozaki, K., Nomura,

N., Egawa, H., Minami, S., Watanabe, Y., Narita, H., and Shiraki, K. (2002). In

vitro and in vivo activities of anti-influenza virus compound T-705. *Antimicrob*

Agents Chemother 46(4), 977-81.

Goldhill, D. H., Langat, P., Xie, H., Galiano, M., Miah, S., Kellam, P., Zambon, M.,

Lackenby, A., and Barclay, W. S. (2019). Determining the Mutation Bias of

Favipiravir in Influenza Virus Using Next-Generation Sequencing. *J Virol* 93(2).

Goldhill, D. H., Te Velthuis, A. J. W., Fletcher, R. A., Langat, P., Zambon, M., Lackenby,

A., and Barclay, W. S. (2018). The mechanism of resistance to favipiravir in

influenza. *Proc Natl Acad Sci U S A* 115(45), 11613-11618.

González-López, C., Gómez-Mariano, G., Escarmís, C., and Domingo, E. (2005).

Invariant aphthovirus consensus nucleotide sequence in the transition to error

catastrophe. *Infection Genetics and Evolution* 5, 366-374.

Gowen, B. B., Wong, M. H., Jung, K. H., Sanders, A. B., Mendenhall, M., Bailey, K. W.,

Furuta, Y., and Sidwell, R. W. (2007). In vitro and in vivo activities of T-705

against arenavirus and bunyavirus infections. *Antimicrob Agents Chemother*

51(9), 3168-3176.

Grande-Pérez, A., Gómez-Mariano, G., Lowenstein, P. R., and Domingo, E. (2005).

Mutagenesis-induced, large fitness variations with an invariant arenavirus

consensus genomic nucleotide sequence. *J. Virol.* 79(16), 10451-9.

Guedj, J., Piorkowski, G., Jacquot, F., Madelain, V., Nguyen, T. H. T., Rodallec, A.,

Gunther, S., Carbonnelle, C., Mentre, F., Raoul, H., and de Lamballerie, X.

(2018). Antiviral efficacy of favipiravir against Ebola virus: A translational study

in cynomolgus macaques. *PLoS Med* 15(3), e1002535.

Jin, Z., Smith, L. K., Rajwanshi, V. K., Kim, B., and Deval, J. (2013). The ambiguous

base-pairing and high substrate efficiency of T-705 (Favipiravir) Ribofuranosyl

5'-triphosphate towards influenza A virus polymerase. *PLoS One* 8(7), e68347.

Lindenbach, B. D. (2009). Measuring HCV infectivity produced in cell culture and in

vivo. *Methods Mol Biol* 510, 329-36.

Loeb, L. A., Essigmann, J. M., Kazazi, F., Zhang, J., Rose, K. D., and Mullins, J. I. (1999).

Lethal mutagenesis of HIV with mutagenic nucleoside analogs. *Proc Natl Acad*

Sci U S A 96(4), 1492-7.

Lorenzo, G., Lopez-Gil, E., Ortego, J., and Brun, A. (2018). Efficacy of different DNA

and MVA prime-boost vaccination regimens against a Rift Valley fever virus

(RVFV) challenge in sheep 12 weeks following vaccination. *Vet Res* 49(1), 21.

Lorenzo, G., Martin-Folgar, R., Hevia, E., Boshra, H., and Brun, A. (2010). Protection

against lethal Rift Valley fever virus (RVFV) infection in transgenic IFNAR(-/-)

mice induced by different DNA vaccination regimens. *Vaccine* 28(17), 2937-44.

Ortega-Prieto, A. M., Sheldon, J., Grande-Perez, A., Tejero, H., Gregori, J., Quer, J.,

Esteban, J. I., Domingo, E., and Perales, C. (2013). Extinction of hepatitis C virus

by ribavirin in hepatoma cells involves lethal mutagenesis. *PLoS ONE* 8(8),

e71039.

Parker, W. B. (2005). Metabolism and antiviral activity of ribavirin. *Virus Res.* 107(2),

165-71.

Qiu, L., Patterson, S. E., Bonnac, L. F., and Geraghty, R. J. (2018). Nucleobases and

corresponding nucleosides display potent antiviral activities against dengue virus

possibly through viral lethal mutagenesis. *PLoS Negl Trop Dis* 12(4), e0006421.

Rolin, A. I., Berrang-Ford, L., and Kulkarni, M. A. (2013). The risk of Rift Valley fever

virus introduction and establishment in the United States and European Union.

Emerg Microbes Infect 2(12), e81.

Sanchez-Campos, S., Dominguez-Huerta, G., Diaz-Martinez, L., Tomas, D. M., Navas-

Castillo, J., Moriones, E., and Grande-Perez, A. (2018). Differential Shape of

Geminivirus Mutant Spectra Across Cultivated and Wild Hosts With Invariant

Viral Consensus Sequences. *Front Plant Sci* 9, 932.

Sangawa, H., Komeno, T., Nishikawa, H., Yoshida, A., Takahashi, K., Nomura, N., and

Furuta, Y. (2013). Mechanism of action of T-705 ribosyl triphosphate against

influenza virus RNA polymerase. *Antimicrob Agents Chemother* 57(11), 5202-8.

Scharton, D., Bailey, K. W., Vest, Z., Westover, J. B., Kumaki, Y., Van Wettere, A.,

Furuta, Y., and Gowen, B. B. (2014). Favipiravir (T-705) protects against peracute

Rift Valley fever virus infection and reduces delayed-onset neurologic disease observed with ribavirin treatment. *Antiviral Res* 104, 84-92.

Sierra, S., Davila, M., Lowenstein, P. R., and Domingo, E. (2000). Response of foot-and-mouth disease virus to increased mutagenesis: influence of viral load and fitness in loss of infectivity. *J Virol* 74(18), 8316-23.

Snell, N. J. (2001). Ribavirin-current status of a broad spectrum antiviral agent. *Expert Opin. Pharmacother.* 2(8), 1317-24.

Takashita, E., Fujisaki, S., Shirakura, M., Nakamura, K., Kishida, N., Kuwahara, T., Shimazu, Y., Shimomura, T., Watanabe, S., and Odagiri, T. (2016). Influenza A(H1N1)pdm09 virus exhibiting enhanced cross-resistance to oseltamivir and peramivir due to a dual H275Y/G147R substitution, Japan, March 2016. *Euro Surveill* 21(24).

Takehara, K., Min, M. K., Battles, J. K., Sugiyama, K., Emery, V. C., Dalrymple, J. M., and Bishop, D. H. (1989). Identification of mutations in the M RNA of a candidate vaccine strain of Rift Valley fever virus. *Virology* 169(2), 452-7.

Vanderlinden, E., Vrancken, B., Van Houdt, J., Rajwanshi, V. K., Gillemot, S., Andrei, G., Lemey, P., and Naesens, L. (2016). Distinct Effects of T-705 (Favipiravir) and Ribavirin on Influenza Virus Replication and Viral RNA Synthesis. *Antimicrob Agents Chemother* 60(11), 6679-6691.

Westover, J. B., Mathis, A., Taylor, R., Wandersee, L., Bailey, K. W., Sefing, E. J., Hickerson, B. T., Jung, K. H., Sheridan, W. P., and Gowen, B. B. (2018). Galidesivir limits Rift Valley fever virus infection and disease in Syrian golden hamsters. *Antiviral Res* 156, 38-45.

Legend to Figures

Figure 1. **Effect of favipiravir treatment on viral yields in cell culture.** Vero cells were

pretreated overnight with the indicated concentrations of drug and then infected with

RVFV 56/74 at a moi of 0.1 PFU/cell. Infection proceeded for 72 hpi in the presence of

the same concentration of favipiravir. Results are shown either as $TCID_{50}$/ml (continuous

line; left Y axis) or as percentages of RVFV titer with respect to the one obtained in

untreated cells (dashed line; right Y axis). Values under the limit of sensitivity of the

assay (estimated to be 10, indicated with a grid line) were arbitrarily represented as 2.

Red line indicates 50%. Error bars denote SD.

Figure 2. **Serial passages of RVFV in Vero cells in the absence or presence of**

favipiravir. A) Viral titers of RVFV after each passage in the absence or presence of

different concentrations of favipiravir (T-705), as indicated. For concentrations higher

than 80 μM, only 3 passages were performed. Viral titers in the supernatant of infected

cells were determined at 72 hpi. Error bars denote SD. B) Lack of detectable infectivity

after blind passages as a criterion for RVFV extinction. Cell culture supernatants from

passages carried out in the presence of favipiravir where no cytopathic effect (cpe) was

observed (Figure 2A, samples whose titres were below the limit of sensitivity of the assay)

were subjected to additional passages (up to five) in the absence of drug. Original samples

are named according to their passage number in the presence of the indicated favipiravir

concentration (in μM). Flattened columns indicate no cpe detection; lifted columns

indicate evident cpe.

Figure 3. **Effect of favipiravir in the specific activity of RVFV.** A) Viral titers of RVFV

after each passage (n=4) in the absence (white columns) or presence (black columns) of

40 μM favipiravir (redrawn from data shown in Figure 2A). B) Quantification by qRT-

PCR (in triplicate) of RNA extracted from the supernatants of untreated (white columns)

or 40 μM favipiravir-treated (black columns) Vero cells after each passage (n=4). C)

Specific infectivity upon passage of RVFV in the absence (open symbols) or presence

(closed symbols) of 40 μM favipiravir. Values correspond to the ratio between infectivity

(panel A) and the amount of viral RNA (panel B). Error bars denote SD. Statistically

significant differences are indicated by asterisks (*$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.005$,

****$p \leq 0.0001$; Student's t-test).

Figure 4. Distribution of mutations among individual clones. Plot of the distribution

of the number of mutations per molecular clone in the RVFV populations passaged in the

absence or presence of 40 μM favipiravir.

Figure 5. Mutational spectrum induced by favipiravir on RVFV. Matrices of

mutation types found in the glycoprotein Gc coding region of RVFV passaged 4 times in

Vero cells in the absence or presence of 40 μM favipiravir. The boxes below each matrix

quantify the mutational bias, according to the ratio shown on the left.

Table 1. Mutant spectrum analysis by molecular cloning and Sanger sequencing of glycoprotein Gc coding region of RVFV passaged 4 times in Vero cells

RVFV population[a]	Number of nucleotides analysed (clones/haplotypes)[b]	Different (total) mutations[c]	Minimum mutation frequency[d]	Maximum mutation frequency[e]	Different synonymous (non-synonymous) mutations	Total synonymous (non-synonymous) mutations
56/74 p4, no drug	31,610 (29/21)	53 (375)	1.7×10^{-3}	1.2×10^{-2}	48 (5)	354 (21)
56/74 p4 + favipiravir	31,610 (29/29)	138 (348)	4.4×10^{-3}	1.1×10^{-2}	85 (53)	284 (64)

[a]The origin and passage history of the viral populations analysed is described in Fig. 2 and Materials and Methods.

[b]The genomic region analysed spans residues 2110-3199 of glycoprotein Gc coding region; the residue numbering is that of the SA-75 genome. The values in parenthesis indicate the number of clones analysed followed by the number of haplotypes (number of different RNA sequences).

[c]The number of different and total mutations were counted relative to the consensus sequence of the corresponding population. Different and total mutations were used to calculate the minimum and maximum mutation frequency, respectively.

[d]Data represent the average number of different mutations per nucleotide in the mutant spectrum relative to the consensus sequence of the corresponding population. The results indicate a significant 2.5-fold increase of minimum mutation frequency associated with replication in the presence of favipiravir ($p<0.0001$; χ^2 test)

[e]Data represent the average number of total mutations per nucleotide in the mutant spectrum relative to the consensus sequence of the corresponding population.

Figure 1

Figure 2

A

B

Figure 3

Figure 4

Mutations/clone

No drug

↗	U	C	G	A
U		15	0	1
C	14		0	0
G	0	1		10
A	1	1	10	

Favipiravir 40 μM

↗	U	C	G	A
U		21	0	1
C	46		0	1
G	0	1		52
A	1	3	12	

$$\frac{[(G \rightarrow A)+(C \rightarrow U)]}{[(A \rightarrow G)+(U \rightarrow C)]}$$

0.96	2.97

Figure 5