Role of sortase-assembled Ebp pili in *Enterococcus faecalis* adhesion to iron oxides and its

impact on extracellular electron transfer

Foo Kiong Ho[1,*], Ling Ning Lam[1,2,3,*], Artur Matysik[1,*], Thomas Dean Watts[1], Jun Jie Wong[1],

Kelvin Kian Long Chong[1], Pei Yi Choo[1], Joe Tolar[4], Pui Man Low[1], Zhi Sheng Chua[1], Jason J.

Paxman[5], Begoña Heras[5], Enrico Marsili[1,^], Caroline M. Ajo-Franklin[4,6,7], Kimberly A.

Kline[1,2,8,**]

[1]Singapore Centre for Environmental Life Sciences Engineering, Nanyang Technological

University, Singapore, Singapore.

[2]School of Biological Sciences, Nanyang Technological University, Singapore, Singapore.

[3]Department of Oral Biology, College of Dentistry, University of Florida, Gainesville, Florida,

USA.

[4]Department of BioSciences, Biomolecular Engineering Rice University, Houston, Texas, USA.

[5]Department of Biochemistry and Chemistry, La Trobe Institute for Molecular Science, La

Trobe University, Bundoora, Melbourne, Australia.

[6]Department of Bioengineering, Rice University, Houston, Texas, USA.

[7]Department of Chemical and Biomolecular Engineering, Rice University, Houston, Texas,

USA.

[8]Department of Microbiology and Molecular Medicine, University of Geneva, Switzerland.

*These authors contributed equally to this work

^Current address: Nottingham Ningbo China Beacons of Excellence Research and Innovation

Institute, University of Nottingham Ningbo, China.

**Correspondence:

Kimberly A. Kline

kimberly.kline@unige.ch

Abstract (214 words)

Enterococcus faecalis sortase-assembled endocarditis and biofilm-associated pili (Ebp) are a virulence factor implicated in enterococcal biofilm-associated infections and gastrointestinal colonization. We previously showed that *E. faecalis* biofilm metabolism is influenced by extracellular electron transfer (EET) under iron-rich conditions, raising the question of whether Ebp pili also play a role in EET. Here, we report a novel role of Ebp pili in *E. faecalis* adhesion to the iron oxides magnetite, goethite and hematite, where the EbpA tip adhesin contributes to this interaction. Adhesion by Ebp pili is conditionally important for EET to iron oxides, as pilus mutants are attenuated in EET under non-static growth conditions. In alignment with the established role of EET in redox homeostasis, we find that EET to ferricyanide supports *E. faecalis* anaerobic growth on glycerol. Further, in an antibiotic-treated mouse gastrointestinal colonization model, we show that *E. faecalis* mutants deficient in EET poorly colonize the intestinal niche. Taken together, our findings suggest that Ebp pili can influence *E. faecalis* metabolic fitness by promoting EET to iron oxides, raising new questions of how Ebp pili shape *E. faecalis* interactions with environmental ecosystems. Furthermore, the important role of EET in *E. faecalis* colonization of the dysbiotic gastrointestinal environment highlights the need for further inquiry into how EET contributes to *E. faecalis* microbial pathogenesis.

Importance (133 words)

In this study, we explored the interplay between extracellular electron transfer (EET) and an *Enterococcus faecalis* biofilm factor, the endocarditis and biofilm-associated pili (Ebp). We demonstrate that Ebp pili have a novel role in adhesion to iron oxides, which consequently promotes EET to iron oxides under non-static conditions. Along with our findings that *E. faecalis* EET can be coupled to anaerobic cell growth, our results point to a potential ecological role of Ebp pili in natural environments, outside of its established function in adhesion to host ligands. We provide the first evidence of the contribution of EET to *E. faecalis* colonization of the antibiotic-treated murine intestinal niche, which adds to the limited experimental evidence linking EET and microbial pathogenesis, as well as highlights the need for further studies of EET in bacterial pathogens.

Keywords

Enterococcus faecalis; pili; biofilm; iron oxides; metal reduction; extracellular electron

transfer; gastrointestinal colonization

Observation

Extracellular electron transfer (EET) is a process in which microorganisms reduce

extracellular substrates, such as iron, and is frequently connected to anaerobic energy

metabolism (1, 2). While EET has been extensively reviewed in the Gram-negative model

species *Geobacter sulfurreducens* and *Shewanella oneidensis* (3-5), the discovery of a flavin-

based EET genetic locus in Gram-positive bacteria, including pathogens and members of the

gut microbiota, raises the question of how microbial EET can influence human health or

disease (6, 7). The gut commensal and opportunistic pathogen *Enterococcus faecalis* also

performs EET (6, 8-11), but the physiological significance of EET in *E. faecalis* is not well-

characterized. As our previous observations showed a relationship between *E. faecalis*

biofilm metabolism and EET (8), this study focuses on investigating whether endocarditis

and biofilm-associated pili (Ebp), an essential factor in *E. faecalis* biofilm formation, play a

role in mediating EET. Given that EET has a prominent impact on the energy metabolism of

another lactic acid bacterium *Lactiplantibacillus plantarum* (12, 13), we also sought to

explore the importance of EET to *E. faecalis* using both *in vitro* and animal models.

Preliminary microscopy observations of *E. faecalis* grown in culture media autoclaved with

ferric chloride showed some cells tightly surrounded by a dense material (**Fig. S1A**). This

material might have been insoluble iron precipitates, inadvertently produced by

autoclaving, as it was reminiscent of iron deposits previously observed in the extracellular

matrix of *E. faecalis* biofilms grown under similar conditions (8). Immunofluorescence

staining showed co-localization of the dense material with Ebp pili in the parent strain, while

the dense material did not associate with the pilus-null Δ*ebpABC* mutant (**Fig. S1A**). As Ebp

pili are associated with adhesion to fibrinogen, collagen and platelets (14, 15), as well as

attachment to abiotic surfaces such as polystyrene (16), we hypothesized that there could

be a similar interaction between Ebp pili and insoluble iron forms. Hence, we assessed the

interaction between Ebp pili and iron by co-incubating *E. faecalis* with the iron oxide

magnetite, enabling the convenient separation of iron-adherent and iron-non-adherent

bacteria using a magnet (17) (**Fig. 1B**).

Heterogenous expression of Ebp pili in *E. faecalis* OG1RF produces subpopulations of

piliated and non-piliated cells (14-16, 18). Through immunofluorescence microscopy and

flow cytometry, we observed that the pilus-expressing cell population selectively adheres to

magnetite (**Fig. 1A**; **Fig. S1B**). The Δ*ebpABC* mutant showed significantly less adherence to

magnetite compared to the parental wild-type strain (**Fig. 1B**; **Fig. S2**). Using the EbpA^{AWAGA}

mutant, which has a mutated metal ion-dependent adhesion site (MIDAS) motif to disrupt

EbpA function without perturbing pilus biogenesis (19), we found that the tip adhesin EbpA

contributes to magnetite binding (**Fig. S2C**). Magnetite binding was also attenuated in the

single deletion mutants of EbpA and major fiber pilin EbpC, while no significant differences

were observed for the deletion mutant in cell wall anchor pilin EbpB (**Fig. S2C**). Findings

from single deletion mutants overall correlate with previously published biofilm formation

phenotypes due to defective pilus biogenesis (20). We were also interested in investigating

whether the adhesion phenotype was similar for goethite and hematite, both of which are

amongst the most common iron oxides found in soils (21). To test for adhesion to other iron

oxides, we separated non-adherent bacteria from mineral-bound bacteria using differential

density centrifugation with a sucrose solution and similarly observed adhesion to goethite

and hematite in a pilus-dependent manner (**Fig. S2D**).

To determine if the adhesin function of Ebp pili influences EET to iron oxides, bacteria were

grown in the presence of iron oxides and the amount of ferrous iron produced and released

into the supernatant was quantified by the ferrozine assay. As the control, we used an EET-

deficient mutant containing an insertional inactivation of the *ndh3* gene. *E. faecalis* Ndh3 is

a NADH dehydrogenase that had previously been characterized to be a component of the

electron transport chain involved in EET, where it oxidizes NADH and transfers electrons to

demethylmenaquinone (6, 9). Ndh3 is distinct from Ndh2, which is likely the NADH

dehydrogenase specific for aerobic respiration (9). Significant differences in iron reduction

were observed between parental and pilus mutant strains when bacterial growth was

conducted under non-static conditions, while these differences were absent or less distinct

under static growth conditions (**Fig. 1C**). Due to the heme auxotrophy of *E. faecalis* and the

absence of heme in the culture media, observations made here were likely not influenced

by the effects of aerobic respiration. Collectively, our findings indicate that Ebp pili play an

accessory role in enhancing EET through the aggregation of *E. faecalis* cells with iron oxides.

Though we did not differentiate between direct and indirect electron transfer, the results

suggest that close contact between bacteria and iron oxides is important for efficient

electron transfer. It is also tempting to speculate that the binding of Ebp pili to iron oxides is

an evolutionary trait that augments bacterial EET in environments, such as soils and

sediments, where iron oxides are commonly present. Ferrous iron produced by EET might

additionally modulate pilus function by competition with or displacement of the native

divalent cation in the MIDAS motif of EbpA (**Fig. S3**).

Previous work in *Listeria monocytogenes* demonstrated that the electron transport chain

involved in EET can route electrons not only to extracellular iron, but also through

extracellular reductases such as fumarate reductase (22). As anaerobic glycerol dissimilation

in *E. faecalis* is dependent on fumarate reductase (23), we predicted that EET could provide

an alternative route to external electron acceptors in place of fumarate. To test if the

membrane-impermeable ferricyanide could act as the electron acceptor for EET during

glycerol dissimilation, we grew *E. faecalis* macrocolonies anaerobically on an agar base

supplemented with ferricyanide. To limit excess fermentation of amino acids, growth

medium was excluded from the agar base and instead, the bacterial inoculum was

resuspended in growth medium prior to spotting on agar plates. We find that the presence

of either fumarate or ferricyanide promotes *E. faecalis* growth on glycerol, whereas growth

was deficient in the absence of an electron acceptor or when the EET-deficient *ndh3*::tn

mutant was used (**Fig. 2A**). Our findings confirmed that *E. faecalis* EET can be coupled to

anaerobic cell growth, which is consistent with metabolic and growth phenotypes

implicated in EET for *L. monocytogenes* and *L. plantarum* (6, 13). It would be particularly

interesting to make further comparisons against *L. plantarum*, as it is a lactic acid bacterium

similar to *E. faecalis* and has been shown to combine features of fermentation and

respiration in a hybrid form of metabolism (13). As EET also promoted *L. monocytogenes*

colonization of the mouse gastrointestinal tract (6), we investigated whether *E. faecalis* EET

plays a similar role. As anticipated, lower numbers of *ndh3*::tn mutant were recovered in an

antibiotic-treated mouse gastrointestinal model of colonization when compared to the

parent strain (**Fig. 2B**).

In summary, we describe a novel role of *E. faecalis* Ebp pili in adherence to iron oxides,

which may promote EET and enhance metabolic fitness in environmental ecosystems where

iron oxides are present (**Fig. 2C**). While we show that EET can be coupled to anaerobic

growth of *E. faecalis*, it is possible that EET may play additional roles, such as in iron uptake

or detoxification (24, 25). Iron availability is a strong driver for colonization of the

gastrointestinal tract by pathogens (26), but it is still unclear how Ebp pili will interact with

different iron forms present in the gastrointestinal tract. Nevertheless, we demonstrate that

mutants deficient in EET poorly colonize the mouse gastrointestinal tract, indicating that EET

contributes to *E. faecalis* outgrowth during gut dysbiosis.

Acknowledgments

This work was supported by the National Research Foundation and Ministry of Education

Singapore under its Research Centre of Excellence Programme and by the Ministry of

Education Singapore under its Tier 2 programme (MOE2014-T2-2-124). Foo Kiong Ho and

Artur Matysik were supported by Singapore Ministry of Health's National Medical Research

Council under its Open Fund Individual Research Grant (MOH-000645) and its Clinical Basic

Research Grant (NMRC/CBRG/0086/2015), respectively. Begoña Heras and Jason Paxman

were supported by the Australian Research Council (ARC) project grant (DP210100673) and

the National Health and Medical Research Council (NHMRC) project grant (GNT1143638).

We thank Hailyn V. Nielsen and Scott Hultgren (Washington University, St Louis, USA) for

providing the pilus mutants, and Gary Dunny (University of Minnesota, USA) for providing

transposon mutants. We also thank Bala Davient and Caroline Manzano for their assistance

with mutant construction.

Data Availability

All data generated or analysed during this study are included in this published article (and its

supplementary information files).

Conflict of Interest

The authors declare no conflict of interest.

References

1. Stevens E, Marco ML. Bacterial extracellular electron transfer in plant and animal

ecosystems. FEMS Microbiol Rev. 2023;47(3).

2. Shi L, Dong H, Reguera G, Beyenal H, Lu A, Liu J, et al. Extracellular electron transfer

mechanisms between microorganisms and minerals. Nat Rev Microbiol. 2016;14(10):651-62.

3. Hu Y, Wang Y, Han X, Shan Y, Li F, Shi L. Biofilm Biology and Engineering of Geobacter

and Shewanella spp. for Energy Applications. Front Bioeng Biotechnol. 2021;9:786416.

4. Zou L, Zhu F, Long ZE, Huang Y. Bacterial extracellular electron transfer: a powerful

route to the green biosynthesis of inorganic nanomaterials for multifunctional applications. J

Nanobiotechnology. 2021;19(1):120.

5. Shi L, Squier TC, Zachara JM, Fredrickson JK. Respiration of metal (hydr)oxides by

Shewanella and Geobacter: a key role for multihaem c-type cytochromes. Mol Microbiol.

2007;65(1):12-20.

6. Light SH, Su L, Rivera-Lugo R, Cornejo JA, Louie A, Iavarone AT, et al. A flavin-based

extracellular electron transfer mechanism in diverse Gram-positive bacteria. Nature.

2018;562(7725):140-4.

7. Wang W, Du Y, Yang S, Du X, Li M, Lin B, et al. Bacterial Extracellular Electron

Transfer Occurs in Mammalian Gut. Anal Chem. 2019;91(19):12138-41.

8. Keogh D, Lam LN, Doyle LE, Matysik A, Pavagadhi S, Umashankar S, et al.

Extracellular Electron Transfer Powers Enterococcus faecalis Biofilm Metabolism. mBio.

2018;9(2).

9.   Hederstedt L, Gorton L, Pankratova G. Two Routes for Extracellular Electron Transfer

in Enterococcus faecalis. J Bacteriol. 2020;202(7).

10.   Pankratova G, Leech D, Gorton L, Hederstedt L. Extracellular Electron Transfer by the

Gram-Positive Bacterium Enterococcus faecalis. Biochemistry. 2018;57(30):4597-603.

11.   Zhang E, Cai Y, Luo Y, Piao Z. Riboflavin-shuttled extracellular electron transfer from

Enterococcus faecalis to electrodes in microbial fuel cells. Can J Microbiol. 2014;60(11):753-

9.

12.   Tolar JG, Li S, Ajo-Franklin CM. The Differing Roles of Flavins and Quinones in

Extracellular Electron Transfer in Lactiplantibacillus plantarum. Appl Environ Microbiol.

2023;89(1):e0131322.

13.   Tejedor-Sanz S, Stevens ET, Li S, Finnegan P, Nelson J, Knoesen A, et al. Extracellular

electron transfer increases fermentation in lactic acid bacteria via a hybrid metabolism.

Elife. 2022;11.

14.   Nallapareddy SR, Singh KV, Sillanpää J, Zhao M, Murray BE. Relative contributions of

Ebp Pili and the collagen adhesin ace to host extracellular matrix protein adherence and

experimental urinary tract infection by Enterococcus faecalis OG1RF. Infect Immun.

2011;79(7):2901-10.

15.   Nallapareddy SR, Sillanpää J, Mitchell J, Singh KV, Chowdhury SA, Weinstock GM, et

al. Conservation of Ebp-type pilus genes among Enterococci and demonstration of their role

in adherence of Enterococcus faecalis to human platelets. Infect Immun. 2011;79(7):2911-

20.

16.   Nallapareddy SR, Singh KV, Sillanpää J, Garsin DA, Höök M, Erlandsen SL, et al.

Endocarditis and biofilm-associated pili of Enterococcus faecalis. J Clin Invest.

2006;116(10):2799-807.

17.   Brown S. Engineered iron oxide-adhesion mutants of the Escherichia coli phage

lambda receptor. Proc Natl Acad Sci U S A. 1992;89(18):8651-5.

18.   Tariq M, Bruijs C, Kok J, Krom BP. Link between culture zeta potential homogeneity

and Ebp in Enterococcus faecalis. Appl Environ Microbiol. 2012;78(7):2282-8.

19.   Nielsen HV, Guiton PS, Kline KA, Port GC, Pinkner JS, Neiers F, et al. The metal ion-

dependent adhesion site motif of the Enterococcus faecalis EbpA pilin mediates pilus

function in catheter-associated urinary tract infection. MBio. 2012;3(4):e00177-12.

20.   Sillanpää J, Chang C, Singh KV, Montealegre MC, Nallapareddy SR, Harvey BR, et al.

Contribution of individual Ebp Pilus subunits of Enterococcus faecalis OG1RF to pilus

biogenesis, biofilm formation and urinary tract infection. PLoS One. 2013;8(7):e68813.

21.   Cornell RM, Schwertmann U. The iron oxides: structure, properties, reactions,

occurrences, and uses. 2nd ed. Weinheim: Wiley-VCH; 2003.

22.   Light SH, Méheust R, Ferrell JL, Cho J, Deng D, Agostoni M, et al. Extracellular

electron transfer powers flavinylated extracellular reductases in Gram-positive bacteria.

Proc Natl Acad Sci U S A. 2019;116(52):26892-9.

23.   Bizzini A, Zhao C, Budin-Verneuil A, Sauvageot N, Giard JC, Auffray Y, et al. Glycerol is

metabolized in a complex and strain-dependent manner in Enterococcus faecalis. J

Bacteriol. 2010;192(3):779-85.

24.   Jeuken LJC, Hards K, Nakatani Y. Extracellular Electron Transfer: Respiratory or

Nutrient Homeostasis? J Bacteriol. 2020;202(7).

25.   Feng H, Xu L, Chen R, Ma X, Qiao H, Zhao N, et al. Detoxification mechanisms of

electroactive microorganisms under toxicity stress: A review. Front Microbiol.

2022;13:1084530.

26.   Kortman GA, Raffatellu M, Swinkels DW, Tjalsma H. Nutritional iron turned inside

out: intestinal stress from a gut microbial perspective. FEMS Microbiol Rev.

2014;38(6):1202-34.

Figure 1. Ebp pilus expression is important for *E. faecalis* adhesion and EET to iron oxides.

(A) Flow cytometry and immunofluorescence microscopy images of *E. faecalis* cells after

magnetic separation using magnetite. Bacterial suspensions of *E. faecalis* OG1RF (left

panels) were mixed with magnetite and then separated into magnetite-bound (middle

panels) and non-bound fractions (right panels) using a magnetic rack. For flow cytometric

analysis, the magnetite-bound bacterial fraction was released from magnetite by

resuspension in 1×PBS containing 100 mM EDTA prior to staining. *E. faecalis* cells were

stained with SYTO 9 DNA stain (green) and labelled for EbpC (red). Histograms show

representative data from three independent experiments. Scale bars represent 20 µm. **(B)**

Schematic diagram (left panel) illustrating how magnetic separation was used to isolate

adherent and non-adherent fractions of bacteria. Culture turbidity (right panel) of non-

adherent bacterial subpopulations after assaying with increasing concentrations of

magnetite. Data points reflect optical density values at 600 nm (OD_{600}) of the non-adherent

fraction, after a magnet was used to remove magnetite and magnetite-bound bacteria.

Horizontal lines represent the mean measurement of three independent experiments and

statistical comparisons represent differences between WT and Δ*ebpABC* at each magnetite

concentration. *$P < 0.05$, **$P < 0.01$ by unpaired Student's *t*-test. **(C)** Iron oxide reduction by

Ebp pilus mutants as assessed by the ferrozine assay under non-static (left panel) or static

(right panel) growth conditions. Horizontal lines represent the mean measurement of three

independent experiments. Statistical comparisons represent differences between WT and

the corresponding pilus mutant. **$P < 0.01$, ***$P < 0.001$ by one-way ANOVA with Tukey's

multiple comparisons.

Figure 2. *E. faecalis* EET contributes to optimal fitness *in vitro* and *in vivo*. (A) Anaerobic

macrocolony growth with glycerol as the carbon source in the presence of either 2.5 mM

fumarate or 2.5 mM ferricyanide as the electron acceptor. After 24-hour growth in an

anaerobic jar, macrocolonies were excised from the agar and CFU enumerated. Data shown

are from five independent experiments and each dot represents the average of three

technical replicates. Horizontal lines represent the median CFU. Statistical comparisons

were assessed by one-way ANOVA with Tukey's multiple comparisons. **$P < 0.01$, ***$P <$

0.001, ns non-significant. **(B)** *E. faecalis* colonization of lower gastrointestinal tract using an

antibiotic-treated mice model of gut colonization. Datapoints represent individual mice and

the dotted line indicates the limit of detection at CFU <40. Data shown are from four

independent experiments with 5 mice per group in each experiment. Horizontal lines

represent the median CFU. Statistical comparisons were assessed by the Mann-Whitney

test. **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$, ns non-significant. **(C)** Proposed model for

accessory role of Ebp pili in promoting adhesion and EET to iron oxides. Efficient EET to iron

oxides under non-static conditions is mediated by Ebp pili, which has a novel role in

adhesion to magnetite, goethite and hematite. Ebp pili are not essential for EET, but

adhesion promotes close contact with iron oxides for more efficient EET. The tip pilin EbpA,

which is responsible for adhesion to host collagen and fibrinogen, contributes to the

adhesion to iron oxides but it is not known if the binding mechanism is similar for these

substrates. EET contributes to the maintenance of redox homeostasis (NAD^+/NADH

balance), which is known to be important for various metabolic processes such as the

anaerobic glycerol dissimilation pathway in *E. faecalis*. While lactate fermentation is the

primary mode of NAD^+ regeneration in *E. faecalis*, experiments in another lactic acid

bacterium *L. plantarum* have demonstrated that EET supplements fermentative pathways to

increase metabolic flux and yield. Lastly, the physiological consequence of EET to iron oxides

would be highly dependent on the thermodynamic favorability of electron transfer in these

environments, but a possibility is that EET serves as a strategy for *E. faecalis* to maintain

competitive fitness by expanding the range of electron acceptors that it is able to utilize.

Components are not drawn to scale. Figure is created using BioRender.