Sortase-assembled pili promote extracellular electron transfer and iron acquisition in

***Enterococcus faecalis* biofilm**

Ling Ning Lam[1,2,7], Jun Jie Wong[1,3], Artur Matysik[1], Jason J. Paxman[4], Kelvin Kian Long

Chong[1,5], Pui Man Low[1], Zhi Sheng Chua[1], Begoña Heras[4], Enrico Marsili[1,6] and Kimberly

A. Kline[1,2,*]

[1]Singapore Centre for Environmental Life Science Engineering, Nanyang Technological

University, 60 Nanyang Drive, Singapore 637551

[2]School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive,

Singapore 637551

[3]Interdisciplinary Graduate School, Nanyang Technological University, Singapore 639798

[4]Department of Biochemistry and Genetics, La Trobe Institute for Molecular Science, La

Trobe University, Science Drive, Bundoora VIC 3083, Melbourne, Australia

[5]Nanyang Technological University Institute for Health Technologies, Interdisciplinary

Graduate School, Nanyang Technological University, 50 Nanyang Drive, Singapore 637553

[6]Department of Chemical and Materials Engineering, Nazarbayev University, 53 Kabanbay

Batryr Avenue, Nur-Sultan 010000, Kazakhstan

[7]Department of Oral Biology, College of Dentistry, University of Florida, 1395 Center Drive,

Gainesville, FL 32610, United States

*Correspondence: kkline@ntu.edu.sg

Keywords: *Enterococcus faecalis* biofilm, iron acquisition, endocarditis and biofilm

associated pili, extracellular electron transfer, virulence

Classification: Biological Sciences, Microbiology

Abstract

Enterococcus faecalis is an opportunistic human pathogen and the cause of biofilm-associated infections of the heart, catheterized urinary tract, wounds, and the dysbiotic gut where it can expand to high numbers upon microbiome perturbations. The *E. faecalis* sortase-assembled endocarditis and biofilm associated pilus (Ebp) is involved in adhesion and biofilm formation *in vitro* and *in vivo*. Extracellular electron transfer (EET) also promotes *E. faecalis* biofilm formation in iron-rich environments, however neither the mechanism underlying EET nor its role in virulence was previously known. Here we show that iron associated with Ebp serve as a terminal electron acceptor for EET, leading to extracellular iron reduction and intracellular iron accumulation. We found that a MIDAS motif within the EbpA tip adhesin is required for interaction with iron, EET, and FeoB-mediated iron uptake. We demonstrate that MenB and Ndh3, essential components of the aerobic respiratory chain and a specialized flavin-mediated electron transport chain, respectively, are required for iron-mediated EET. In addition, using a mouse gastrointestinal (GI) colonization model, we show that EET is essential for colonization of the GI tract, and Ebp is essential for augmented *E. faecalis* GI colonization when dietary iron is in excess. Taken together, our findings show that pilus mediated capture of iron within biofilms enables EET-mediated iron acquisition in *E. faecalis*, and that these processes plays an important role in *E. faecalis* expansion in the GI tract.

Significance

Understanding enterococcal biofilm development is the first step towards improved therapeutics for the often antimicrobial resistant infections caused by these bacteria. Here we report a role for *Enterococcus faecalis* endocarditis and biofilm associated pili (Ebp) in mediating iron-dependent biofilm growth and contributing to extracellular electron transfer

(EET) which in turn promotes iron acquisition. Furthermore, we characterize the mechanisms

underlying electron transfer in the *E. faecalis* biofilm. Our findings support a model in which

E. faecalis use EET to drive the reduction of pilus-associated ferric iron, leading to iron

acquisition in *E. faecalis* biofilm, and contributing to enterococcal virulence in the GI tract.

Introduction

Enterococcus faecalis is an important human opportunistic pathogen that causes a variety of

diseases including endocarditis, urinary tract infections (UTI), bacteremia, wound infection,

and medical device-associated infections [1]. Many of these infections are polymicrobial and

biofilm-associated, rendering them more tolerant to antimicrobial and immune clearance, and

contributing to their persistent nature [1]. Therefore, a detailed understanding of enterococcal

biofilm development is a critical step towards advancing new therapeutics for treatment of

these often antimicrobial resistant infections [1,2].

One important factor contributing to *E. faecalis* biofilm formation and virulence is the

sortase-assembled endocarditis and biofilm associated pilus (Ebp) [3-7]. The importance of Ebp

to *E. faecalis* is supported by the presence of its coding sequence in the core *E. faecalis*

genome [5,8]. The Ebp fiber is primarily composed of the major pilin subunit EbpC, along with

two dispensable minor pilin subunits EbpB and EbpA [5], and these pilin subunits are

covalently assembled by Sortase C on the cell membrane, prior to attachment to the cell wall

by Sortase A [6,9]. EbpA is the tip adhesin of the pilus, and its N terminus encompasses a metal

ion-dependent adhesion site (MIDAS) motif. In eukaryotes, the MIDAS motif is important

for ligand binding [10] and coordinates divalent cations, most often Mg^{2+} [11,12], for cell adhesion

and interaction with extracellular matrix (ECM) proteins [12]. The MIDAS motif within the N-

terminal domain of *E. faecalis* EbpA contributes to *in vitro* biofilm formation, adherence to

fibrinogen, and bladder colonization in a mouse catheter-associated urinary tract infection

(CAUTI) model [13-15]. Sortase-assembled pili are conserved in many Gram-positive bacteria,

where they often contribute to adhesion, biofilm formation, modulation of host immune

response, and virulence [16,17]. The adhesive tips of these pilins (PilA of *Streptococcus*

agalactiae [18], RrgA of *Streptococcus pneumoniae* [19], and AP-1 of *Streptococcus pyogenes* [20])

include a conserved von Willebrand factor A (vWFA) domain containing a MIDAS motif,

which are essential for adhesion. However, the specificity and affinity for metal binding to

the MIDAS motif in bacteria, including *E. faecalis,* has not been characterized.

Previously, we demonstrated that *E. faecalis* is electrogenic, and that iron promotes both

extracellular electron transfer (EET) and biofilm growth in *E. faecalis* [21]. Several mechanisms

for EET have been described in both Gram-positive and Gram-negative bacteria [22,23]. In some

bacteria, microbial nanowires, composed of extracellular filamentous protein fibers and/or of

cytochrome-associated fibers, have been proposed to aid EET by directly connecting the

bacterial cell surface to extracellular metal oxides, or alternatively, by interacting with

extracellular soluble redox mediators [23,24]. Biochemical and structural analysis have recently

demonstrated that the conductive filaments of *Geobacter sulfurreducens* consist of

polymerized chains of the hexaheme cytochrome OmcS [25]. Gram positive bacteria have

evolved a highly conserved mechanism for EET, described in *L. monocytogenes*, which

involves a specialized respiratory complex that channels electrons through discrete

membrane-localized quinone pools to flavin intermediates, for subsequent delivery to

terminal extracellular electron acceptors [26]. In *E. faecalis,* electroactivity requires the ability

to synthesize L-lactate dehydrogenase (LDH) [21] and demethylmenaquinone (DMK) [27],

important for catalyzing redox reactions and electron transfer during respiration, respectively

[28].

In this study, we sought to understand the mechanism and physiological role of EET in *E.*

faecalis. We show that EET in *E. faecalis* makes use of both DMK and the conserved flavin

EET pathway described in *L. monocytogenes* for electron delivery to extracellular pilus-

associated iron deposits. Iron induces the expression of Ebp pili, increasing the interaction

sites for iron, which include the MIDAS motif of the EbpA tip adhesin, which is required for

pilus association with iron as well as for iron-augmented biofilm formation. In addition, we

show that EET drives the reduction of iron leading to FeoB-mediated uptake of iron into the

E. faecalis biofilm cells. Finally, we demonstrate that a high iron diet promotes *E. faecalis*

mouse gastrointestinal (GI) colonization in an *ebp*-dependent manner, and EET mutants are

attenuated in the GI tract. Together, these findings suggest a model in which iron promotes

pilus-mediated biofilm formation. We propose that the association of iron with sortase-

assembled Ebp sequesters iron close to the cell surface within biofilms where it can serve as

an extracellular electron acceptor for EET, resulting in the generation of ferrous iron for

subsequent uptake into the cell. This new paradigm for iron acquisition in which pili and

aggregates of piliated cells serve as iron sinks to promote biofilm formation and intracellular

iron accumulation, significantly advances our mechanistic understanding of both biofilm

formation and the pathogenesis of *E. faecalis*.

Results

Iron-augmented *E. faecalis* biofilm formation requires pilus expression

We previously demonstrated that *E. faecalis* biofilm growth in media supplemented with iron

resulted in two-fold more biomass [21]. To test the hypothesis that Ebp contributes to iron-

augmented biofilm, we performed biofilm assays in growth media supplemented with 2 mM

ferric chloride ($FeCl_3$), and quantified total adherent biofilm biomass. In iron-supplemented

media, deletion mutants of the entire Ebp pilus (Δ*ebpABC)*, pilus tip (Δ*ebpA*) and pilus fiber

(Δ*ebpC*) displayed more than a 50% reduction in biofilm growth compared to the wild type

control **(Figure 1a)** and mutant biofilm levels in iron-supplemented media were most similar

to wild type biofilm in the absence of iron augmentation **(Figure S1a)**. Biofilm formation by

the pilus null Δ*ebpABC* mutant was restored to wild type levels upon genetic

complementation with the full locus (p*ebpABCsrtC*). A Δ*ebpB* deletion mutant also displayed

significantly reduced biofilm growth as compared to the wild type control in iron-

supplemented media; however, it was not as strongly attenuated compared to the other pilus

mutants **(Figure 1a)**. These findings are consistent with previous reports in which deletion of

ebpA and *ebpABC* resulted in significantly reduced biofilm formation in normal media

(Figure S1a, S2b) [6,29]. These results indicate that the fully assembled Ebp pilus fiber

contributes substantially to iron-augmented biofilm formation with a key role for the EbpA

pilus tip **(Figure 1a, S1b)**. We next tested the contribution of the MIDAS motif within EbpA

to iron-mediated biofilm growth. Using a MIDAS mutant (*ebpA^{AWAGA}*), in which the native

motif (Asp^{275}-Trp-Ser^{277}-Gly-Ser^{279}) is mutated to (Ala^{275}-Trp-Ala^{277}-Gly-Ala^{279}) [13], we

observed that it was as attenuated as the *ebpABC* null mutant for biofilm formation in iron-

supplemented media **(Figure 1a, S1a)**. Collectively, these data demonstrate that both the

MIDAS motif of the EbpA tip adhesin, as well as an intact EbpC pilus fiber, are essential for

iron-mediated biofilm growth.

Iron induces pilus expression and drives aggregation of pilus-expressing cells in *E.*

***faecalis* biofilm**

Iron-augmented *E. faecalis* biofilms have an altered 3-D structure [21]. To determine whether

pilus expression contributes to the biofilm ultrastructure, we used immunofluorescence

microscopy to visualize pilus-expressing cells within biofilms. Augmented biofilm biomass

after growth in iron-supplemented media, as previously reported, was not apparent for the

$\Delta ebpABC$ or the $ebpA^{AWAGA}$ mutants **(Figure 1b)** [21]. In addition, pilus-expressing cells tended

to aggregate in areas of higher cell density in biofilms grown in iron-supplemented media

(Figure 1c). The abundance of piliated cell aggregates within the biofilm suggested that

pilus expression may be induced in the presence of iron. We therefore quantified the number

of biofilm cells expressing pili and observed a significantly larger pilus-expressing

population in biofilms grown in ferric iron-supplemented media, but not in the presence of

alternative iron sources or with other cationic metals **(Figure 1d)**. By contrast, we observed

no significant difference in planktonically grown pilus-expression in response to iron **(Figure**

1d). Together these results show that ferric iron induces pilus expression, which in turn

increases the biofilm structure and mass.

Iron colocalizes with pili on the cell surface and is dependent on the EbpA MIDAS motif

While monitoring *E. faecalis* populations for piliation in iron-supplemented media, we

observed dense deposits at the polar hemispheres of *E. faecalis* cells which co-localized with

sites of EbpC deposition, suggesting that these dense deposits may be iron interacting with

pili **(Figure 2a, Figure S2a-b)**. We observed these dense deposits less frequently in the

$ebpA^{AWAGA}$ mutant, even though this mutant strain expresses a similar proportion of piliated

cells as WT **(Figure S2c),** and the dense deposits were completely absent in $\Delta ebpABC$

deletion mutant **(Figure 2a, Figure S2a-b)**, suggesting that pili may be interacting with ferric

iron via the MIDAS motif of EbpA.

To confirm the interaction between the EbpA MIDAS motif and iron, we extracted Ebp from

E. faecalis wild type and $ebpA^{AWAGA}$ biofilms grown in normal or iron supplemented media,

and in parallel performed control extractions from $\Delta ebpABC$ biofilms. Pili were isolated

under native conditions and analyzed for iron content by inductive-coupled plasma mass

spectrometry (ICP-MS). Pilus extracts from wild type cells grown in the iron-supplemented

media were associated with nearly three times more iron than wild type pilus extracts from

biofilms grown in control media. Ebp association with iron was dependent on the EbpA

MIDAS motif because pilus preps from *ebpA^AWAGA* were similar to background iron levels in

control Δ*ebpABC* extracts **(Figure 2b)**.

To better understand how the EbpA MIDAS motif could interact with ferric iron, we

modelled the EbpA structure based upon the crystal structures of its Mg bound homologs, *S.*

agalactiae PilA GBS104 (PDB: 3txa) [30] and *S. pneumoniae* RrgA (PDB: 2ww8) [19]. The

overall EbpA structural model (N177-P620) showed the expected vWFA structural fold

encompassing the MIDAS motif, consistent with other pili tip adhesins **(Figure 2c-d)**. The

model supports an interaction of the MIDAS motif and associated metal binding residues

with ferric iron **(Figure 2e)**. D275 would make a water mediated contact with the iron, the

hydroxyl oxygen atoms of S277 and S279 and the carboxylate oxygen of D378 would make

direct ionic bonds with iron and the classic 6-coordination of iron would be completed by 2

water molecules. vWFA domains are known to undergo conformational changes from closed

to open states, whereby iron bonding would be shifted from D378 to the nearby T350 [31]. The

metal binding residues were confirmed by an alignment of the EbpA, GBS104 and RrgA

sequences, which despite low sequence identity, showed strict conservation of the MIDAS

motif D275, S277, S279, T350 and D378 residues **(Figure S3)**.

Pili and the respiration-associated electron transport chain contribute to extracellular

electron transfer (EET)

The association of Ebp with iron, coupled with the ability of *E. faecalis* to undergo iron-

mediated EET [21], suggested that pili themselves may be involved in the EET. To address this,

we measured the instantaneous current output by chronoamperometry and the cumulative

charge production in biofilms grown in iron-supplemented media for 20 hours. We observed

a reduction in cumulative charge production in both the $\Delta ebpABC$ and $ebpA^{AWAGA}$ MIDAS

mutants when grown in 0.5 mM iron-supplemented media **(Figure 3a)**. At higher iron

concentrations, we did not observe significant differences in cumulative charge production

over 20 hours **(Figure S4a)**. The current output in the first few hours was similar for all

strains **(Figure S4b)**, but both pilin mutants exhibited an inability to sustain current output as

compared to the wild- type control at later time points **(Figure S4a-b)**. We therefore

conclude that pili contribute to sustained EET, but that other mechanisms also contribute to

E. faecalis EET in the absence of the pili.

To identify additional pathways, in addition to pili, that contribute to iron-mediated EET, we

examined the *E. faecalis* homolog of the *L. monocytogenes* EET-specific NADH

dehydrogenase (*ndh3*) for flavin-mediated EET [26], as well as demethylmenaquinone (*menB*)

for quinone-mediated EET [28]. We hypothesized that these electron transfer pathways may

work together to promote EET. Indeed, we observed attenuated charge production in both

ndh3::Tn and *menB::Tn* mutants [32,33] which was restored upon genetic complementation

(Figure 3b), as well as in mutants predicted to work in concert with Ndh3 in the conserved

EET pathway **(Figure S5)**. Importantly, charge production was further reduced in the

ndh3::Tn ΔmenB double mutant compared to either single mutant **(Figure 3b),** suggesting

that DMK and Ndh3 work in non-redundant concert for efficient EET**.** Together, these

findings demonstrate that both respiratory electron transport conduits and pili contribute to

iron-mediated EET in *E. faecalis* biofilm.

Genes encoding Ebp and the FeoB ferrous iron transporter are upregulated in *E. faecalis* biofilm

To further understand how iron promotes *E. faecalis* biofilm, we performed RNA sequencing and found a total of 90 genes that were differentially regulated upon growth in iron-supplemented media **(Table S1).** As expected, the pilus genes *ebpA* (OG1RF_RS04555) and *ebpB* (OG1RF_RS04560) were induced 1.68-fold and 1.55-fold respectively, which we validated by qRT-PCR **(Figure S6)**. Surprisingly, the only other iron-associated gene that showed both significant and differential expression was *feoB* (OG1RF_RS01950) encoding a predicted ferrous iron transporter, which was up-regulated 1.38-fold **(Table S1).** Importantly, this suggested that *E. faecalis* biofilm grown in iron-supplemented media may be importing ferrous iron. This finding was consistent with our earlier study that showed *E. faecalis* intracellular iron concentrations were increased in biofilm cells grown in iron-supplemented media [21]. The simultaneous up-regulation of *ebpABC* and *feoB* as the sole iron-responsive transporter, along with our observations that Ebp co-purify with iron and are required for EET suggested that pilus-associated iron may be reduced during EET and subsequently taken up by the ferrous iron transporter FeoB.

Extracellular ferric iron reduction replenishes ferrous iron pool for uptake by *E. faecalis* biofilm cells

Since ferric iron reduction to ferrous iron requires EET and is facilitated by both Ebp pili and the respiration electron transport chain, we predicted that an inability to perform EET would coincide with attenuation in ferric iron reduction and subsequent diminished extracellular ferrous iron pools available for uptake via FeoB. We therefore performed ICP-MS to determine the intracellular iron content in pilus mutants and EET-associated mutants grown in iron-supplemented media. Since Ldh1 was previously shown to be important for *E.*

faecalis EET [21], we predicted that the absence of *ldh1* may also lead to reduced intracellular

iron accumulation. Consistent with these predictions, we observed significantly reduced

intracellular iron for all pilus deletion mutants which was restored upon complementation, as

well as reduced intracellular iron in the *ebpA^{AWAGA}* MIDAS and the *ldh1* mutants.

Furthermore, we observed significantly reduced intracellular iron in the *ndh3::Tn ΔmenB*

mutant **(Figure 4a)**. As predicted, the absence of *feoB* also led to reduced intracellular iron,

in a genetically complementable manner **(Figure 4b)**.

If intracellular ferrous iron accumulation is a direct consequence of EET catalyzed ferric iron

reduction, EET mutants should display reduced ability for ferric iron reduction. We tested

this using a ferrozine assay, which reacts with ferrous iron to form a stable colored product.

With a functional EET mechanism, we predict the presence of increased ferrous iron pools.

As expected, we observed a significant decrease in ferrous iron pools in *ldh1, ndh3* and *menB*

mutants **(Figure 4c)**. Collectively, these data show that EET mutants are attenuated for ferric

iron reduction and therefore intracellular iron accumulation.

Pili and respiration-associated electron transfer systems contribute to colonization of

mouse GI tract

To understand how iron-augmented biofilms and EET impact *E. faecalis* pathogenesis, we

examined the role of dietary iron in an antibiotic-treated mouse gastrointestinal (GI)

colonization model. We fed mice a diet containing normal amounts of iron (control; 200

mg/kg ferric chloride) or excess iron (high iron; 2000 mg/kg ferric chloride), for 3 weeks

prior to antibiotic exposure and subsequent ingestion of *E. faecalis* in the drinking water

(Figure 5a). Twenty-four hours after inoculation, we recovered significantly more *E. faecalis*

from the mouse colon in mice fed the high iron diet as compared to the control group and this

increase was dependent on Ebp expression **(Figure 5b)**. Mice fed with high iron diet had

increased levels of colon tissue-associated iron, which was not apparent in the cecum or small

intestine **(Figure S7a)**. Accordingly, no iron-dependent colonization differences were

observed in those compartments **(Figure S7b)**. These data suggest that dietary iron promotes

E. faecalis Ebp-dependent colonization in the mouse colon. We next asked whether EET also

provides a colonization advantage for *E. faecalis* in the GI tract. Indeed, both Ndh3 and

MenB are required for GI colonization, independent of dietary manipulation **(Figure 5c,**

Figure S7c). Together, these findings demonstrate that both the pili and EET are important

for GI colonization.

Discussion

We previously showed that *E. faecalis* performs EET in the presence of extracellular iron,

which was associated with ATP production, and that iron supplementation augments biofilm

growth [21]. In this study, we sought to understand the mechanism underlying EET and its

physiological role in *E. faecalis*. We show that iron induces expression of the genes encoding

the sortase-assembled Ebp, which in turn associate with iron via the MIDAS motif of the

EbpA tip adhesin, and promote biofilm formation. We also show that EET is dependent on

both DMK in the electron transport chain, and a specialized flavin-associated respiratory

complex, leading to reduction of extracellular iron and FeoB-mediated uptake of iron (**Figure**

6). Furthermore, we demonstrate that pili are important for colonization in the iron-loaded

mouse GI tract.

Many microorganisms have the capacity to generate electrical current via dissimilatory metal

reduction [22,34,35]. In *E. faecalis*, it was previously known that EET relies upon DMK, part of

its minimal electron transport chain [27], as well as Ldh1, involved in cellular redox

homeostasis [36]. Here we confirm that *menB*, required for DMK synthesis [37], contributes to

EET, along with a recently reported flavin-associated alternative membrane-bound electron

transport chain [26]. Since mutation of either pathway results in decreased charge production

and ferric iron reduction, whereas mutation of both pathways completely abrogates both

activities, we suggest that electrons flow through both respiratory complexes for maximally

efficient EET and ferric iron reduction. While the precise mechanism for electron transfer

outside of the *E. faecalis* cell may vary depending on the environmental niche, like *L.*

monocytogenes [26], *E. faecalis* is a flavin auxotroph [38], necessitating the presence of this

soluble electron shuttle in the vicinity of the microbes. Indeed, addition of riboflavin to *E.*

faecalis biofilms growing in microbial fuel cells augments current production [39]. Our work

also demonstrates that *E. faecalis* can use iron as a terminal acceptor for EET, which is likely

relevant both in the outside environment where Enterococci can persist for long periods of

time [40], as well as in the GI tract where less than 15% of dietary iron is absorbed at the

duodenum, leaving the majority destined for excretion [41] and potentially available for EET. In

addition, there are likely alternative terminal electron acceptors available for EET in the GI

tract, such as host associated iron, host-derived nitrate, or microbially-derived molecules such

as humic substances [42].

We previously demonstrated that *E. faecalis* biofilms, but not planktonic cells, grow better

upon iron supplementation [21]. We further showed that iron deposits accumulate in the *E.*

faecalis biofilm matrix [21], which is composed of extracellular DNA [43,44] as well as, most

likely, sortase substrates including Ebp that are continuously shed from wild type cells [6,9].

Here we show that the same pili are required for the formation of biofilm matrix-associated

iron deposits, and are both essential for iron-augmented biofilm formation and necessary for

EET. We speculate that Ebp contribute to EET in *E. faecalis* by sequestering iron as terminal

electron acceptors in close proximity to the cells, either as surface attached pili or within the

tightly packed biofilm matrix, for efficient EET.

Both pilus associated EET and biofilm formation depend on an intact MIDAS motif within

the EbpA pilus tip adhesin of a fully polymerized pilus. Ebp also co-purifies with iron in a

MIDAS-dependent manner, demonstrating for the first time that the MIDAS motif

contributes to the association of a protein with iron. Indeed, our structural model of EbpA

aligns well with other pilus adhesin tips such as *S. agalactiae* GBS104 and *S. pneumoniae*

RrgA, with a vWFA fold and MIDAS motif configured for binding cationic metals [76,77].

MIDAS motifs have been shown to bind a range of cationic metals, but in the case of EbpA

this can include Fe^{3+} due to the local available concentrations in iron-rich environments.

Apart from binding metals, MIDAS motifs also form part of substrate binding sites, whereby

iron binding to this site could influence the interaction of EbpA with biofilm factors to

promote an increase in biofilm mass [78]. Collectively, our data confirm that iron binding at the

MIDAS motif has a dual role as both a terminal electron acceptor for EET and to promote

biofilm formation. Moreover, vWFA domains can undergo conformational changes to open

and closed states [78], so it is tempting to speculate that such a conformational change may

have a role in the interaction with biofilm promoting factors and/or the release of iron after its

reduction to Fe^{2+}.

Consistent with an important role for Ebp pili in an iron-rich environment, we show that the

ebp operon is transcriptionally induced when iron is supplemented. Ebp are displayed on the

surface of a subset of cells in any given *E. faecalis* population [5,6,9,29,45], and this population

can be increased in the presence of serum [5] and bicarbonate [46]. Here we add another

environmental signal, iron abundance, that can influence Ebp expression. Several intrinsic

factors can impact *ebp* gene expression including transcriptional regulators, EbpR [47], AhrC

(also annotated as ArgR3) and ArgR2 [48], the FsrB quorum sensing peptide [49] and the RNA

processing enzyme RNase J2 [50,51]. While none of these factors were differentially regulated

upon iron supplementation which may suggest uncharacterized regulatory pathways are

responsible for iron-regulated *ebp* transcription, we cannot rule out the possibility that excess

iron impacts the function of these known regulators of *ebp* expression.

In addition to *ebp*, the only iron-regulated *E. faecalis* gene annotated as iron-associated was

feoB, involved in ferrous iron uptake. Our discoveries that 1) Ebp expressing cells aggregate

within the biofilm, 2) *E. faecalis* accumulate intracellular iron in an Ebp-, EbpA MIDAS-,

and FeoB-dependent manner, and that 3) intracellular iron accumulation is abrogated in the

absence of EET and ability for ferric iron reduction, together suggest the presence of spatially

segregated areas within the biofilm that may be especially enriched for ferric iron

sequestration and reduction, subsequent ferrous iron acquisition, leading to biofilm growth

and restructuring. Our observation that *E. faecalis* can combine dissimilatory metal reduction

with iron acquisition is similar to early reports for *Shewanella putrefaciens* [52]. However, the

precise mechanism by which iron acquisition contributes to *E. faecalis* biofilm formation is

currently under investigation.

Enterococci are common but minor members of the GI microbiome. However, antibiotic

mediated dysbiosis favors overgrowth of Enterococci, harbouring both intrinsic and acquired

antibiotic resistance, in the lower GI tract [53-56]. In addition, their inherent tolerance to

oxidative stress [57] and high concentration of metals [21,58,59] can provide a selective advantage

for *E. faecalis* to colonize and bloom in niches unfavorable to other microbes. Our

observation that *E. faecalis* can both use EET and take advantage of iron-overload in the GI

tract provides yet another mechanism by which *E. faecalis* can thrive in this niche.

Collectively, the findings of this study demonstrate that Ebp and EET facilitate iron reduction

for its subsequent uptake, biofilm augmentation, and virulence in the antibiotic treated gut.

Given the strong conservation of sortase-assembled pili containing MIDAS motifs in their tip

adhesins among Gram positive pathogens, along with the conservation of genes encoding

EET components in the same organisms, we propose that pilus-mediated metal capture and

acquisition may be a common mechanism used by microbes to acquire limiting nutrients in a

diversity of niches. This work therefore raises the possibility for new therapeutic strategies

for *E. faecalis*, and potentially many other pathogens, aimed inhibiting extracellular electron

flow, iron reduction and acquisition, and EET-associated ATP production.

Materials and Methods

Bacterial Strains and Growth Conditions. *E. faecalis* was grown in Brain Heart Infusion

broth (BHI; Becton, Dickinson and Company, Franklin Lakes, NJ) and cultured at 37^oC

under static or shaking (200rpm) conditions, as indicated. Preparation of inocula for biofilm

and planktonic growth assays was performed as previously described following growth in

Tryptic Soy Broth (TSB) or agar, supplemented with 0.175% glucose (TSBG) (Oxoid Inc.,

Ontario, Canada) [21]. Bacterial strains used in this study are listed in **Table S2.** Where

appropriate, strains harbouring pGCP123 plasmids were grown in 1000 µg/mL kanamycin

(Sigma Aldrich, USA) and strains harbouring pMSP3535 plasmids were grown in 100 µg/mL

erythromycin unless stated otherwise. Metals were filtered sterilized and supplemented

during medium preparation in autoclaved TSBG media. For experiments using ferric chloride

only, metal is supplemented in TSBG media and autoclaved together. Ferric citrate hydrate

≥98%, magnesium chloride anhydrous ≥98%, copper chloride dihydrate ≥99%, ferrous

sulphate heptahydrate ≥99%, ferric sulphate hydrate ≥97%, ferric chloride anhydrous ≥99%,

heme ≥90%, and the chelator 2,2'dipyridyl ≥99% were all supplied by Sigma Aldrich, St

Louis, MO, USA. Manganese chloride tetrahydrate and zinc chloride were supplied by Merk

Millipore, Singapore.

General cloning techniques. Both *menB* and *ndh3* nucleotide sequences are based on the *E.*

faecalis OG1RF genome obtained from BioCyc [60]. The Wizard genome DNA purification kit

(Promega Corp., Madison, WI) was used for isolation of bacterial genomic DNA (gDNA),

and Monarch® Plasmid miniprep Kit (New England BioLabs, Ipswitch, MA) was used for

purification of plasmid for gene expression and construction of deletion mutant. The

Monarch® DNA Gel Extraction Kit (New England BioLabs, Ipswitch, MA) was used to

isolate PCR products during extension overlay PCR. In-Fusion HD Cloning Kit (TaKara Bio,

USA) was used for fast, directional cloning of DNA fragments into vector for both

expression vector and in-frame deletion vector. All plasmids used in the study are listed in

Table S3. T4 DNA ligase and restriction endonucleases were purchased from New England

BioLabs (Ipswitch, MA). Colony PCR was performed using Taq DNA polymerase (Thermo

Fisher Scientific, Waltham, MA, USA) and PCR of gene of interest for plasmid construction

was performed using Phusion DNA polymerase (Thermo Fisher Scientific, Waltham, MA,

USA). Ligations were transformed into *E. coli* Dh5α cells. Plasmids derived in this study

were confirmed by sequencing of purified plasmid.

Strain construction. To construct *menB* and *ndh3* complementation plasmids, primers

(menB_F' and menB_R' for *menB*, or ndh3_F' and ndh3_R' for *ndh3*; **Table S3)** were

designed with BamHI restriction site or SpeI restriction sites flanking the gene of interest, to

generate DNA fragments as templates for In-Fusion cloning (Takara Bio USA Inc.) using

primers (menB_F'_Infusion and menB_R'_Infusion for *menB*, or ndh3_F'_Infusion and

ndh3_R'_Infusion for *ndh3*) with at least 15 bp complementary sequence for ligation into the

nisin-inducible vector pMSP3535 digested with the same restriction enzymes. Both

pMSP3535::*menB* and pMSP3535::*ndh3* plasmids were generated in *E. coli* Dh5α, verified

by sequencing, and transformed into *E. faecalis* as described previously [13]. Deletion of the

menB coding sequence from the OG1RF chromosome was accomplished by allelic

replacement using pGCP213 temperature sensitive shuttle vector as described previously [13].

The deletion allele was constructed by extension overlap PCR, consisting of the upstream

(IFD_menB_Frag2_F' and IFD_menB_Frag2_R'; **Table S3)** and downstream

(IFD_menB_Frag1_F' and IFD_menB_Frag1_R'; **Table S3)** of the *menB* coding sequence,

and introduced into pGCP213 using in-fusion cloning with IFD_menB_Frag2_F' and

IFD_menB_Frag1_R' primers which has at least 15 bp complementary. Deletion of *menB*

was performed in *ndh3*::Tn parent strain to generate a double mutant. Transformants were

PCR screened for in-frame deletion as previously described [13].

Immunofluorescence Imaging and 3D Reconstruction of Biofilm. Bacterial cultures were

normalized to OD_{600} 0.7 and diluted 1000x prior inoculating into each well of μ-Slide 8 well

glass bottom slides (ibiTreat coated) (ibidi Inc., USA) containing 40% v/v TSBG for

incubation at 37°C under static conditions for 24 hours. After incubation, adherent biofilm

was fixed in 4% paraformaldehyde (Sigma Aldrich, USA)-PBS for 30 minutes, blocked with

3% bovine serum album (Sigma Aldrich, USA) (BSA)-PBS for 30 minutes, and incubated

with primary antibody (Rabbit Anti-Group D antigen, Thermo Scientific Singapore) or

Guinea Pig Anti-EbpC [45] at 1:500 dilution for 30 minutes. The biofilm was then washed with

3% BSA-PBS and incubated with secondary antibody (IgG horseradish peroxidase-

conjugated anti-rabbit or anti guinea-pig, Thermo Scientific Singapore) at 1:500 dilution for

30 minutes and washed with 3% BSA-PBS. Biofilms were hydrated with PBS prior to

imaging. Biofilm morphology, biomass, and cell distribution were analysed by confocal laser

scanning microscopy (CLSM). Images were acquired using LSM780 confocal microscope

(Zeiss, Germany) equipped with 63x/1.4 Oil DIC M27 and controlled by ZEN software.

Samples were illuminated with 488 nm and 561nm Argon laser line; GFP emitted

fluorescence was collected in the 493-580 nm range and RFP emitted fluorescence was

collected in the 568-712 nm range. Optical sections (134.95x134.95 μm) were collected

every 0.637 μm through the entire biofilm thickness and signal from each section was

averaged 4 times. Fiji software was used for further processing (levels adjustment, stack

resliced). To visualize the biofilm spatial organization coordinates representing cocci position

were plotted as spheres with cell-size diameter as described previously[21]. For each cell, local

density was calculated as number of neighbour cells within 4μm radius, normalized and

visualized using colour gradient.

Biofilm Assay. Bacterial cultures were normalized as previously described [21] and inoculated

in TSBG in a 96-well flat bottom transparent microtiter plate (Thermo Scientific, Waltman,

MA, USA), and incubated at 37°C under static conditions for 5 days unless specified

otherwise. Strains harboring pGCP123 complementation plasmids were grown in the

presence of kanamycin prior to the biofilm assay, but kanamycin was not added to the biofilm

assay itself because we found that kanamycin precipitates in the presence of excess iron

leading to aberrant biofilm formation (data not shown). Adherent biofilm biomass was

stained using 0.1% w/v crystal violet (Sigma-Aldrich, St Louis, MO, USA) at 4°C for 30

minutes. The microtiter plate was washed twice with PBS followed by crystal violet

solubilization with ethanol:acetone (4:1) for 45 minutes at room temperature. Quantification

of adherent biofilm biomass was measured by absorbance at OD_{595nm} using a Tecan Infinite

200 PRO spectrophotometer (Tecan Group Ltd., Männedorf, Switzerland).

Planktonic Growth Assay. Bacterial cultures were normalized as previously described [21]

and serially diluted by a dilution factor of 200. Diluted cultures were inoculated into the

media at a ratio of 1:25, which is 8 µL of the inoculum in 200 µL of media, incubated at $37^{o}C$

for 18 hours, and absorbance at OD_{600nm} was measured using a Tecan Infinite 200 PRO

spectrophotometer (Tecan Group Ltd., Männedorf, Switzerland) at 15 minute intervals.

Population Analysis of Pilus Expression. Biofilms were grown in 6 well tissue culture

plates at $37^{o}C$ under static conditions for 24 hours in TSBG or TSBG supplemented with

metals where appropriate, and planktonic cultures were grown in 50-ml falcon tube under

shaking conditions (200 rpm). Immunoblotting was performed as described previously [61] with

some modification. Spent media was removed, and the adherent biofilm was suspended in

PBS. A cell scraper was used to dislodge the biofilm, and the dislodged cells were

centrifuged at 14,000 rpm for 2 minutes at room temperature to remove the supernatant.

Biofilm cells were suspended in PBS and normalized to OD_{600nm} 1.0. Normalized

suspensions were fixed in 4% paraformaldehyde (Sigma Aldrich, USA)-PBS for 20 minutes

at $4^{o}C$, centrifuged at 14,000 rpm for 2 minutes. and the supernatant then discarded. The

pellets were washed with PBS, centrifuged at 14,000 rpm for 2 minutes, and the supernatant

discarded again. Immunofluorescence microscopy was performed as described previously [61]

with the following modifications: 1:500 dilution of guinea pig anti-EbpC (Thermo Scientific,

Singapore) in PBS-3% bovine serum albumin, and 1:500 dilution of goat Anti-guinea pig

AlexaFluor-568 (Invitrogen Inc., USA) incubation for 30 minutes. After washing with PBS

twice, resuspended cells were applied to poly-L-lysine slides (Polysciences Inc., USA), dried

for 5 minutes, mounted with Vectashield®, and sealed with a cover slip. Images were

acquired using an inverted Epi-fluorescence microscope (Zeiss Axio observer Z1, Germany)

equipped with an EC Plan-Neofluar 100x/1.3 Oil objective and controlled by ZEN software.

For each biological sample, 5 images were acquired and the percentage of the cell population

expressing pili was quantified. At least 300 bacterial cells per strain per experiment were

scored for pilus expression. The metal concentrations used during the pilus expression

experiments were based on the highest concentration in which *E. faecalis* cells did not exhibit

growth inhibition (data not shown).

EbpA Structural Modelling. The EbpA amino acid sequence was submitted to the structural

homology-modelling server Swiss-Model (https://swissmodel.expasy.org) [62].   Template

structures for structural modelling were automatically selected based upon sequence identity.

The initial template search derived 723 templates that were filtered down to 50 templates,

whereby 14 models were constructed and ranked based on sequence similarity and coverage.

The top solution was based upon the PDB 3txa (GBS104 from *Streptococcus agalactiae*) that

showed 34% sequence similarity and 38% sequence coverage. The final EbpA model

included N177-P620. The $Fe3^+$ was created with Sketcher [63] and placed into position using

the MIDAS bound magnesium of 3txa and 2ww8 as a guide with the program COOT [64]. The

two conserved water molecules from both 3txa and 2ww8 were fitted into EbpA by a similar

means. Final alignments, analysis and structural figures were created with PyMOL [65].

Quantitative Real time PCR (qRT-PCR) and RNA sequencing. Biofilm cells were

prepared as described above for TEM. After washing, the biofilm cell pellet was incubated

with lysozyme from chicken egg white (10mg/ml) (Sigma Aldrich, USA) for 30 minutes at

37^oC to remove the cell wall and centrifuged at 14,000 rpm for 2 minutes at room

temperature to remove supernatant prior to cell lysis. RNA extraction was performed in a

Purifier® filtered PCR enclosure using the PureLink™ RNA mini kit (Invitrogen, USA)

according to the manufacturer's instructions. RNA purification and removal of DNA was

performed using TURBO DNA-free™ kit (Thermo Fisher, USA) and Agencourt®

RNAClean® XP Kit (Beckman Coulter, USA). Measurement of RNA yield and quality was

performed using Qubit® RNA HS assay kit (Thermo Fisher, USA) and RNA ScreenTape

System and 2200 TapeStation (Agilent, USA). Synthesis of cDNA was performed using

SuperScript III First-strand (Invitrogen, USA). Quantitative real-time PCR using cDNA was

performed using KAPA SYBR fast qPCR master mix kit (Sigma Aldrich, USA) and Applied

Biosystems StepOne Plus Real-Time PCR system. The expression of *ebpC*, *ebpR*, *srtA*, *srtC,*

argR3 and *gyrA* were analysed using primer pairs listed in **Table S2**. For each primer set, a

standard curve was established using genomic DNA from *E. faecalis* OG1RF. Normalized

concentrations of cDNA were used to determine the relative fold change in gene expression

as compared to *E. faecalis* OG1RF biofilm grown in TSBG. For RNA sequencing, ribosomal

RNA depletion was performed after RNA purification using Ribo-Zero™ rRNA removal kit

(Illumina, USA). cDNA library synthesis was performed using NEBNext RNA First-strand

and NEBNext Ultra directional RNA Second-strand synthesis module (New England BioLab,

US). Transcriptome library preparation was performed using 300bp paired end Illumina

sequencing.

Pilus Extraction and Purification. Cell surface protein extracts from biofilms were prepared

as described previously [5] with minor modifications. Biofilms were grown cells dislodged as

described above for TEM. Dislodged biofilm cells were centrifuge at 3300 g for 4 minutes at

15°C, washed with Tris buffer (10 mM Tris-HCl pH 8, 1 mM EDTA, 1 mM DTT) and

suspended in protoplast buffer with Pierce™ protease inhibitors (Thermo Scientific, USA).

Mutanolysin from *Streptomyces globisporus* ATCC 21553 (Sigma Aldrich, USA) and

benzonase nuclease (Sigma Aldrich, USA) were added and incubated at 37°C for 6 hours

with shaking (200 rpm). Following incubation, biofilm cells were centrifuged at 3300 g for 4

minutes, 15°C to obtain the supernatant. Supernatants were filtered through a 0.45μm Supor®

membrane (Sigma Aldrich, USA) and loaded into Corning Spin-X UF concentrators (MWCO

100kDa) (Sigma Aldrich, USA), centrifuged for 20 minutes, 15°C at 6000 g, and dialyzed

using Tris buffer using Spectra/Por® Float-A-Lyzer® G2 dialysis device (MWCO 300kDa)

(Spectrum Labs, USA). Pilus extracts were collected, quantified using Nanodrop 2000

Spectrophotometer (Thermofisher, USA), and normalized to 5 μg per sample. Samples were

mixed with an equal volume of NuPAGE® Tris-glycine native sample buffer (Novex®) and

loaded on to NuPAGE™ 3-8% Tris-acetate gel (Thermofisher, USA). Gel blots were run at

150 V for 3 hours. Western blotting was performed as described previously [6] with minor

modifications. Membranes were blocked in 0.05% v/v Tween-3% v/v bovine serum albumin-

PBS (Sigma Aldrich, USA) overnight at 4 $^{\circ}$C and washed twice with 0.05% v/v Tween

(Sigma Aldrich, USA)-PBS prior to incubation with primary and secondary antibodies, and

prior to incubation with detection substrate. Primary antibody (guinea pig Anti-EbpC [45] or

rabbit Anti-SecA [6]) diluted 1:1500, and secondary antibody (IgG guinea pig or rabbit

conjugated horseradish peroxidase) diluted 1:4000 were used. Protein was detected using

SuperSignal® west femto maximum sensitivity substrate kit (Thermo Scientific, USA).

Inductive-Coupled Plasma Mass Spectrometry (ICP-MS). Pilus extracts, normalized to

5μg per sample, were run under native condition as described above. Using the western blot

prepared gel as a reference, a separate identical NuPAGE™ Tris-acetate gel (Thermofisher,

USA) was cut to isolate pilus extract ladder above a molecular weight of 100 kDa. At a ratio

of 2:1, 70% nitric acid (Sigma Aldrich, USA) and 30% hydrogen peroxide (Sigma Aldrich,

USA) was added to the excised gel slice containing the pilus protein, and left under room

temperature for 3 days to allow complete digestion. The digested samples were diluted with

LC-MS grade water and filtered using 0.2 μm membrane, prior to analysis using ICP-MS.

Analysis of trace metals in samples was performed using ICP-MS model Elan-DRCe,

Meinhard Nebulizer model TR-30-C3 (Perkin Elmer; Model: N8122006 (Elan Standard

Torch)).

Electrochemical Assay and Analysis. Screen printed electrodes (SPE) (model DRP-C110;

DropSens, Spain) consisting of a 4 mm diameter carbon working electrode, carbon counter

electrode, and a Ag pseudo-reference electrode were controlled by a multichannel

potentiostat (VSP, Bio-Logic, France) in an electrochemical cell of 9 mL working volume

sealed with a Teflon cap. Chronoamperometry was performed as previously described with

minor modifications [21]. All electrochemical experiments were conducted at 37^{o}C in TSBG

medium supplemented with 1 mM $FeCl_3$ unless otherwise stated.

Ferrozine assay of ferric iron reductase activity. *E. faecalis* cells were normalized to

OD_{600nm} 0.7, diluted 1:200 in PBS and resuspended in TSBG supplemented with 2 mM iron

(III) chloride as previously described [21] and then supplemented with 0.5 mM ferrozine.

Experiments were performed as previously described with minor modification [26]. $OD592_{nm}$

measurements were made every 2 mins for up to 7 hours on *E. faecalis* biofilms grown

statically. Corresponding change in optical density measurements correlates with color

change due to ferrous iron binding to ferrozine after ferric iron is reduced to ferrous iron.

Mouse Gastrointestinal Tract (GI) Infection Model. Three week old male C57BL/6NTac

mice were administered ampicillin (VWR, USA) in their drinking water (1 g/L) for 5 days as

previously described [53,55]. Mice were then given one day of recovery from antibiotic

treatment prior to administration of approximately 1-5 x 10^8 CFU/ml *E. faecalis* (OD_{600nm}

0.5) in the drinking water for 3 days as previously described [66]. Before and after infection,

mice were monitored for signs of disease and weight loss. All animal experiments were

approved and performed in compliance with the Nanyang Technological University

Institutional Animal Care and Use Committee (IACUC). For dietary manipulation,

customized synthesized diets (C1038 iron deficient diet with 200 mg/kg ferric chloride as

control diet, C1038 iron deficient diet with 2000 mg/kg ferric chloride as high iron diet)

(Altromin, Germany) were given to mice prior to infection and throughout infection. Mice

were kept on customized diets for 3 weeks before administrating ampicillin as previously

described with minor modification in the synthesized diet [67,68]. At the indicated timepoints,

the small intestine, colon, and cecum were harvested. Tissue samples were homogenised in

PBS, serial diluted in PBS, and spot-plated on BHI agar with 10 mg/L colistin, 10 mg/L

nalidixic acid, 100 mg/L rifampicin, 25 mg/L fusidic acid for CFU enumeration. All

antibiotics were obtained from Sigma Aldrich, USA.

Authors Contributions

L.N.L and K.A.K. conceptualized the study. L.N.L, E.M and K.A.K designed the

experiments, analyzed the data and prepared the manuscript. L.N.L performed biofilm

experiments, immuno-fluorescence assays, ICP-MS, and analyzed the data. A.M analyzed the

confocal data and generated 3D reconstruction models. LN.L, P.M.L., Z.S.C, and E.M

performed the electrochemistry experiments and analyzed the data. L.N.L and J.J.W

performed mouse GI experiments and analyzed the data. L.N.L prepared the RNA samples

for RNA sequencing and K.K.L.C analyzed the data. J.P. and B.H. performed modeling

experiments. All authors reviewed the manuscript.

Acknowledgments

This work was supported by the National Research Foundation and Ministry of Education

Singapore under its Research Centre of Excellence Programme and by the Ministry of

Education Singapore under its Tier 2 programme (MOE2014-T2-2-124). Artur Matysik was

supported by the National Medical Research Council under its Clinical Basic Research Grant

(NMRC/CBRG/0086/2015). We thank Wandy Betty (Washington University, St Louis,

USA) for performing TEM, Hailyn V. Nielsen and Scott Hultgren (Washington University,

St Louis, USA) for providing the pilus mutants, and Gary Dunny (University of Minnesota)

for providing transposon mutants. Finally, we thank Sam Light and Dan Portnoy (University

of California, Berkeley) for sharing unpublished data and for critical discussions of the

manuscript.

References

1   Ch'ng, J. H., Chong, K. K. L., Lam, L. N., Wong, J. J. & Kline, K. A. Biofilm-associated infection
by enterococci. *Nature reviews. Microbiology* **17**, 82-94, doi:10.1038/s41579-018-0107-z
(2019).
2   O'Driscoll, T. & Crank, C. W. Vancomycin-resistant enterococcal infections: epidemiology,
clinical manifestations, and optimal management. *Infection and drug resistance* **8**, 217-230,
doi:10.2147/idr.s54125 (2015).
3   Nallapareddy, S. R. *et al.* Conservation of Ebp-type pilus genes among Enterococci and
demonstration of their role in adherence of *Enterococcus faecalis* to human platelets.
Infection and immunity **79**, 2911-2920, doi:10.1128/iai.00039-11 (2011).
4   Nallapareddy, S. R., Singh, K. V., Sillanpaa, J., Zhao, M. & Murray, B. E. Relative contributions
of Ebp Pili and the collagen adhesin ace to host extracellular matrix protein adherence and
experimental urinary tract infection by *Enterococcus faecalis* OG1RF. *Infection and immunity*
79, 2901-2910, doi:10.1128/iai.00038-11 (2011).
5   Nallapareddy, S. R. *et al.* Endocarditis and biofilm-associated pili of *Enterococcus faecalis*.
The Journal of clinical investigation **116**, 2799-2807, doi:10.1172/jci29021 (2006).
6   Nielsen, H. V. *et al.* Pilin and sortase residues critical for endocarditis- and biofilm-associated
pilus biogenesis in *Enterococcus faecalis*. *Journal of bacteriology* **195**, 4484-4495,
doi:10.1128/jb.00451-13 (2013).
7   Singh, K. V., Nallapareddy, S. R. & Murray, B. E. Importance of the ebp (endocarditis- and
biofilm-associated pilus) locus in the pathogenesis of *Enterococcus faecalis* ascending urinary
tract infection. *J Infect Dis* **195**, 1671-1677 (2007).

8   Sillanpää, J., Xu, Y., Nallapareddy, S. R., Murray, B. E. & Höök, M. A family of putative
MSCRAMMs from *Enterococcus faecalis*. *Microbiology* **150**, 2069-2078,
doi:doi:10.1099/mic.0.27074-0 (2004).
9   Kline, K. A. *et al.* Mechanism for sortase localization and the role of sortase localization in
efficient pilus assembly in *Enterococcus faecalis*. *Journal of bacteriology* **191**, 3237-3247,
doi:10.1128/JB.01837-08 (2009).
10   Plow, E. F., Haas, T. A., Zhang, L., Loftus, J. & Smith, J. W. Ligand Binding to Integrins. *Journal*
of Biological Chemistry **275**, 21785-21788, doi:10.1074/jbc.R000003200 (2000).
11   Bergelson, J. M. & Hemler, M. E. Integrin-ligand binding. Do integrins use a 'MIDAS touch' to
grasp an Asp? *Current biology : CB* **5**, 615-617 (1995).
12   Whittaker, C. A. & Hynes, R. O. Distribution and evolution of von Willebrand/integrin A
domains: widely dispersed domains with roles in cell adhesion and elsewhere. *Molecular*
biology of the cell **13**, 3369-3387, doi:10.1091/mbc.e02-05-0259 (2002).
13   Nielsen, H. V. *et al.* The Metal Ion-Dependent Adhesion Site Motif of the *Enterococcus*
faecalis EbpA Pilin Mediates Pilus Function in Catheter-Associated Urinary Tract Infection.
mBio **3**, e00177-00112, doi:10.1128/mBio.00177-12 (2012).
14   Flores-Mireles, A. L., Pinkner, J. S., Caparon, M. G. & Hultgren, S. J. EbpA vaccine antibodies
block binding of *Enterococcus faecalis* to fibrinogen to prevent catheter-associated bladder
infection in mice. *Science translational medicine* **6**, 254ra127-254ra127,
doi:10.1126/scitranslmed.3009384 (2014).
15   Schembri, M. A., Lo, A. W. & Ulett, G. C. Blocking the 'MIDAS' touch of *Enterococcus faecalis*.
Annals of translational medicine **3**, S3-S3, doi:10.3978/j.issn.2305-5839.2015.03.32 (2015).
16   Proft, T. & Baker, E. N. Pili in Gram-negative and Gram-positive bacteria - structure, assembly
and their role in disease. *Cellular and molecular life sciences : CMLS* **66**, 613-635,
doi:10.1007/s00018-008-8477-4 (2009).
17   Danne, C. & Dramsi, S. Pili of Gram-positive bacteria: roles in host colonization. *Research in*
Microbiology **163**, 645-658, doi:https://doi.org/10.1016/j.resmic.2012.10.012 (2012).
18   Konto-Ghiorghi, Y. *et al.* Dual Role for Pilus in Adherence to Epithelial Cells and Biofilm
Formation in *Streptococcus agalactiae*. *PLOS Pathogens* **5**, e1000422,
doi:10.1371/journal.ppat.1000422 (2009).
19   Izore, T. *et al.* Structural basis of host cell recognition by the pilus adhesin from
Streptococcus pneumoniae. *Structure (London, England : 1993)* **18**, 106-115,
doi:10.1016/j.str.2009.10.019 (2010).
20   Becherelli, M. *et al.* The ancillary protein 1 of *Streptococcus pyogenes* FCT-1 pili mediates cell
adhesion and biofilm formation through heterophilic as well as homophilic interactions.
Molecular microbiology **83**, 1035-1047, doi:10.1111/j.1365-2958.2012.07987.x (2012).
21   Keogh, D. *et al.* Extracellular Electron Transfer Powers *Enterococcus faecalis* Biofilm
Metabolism. *mBio* **9**, doi:10.1128/mBio.00626-17 (2018).
22   Shi, L. *et al.* Extracellular electron transfer mechanisms between microorganisms and
minerals. *Nature reviews. Microbiology* **14**, 651-662, doi:10.1038/nrmicro.2016.93 (2016).
23   Weber, K. A., Achenbach, L. A. & Coates, J. D. Microorganisms pumping iron: anaerobic
microbial iron oxidation and reduction. *Nature reviews. Microbiology* **4**, 752-764,
doi:10.1038/nrmicro1490 (2006).
24   Sure, S. K., Ackland, L. M., Torriero, A. A., Adholeya, A. & Kochar, M. Microbial Nanowires: An
Electrifying Tale. *Microbiology*, doi:10.1099/mic.0.000382 (2016).
25   Wang, F. *et al.* Structure of Microbial Nanowires Reveals Stacked Hemes that Transport
Electrons over Micrometers. *Cell* **177**, 361-369.e310, doi:10.1016/j.cell.2019.03.029 (2019).
26   Light, S. H. *et al.* A flavin-based extracellular electron transfer mechanism in diverse Gram-
positive bacteria. *Nature* **562**, 140-144, doi:10.1038/s41586-018-0498-z (2018).

27   Pankratova, G., Leech, D., Gorton, L. & Hederstedt, L. Extracellular Electron Transfer by the
Gram-Positive Bacterium *Enterococcus faecalis*. *Biochemistry* **57**, 4597-4603,
doi:10.1021/acs.biochem.8b00600 (2018).
28   Ramsey, M., Hartke, A. & Huycke, M. in *Enterococci: From Commensals to Leading Causes of*
Drug Resistant Infection   (eds M. S. Gilmore, D. B. Clewell, Y. Ike, & N. Shankar)
(Massachusetts Eye and Ear Infirmary, 2014).
29   Sillanpaa, J. *et al.* Contribution of individual Ebp Pilus subunits of *Enterococcus faecalis*
OG1RF to pilus biogenesis, biofilm formation and urinary tract infection. *PloS one* **8**, e68813,
doi:10.1371/journal.pone.0068813 (2013).
30   Krishnan, V. *et al.* Structure of *Streptococcus agalactiae* tip pilin GBS104: a model for GBS pili
assembly and host interactions. *Acta crystallographica. Section D, Biological crystallography*
69, 1073-1089, doi:10.1107/S0907444913004642 (2013).
31   Shimaoka, M., Takagi, J. & Springer, T. A. Conformational regulation of integrin structure and
function. *Annual review of biophysics and biomolecular structure* **31**, 485-516,
doi:10.1146/annurev.biophys.31.101101.140922 (2002).
32   Kristich, C. J. *et al.* Development and use of an efficient system for random mariner
transposon mutagenesis to identify novel genetic determinants of biofilm formation in the
core *Enterococcus faecalis* genome. *Applied and environmental microbiology* **74**, 3377-3386,
doi:10.1128/aem.02665-07 (2008).
33   Dale, J. L. *et al.* Comprehensive Functional Analysis of the *Enterococcus faecalis* Core
Genome Using an Ordered, Sequence-Defined Collection of Insertional Mutations in Strain
OG1RF. *mSystems* **3**, doi:10.1128/mSystems.00062-18 (2018).
34   Logan, B. E., Rossi, R., Ragab, A. a. & Saikaly, P. E. Electroactive microorganisms in
bioelectrochemical systems. *Nature Reviews Microbiology*, doi:10.1038/s41579-019-0173-x
(2019).
35   Logan, B. E. Exoelectrogenic bacteria that power microbial fuel cells. *Nature Reviews*
Microbiology **7**, 375, doi:10.1038/nrmicro2113 (2009).
36   Rana, N. F. *et al.* Redox balance via lactate dehydrogenase is important for multiple stress
resistance and virulence in *Enterococcus faecalis*. *Infection and immunity* **81**, 2662-2668,
doi:10.1128/IAI.01299-12 (2013).
37   Huycke, M. M. *et al.* Extracellular superoxide production by *Enterococcus faecalis* requires
demethylmenaquinone and is attenuated by functional terminal quinol oxidases. *Molecular*
microbiology **42**, 729-740 (2001).
38   Vitreschak, A. G., Rodionov, D. A., Mironov, A. A. & Gelfand, M. S. Regulation of riboflavin
biosynthesis and transport genes in bacteria by transcriptional and translational attenuation.
Nucleic acids research **30**, 3141-3151 (2002).
39   Zhang, E., Cai, Y., Luo, Y. & Piao, Z. Riboflavin-shuttled extracellular electron transfer from
Enterococcus faecalis to electrodes in microbial fuel cells. *Canadian journal of microbiology*
60, 753-759, doi:10.1139/cjm-2014-0389 (2014).
40   Byappanahalli, M. N., Nevers, M. B., Korajkic, A., Staley, Z. R. & Harwood, V. J. Enterococci in
the environment. *Microbiology and molecular biology reviews : MMBR* **76**, 685-706,
doi:10.1128/MMBR.00023-12 (2012).
41   Benito, P. & Miller, D. Iron absorption and bioavailability: An updated review. *Nutrition*
Research **18**, 581-603, doi:https://doi.org/10.1016/S0271-5317(98)00044-X (1998).
42   Saunders, S. H. & Newman, D. K. Extracellular Electron Transfer Transcends Microbe-Mineral
Interactions. *Cell Host & Microbe* **24**, 611-613,
doi:https://doi.org/10.1016/j.chom.2018.10.018 (2018).
43   Guiton, P. S. *et al.* Contribution of autolysin and Sortase a during *Enterococcus faecalis* DNA-
dependent biofilm development. *Infection and immunity* **77**, 3626-3638,
doi:10.1128/iai.00219-09 (2009).

44 Thomas, V. C., Thurlow, L. R., Boyle, D. & Hancock, L. E. Regulation of autolysis-dependent
extracellular DNA release by *Enterococcus faecalis* extracellular proteases influences biofilm
development. *Journal of bacteriology* **190**, 5690-5698, doi:10.1128/jb.00314-08 (2008).
45 Afonina, I., Lim, X. N., Tan, R. & Kline, K. A. Planktonic Interference and Biofilm Alliance
between Aggregation Substance and Endocarditis- and Biofilm-Associated Pili in
Enterococcus faecalis. *Journal of bacteriology* **200**, doi:10.1128/jb.00361-18 (2018).
46 Bourgogne, A., Thomson, L. C. & Murray, B. E. Bicarbonate enhances expression of the
endocarditis and biofilm associated pilus locus, ebpR-ebpABC, in *Enterococcus faecalis*. *BMC*
Microbiol **10**, 17, doi:10.1186/1471-2180-10-17 (2010).
47 Bourgogne, A. *et al.* EbpR is important for biofilm formation by activating expression of the
endocarditis and biofilm-associated pilus operon (ebpABC) of *Enterococcus faecalis* OG1RF.
Journal of bacteriology **189**, 6490-6493, doi:10.1128/JB.00594-07 (2007).
48 Manias, D. A. & Dunny, G. M. Expression of Adhesive Pili and the Collagen-Binding Adhesin
Ace Is Activated by ArgR Family Transcription Factors in *Enterococcus faecalis*. *Journal of*
bacteriology **200**, doi:10.1128/jb.00269-18 (2018).
49 Bourgogne, A., Hilsenbeck, S. G., Dunny, G. M. & Murray, B. E. Comparison of OG1RF and an
Isogenic fsrB Deletion Mutant by Transcriptional Analysis: the Fsr System of *Enterococcus*
faecalis Is More than the Activator of Gelatinase and Serine Protease. *Journal of bacteriology*
188, 2875-2884, doi:10.1128/JB.188.8.2875-2884.2006 (2006).
50 Gao, P. *et al. Enterococcus faecalis* rnjB is required for pilin gene expression and biofilm
formation. *Journal of bacteriology* **192**, 5489-5498, doi:10.1128/jb.00725-10 (2010).
51 Gao, P. *et al.* Functional studies of *E. faecalis* RNase J2 and its role in virulence and fitness.
PloS one **12**, e0175212, doi:10.1371/journal.pone.0175212 (2017).
52 Glasauer, S., Langley, S. & Beveridge, T. J. Intracellular iron minerals in a dissimilatory iron-
reducing bacterium. *Science (New York, N.Y.)* **295**, 117-119, doi:10.1126/science.1066577
(2002).
53 Ubeda, C. *et al.* Vancomycin-resistant *Enterococcus* domination of intestinal microbiota is
enabled by antibiotic treatment in mice and precedes bloodstream invasion in humans. *The*
Journal of clinical investigation **120**, 4332-4341, doi:10.1172/JCI43918 (2010).
54 Caballero, S. *et al.* Distinct but Spatially Overlapping Intestinal Niches for Vancomycin-
Resistant *Enterococcus faecium* and Carbapenem-Resistant *Klebsiella pneumoniae*. *PLOS*
Pathogens **11**, e1005132, doi:10.1371/journal.ppat.1005132 (2015).
55 Caballero, S. *et al.* Cooperating Commensals Restore Colonization Resistance to Vancomycin-
Resistant *Enterococcus faecium*. *Cell Host Microbe* **21**, 592-602.e594,
doi:10.1016/j.chom.2017.04.002 (2017).
56 La Rosa, S. L. *et al.* In vivo assessment of growth and virulence gene expression during
commensal and pathogenic lifestyles of luxABCDE-tagged *Enterococcus faecalis* strains in
murine gastrointestinal and intravenous infection models. *Applied and environmental*
microbiology **79**, 3986-3997, doi:10.1128/aem.00831-13 (2013).
57 Riboulet, E. *et al.* Relationships between oxidative stress response and virulence in
Enterococcus faecalis. *Journal of molecular microbiology and biotechnology* **13**, 140-146,
doi:10.1159/000103605 (2007).
58 Aktan, Y., Tan, S. & Icgen, B. Characterization of lead-resistant river isolate *Enterococcus*
faecalis and assessment of its multiple metal and antibiotic resistance. Vol. 185 (2012).
59 Colomer-Winter, C., Gaca, A. O. & Lemos, J. A. Association of Metal Homeostasis and
(p)ppGpp Regulation in the Pathophysiology of *Enterococcus faecalis*. *Infection and*
immunity **85**, doi:10.1128/iai.00260-17 (2017).
60 Karp, P. D. *et al.* The BioCyc collection of microbial genomes and metabolic pathways.
Briefings in bioinformatics, doi:10.1093/bib/bbx085 (2017).

61   Kline, K. A. *et al.* Mechanism for Sortase Localization and the Role of Sortase Localization in
Efficient Pilus Assembly in *Enterococcus faecalis. Journal of bacteriology* **191**, 3237-3247,
doi:10.1128/jb.01837-08 (2009).
62   Waterhouse, A. *et al.* SWISS-MODEL: homology modelling of protein structures and
complexes. *Nucleic Acids Res* **46**, W296-w303, doi:10.1093/nar/gky427 (2018).
63   Winn, M. D. *et al.* Overview of the CCP4 suite and current developments. *Acta Crystallogr D*
Biol Crystallogr **67**, 235-242, doi:10.1107/s0907444910045749 (2011).
64   Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta Crystallogr D*
Biol Crystallogr **60**, 2126-2132, doi:10.1107/s0907444904019158 (2004).
65   L DeLano, W. *The PyMOL Molecular Graphics System (2002) DeLano Scientific, Palo Alto, CA,*
USA. http://www.pymol.org. (2002).
66   Banla, I. L. *et al.* Modulators of *Enterococcus faecalis* Cell Envelope Integrity and
Antimicrobial Resistance Influence Stable Colonization of the Mammalian Gastrointestinal
Tract. *Infection and immunity* **86**, e00381-00317, doi:10.1128/IAI.00381-17 (2017).
67   Zackular, J. P. *et al.* Dietary zinc alters the microbiota and decreases resistance to *Clostridium*
difficile infection. Nat Med **22**, 1330-1334, doi:10.1038/nm.4174 (2016).
68   Juttukonda, L. J. *et al.* Dietary Manganese Promotes Staphylococcal Infection of the Heart.
Cell Host Microbe **22**, 531-542.e538, doi:10.1016/j.chom.2017.08.009 (2017).

Figure 1. Contribution of pilus to iron mediated *E. faecalis* biofilm growth. (a) *E.*

faecalis biofilm biomass quantification grown for 120 hours using crystal violet staining.

N=9 repeated on non-consecutive days. **(b)** Volume viewer of CSLM images of *E. faecalis*

biofilm at 24 hr for control (40% TSBG) and iron-supplemented (40% TSBG, 0.2 mM FeCl$_3$)

at 63x magnification. *E. faecalis* cells are labeled with antibody against Group D antigen. **(c)**

Overlap of cell aggregation density map with the piliated cell population density map using

3D reconstruction of CLSM z-stack images of iron-supplemented *E. faecalis* biofilm

841   compared with control. N=3, repeated on non-consecutive days. **(d)** Quantification of

842   immuno-fluorescent labelled anti-EbpC *E. faecalis* cells from 6-well microtiter plate 24 hour

843   biofilm or planktonic culture grown at 37 ^{o}C in control and metal supplemented media. N>3.

844   Statistical significance was determined by two-way ANOVA. Error bars represent standard

845   error of the mean (SEM).

Figure 2. Evidences for EbpA-iron interaction. (a) Representative immunofluorescent

images from 24 hour *E. faecalis* biofilm (wild type, Δ*ebpABC* and *ebpA^{AWAGA}*) labeled with

EbpC antibody-conjugated fluorescent probe, grown from 6-well microtiter plate biofilm in

the presence of iron, N=3. **(b)** ICP-MS analysis of iron (ppb) from pilus preps isolated under

native condition. N=6. Statistical analysis is performed using two-way ANOVA Fisher LSD

test. Error bars represent SEM. Black bar indicates 500nm. **(c)** Superimposition of EbpA

structural model (N177-P620) (blue) with that of the *S. agalactiae* GBS104 (grey) and *S.*

pneumoniae RrgA (pink). In agreement with other pilin adhesins, EbpA tip folds into a

vWFA domain comprising of a 5 stranded β-sheet surrounded by 7 α-helices. The MIDAS

motif is located at the N-terminus of the β-sheet. Only a partial model was obtained for the 2

extended arms of this domain. The EbpA tip is then followed by an IgG fold domain. **(d)**

Closeup view of the superimposition of the MIDAS motif from EbpA (blue), GBS104 (grey)

and RrgA (pink) showing conservation of D275, S277, S279, T350 and D378 residues (EbpA

numbering). Bound Mg^{2+} ions (grey/magenta) in GBS104 and RrgA and modelled

Fe^{3+}(orange) in EbpA are shown as spheres. **(e)** The Fe^{3+} is coordinated by the EbpA MIDAS

motif by ~2Å direct interactions with S277, S279, D378 and a water mediated interaction

with D275 along with interactions with two additional water molecules (water molecules

shown as blue spheres). Upon a closed to open conformational change, bonding could shift

from D378 to T350.

Figure 3. Extracellular electron transfer in *E. faecalis* biofilm. (a) Cumulative charge production (over 20 hours) in *E. faecalis* pilin mutants

grown in the presence of 0.5 mM iron and **(b)** EET-associated mutants grown in the presence of 2 mM iron on a screen-printed carbon mini-

electrode for 20 hours in a closed static system at 37°C. Experiments were repeated on non-consecutive days, N>3. Statistical analysis is

performed using one-way ANOVA with Bonferroni multiple comparison test. Error bar represents standard error margin (SEM).

Figure 4: Intracellular iron accumulation is dependent on the ability for extracellular ferric iron reduction. (a-b) ICP-MS analysis of intracellular iron (ppb) in *E. faecalis* biofilm cells grown in iron-supplemented media. N=9. **(c)** Quantification of ferric iron reduction activity by *E. faecalis*. Statistical significance was determined by one-way ANOVA. Error bars represent standard error margin (SEM).

Figure 5. *E. faecalis* **colonization in lower gastrointestinal tract. (a)** Schematic of diet

manipulation in GI colonization model. **(b)** Bacterial titers of pilus null mutant (N=3, n=5)

and EET-associated mutants (for WT n=30, for mutants n=15) recovered from mouse colons,

repeated on non-consecutive days. Statistical significance was determined by Mann-Whitney

test. Black line indicates the median. Dotted line indicates limitation of detection at CFU <40.

Figure 6. Model for the role of iron in *E. faecalis* biofilm. Biofilm growth in excess iron induces expression of the genes encoding the sortase-assembled Ebp, which in turn associate with iron via the MIDAS motif of the EbpA tip adhesin. EET is dependent on both DMK in the electron transport chain, as well as a specialized flavin-associated respiratory chain, leading to reduction of extracellular iron and FeoB-mediated uptake of reduced free ferrous iron.

Figure S1. Pili contribution to biofilm formation. (a) *E. faecalis* biofilm total biomass quantification and **(b)** fold change calculations for the

same data shown in Fig 1A and Fig S1A after growth for 120 hours grown in non-supplemented and iron-supplemented media in 96-well

922   microtiter plate as measured by crystal violet staining. N=9 repeated on non-consecutive days. Statistical significance was determined by two-

923   way ANOVA. Error bars represent the standard error of the mean (SEM).

Figure S2. Antibody specificity against cell surface expressed pili. Representative

immunofluorescent images of *E. faecalis* and pilus mutants **(a)** cropped and **(b)** at original

63x magnification from biofilm grown in 6-well microtiter plates for 24 hours at 37 degrees

in non-supplemented media or both non-supplemented and iron-supplemented media. N=6,

with 5 images per biological replicate, capturing approximately 100-300 cells per images,

repeated on non-consecutive days. **(c)** Quantification of immuno-fluorescent labelled anti-

EbpC *E. faecalis* cells from 6-well microtiter plate 24hr biofilm grown at 37 degrees in

control and metal supplemented media. Statistical significance was determined by two-way

ANOVA with Bonferroni multiple comparison test. Error bars represent the standard error of

the mean (SEM).

```
EbpA     MITDENDKTNINIELNLLNQTEQPLQREIQLKNAQFMDTAVIEKDGYSYQVTNGTLYLTL      60
GBS104   ----------------------------MKKRQKIWRGLSVTLLILSQI---P--FGI      25
RrgA     ----------------------MLNRETHMKKVRKIFQKAVAGLCCISQL---TAFSSI      34
                                     :*: : :                *:         :

EbpA     DAQVK----KPVQLSLAVEQSSLQTAQPPKLLYE-NNEYDVSVTSEKITVEDS---AKES     112
GBS104   LVQGETQDTNQALGKVIVKKTGDNATPLGKATFVLKNDNDKSETSHE-TVEGSGEATFEN      84
RrgA     VALAETPETSPAIGKVVIKETGEGGALLGDAVFELKNNTDGTTVSQR-TEAQTGEAIFSN      93
              . :    . .  .: ::::.   :   . : :*: * : .*.. *    :     ..

EbpA     TEPEKITVPENTK-----ETNKNDSAPEKTEQPTATEEVTNPFAEARMAPATLRANLALP     167
GBS104   IKPGDYTLREETAPIGYKKTDKTWKVKVADNGATIIEGMDADKAE-----------KRKE     133
RrgA     IKPGTYTLTEAQPPVGYKPSTKQWTVEVEKNGRTTVQGE---QVE-----------NREE     139
            :*    *: *          : *  ..    :  * :          .*

EbpA     LIAPQYTTDNSGTYPTANWQPTGNQNVLNHQGNKDGGAQWDGQTSWNGDPTNRTNSYTEY     227
GBS104   VLNAQYP--KSAIYEDTK----ENYPLVNVEGSKVG-EQYKALNPINGKDGRREIAE---     183
RrgA     ALSDQYP--QTGTYPDVQ----TPYQIIKVDGSEKN-GQHKALNPN---PYERVIPE---     186
            :   **   ::.  *  .:         ::: :*.:.   *  ...        .*

EbpA     GGTGDQADYAIRKYARETTTPGLFDVYLNVRGNV----QKEITPLDLVLVVDWSGSMNEN     283
GBS104   -G-----WLSKKITGVNDLDKNKYKIELTVEGKTTVETKELNQPLDVVVLLDNSNSMNNE     237
RrgA     -G-----TLSKRIYQVNNLDDNQYGIELTVSGKTVYEQKDKSVPLDVVILLDNSNSMSNI     240
          *         : :   :     . : : *.* *:.     :.    ***:*:::* *.**.:

EbpA     NRIG-----EVQKGVNRFVDTLADSGITNNINMGYVGYSSDGYN----------------     322
GBS104   RANNSQRALKAGEAVEKLIDKITSNKD---NRVALVTYASTIFDGTEATVSKGVADQNGK     294
RrgA     RNKNARRAERAGEATRSLIDKITSDSE---NRVALVTYASTIFDGTEFTVEKGVADKNGK     297
          .  .      ..  :... ::*.::..       .:. * *:*  ::

EbpA     ----------------NNAIQM---------GPFDTVKN-------PIKNITPSSTRGGT     350
```

```
998    GBS104    ALNDSVSWDYHKTTFTATTHNYSYLNLTNDANEVNILKSRIPKEAEHINGDRTLYQFGAT    354
999    RrgA      RLNDSLFWNYDQTSFTTNTKDYSYLKLTNDKNDIVELKNKVPTEAEDHDGNRLMYQFGAT    357
1000                       .: :              . .   :*.          ..        *.*

1001

1002   EbpA      FTQKALRDAGDMLATP--NGHKKVIVLLTDGVPTFSYKVSRVQTEADGRFYGTQFTNRQD    408
1003   GBS104    FTQKALMKANEILETQSSNARKKLIFHVTDGVPTMSYAINFNPY-ISTS-YQNQFN----    408
1004   RrgA      FTQKALMKADEILTQQARQNSQKVIFHITDGVPTMSYPINFNHATFAPS-YQNQLN----    412
1005             ****** .*.::*       :   :*:*. :******:** :.          * .*:.

1006

1007   EbpA      QPGSTSYISGSYNAPDQNNISKRINSTFIATI--GEAMALKQRGIEIHGLGIQLQSDPRA    466
1008   GBS104    -----SFLN---KIPDRSGI---LQEDFIING--DDYQIVKGD-----GESFKLFSD---    447
1009   RrgA      -----AFFS---KSPNKDGI---LLSDFITQATSGEHTIVRGD-----GQSYQMFTD---    453
1010               :::.    :  *::..*    :  . **       .:    ::          * . :: :*

1011

1012   EbpA      NLSKQQVEDKMREMVSADENGDLYYESADYAPDISDYLAKKAVQISGTVVNGKVVDPIAE    526
1013   GBS104    ----------------------------------------RKVPVTGG--------TTQA    459
1014   RrgA      ----------------------------------------KTVYEKGA--------P--A    463
1015                                                      :  *   .*

1016

1017   EbpA      PFKYEPNTLSMKSVGPVQVQTLPEVSLTGATINSNEIYLGKGQEIQIHYQVRIQT-----    581
1018   GBS104    AYRVPQNQLSVMS--------NEGY-----AINSGYIYLWRDYN-WVYPFDPKTKKVSA    505
1019   RrgA      AFPVKPEKYSEMK--------AAGYAVIGDPINGGYIWLNWRESI-LAYPFNSNTA----    510
1020               :    :  *  .                  **.. *:*   :       * . :*

1021

1022   EbpA      ---ESENFKPDFWYQMNGRTTFQPLATAPEKVDFGVPSGKAPGVKLNVKKIWEEYDQDPT    638
1023   GBS104    TKQIKTHGEPTTLYFNGN-I--RPKGYDI--FTVGIGVNGDPGATPLE---AEKFMQSIS    557
1024   RrgA      --KITNHGDPTRWYYNGN-I--APDGYDV--FTVGIGINGDPGTDEAT---ATSFMQSIS    560
1025                .  : .*    *  ..      *  .    . .*:  .  **.         .: *. :

1026

1027   EbpA      SRPDNVIYEISRKQVTDTANWQTGYIKLSKPENDTSNSWERKNVTQLSKTADESYQEVLG    698
1028   GBS104    SKTEN-------------------------YTNVDDTNKIYDELNKY----FKTIVE    585
1029   RrgA      SKPEN-------------------------YTNVTDTTKILEQLNRY----FHTIVT    588
```

```
1030                    *: :*                                *.       *    :*.:      :: ::

1031

1032    EbpA      LPQYNNQGQAFNYQTTRELAVPGYS---QEKIDDTTWKNTKQFKPLDLKVIKNSSSGEKN    755

1033    GBS104    ---------------EKHSIVDGNVTDPMGEMIEFQLKNGQSFTHDDYVLVGNDGSQLKN    630

1034    RrgA      ---------------EKKSIENGTITDPMGELIDLQLGTDGRFDPADYTLTANDGSRLEN    633

1035                               :.    *         :: :      .    *    *   :   *..*   :*

1036

1037    EbpA      LVGAVFELSGKNVQTTLVDNKDGSYSLPKDVRLQKGERYTLTEVKAPAGHELGKKTTWQI    815

1038    GBS104    GV----ALGGPNSDGGILKDVTVTYD----------------------------------    652

1039    RrgA      GQ----AVGGPQNDGGLLKNAKVLYD----------------------------------    655

1040                  :.* : :  ::.:     *.

1041

1042    EbpA      EVNEQGKVSIDGQEVTTTNQVIPLEIENKFSSLPIRIRKYTMQNGKQVNLAEATFALQRK    875

1043    GBS104    ---------------------------KTSQTIKINHLNLGSGQKVVLTY---DVRLK    680

1044    RrgA      ---------------------------TTEKRIRVTGLYLGTDEKVTLTY---NVRLN    683

1045                                       :.   *::      :  ..::* *:       :: :

1046

1047    EbpA      NA---QGSYQTVATQKTDTAGLSYFKISEPGEYRMVEQSGPLGYDTLAGNYEFTVDKYGE    932

1048    GBS104    DNYISNKFYNTNNR-----TTLSPKSEKEPNTI----------------RDFPIPKIR-    717

1049    RrgA      DEFVSNKFYDTNGR-----TTLHPKE-VEQNTV----------------RDFPIPKIR-    719

1050              :    :   *:*         : *   . * .                        :*  : *

1051

1052    EbpA      IHYAGKNIEENAPEWTLTHQNHLKPFDLTVHKKADNQTPLKGAKFRLTGPDTDIEL----    988

1053    GBS104    -------DVREFPVLTISNQKKMGEVEFIKVNKDKHSESLLGAKFQLQIEK-DF---SGY    766

1054    RrgA      -------DVRKYPEITISKEKKLGDIEFIKVNKNDK-KPLRGAVFSLQKQHPDYPDIYGA    771

1055                     .:  *   *:::::::  .::    :* .:     * **  * *    . *

1056

1057    EbpA      -------PKDGKETDTFVFENLKPGKYVLTETFTPEGYQGLKEPIE-LIIREDGSVTID    1039

1058    GBS104    KQFVPEGSDVTTKNDGKIYFKALQDGNYKLYEISSPDGYIEVKTKPVVTFTIQNGEVTNL    826

1059    RrgA      IDQNGTYQNVRTGEDGKLTFKNLSDGKYRLFENSEPAGYKPVQNKPIVAFQIVNGEVRDV    831

1060                  :  ..:  *:  *.  *:*  *  *    *  **   ::          :     :*.*

1061
```

```
1062   EbpA       GEKV-AD----VLISGEKNNQITLDVTNQAKVPLPETGGIGRLWFYLIAISTFVIAGVYL    1094

1063   GBS104     KADPNANKNQIGYLEGNG-KHLITNTPKRPPGVFPKTGGIGTIVYILVGSTFM-ILTICS    884

1064   RrgA       TSIV-PQDIPAGYEFTND-KHYITNEPIPPKREYPRTGGIGMLPFYLIGCMMM-GGVLLY    888

1065                  :         :  ::   :            *.***** : : *:.    :      :

1066

1067   EbpA       FIRRPEGSV    1103

1068   GBS104     FRRKQL---    890

1069   RrgA       TRKHP----    893

1070               ::
```

Figure S3. Clustal alignment of EbpA homologs

Alignment of the EbpA sequence with *S. agalactiae* GBS104 (Q8E0S5) and *S. pneumoniae* RrgA

(A0A0H2UNT6), showing conservation of the EbpA MIDAS motif and associated metal binding

residues.

1076

Figure S4. Current output and cumulative charge production of pilin mutants in a

dosage dependent manner. (a) Cumulative charge production over 20 hours and **(b)** current

output in *E. faecalis* mutants grown on a screen-printed carbon mini-electrode over 20 hr in

closed static system at 37°C at 0.5 mM, 1 mM and 2 mM iron concentrations. Statistical

analysis is performed using one-way ANOVA with Bonferroni multiple comparison test,

N>3. Error bars represent the standard error of the mean (SEM).

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

Figure S5. Respiratory electron transport mechanisms in *E. faecalis*. (a) Gene homologs

of EET specialized respiratory electron transport chain and menaquinone biosynthesis genes

required for DMK. **(b)** Cumulative charge production over 20 hours by *E. faecalis* mutants of

EET specialized gene locus and menaquinone biosynthesis genes grown on a screen-printed

carbon mini-electrode over 20 hr in closed static system at 37°C at 2 mM iron concentration,

1112   N=3. Statistical analysis is performed using one-way ANOVA with Bonferroni multiple

1113   comparison test. Error bars represent the standard error of the mean (SEM).

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139 **Figure S6. Regulation of pilus-associated genes in response to iron.** qRT-PCR of RNA

1140 transcript abundance of genes associated with pilus biogenesis. N=6, repeated on non-

1141 consecutive days. N=9. Statistical significance was determined by one-way ANOVA. Error

1142 bars represent the SEM.

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

Figure S7. *E. faecalis* **pilin mutant colonization in mouse lower gastrointestinal tract. (a)** ICP-MS analysis of iron in the mouse tissues after feeding on normal or high iron diets, N=2, n=5. **(b)** Bacterial titers of pilin mutants recovered from the mouse cecum and small intestine, N=3, n=5. **(c)** Bacterial titers of EET associated mutants recovered from the mouse cecum and small intestine, N=3, n=5. Statistical significance was determined by Mann-Whitney test. Black line indicates median. Dotted line indicates limitation of detection of CFU <40.

Supplementary Table 1. RNA-sequencing results.

Old Gene Locus	New Gene Locus	RF-Bio1	RF-Bio2	RF-Bio3	RF_Fe-Bio1	RF_Fe-Bio2	RF_Fe-Bio3	logFC	logCPM	P-Value	FDR	Annotation
OG1RF_10072	OG1RF_RS00365	8511.8527	6488.9180	7624.4051	2705.9691	2800.5727	3142.9526	-1.3872	12.3487	0.0003	0.0140	transcriptional regulator (stress response regulator Gls24)
OG1RF_10073	OG1RF_RS00370	3214.8225	2303.0033	4763.1714	1013.7927	1268.0472	1233.1796	-1.5480	11.1690	0.0002	0.0111	membrane protein (transglycosylase associated protein)
OG1RF_10171	OG1RF_RS00860	1352.7718	926.4525	639.5307	3502.8808	2184.6641	2369.1118	1.4632	10.8424	0.0007	0.0268	preprotein translocase subunit SecY
OG1RF_10172	OG1RF_RS00865	411.1365	292.5639	283.1256	1160.0613	713.7294	798.6700	1.4329	9.2682	0.0005	0.0212	adenylate kinase
OG1RF_10269	OG1RF_RS01485	79.5748	82.5180	73.2796	161.3998	173.9036	240.0148	1.2876	7.1469	0.0017	0.0468	esterase
OG1RF_10296	OG1RF_RS01630	10.6100	18.7541	19.9853	65.5687	68.8368	64.1419	2.0227	5.5802	0.0001	0.0046	PTS mannitol transporter subunit IICB
OG1RF_10338	OG1RF_RS01840	1249.3246	465.1016	886.0165	254.7091	396.7176	326.9167	-1.4098	9.2303	0.0019	0.0500	osmotically inducible protein C
OG1RF_10340	OG1RF_RS01850	29.1774	7.5016	13.3236	113.4843	54.3449	78.6255	2.2048	5.8090	0.0000	0.0036	PTS sorbose transporter subunit IIC
OG1RF_10360	OG1RF_RS01950	172.4121	97.5213	109.9193	393.4121	226.4370	380.7132	1.3840	7.8873	0.0014	0.0405	ferrous iron transport protein B
OG1RF_10396	OG1RF_RS02135	1726.7735	1425.3115	1122.5096	522.0276	280.7819	403.4732	-1.8269	9.8397	0.0000	0.0017	hypothetical protein
OG1RF_10541	OG1RF_RS02845	198.9370	225.0492	369.7287	65.5687	94.1978	120.0074	-1.4948	7.5083	0.0007	0.0268	membrane protein (transglycosylase-associated protein)
OG1RF_10544	OG1RF_RS02860	140.5822	56.2623	99.9267	494.2870	278.9704	289.6731	1.8213	7.8679	0.0001	0.0057	glucuronyl hydrolase
OG1RF_10545	OG1RF_RS02865	98.1423	30.0066	39.9707	310.1903	206.5105	192.4257	2.0373	7.2614	0.0000	0.0029	beta-galactosidase
OG1RF_10549	OG1RF_RS02885	15.9150	0.0000	3.3309	83.2218	32.6069	37.2437	2.7733	5.1351	0.0001	0.0052	PTS fructose transporter subunit IID
OG1RF_10551	OG1RF_RS02895	100.7948	138.7803	376.3905	30.2625	83.3288	53.7964	-1.8588	7.0590	0.0009	0.0315	GTP cyclohydrolase
OG1RF_10565	OG1RF_RS02965	233.4195	82.5180	66.6178	1033.9677	364.1107	662.1098	2.4125	8.6951	0.0000	0.0018	PTS cellbiose transporter subunit IIC
OG1RF_10567	OG1RF_RS02975	132.6247	30.0066	26.6471	441.3277	144.9197	277.2585	2.1551	7.5121	0.0004	0.0184	hypothetical protein (outer surface protein)
OG1RF_10585	OG1RF_RS03065	34.4824	22.5049	9.9927	85.7437	52.5334	84.8328	1.6825	5.7758	0.0011	0.0342	hypothetical protein (M protein trans-acting positive regulator)
OG1RF_10627	OG1RF_RS03265	2503.9542	3803.3312	5176.2016	10014.3554	12850.7521	8520.5260	1.4507	12.8041	0.0006	0.0248	bifunctional acetaldehyde-CoA/alcohol dehydrogenase
OG1RF_10838	OG1RF_RS04395	387.2641	457.6000	419.6920	12.6094	19.9265	8.2764	-4.8996	7.7734	0.0000	0.0000	manganese transporter
OG1RF_10839	OG1RF_RS04400	82.2273	86.2689	93.2649	2.5219	0.0000	6.2073	-4.8069	5.5680	0.0000	0.0000	universal stress protein UspA
OG1RF_10849	OG1RF_RS04450	320.9518	150.0328	266.4711	1094.4926	561.5637	678.6626	1.6552	9.0194	0.0002	0.0120	galactokinase
OG1RF_10851	OG1RF_RS04460	400.5266	315.0689	269.8020	1205.4550	570.6212	993.1648	1.4870	9.3048	0.0006	0.0221	galactose-1-phosphate uridylyltransferase
OG1RF_10859	OG1RF_RS04500	289.1218	123.7770	176.5371	991.0959	445.6280	639.3498	1.8061	8.8178	0.0001	0.0077	excinuclease ABC subunit A (Y family DNA-directed DNA polymerase)
OG1RF_10860	OG1RF_RS04505	55.7024	26.2557	53.2942	216.8810	101.4438	167.5966	1.8143	6.7839	0.0001	0.0078	hypothetical protein

OG1RF_10861	OG1RF_RS04510	34.4824	15.0033	36.6398	98.3530	72.4598	111.7310	1.6747	6.0889	0.0004	0.0182	hypothetical protein
OG1RF_10862	OG1RF_RS04515	119.3622	75.0164	99.9267	257.2310	231.8715	285.5349	1.3836	7.5312	0.0006	0.0248	hypothetical protein
OG1RF_10870	OG1RF_RS04555	132.6247	157.5344	136.5665	360.6278	507.2189	502.7897	1.6850	8.2613	0.0000	0.0027	cell wall surface anchor protein (ebpA)
OG1RF_10871	OG1RF_RS04560	161.8021	311.3180	236.4931	491.7651	833.2881	755.2190	1.5567	8.8831	0.0003	0.0149	cell wall surface anchor protein (ebpB)
OG1RF_10874	OG1RF_RS04575	42.4399	71.2656	86.6031	7.5656	19.9265	18.6218	-2.0480	5.4664	0.0001	0.0061	DUF378 domain-containing protein
OG1RF_10910	OG1RF_RS04755	58.3549	176.2885	266.4711	17.6531	47.0989	2.0691	-2.8636	6.5904	0.0001	0.0052	CBS domain-containing protein
OG1RF_10911	OG1RF_RS04760	84.8798	183.7902	229.8313	22.6969	41.6644	24.8291	-2.4465	6.6443	0.0000	0.0002	2,3,4,5-tetrahydropyridine-2,6-dicarboxylate N-acetyltransferase
OG1RF_10912	OG1RF_RS04765	92.8373	93.7705	119.9120	17.6531	39.8529	24.8291	-1.8562	6.0907	0.0001	0.0041	acetyldiaminopimelate deacetylase
OG1RF_10993	OG1RF_RS05165	7.9575	30.0066	46.6324	2.5219	1.8115	2.0691	-3.5309	4.2002	0.0001	0.0075	spermidine/putrescine ABC transporter substrate-binding protein
OG1RF_10995	OG1RF_RS05175	23.8724	22.5049	29.9780	5.0437	3.6230	4.1382	-2.5126	4.2291	0.0002	0.0082	amidohydrolase
OG1RF_11012	OG1RF_RS05265	5.3050	3.7508	3.3309	20.1750	23.5494	24.8291	2.3575	4.2929	0.0018	0.0475	DUF5067 domain-containing protein
OG1RF_11036	OG1RF_RS05385	1562.3189	978.9639	769.4354	363.1496	461.9315	333.1240	-1.5149	9.5479	0.0004	0.0164	ATPase P (E1-E2 family cation-transporting ATPase)
OG1RF_11050	OG1RF_RS05450	87.5323	191.2918	173.2062	42.8718	67.0253	55.8655	-1.4239	6.7270	0.0018	0.0475	DNA replication protein DnaI
OG1RF_11058	OG1RF_RS05490	1501.3115	3158.1902	2081.8057	423.6746	742.7133	643.4880	-1.8948	10.4789	0.0000	0.0019	hypothetical protein
OG1RF_11059	OG1RF_RS05495	273.2069	675.1475	516.2878	95.8312	141.2967	142.7674	-1.9372	8.2731	0.0000	0.0020	tail protein
OG1RF_11060	OG1RF_RS05500	713.5208	1462.8197	959.2961	216.8810	355.0532	347.6077	-1.7653	9.4052	0.0001	0.0041	structural protein
OG1RF_11061	OG1RF_RS05505	1042.4301	2085.4557	1488.9074	421.1527	514.4648	562.7934	-1.6212	9.9964	0.0001	0.0065	hypothetical protein
OG1RF_11062	OG1RF_RS05510	363.3917	510.1115	393.0449	148.7905	126.8047	215.1857	-1.3647	8.2088	0.0009	0.0315	holin
OG1RF_11063	OG1RF_RS05515	572.9387	1211.5148	1009.2594	199.2279	251.7979	252.4294	-1.9852	9.1916	0.0000	0.0006	N-acetylmuramoyl-L-alanine amidase
OG1RF_11167	OG1RF_RS06055	106.0998	82.5180	99.9267	274.8841	201.0760	244.1530	1.3095	7.4482	0.0012	0.0362	alanine--tRNA ligase
OG1RF_11246	OG1RF_RS06450	18647.0316	2726.8459	3350.8744	2047.7604	1758.9626	2296.6935	-2.0184	12.3283	0.0014	0.0406	type I glyceraldehyde-3-phosphate dehydrogenase
OG1RF_11269	OG1RF_RS06565	1973.4554	4403.4623	7431.2136	1412.2486	1068.7826	1154.5540	-1.9250	11.5063	0.0001	0.0060	transcriptional regulator (DNA-binding protein HU)
OG1RF_11379	OG1RF_RS07115	1729.4260	1331.5410	1462.2603	615.3369	579.6787	655.9026	-1.2893	10.0579	0.0008	0.0274	oxidoreductase
OG1RF_11400	OG1RF_RS07215	156.4971	101.2721	113.2502	469.0683	210.1335	343.4695	1.4507	7.8997	0.0011	0.0343	adenine phoshoribosyltransferase
OG1RF_11437	OG1RF_RS07400	156.4971	273.8098	406.3685	98.3530	105.0668	88.9710	-1.5094	7.5769	0.0008	0.0279	cold-shock protein
OG1RF_11489	OG1RF_RS07650	2697.5862	1155.2525	1785.3566	52.9593	32.6069	35.1746	-5.5472	9.9098	0.0000	0.0000	phosphoribosylamine--glycine ligase
OG1RF_11490	OG1RF_RS07655	1917.7530	933.9541	909.3327	47.9156	23.5494	55.8655	-4.8856	9.3450	0.0000	0.0000	bifunctional phosphoribosylaminoimidazolecarboxamide formyltransferase/inosine monophosphate cyclohydrolase
OG1RF_11491	OG1RF_RS07660	400.5266	157.5344	173.2062	20.1750	1.8115	10.3455	-4.5388	7.0238	0.0000	0.0000	phosphoribosylglycinamide formyltransferase

OG1RF_11492	OG1RF_RS07665	270.5544	168.7869	179.8680	10.0875	5.4345	4.1382	-4.9605	6.7653	0.0000	0.0000	phosphoribosylformylglycinamidine cyclo-ligase
OG1RF_11493	OG1RF_RS07670	374.0016	206.2951	256.4785	25.2187	10.8690	12.4146	-4.1262	7.2284	0.0000	0.0000	amidophosphoribosyltransferase
OG1RF_11494	OG1RF_RS07675	631.2935	266.3082	266.4711	7.5656	19.9265	22.7600	-4.4845	7.6805	0.0000	0.0000	phosphoribosylformylglycinamidine synthase II
OG1RF_11495	OG1RF_RS07680	196.2845	78.7672	99.9267	12.6094	3.6230	14.4837	-3.6189	6.1465	0.0000	0.0000	phosphoribosylformylglycinamidine synthase subunit PurQ
OG1RF_11496	OG1RF_RS07685	18.5675	22.5049	19.9853	0.0000	0.0000	0.0000	-6.0205	3.7993	0.0000	0.0003	phosphoribosylformylglycinamidine synthase
OG1RF_11497	OG1RF_RS07690	76.9223	18.7541	3.3309	7.5656	0.0000	2.0691	-3.3788	4.4932	0.0015	0.0432	phosphoribosylaminoimidazolesuccinocarboxamide synthase
OG1RF_11498	OG1RF_RS07695	127.3197	37.5082	33.3089	27.7406	5.4345	6.2073	-2.3806	5.4484	0.0015	0.0424	5-(carboxyamino)imidazole ribonucleotide synthase
OG1RF_11506	OG1RF_RS07735	209.5470	311.3180	469.6554	88.2655	157.6001	111.7310	-1.4594	7.8308	0.0012	0.0358	hypothetical protein (putative lipoprotein)
OG1RF_11519	OG1RF_RS07800	34.4824	93.7705	66.6178	10.0875	18.1150	4.1382	-2.5097	5.3395	0.0000	0.0031	universal stress protein A
OG1RF_11564	OG1RF_RS08025	259.9444	367.5803	906.0018	100.8749	110.5013	82.7637	-2.3767	8.2599	0.0000	0.0005	50S ribosomal protein L19
OG1RF_11576	OG1RF_RS08085	37.1349	48.7607	109.9193	22.6969	14.4920	18.6218	-1.8012	5.4978	0.0016	0.0434	membrane protein
OG1RF_11584	OG1RF_RS08125	31.8299	56.2623	66.6178	174.0092	146.7312	167.5966	1.6689	6.8297	0.0002	0.0096	C4-dicarboxylate ABC transporter (C4-dicarboxylate anaerobic carrier)
OG1RF_11677	OG1RF_RS08600	135.2772	150.0328	139.8973	7.5656	12.6805	6.2073	-3.9282	6.2738	0.0000	0.0000	ABC transporter ATP-binding protein
OG1RF_11678	OG1RF_RS08605	108.7522	93.7705	66.6178	7.5656	28.9839	8.2764	-2.5118	5.7973	0.0000	0.0004	membrane protein
OG1RF_11679	OG1RF_RS08610	907.1529	2096.7082	2891.2117	27.7406	27.1724	16.5527	-6.3493	9.9577	0.0000	0.0000	manganese ABC transporter substrate-binding protein
OG1RF_11749	OG1RF_RS08955	973.4652	1204.0131	1825.3272	602.7275	489.1039	546.2406	-1.2886	9.8813	0.0016	0.0442	hypothetical protein
OG1RF_11752	OG1RF_RS08970	42.4399	11.2525	19.9853	131.1374	56.1564	74.4874	1.7552	5.9621	0.0011	0.0342	hypothetical protein
OG1RF_11753	OG1RF_RS08975	458.8814	603.8820	649.5234	1621.5640	1345.9415	1309.7360	1.3213	9.9727	0.0007	0.0268	PTS maltose transporter subunit IIBC
OG1RF_11774	OG1RF_RS09080	63.6599	45.0098	83.2722	292.5372	271.7244	256.5676	2.0848	7.4603	0.0000	0.0002	sugar ABC transporter substrate-binding protein
OG1RF_11775	OG1RF_RS09085	74.2698	41.2590	96.5958	378.2809	239.1175	246.2221	2.0099	7.5416	0.0000	0.0014	glucuronyl hydrolase
OG1RF_11776	OG1RF_RS09090	236.0720	367.5803	546.2658	1518.1672	1068.7826	1222.8341	1.7293	9.7030	0.0001	0.0050	hypothetical protein
OG1RF_11778	OG1RF_RS09100	400.5266	247.5541	326.4271	781.7805	661.1960	989.0266	1.3152	9.1666	0.0012	0.0367	LacI family transcriptional regulator
OG1RF_11794	OG1RF_RS09180	119.3622	26.2557	43.3016	264.7966	210.1335	184.1493	1.7648	7.2150	0.0005	0.0212	hypothetical protein
OG1RF_11860	OG1RF_RS09520	193.6320	135.0295	229.8313	22.6969	32.6069	28.9673	-2.7058	6.7797	0.0000	0.0000	guanosine monophosphate reductase
OG1RF_11976	OG1RF_RS10110	599.4636	303.8164	373.0596	2080.5448	746.3363	1390.4307	1.7199	9.8493	0.0003	0.0140	PTS beta-glucoside transporter subunit EIIBCA
OG1RF_12046	OG1RF_RS10465	448.2715	933.9541	1515.5545	146.2686	447.4395	258.6367	-1.7602	9.2931	0.0009	0.0315	adapter protein MecA
OG1RF_12070	OG1RF_RS10590	39.7874	82.5180	196.5225	25.2187	38.0414	18.6218	-1.9329	6.1199	0.0012	0.0373	hypothetical protein
OG1RF_12078	OG1RF_RS10630	18.5675	26.2557	29.9780	78.1780	65.2139	72.4183	1.5324	5.7721	0.0011	0.0342	AraC family transcriptional regulator
OG1RF_12143	OG1RF_RS10965	68.9648	157.5344	149.8900	55.4812	27.1724	37.2437	-1.6493	6.4149	0.0007	0.0265	DUF3042 domain-containing protein

OG1RF_12243	OG1RF_RS11490	124.6672	161.2852	129.9047	388.3684	260.8554	492.4442	1.4575	8.0563	0.0005	0.0217	D-ribose pyranase
OG1RF_12244	OG1RF_RS11495	302.3843	397.5869	356.4051	859.9585	722.7869	1051.0994	1.3189	9.2800	0.0009	0.0315	ribokinase
OG1RF_12280	OG1RF_RS11675	10.6100	15.0033	9.9927	100.8749	47.0989	57.9346	2.5045	5.5431	0.0000	0.0020	allantoin permease
OG1RF_12310	OG1RF_RS11830	95.4898	75.0164	99.9267	10.0875	52.5334	16.5527	-1.7225	5.9557	0.0013	0.0397	peptidase
OG1RF_12328	OG1RF_RS11925	50.3974	153.7836	303.1109	20.1750	50.7219	26.8982	-2.3421	6.6853	0.0001	0.0080	hypothetical protein
OG1RF_12393	OG1RF_RS12260	164.4546	183.7902	343.0816	73.1343	67.0253	47.5891	-1.8762	7.2214	0.0000	0.0031	hypothetical protein
OG1RF_12425	OG1RF_RS12410	503.9738	341.3246	346.4125	1684.6108	912.9939	1247.6632	1.6862	9.7253	0.0001	0.0044	glycosyl hydrolase
OG1RF_12461	OG1RF_RS12570	47.7449	45.0098	46.6324	110.9624	481.8579	157.2511	2.4236	7.2917	0.0000	0.0023	murein hydrolase regulator LrgA

 Supplementary Table 2. Strains used in this study.

	Strain name	Relevant characteristics	Plasmid Source	Plasmid made	References
***E. faecalis* laboratory strain**	*E. faecalis* OG1RF wild type	Laboratory strain, Rif^R, Fus^R	-	-	[1]
***E. faecalis* markerless, in-frame chromosomal mutants**	OG1RF Δ*ebpABC*	*ebpABC* triple deletion mutant; Rif^R, Fus^R	-	-	[2,3]
	OG1RF Δ*ebpA*	*ebpA* single deletion mutant; Rif^R, Fus^R	-	-	[2,3]
	OG1RF Δ*ebpB*	*ebpB* single deletion mutant; Rif^R, Fus^R	-	-	[2,3]
	OG1RF Δ*ebpC*	*ebpC* single deletion mutant; Rif^R, Fus^R	-	-	[2,3]
	OG1RF *ndh3*::Tn Δ*menB*	*menB* in-frame deletion with insertional mutation of *ndh3*	pGCP213	pGCP213::*delta_menB*	This study
***E. faecalis* chromosomal pilin mutant**	OG1RF ebpA^{AWAGA} (MIDAS)	ebpA allelic replacement with ebpA^{AWAGA} (coding for Ala^{315}-Trp-Ala^{317}-Gly-Ala^{319}); Rif^R, Fus^R	-	-	[2,3]
***E. faecalis* transposon mutants**	OG1RF *menB*::Tn	Rif^R, Fus^R, Cm^R (Insertional position 5' – 3': 343473)	-	-	[4,5]
	OG1RF *ndh3*::Tn	Rif^R, Fus^R, Cm^R (Insertion position 5' – 3': 2658965)	-	-	[4,5]
	OG1RF *ldh1*::Tn	Rif^R, Fus^R, Cm^R (Insertion position 5' – 3': 197152)	-	-	[4-6]
	OG1RF_RS12785-12815::Tn	Rif^R, Fus^R, Cm^R (*RS12785* Insertion position 5' – 3': 2654550) (*RS12795* Insertion position 5' – 3': 2657169) (*RS12800* Insertion position 5' – 3': 2658965) (*RS12805* Insertion position 5' – 3': 2660151) (*RS12810* Insertion position 5' – 3': 2660397) (*RS12815* Insertion position 5' – 3': 2660975)	-	-	[4,5]
	OG1RF_RS01800-01825::Tn	Rif^R, Fus^R, Cm^R (*RS01800* Insertion position 5' – 3': 343437) (*RS01805* Insertion position 5' – 3': 343749) (*RS01820* Insertion position 5' – 3': 348095) (*RS01825* Insertion position 5' – 3': 349226)	-	-	[4,5]
***E. faecalis* strains with empty vector**	OG1RF Δ*ebpABC*	*ebpABC* triple deletion mutant; Rif^R, Fus^R, Kan^R	pGCP123	-	[2,3]
	OG1RF wild type	Laboratory strain, Rif^R, Fus^R, Kan^R	pGCP123	-	This study

	OG1RF *menB*::Tn	Rif[R], Fus[R], Erm[R]	pMSP3535	-	This study
	OG1RF *ndh3*::Tn	Rif[R], Fus[R], Erm[R]	pMSP3535	-	This study
	OG1RF *ldh1*::Tn	Rif[R], Fus[R], Erm[R]	pMSP3535	-	[6]
E. faecalis **mutants**	OG1RF *ΔebpABC*	Complement mutation; Rif[R], Fus[R] ,Kan[R]	*gcp123::ebpABCsrtC*		[2,3]
with gene	OG1RF *menB*::Tn	Complement mutation; Rif[R], Fus[R] ,Erm[R]	pMSP3535	pMSP3535::*menB*	This study
complementation	OG1RF *ndh3*::Tn	Complement mutation; Rif[R], Fus[R] ,Erm[R]	pMSP3535	pMSP3535::*ndh3*	This study
	OG1RF *ldh1*::Tn	Complement mutation; Rif[R], Fus[R] ,Erm[R]	pMSP3535	pMSP3535::*ldh1*	[6]

1171

1172

1   Bourgogne, A. *et al.* Large scale variation in *Enterococcus faecalis* illustrated by the genome analysis of strain OG1RF. *Genome biology* **9**, R110,
doi:10.1186/gb-2008-9-7-r110 (2008).

2   Nielsen, H. V. *et al.* Pilin and sortase residues critical for endocarditis- and biofilm-associated pilus biogenesis in *Enterococcus faecalis*. *Journal of*
bacteriology **195**, 4484-4495, doi:10.1128/jb.00451-13 (2013).

3   Nielsen, H. V. *et al.* The metal ion-dependent adhesion site motif of the *Enterococcus faecalis* EbpA pilin mediates pilus function in catheter-associated
urinary tract infection. *mBio* **3**, e00177-00112, doi:10.1128/mBio.00177-12 (2012).

4   Kristich, C. J. *et al.* Development and use of an efficient system for random mariner transposon mutagenesis to identify novel genetic determinants
of biofilm formation in the core *Enterococcus faecalis* genome. *Appl Environ Microbiol* **74**, 3377-3386, doi:10.1128/aem.02665-07 (2008).

5   Dale, J. L. *et al.* Comprehensive Functional Analysis of the *Enterococcus faecalis* Core Genome Using an Ordered, Sequence-Defined Collection of
Insertional Mutations in Strain OG1RF. *mSystems* **3**, doi:10.1128/mSystems.00062-18 (2018).

6   Keogh, D. *et al.* Extracellular Electron Transfer Powers *Enterococcus faecalis* Biofilm Metabolism. *mBio* **9**, doi:10.1128/mBio.00626-17 (2018).

1184

1185

1186

1187

1188

1189

1190 **Supplementary Table 3. Primers used in this study.**

	Primer name	Primer sequence (5' → 3')	Restriction sites[a]	References
Primers used to generate complement plasmids	ndh3_F'	GGATCCAAGGAGCGGATTGTCAA	BamHI	This study
	ndh3_R'	ACTAGTGCTCGTAGTTGTCAATCGAT	SpeI	This study
	ndh3_F'_Infusion	GACTCTGCATGGATCCGGATCCAAGGAGCGGAT	BamHI	This study
	ndh3_R'_Infusion	GCAGCCCGGGACTAGTACTAGTGCTCGTAGTTGTCAA	SpeI	This study
	menB_F'	GGATCCTTTCTCTAAAGTGCCGAT	BamHI	This study
	menB_R'	ACTAGTTGTTTATTTAACCAAGTCATTGG	SpeI	This study
	menB_F'_Infusion	GACTCTGCATGGATCCGGATCCTTTCTCTAAAGTGCC	BamHI	This study
	menB_R'_Infusion	GCAGCCCGGGACTAGTACTAGTTGTTTATTTAACCAAGTCATT	SpeI	This study
Primers used to generate chromosomal deletion strain	IFD_menB_Frag1_F'	TCTCTAAAACCAATGACTTGGTTAAATAAAC	-	This study
	IFD_menB_Frag1_R'	GGCGGCCGTTACTAGTCAAATGCGTATTTAAAAATAATAACTCTTTCC	SpeI	This study
	IFD_menB_Frag2_F'	TACCGAGCTCGGATCCAAGATGCCGTTTAAGCCTC	BamHI	This study
	IFD_menB_Frag2_R'	CATTGGTTTTAGAGAAAAAGTCGCTTGAATTTTAAGA	-	This study
Primers used in qRT-PCR	gyrA_F'	TGTTCGTCGGGATGTGAGTG		[1,2]
	gyrA_R'	GGTACGCCTTTTTCGATGGC		[1,2]
	ebpR_F'	TAAACAGCGTTGGGGCGAAA		[1]
	ebpR_R'	TGGGTGGTCGTTGACGTTTT		[1]
	ebpC_F'	CGGTCATACCGACGACCAAA		[1]
	ebpC_R'	TGTCACATCGCCATCGACTT		[1]
	srtA_F'	TCGTACGCCGTTAGCAAGTT		This study
	srtA_R'	TTCATCACCGCTTCTGTGCT		This study
	srtC_F'	ACACATGCGGTCATTTCAGG		This study
	srtC_R'	GCGTCTTCCCATTGACTTCG		This study
	argR3_F'	CCACTATTTACAAGAAAAGGGCGT		This study
	argR3_R'	CGCGAGAAATCGTTGCTTGT		This study

1191 [a] Restriction sites are underlined in the primer sequence.

1192

1193   1   Afonina, I., Lim, X. N., Tan, R. & Kline, K. A. Planktonic Interference and Biofilm Alliance between Aggregation Substance and Endocarditis- and Biofilm-
1194   Associated Pili in *Enterococcus faecalis*. *Journal of bacteriology* **200**, doi:10.1128/jb.00361-18 (2018).
1195   2   Bourgogne, A. *et al.* EbpR is important for biofilm formation by activating expression of the endocarditis and biofilm-associated pilus operon (ebpABC)
1196   of *Enterococcus faecalis* OG1RF. *Journal of bacteriology* **189**, 6490-6493, doi:10.1128/jb.00594-07 (2007).

1197