ADDITIVE GENETIC VARIANCE FOR LIFETIME FITNESS AND THE

CAPACITY FOR ADAPTATION IN THE WILD

Mason W. Kulbaba[1], Seema N. Sheth[1,2], Rachel E. Pain[1], Vincent M. Eckhart[3], Ruth G. Shaw[1]

[1] Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, MN, USA.

[2] Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC,

USA. [3] Department of Biology, Grinnell College, Grinnell, IA, USA

Abstract

The immediate capacity for adaptation under current environmental conditions is directly proportional to the additive genetic variance for fitness, $V_A(W)$. Mean fitness, $\overline{W}$, is predicted to change at the rate $\frac{V_A(W)}{\overline{W}}$, according to Fisher's Fundamental Theorem of Natural Selection. Despite ample research evaluating degree of local adaptation, direct assessment of $V_A(W)$ and the capacity for ongoing adaptation is exceedingly rare. We estimated $V_A(W)$ and total lifetime fitness in three pedigreed populations of the annual *Chamaecrista fasciculata*, over three consecutive years in the wild. Contrasting with prevailing expectations of minimal $V_A(W)$, we found significant $V_A(W)$ in all populations and years, predicting increased mean fitness (0.83 to 6.12 seeds per individual). Further, we identified two instances predicting "evolutionary rescue", where selection on standing $V_A(W)$ was expected to increase fitness of declining populations ($\overline{W} < 1.0$) to levels consistent with population sustainability and growth. Within populations, interannual differences in the genetic expression of fitness were striking. Significant genotype-by-year interactions reflected modest correlations between breeding values across years (all $r < 0.490$), indicating temporally variable selection contributes to maintaining $V_A(W)$. By directly estimating $V_A(W)$ and total lifetime $\overline{W}$, our study presents an experimental approach for future studies of adaptive capacity in the wild.

Introduction

A population's mean absolute fitness, in terms of the per capita contribution of offspring, is a measure of its degree of adaptation and corresponds to its growth rate. The change in mean population fitness from one generation to the next represents the rate of adaptation. When the timing of abrupt environmental change is documented, rapid genetic differentiation and adaptation to changes in local conditions have been observed in response to edaphic conditions (Antonovics and Bradshaw 1970; Al-Hiyaly et al. 1993), herbicides (Vigueira et al. 2013; Baucom 2019), predation (Fisk et al. 2007), and climate (Franks et al. 2007; Geerts et al. 2015) across a variety of taxa. These examples of relatively rapid local adaptation reveal a propensity for adaptation to environmental differences (but see Leimu and Fischer 2008) from standing genetic variation (Barrett and Schluter 2008). However, such studies are limited to characterizing current adaptation in response to past selection and do not evaluate the potential for ongoing adaptation. Predicting future adaptation under ongoing natural selection, not only whether to expect adaptation but also the rate of fitness increase, requires direct, quantitative estimates of standing additive genetic variance for fitness, $V_A(W)$. Studies that take such a direct and predictive approach are exceedingly rare (but see Etterson 2004a,b; Sheth et al. 2018), limiting our understanding of the adaptive process in the wild.

The immediate capacity for adaptation is proportional to the magnitude of a population's additive genetic variance for fitness (Fisher 1930; Ewens 2004). Fisher's Fundamental Theorem of Natural Selection (Fisher 1930) predicts that the rate of adaptation under prevailing environmental conditions as the ratio of $V_A(W)$ to mean population fitness, $\overline{W}$. With fitness evaluated as an individuals' lifetime contribution of offspring to the population, this ratio predicts the genetically based change, from one generation to the next, in the per capita rate of

population growth. Such predictions are vital to understanding the adaptive process, by first

determining the possibility for any change in mean fitness, and second, the expected magnitude

of fitness increase in the wild. In addition, estimates of the magnitude of $V_A(W)$ are critical for

predicting the genetically-based response to shifting patterns of natural selection that accompany

environmental change. Given the elevated threat to population persistence due to rapid change in

climate and other aspects of the environment, quantifying the opportunity and capacity for

adaptation, and population stability, has become especially pressing.

Despite their importance, very few direct estimates of $V_A(W)$ exist (but see Etterson

2004a,b; Winn 2004; Sheth et al. 2018). Several reasons account for this paucity of empirical

research. First, population mean fitness is often thought to be at or near optimum due to incessant

selection, which would tend to fix or eliminate alleles having directional effects on fitness.

Accordingly, $V_A(W)$, and therefore the capacity for adaptation, is conjectured to be reduced to

low or negligible levels (Mazer 1987; Barton and Keightley 2002). However, even with

consistent and strong selection at individual loci, fixation requires hundreds of generations (e.g.,

Fig. 6.3 in Hartl and Clark 1997; Messer et al. 2016). The continuing response to selection in the

Illinois corn selection experiment (Moose et al. 2004), even after 100 generations, demonstrates

the persistence of variation supporting a response to selection. However, the resilience of

variation under selection in nature remains an open empirical question. Second, numerous

studies have found that estimates of narrow-sense heritability $\left(\frac{V_A}{V_P}\right)$ for individual fitness

components tend to be lower than for morphological traits (Roff and Mousseau 1987; McFarlane

et al. 2014). Because environmental variation inherent to fitness components is compounded

with environmental variance of underlying morphological traits, high environmental variance,

rather than low additive genetic variance, may explain low heritability of fitness components

(Price and Schluter 1991). Nevertheless, this empirical generalization about individual

components of fitness contributed to the expectation that additive genetic variance for total

fitness is likely to be modest. This expectation may, however, be inaccurate because it does not

account for the dependence of sequential fitness components through the life-span. Third, when

pedigrees are not experimentally determined and may be uncertain (e.g. Foerster et al. 2007),

logistical challenges encumber efforts to both reconstruct relatedness of individuals and

determine total lifetime fitness. Uncertainty in the assignment of relatedness casts doubt on the

estimates of quantitative genetic parameters (Thomas et al. 2000). Finally, confounding between

genetic and environmental influences, likely in observational studies, can also bias $V_A(W)$

estimates (Pemberton 2010), and therefore inferences drawn from them.

Whereas direct estimates of $V_A(W)$ in the wild are rare, theory bearing on whether

substantial genetic variation could be maintained is ambiguous. In particular, early theory

addressing the role of environmental variation, particularly temporal variability, indicated that

rather restrictive conditions are required to maintain genetic variation within populations

(Felsenstein 1976; Hedrick 1986; Gillespie and Turelli 1989). More recently, models focusing on

$V_A(W)$ under environmentally varying selection have demonstrated maintenance of substantial

$V_A(W)$ (Zhang 2012; Shaw and Shaw 2014; see also Yi and Dean 2013).

Predicting the rate of adaptation based on mean fitness and $V_A(W)$ is not only vital to

understanding ongoing adaptation in wild populations, but also in assessing whether standing

levels of $V_A(W)$ suffice for contemporary natural selection in declining populations (i.e. $\overline{W} < 1$)

to halt further decline (Bradshaw 1991). Such "evolutionary rescue" (Gomulkiewicz and Holt

1995) that increases fitness to 1 (i.e. individuals demographically replacing themselves) or

greater depends on current standing $V_A(W)$. However, a sufficient baseline of $V_A(W)$ for such

fitness increase is unknown, particularly in wild populations that undergo environmental

fluctuations (Carlson et al. 2014). Given the increasing rate of change in climate and other

aspects of environment and their effects on natural populations (Shaw and Etterson 2012;

Neumann et al. 2017), it is imperative to clarify the efficacy of selection in maintaining and

enhancing population fitness *in situ.*

Direct estimation of $V_A(W)$ is not a trivial endeavor, requiring sound experimental design

and statistical inference. Designs for organisms that facilitate estimates of complete lifetime

fitness expressions from fully pedigreed individuals, and avoid confounding environmental and

genetic effects are essential (Shaw 2019). Further, statistical tools are needed that can

accommodate fitness measures that seldom approach conventional statistical distributions, and

incorporate the dependent nature of sequential life-history fitness expressions that culminate in

total lifetime fitness (Shaw et al. 2008). Aster models incorporate such consecutive fitness

expressions, including their unconventional distributions, to estimate total mean lifetime fitness

(Geyer et al. 2007), and random parental effects (Geyer et al. 2013) to estimate the additive

genetic effects for lifetime fitness.

In this study, we estimate the capacity for adaptation under contemporary natural

selection in wild populations of the annual legume *Chamaecrista fasciculata* growing in their

native locations. Using Fisher's Fundamental Theorem (Fisher 1930) and aster modelling to

analyze records of lifetime fitness (Geyer et al. 2007, 2013), we directly estimate $V_A(W)$ and $\overline{W}$

to quantitatively predict the rate of adaptation. To assess how populations differ in these

attributes, and how each differs among years, we conducted our experiment in three populations

in three successive years. Our experimental design allowed us to study the rate of future

adaptation and to characterize population-specific expression of $V_A(W)$ and temporal variation in

the expression of $V_A(W)$, and therefore variation in the capacity for adaptation, in three

populations over three years.

Material and Methods

Study system and pedigree generation

We estimated $V_A(W)$ and $\overline{W}$ as the basis for predicting the generational change in mean

fitness in three populations (Fig. 1) of the annual legume *Chamaecrista fasciculata,* each

growing in its home site. The range of *C. fasciculata* spans from the prairies of the Northern

Great Plains in Minnesota to Central Mexico and to the eastern seaboard of North America

(Irwin and Barneby 1982). *C. fasciculata* produces hermaphroditic, enantiostylous flowers that

are buzz-pollinated (Lee and Bazzaz 1982) by bumblebees (*Bombus* spp). Its experimental

tractability has made it the focus of previous studies of gene flow (Fenster 1991a,b), quantitative

genetics (Kelly 1993; Etterson 2004b), selection on traits (Etterson 2004a), local adaptation

(Galloway and Fenster 2000), adaptation to climate change (Etterson and Shaw 2001), evolution

of range limits (Stanton-Geddes et al. 2012, 2013), and genotype-by-environment interactions

(Sheth et al. 2018).

Genetic design

In the fall of 2012, 2-3 fruits from 200 maternal plants were collected from a population

of *C. fasciculata at* Kellogg-Weaver Dunes in Minnesota (44°15'43" N, 91°55'15" W; hereafter

"McCarthy Lake"). Similarly the following fall, fruits from 200 maternal plants were collected

from populations of *C. fasciculata* at Grey Cloud Dunes Scientific and Natural Area in

Minnesota (44°46'32"N, 93°01'38"W; hereafter "Grey Cloud"), and 100 plants from the Conard

Environmental Research Area in Iowa (41°40'44.2"N 92°51'24.9"W; hereafter "CERA").

To obtain pedigreed populations of seeds representing each of these populations, six

seeds from each maternal plant were surface sterilized using an 8.9% bleach solution followed

by a 70% ethanol rinse. We used 100 grit sandpaper to scarify the seeds, imbibed them in sterile

water for three days, and then planted them in small peat pots. A total of 167 Grey Cloud, 196

McCarthy Lake, and 84 CERA individuals were used to generate pedigreed plants. One seedling

per individual was used to represent each distinct family and was transplanted into a large tree

pot with 1 teaspoon of Osmocote ® slow release fertilizer (14:14:14) after producing five true

leaves. Plants were grown in the University of Minnesota Plant Growth Facilities greenhouse

under 16:8-h photoperiod. Hand crosses were made according to a nested paternal half-sibling

design (North Carolina Design I; Comstock and Robinson 1948), with independent sets of three

dams randomly assigned to sires. Hand crosses were performed daily, and all non-pollinated

flowers were removed. After accounting for mortality in the greenhouse and planting logistics,

the final pedigree consisted of 42 paternal half-sibling families and 124 maternal full-sibling

families at Grey Cloud, 48 paternal half-sibling families and 132 maternal full-sibling families at

McCarthy Lake, and 21 paternal half-sibling families and 60 maternal full-sibling families at

CERA.

Field planting design

In October-November of each of three consecutive years (2014-2016), at each site of

origin, pedigreed seeds from each respective population were planted into restored tallgrass

prairie. To maintain the environment for the experimental populations as realistic as possible,

vegetation was left in place and only mown (or burned in the case of CERA in 2016 and

McCarthy Lake in 2014) in the fall to facilitate planting. Within a randomly chosen subset of

rows within blocks, seeds were planted at randomly chosen positions according to the following

protocol. Seeds from each full-sibling family from the greenhouse crosses were combined and

haphazardly distributed into envelopes, such that each envelope contained five seeds. We

randomly assigned each envelope to a position along 50 m rows at least two meters apart in a

randomized block design (eight blocks at Grey Cloud and McCarthy Lake, and four blocks at

CERA). Planting positions within rows were marked by nails at two-meter intervals (i.e. 25

marked positions per row). At each position centered at the nail, seeds were planted at 10 cm

spacing, approximately 1 cm under the soil surface.

Fitness surveys

We regularly recorded fitness components for each planted seed throughout the growing

season of the following year, including emergence, survival to flowering, and reproductive

output (Table 1). A plant was considered present if cotyledons had emerged, and survival to

flowering was scored based on the presence of open flowers or evidence of flowering (e.g.,

wilted flowers, pedicels, or fruit). At each late-season census (i.e. after flowering), we recorded

the number of ripe fruit collected, number of fully elongated but immature fruits collected,

number of fruits collected off the ground in the immediate vicinity, number of immature fruits,

and pedicels from dehisced fruits remaining on the plant (see Sheth et al. 2018). Herbivory,

ranging from light browsing to consumption of the entire plant, or severing of the stem near the

soil surface, was recorded throughout the growing season. When herbivory occurred early in the

growing season (i.e. June-July), plants sometimes recovered and eventually reproduced.

Therefore, we retained these plants in censuses after herbivory to account for this late season

seed production. Because individuals ranged widely in number of fruits set (1-68), that

explosively dehisce at maturity, we were unable to collect all mature fruits or count all seeds

each plant produced. Therefore, seed counts were obtained on a subsample of fruits produced.

Subsampling was accounted for in the statistical analyses (see below).

Statistical Analyses

We used aster models (Geyer et al. 2007; Shaw et al. 2008) to estimate mean lifetime

fitness, as mean number of seeds produced per seed planted, and the additive genetic variance for

lifetime fitness for each population in each year. Unconditional aster analyses, as implemented in

the package "aster" (Geyer 2018), were used in the R 3.5.1 environment (R Core Team 2018).

The graphical model for lifetime fitness (Fig. 1) included seedling emergence and survival of

plants to flowering, which were modeled as Bernoulli variables, and the number of fruits and

seeds produced, which were modeled with Poisson distributions. To account for subsampling of

fruits and scale up to total seed production, we followed (Stanton-Geddes et al. 2012),

incorporating in our graphical models an additional node representing the fraction of fruits

sampled.

For each population in each year, we conducted separate aster analyses to estimate

$V_A(W)$, including as predictors parental (genetic) effects and block. Only parental effects were

treated as random, as in Sheth et al. (2018), following advice to minimize the use of random

factors in GLMM, including aster models (Bolker et al. 2009; Geyer et al. 2013). From separate

analyses, also including blocks as fixed effects, we estimated $\bar{W}$. In addition, to test whether

parental genetic effects differed among years, we jointly analyzed the data for cohorts grown in

all three years, including year as a fixed factor and parental (genetic) effects and the interaction

between parental effects and year (genotype-by-year) as random factors. We tested fixed factors

with likelihood ratio tests of the full model vs. models excluding each of these factors. For

estimates of additive genetic variance for lifetime fitness, we constructed models that equated

dam and sire effects (hereafter parental effects) after confirming statistical significance of sire

and dam effects separately (see Sheth et al. 2018).

For each population-year combination, mean fitness and standard error were represented

by the median estimate across all blocks. To obtain estimates of $V_A(W)$ on the measurement

scale (i.e. number of seeds produced per seed planted), we performed non-linear transformations

of estimates of $V_A(W)$ from the canonical parameter scale to the mean-value parameter scale

using a mapping function that adds a component of a fixed effect (see Table 3 for relevant

blocks) to each random parental effect (Geyer and Shaw 2013; de Villemereuil et al. 2016). The

resulting values represent breeding values on the mean-value parameter (i.e. measurement) scale

and are used to estimate $V_A(W)$. Predicted changes in fitness from Fisher's Fundamental

Theorem of Natural Selection $\left(\frac{V_A(W)}{\overline{W}}\right)$ were added to the estimates of mean fitness ($\overline{W}$) to

generate the predicted mean fitness of the progeny generation. Apart from the possibility of seed

dormancy, our fitness records cover the entire life history of individual plants, and therefore

represent estimates of total absolute fitness over individual life spans.

To compare the genetic effects on fitness (breeding values) between years, and the

potential for temporal fitness trade-offs, we obtained breeding values for each sire in each year

and plotted each sire's value in one year against its breeding value in another for all three

combinations of years. We also calculated Pearson's product-moment correlation coefficients

between breeding values expressed in different years. We emphasize that these coefficients do

not account for the uncertainty associated with estimates of breeding values; we therefore present

these correlation coefficients as a coarse description of association between breeding values

expressed in different years. All data and scripts associated with this study are available at

https://github.com/mason-kulbaba/adaptive-capacity.git

Results

Expression of additive genetic variance for lifetime fitness

We found highly significant additive genetic variance for lifetime fitness, $V_A(W)$, in all three years for all three populations, as reflected by significant parental effects in all models (Table 2); block effects were also significant in all cases, indicating spatial variation in expression of fitness. Estimates of $V_A(W)$ varied widely among years and populations, with the smallest and largest estimates of $V_A(W)$ found in consecutive years for the Grey Cloud population ($V_A(W) = 0.631$ in 2016, and 6.491 in 2017). The findings of significant and substantial estimates of $V_A(W)$ (Table 3) indicate ample basis for the genetic response to natural selection based on standing $V_A(W)$. Probability distributions of breeding values for each of the nine cases are displayed in Fig. 2.

Significant genotype-by-year interactions for each population (Table 2) indicated that the expression of parental genetic effects varied among years. Year-specific breeding values within each population revealed large differences between years in family-specific expressions of total lifetime fitness (Fig. 3). Breeding values exhibited small to moderate correlations between years, with two cases of modest negative correlations: 2015 vs. 2016 in McCarthy Lake and 2016 vs. 2017 in CERA (Fig. 3). The findings of significant genotype-by-year interactions indicate that families contributing disproportionately to the next generation often differ among years.

Mean fitness and predicted changes in mean fitness

As with estimates of $V_A(W)$, estimates of mean fitness and predicted rates of adaptation varied among populations and years (Table 3; Fig. 4). Mean fitness was highest in the McCarthy Lake site in 2015 (mean and standard error: 3.528 ± 1.227) and smallest in Grey Cloud in 2016

(0.640 ± 0.129). This and one other instance (CERA mean and standard error: 0.725 ± 0.112)

indicating declining populations (i.e. $\overline{W} < 1$) were found in the same year.

The predicted change in mean fitness given by Fisher's Fundamental Theorem $\left(\frac{V_A\,(W)}{\overline{W}}\right)$,

varied with the magnitude of both $\overline{W}$ and $V_A(W)$. The largest increase in mean fitness was

predicted for the progeny of Grey Cloud plants after 2017 (mean increase of 6.122 seeds per

plant) and the smallest for progeny of McCarthy Lake plants after 2016 (mean increase of 0.834

seeds per plant). We identified two predictions of evolutionary rescue for the progeny

generations of 2016 in Grey Cloud and CERA. This predicted increase in mean fitness is

consistent with population sustainability and growth of the progeny generations (Table 3; Fig. 4).

Discussion

The immediate capacity for population-level adaptation is strongly dependent on the

presence and magnitude of additive genetic variance for lifetime fitness (Fisher 1930; Ewens

2004). Our study detected significant and substantial capacity for immediate adaptation through

current natural selection on standing levels of $V_A(W)$ in three populations in three years. Among

these nine cases, we detected two instances of declining populations, and in those cases obtained

predictions of evolutionary rescue increasing mean fitness to levels consistent with maintaining

or increasing population size. Genotype-by-year interactions indicated that genetic effects on

fitness differed among years. Estimates of the magnitude of $V_A(W)$ and predicted change in

mean fitness also varied among years, suggesting that the capacity for adaptation is strongly

influenced by temporal environmental variation. Below, we discuss the importance of these

findings and consider how our results from direct study of the adaptive process in the wild

illuminate the potential for ongoing adaptation. We also relate our findings to the potential for

evolutionary rescue of declining populations. Finally, we describe how our experimental and

analytical approaches overcome obstacles commonly associated with directly estimating

adaptive capacity in the wild, and we advocate for their broader implementation.

Expression of additive genetic variance for lifetime fitness

Our detection of prevalent and substantial additive genetic variance for lifetime fitness,

characterized as the number of seeds produced per seed planted, indicates differential genetic

contributions to the progeny generation and, via the Fundamental Theorem of Natural Selection,

abundant capacity for ongoing adaptation under prevailing natural selection. We estimated

significant $V_A(W)$ in the home site of all three populations in all three years of study (Fig. 2),

albeit at varying magnitudes. These results contrast with those of several studies that conclude

limited capacity for adaptation, through estimates of heritability of fecundity (e.g. Kruuk et al.

2000), or other individual fitness components (Mousseau and Roff 1987; Matos et al. 2000;

McCleery et al. 2004; Teplitsky et al. 2009). We suggest that these studies can be reconciled

with our findings in part by recognizing that even when additive genetic variances of individual

fitness components are modest, estimates of $V_A(W)$ may be considerably larger due to the

compounding of fitness components through an individual's life-history (Shaw et al. 2008; Shaw

and Geyer 2010). This compounding plays out over the entire life-history, regardless of the

extent to which there are genetically based trade-offs between individual components of fitness

(e.g. Rose and Charlesworth 1981). Our analysis of total lifetime fitness through aster modeling

enabled us to account for the dependent nature of sequential fitness components in our estimates

of total fitness.

Our finding of prevalent interactions between genotype and year reflects two aspects of

environmental dependence of genetic expression. First, correlations between years of family-

specific breeding values for lifetime fitness were generally low and, in two cases, slightly

negative (Fig. 3). Thus, genotypic contributions of offspring in one year are not predictive of

their contributions in another year. Interestingly, the two strongest genetic correlations between

years were not for consecutive years. These modest between-year genetic correlations suggest

that the response to selection may not be accompanied by directional change in the frequency of

the same alleles over years. Second, for each population, estimates of $V_A(W)$ differed strikingly

among years (Table 3), though we acknowledge substantial uncertainty in the estimates.

Nevertheless, these differences in estimated $V_A(W)$ suggest that the immediate capacity for

ongoing adaptation varies among years. Almost certainly, many differences, whether slight or

major, distinguished the environments among years; which of these influenced genetic disparities

in fitness remains unknown.

325   Whereas we found considerable $V_A(W)$ for multiple populations, explanations for the

maintenance of such potential variation has long been elusive (e.g. Burt 1995). One potential

explanation is that trade-offs between individual fitness components maintain additive genetic

variation (Barton and Keightley 2002). Previous work on *C. fasciculata* revealed positive

correlations of breeding values between fitness components within and between two

environments (Sheth et al. 2018). This result suggests that trade-offs (i.e. negative genetic

correlations) do not account for the maintenance of observed levels of $V_A(W)$.

332   Temporal environmental variation has long been conjectured to maintain genetic

variation, but the theoretical conditions that predict this outcome are restrictive (Felsenstein

1976). Experiments manipulating environmental variation have led to conflicting conclusions on

the role of environmental variation in the maintenance of genetic variation (Mackay 1980;

Yeaman et al. 2010; Huang et al. 2015). Through estimates of total lifetime fitness, instead of

individual traits or components of fitness, our results have documented extensive temporal

variation in the expression of breeding values within populations (Fig. 2; Fig. 3). Given the short

timeframe of our study (i.e. single generation replicated over three years), the impact of novel

mutations on lifetime fitness would be negligible. Additional phenomena may also contribute to

the conservation of genetic variation (e.g., marginal overdominance; Levene 1953; Gillespie

1984), but the differential and variable genetic contribution of families to the following progeny

generation across years (Fig. 3) is consistent with a role for temporal environmental variation in

maintaining $V_A(W)$. These annual variations may contribute to the maintenance of the

appreciable levels of observed $V_A(W)$ in our study.

Mean fitness and predicted changes in mean fitness

Similar to estimates of $V_A(W)$, mean fitness and the predicted rate of adaptation across

generations varied with population and year. Predictions based on Fisher's Fundamental

Theorem $\left(\frac{V_A(W)}{\overline{W}}\right)$ indicated that even the more modest estimates of $V_A(W)$ (e.g., Grey Cloud

and CERA in 2016; Table 3) were sufficient to increase mean fitness under contemporary natural

selection. However, when estimates of $V_A(W)$ were large (e.g. Grey Cloud in 2017),

contemporary selection was predicted to drastically increase fitness to several times that of the

parental generation (Table 3, Fig. 4).

Our study identified two cases where evolutionary rescue was predicted to increase mean

fitness of declining populations to levels consistent with sustainability and even population

growth. Mean fitness of both the Grey Cloud and CERA populations during the 2016 growing

season was less than 1.0, indicating declining numbers of individuals to the following year. The

prediction of change in mean fitness, however, is that fitness in the progeny generation would

increase to more than double the current mean fitness. Realization of this prediction would not

only prevent further population decline but would elevate seed production sufficiently to result in

population growth. Whereas evolutionary rescue has been directly demonstrated in controlled

laboratory systems (Bell and Gonzalez 2009, 2011), examples from wild systems are lacking.

Beginning with Gomulkiewicz and Holt (1995), mathematical modeling efforts have described

the likelihood of evolutionary rescue under various demographic and environmental scenarios,

but the promise of evolutionary rescue of declining and threatened populations remains unclear

(Bell 2017). Therefore, along with Gomulkiewicz and Shaw (2012) we advocate for further

empirically-based predictions of adaptation, as presented in our study, to elucidate the potential

of evolutionary rescue in the wild.

Our assessments of fitness encompass components of fitness expressed across the entire

life span of individuals, from each seed planted through to the seeds it produced; these fitness

evaluations are thus uncommonly complete. Nevertheless, some aspects of fitness are not

included. For example, this study did not account for fitness realized through siring of seeds

(male fecundity). In a companion study, however, Kulbaba and Shaw (in review) found strongly

positive genetic correlations between lifetime fitness measured as maternal contributions (seeds

set) and lifetime fitness measured through paternal contributions (seeds sired), regardless of

population density and genetic relatedness among individuals. We therefore expect paternal-

specific fitness to scale proportionately with maternal-specific fitness. A second omission from

our fitness estimates is the potential for a delayed contribution to fitness due to seed dormancy.

Partial surveys of germination in later years found low emergence from seeds originally planted,

representing a modest proportion of each cohort. Moreover, in increasing populations, (i.e. $\overline{W} >$

1), as we found in most cases, delays in an individual's life-cycle are expected to have a

discounting effect on its contributions to fitness (Roughgarden 1996, Eqs. 18.31, p. 331). On the

temporal scale of adaptation of annual populations between consecutive years, which is our

present focus, it is valid to neglect such delays. To address ongoing adaptation on longer time

scales, further studies that include instances of delayed germination in estimates of fitness, with

appropriate discounting, as in Eck et al. (2015) would be worthwhile. This is especially so, in

view of our finding of weak correspondence of genetic selection between consecutive years (Fig.

3).

Our study has generated precise quantitative predictions of fitness increase for declining

populations. However, correspondence between predicted and realized fitness changes across

generations remains to be assessed. We are currently evaluating this correspondence to address

the accuracy of predictions of the rate of adaptation from Fisher's Fundamental Theorem for

these populations.

Conclusions

Whereas numerous authors invoke relentless selection to explain limited heritability of

total fitness, or traits closely related to fitness (Kruuk et al. 2000; Coltman et al. 2005;

McFarlane et al. 2014; de Villemereuil et al. 2019), our results demonstrate the persistence of

high $V_A(W)$ despite contemporary selection. Significant interactions between genotype and year

reflected modest correlations between years of family-specific breeding values for lifetime

fitness, as well as estimates of $V_A(W)$ that varied widely across years. Along with estimates of

$V_A(W)$, mean expected fitness varied among sites and years. Using Fisher's Fundamental

Theorem, we predicted the rate of adaptation, and two instances of evolutionary rescue from

standing $V_A(W)$ under contemporary natural selection. Our results reveal the power and utility of

Fisher's Fundamental Theorem to identify the capacity for adaptation, and to generate

quantitative predictions for fitness increase in the wild.

Our experimental approach in combination with aster analyses for total lifetime fitness

provided a direct estimate of $V_A(W)$ and fitness change (Shaw 2019). Many studies concerning

the potential for adaptation rely on indirect estimates of fitness, and often focus on one or a few

individual traits or fitness components, thus omitting the role of multiple life-history fitness

expressions. Our approach to evaluating total lifetime fitness with aster models provides a more

complete picture of fitness and $V_A(W)$. We recommend our empirical and analytical strategy for

future studies of other organisms having different life-histories to more fully describe the

capacity for ongoing adaptation.

Acknowledgements

We thank land managers: Larissa Mottl, Don Ramsden, Karen Schik, and Elizabeth Hill

for site preparation. Amber Nashoba, Sam Weaver, Carlee Steppe, Becca Tucker, Bailey

Choudhury, Linley Davidson, William Peterson, Sarah Jordan, Greg Margida, Sam Sokolosky,

Aidan Healey, Christine Solomon, Tim Burnette, Ana Pulak, and members of the Grinnell

College BIO 305 (Evolution of the Iowa Flora) and BIO 368 (Ecology) classes supplied critical

greenhouse and field assistance. We are grateful to Charles Geyer and Frank Shaw for guidance

with aster modeling. This work was funded by a National Science Foundation grant (DEB

1257462) to RGS. USDA National Institute of Food and Agriculture Hatch project 1016272

supported SNS during the writing stages of this work.

Literature Cited

428   Al-Hiyaly, S. A. K., T. McNeilly, A. D. Bradshaw, and A. M. Mortimer. 1993. The effect of zinc

429   contamination from electricity pylons. Genetic constraints on selection for zinc tolerance.

430   Heredity 70:22–32.

431   Antonovics, J., and A. D. Bradshaw. 1970. Evolution in closely adjacent plant populations VIII.

432   Clinal patterns at a mine boundary. Heredity 25:349–362.

433   Barton, N. H., and P. D. Keightley. 2002. Understanding quantitative genetic variation. Nature

434   Reviews Genetics 3:11–21.

435   Baucom, R. S. 2019. Evolutionary and ecological insights from herbicide resistant weeds: what

436   have we learned about plant adaptation, and what is left to uncover? New Phytologist,

437   doi: 10.1111/nph.15723.

438   Bell, G. 2017. Evolutionary rescue. Annual Review of Ecology, Evolution, and Systematics

439   48:605–627.

440   Bell, G., and A. Gonzalez. 2011. Adaptation and evolutionary rescue in metapopulations

441   experiencing environmental deterioration. Science 332:1327–1330.

442   Bell, G., and A. Gonzalez. 2009. Evolutionary rescue can prevent extinction following

443   environmental change. Ecology Letters 12:942–948.

444   Bolker, B. M., M. E. Brooks, C. J. Clark, S. W. Geange, J. R. Poulsen, M. H. H. Stevens, and J.-

445   S. S. White. 2009. Generalized linear mixed models: a practical guide for ecology and

446   evolution. Trends in Ecology & Evolution 24:127–135.

Bradshaw, A. D. 1991. The Croonian Lecture, 1991. Genostasis and the limits to evolution.

Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences

333:289–305.

Burt, A. 1995. The evolution of fitness. Evolution 49:1–8.

Carlson, S. M., C. J. Cunningham, and P. A. H. Westley. 2014. Evolutionary rescue in a

changing world. Trends in Ecology & Evolution 29:521–530.

Coltman, D. W., P. O'Donoghue, J. T. Hogg, and M. Festa-Bianchet. 2005. Selection and genetic

(co)variance in bighorn sheep. Evolution 59:1372–1382.

Comstock, R. E., and H. F. Robinson. 1948. The components of genetic variance in populations

of biparental progenies and their use in estimating the average degree of dominance.

Biometrics 4:254–266.

de Villemereuil, P., A. Rutschmann, K. D. Lee, J. G. Ewen, P. Brekke, and A. W. Santure. 2019.

Little adaptive potential in a threatened passerine bird. Current Biology. doi:

10.1016/j.cub.2019.01.072.

de Villemereuil, P., H. Schielzeth, S. Nakagawa, and M. Morrissey. 2016. General methods for

evolutionary quantitative genetic inference from generalized mixed models. Genetics

204:1281–1294. Genetics.

Eck, D. J., R. G. Shaw, C. J. Geyer, and J. G. Kingsolver. 2015. An integrated analysis of

phenotypic selection on insect body size and development time. Evolution 69:2525–2532.

Etterson, J. R. 2004a. Evolutionary potential of *Chamaecrista fasciculata* in relation to climate

change. I. Clinal patterns of selection along an environmental gradient in the great plains.

Evolution 58:1446–1458.

Etterson, J. R. 2004b. Evolutionary potential of *Chamaecrista fasciculata* in relation to climate

change. II. Genetic architecture of three populations reciprocally planted along an

environmental gradient in the great plains. Evolution 58:1459–1471.

Etterson, J. R., and R. G. Shaw. 2001. Constraint to Adaptive Evolution in Response to Global

Warming. Science 294:151–154.

Ewens, W. J. 2004. Mathematical Population Genetics. 2nd ed. Springer.

Felsenstein, J. 1976. The theoretical population genetics of variable selection and migration.

Annual Review of Genetics 10:253–280.

Fenster, C. B. 1991a. Gene flow in *Chamaecrista fasciculata* (Leguminosae) I. Gene dispersal.

Evolution 45:398–409.

Fenster, C. B. 1991b. Gene flow in *Chamaecrista fasciculata* (Leguminosae) II. Gene

establishment. Evolution 45:410–422.

Fisher, R. A. 1930. The Genetical Theory of Natural Selection. Oxford: Claredon Press.

Fisk, D. L., L. C. L. IV, R. A. Knapp, and M. E. Pfrender. 2007. Rapid evolution in response to

introduced predators I: rates and patterns of morphological and life-history trait

divergence. BMC Evolutionary Biology 11.

Foerster, K., T. Coulson, B. C. Sheldon, J. M. Pemberton, T. H. Clutton-Brock, and L. E. B.

Kruuk. 2007. Sexually antagonistic genetic variation for fitness in red deer. Nature

447:1107–1110.

Franks, S. J., S. Sim, and A. E. Weis. 2007. Rapid evolution of flowering time by an annual plant

in response to a climate fluctuation. PNAS 104:1278–1282.

Galloway, L. F., and C. B. Fenster. 2000. Population differentiation in an annual legume: local

adaptation. Evolution 54:1173–1181.

Geerts, A. N., J. Vanoverbeke, B. Vanschoenwinkel, W. Van Doorslaer, H. Feuchtmayr, D.

Atkinson, B. Moss, T. A. Davidson, C. D. Sayer, and L. De Meester. 2015. Rapid

evolution of thermal tolerance in the water flea *Daphnia*. Nature Climate Change 5:665–

668.

Geyer, C. J. 2018. aster: Aster Models. R package version 0.9.1.1. https://cran.r-

project.org/package=aster.

Geyer, C. J., C. E. Ridley, R. G. Latta, J. R. Etterson, and R. G. Shaw. 2013. Local adaptation

and genetic effects on fitness: Calculations for exponential family models with random

effects. The Annals of Applied Statistics 7:1778–1795.

Geyer, C. J., and R. G. Shaw. 2013. Aster models with random effects and additive genetic

variance for fitness. Accessed June 24, 2018.

https://conservancy.umn.edu/handle/11299/152355.

Geyer, C. J., S. Wagenius, and R. G. Shaw. 2007. Aster models for life history analysis.

Biometrika 94:415–426.

Gillespie, J. H. 1984. Pleiotropic overdominance and the maintenance of genetic variation in

polygenic characters. Genetics 107:321–330.

Gomulkiewicz, R., and R. D. Holt. 1995. When does evolution by natural selection prevent

extinction? Evolution 49:201–207.

Gomulkiewicz, R., and R. G. Shaw. 2012. Evolutionary rescue beyond the models. Philosophical

Transactions of the Royal Society B: Biological Sciences 368: 20120093.

Hartl, D. L., and A. G. Clark. 1997. Principles of Population Genetics. 3rd ed. Sinauer.

Huang, Y., J. R. Stinchcombe, and A. F. Agrawal. 2015. Quantitative genetic variance in

experimental fly populations evolving with or without environmental heterogeneity:

quantitative genetic variance. Evolution 69:2735–2746.

Irwin, H., and R. Barneby. 1982. The American Cassiinaea synoptical revision of Leguminosae

tribe Cassieae subtribe Cassiinae in the New World. Memoirs of the New York Botanical

Garden 35:1–918.

Kelly, C. A. 1993. Quantitative genetics of size and phenology of life-history traits in

Chamaecrista fasciculata. Evolution 47:88–97.

Kruuk, L. E. B., T. H. Clutton-Brock, J. Slate, J. M. Pemberton, S. Brotherstone, and F. E.

Guinness. 2000. Heritability of fitness in a wild mammal population. Proceedings of the

National Academy of Sciences 97:698–703.

Kulbaba, M. W., and R. G. Shaw. In Review. Lifetime fitness through female and male function:

the influence of density and genetically effective population size. American Naturalist.

Lee, T. D., and F. A. Bazzaz. 1982. Regulation of fruit and seed production in an annual legume,

Cassia fasciculata. Ecology 63:1363–1373.

Levene, H. 1953. Genetic equilibrium when more than one ecological niche is available. The

American Naturalist 87:331–333.

Mackay, T. F. C. 1980. Genetic variance, fitness, and homeostasis in varying environments: an

experimental check of the theory. Evolution 34:1219–1222.

Matos, Rose, M. T. R. Pite, Rego, and Avelar. 2000. Adaptation to the laboratory environment in

Drosophila subobscura. Journal of Evolutionary Biology 13:9–19.

Mazer, S. J. 1987. The Quantitative genetics of lifehistory and fitness components in *Raphanus*

raphanistrum L. (Brassicaceae): ecological and evolutionary consequences of seed-

weight variation. The American Naturalist 130:891–914.

McCleery, R. H., R. A. Pettifor, P. Armbruster, K. Meyer, B. C. Sheldon, and C. M. Perrins.

2004. Components of variance underlying fitness in a natural population of the great tit

Parus major. The American Naturalist 164:E62–E72.

McFarlane, S. E., J. C. Gorrell, D. W. Coltman, M. M. Humphries, S. Boutin, and A. G.

McAdam. 2014. Very low levels of direct additive genetic variance in fitness and fitness

components in a red squirrel population. Ecology and Evolution 4:1729–1738.

Messer, P. W., S. P. Ellner, and N. G. Hairston. 2016. Can population genetics adapt to rapid

evolution? Trends in Genetics 32:408–418.

Moose, S. P., J. W. Dudley, and T. R. Rocheford. 2004. Maize selection passes the century mark:

a unique resource for 21st century genomics. Trends in Plant Science 9:358–364.

Mousseau, T. A., and D. A. Roff. 1987. Natural selection and the heritability of fitness

components. Heredity 59:181–197.

Neumann, M., V. Mues, A. Moreno, H. Hasenauer, and R. Seidl. 2017. Climate variability drives

recent tree mortality in Europe. Global Change Biology 23:4788–4797.

Pemberton, J. M. 2010. Evolution of quantitative traits in the wild: mind the ecology.

Philosophical Transactions of the Royal Society B: Biological Sciences 365:2431–2438.

R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Roff, D. A., and T. A. Mousseau. 1987. Quantitative genetics and fitness: lessons from

Drosophila. Heredity 58:103–118.

Rose, M. R., and B. Charlesworth. 1981. Genetics of life history in *Drosophila melanogaster*. II.

exploratory selection experiments. Genetics 97:187–196.

Roughgarden, J. 1996. The Theory of Population Genetics and Evolutionary Ecology: An

Introduction. Prentice Hall.

Sæther, B.-E., and S. Engen. 2015. The concept of fitness in fluctuating environments. Trends in

Ecology & Evolution 30:273–281.

Shaw, R. G. 2019. From the past to the future: considering the value and limits of evolutionary

prediction. The American Naturalist 193:1–10.

Shaw, R. G., and J. R. Etterson. 2012. Rapid climate change and the rate of adaptation: insight

from experimental quantitative genetics. New Phytologist 195:752–765.

Shaw, R. G., and C. J. Geyer. 2010. Inferring fitness landscapes. Evolution 64:2510–2520.

Shaw, R. G., C. J. Geyer, S. Wagenius, H. H. Hangelbroek, and J. R. Etterson. 2008. Unifying

life-history analyses for inference of fitness and population growth. The American

naturalist 172:E35-47. The University of Chicago Press .

Shaw, R. G., and F. H. Shaw. 2014. Quantitative genetic study of the adaptive process. Heredity

112:13–20.

Sheth, S. N., M. W. Kulbaba, R. E. Pain, and R. G. Shaw. 2018. Expression of additive genetic

variance for fitness in a population of partridge pea in two field sites. Evolution 72:2537–

2545.

Stanton-Geddes, J., R. G. Shaw, and P. Tiffin. 2013. Insights from population genetics for range

limits of a widely distributed native plant. American Journal of Botany 100:744–753.

Stanton-Geddes, J., P. Tiffin, and R. G. Shaw. 2012. Role of climate and competitors in limiting

fitness across range edges of an annual plant. Ecology 93:1604–1613.

Teplitsky, C., J. A. Mills, J. W. Yarrall, and J. Merilä. 2009. Heritability of fitness components in

a wild bird population. Evolution 63:716–726.

Thomas, S. C., J. M. Pemberton, and W. G. Hill. 2000. Estimating variance components in

natural populations using inferred relationships. Heredity 84:427–436.

Vigueira, C. C., K. M. Olsen, and A. L. Caicedo. 2013. The red queen in the corn: agricultural

weeds as models of rapid adaptive evolution. Heredity 110:303–311.

Winn, A. A. 2004. Natural selection, evolvability and bias due to environmental covariance in

the field in an annual plant: bias in phenotypic selection gradients. Journal of

Evolutionary Biology 17:1073–1083.

Yeaman, S., Y. Chen, and M. C. Whitlock. 2010. No effect of environmental heterogeneity on

the maintenance of genetic variation in wing shape in *Drosophila melanogaster*:

heterogeneity and maintenance of genetic variation. Evolution 64:3398–3408.

Yi, X., and A. M. Dean. 2013. Bounded population sizes, fluctuating selection and the tempo and

mode of coexistence. Proceedings of the National Academy of Sciences 110:16945–

16950.

Zhang, X.-S. 2012. Fisher's geometrical model of fitness landscape and variance in fitness

within a changing environment. Evolution 66:2350–2368.

Table 1. Range of dates for collection of fitness expression data collection, for three sites of

Chamaecrista fasciculata, over three consecutive years.

	Year		
Site	2015	2016	2017
Grey Cloud	June 1 – October 1	June 10 – October 7	June 5 – October 3
McCarthy Lake	May 27 – September 25	June 7 – October 11	June 8 – October 5
CERA	May 10 – September 23	June 2 – September 29	June 1 – October 23

602 Table 2. Summary from aster models testing the fixed effects of block and year, the random

603 factors of combined parental effects, and the interaction between parental effects and year on

604 individual lifetime fitness. Statistical significance of predictor variables was assessed using

605 likelihood ratio tests, and random parental effects were assessed from summary output of aster

606 analyses.

Model	Test df	Test deviance	P
Grey Cloud – *G x Year* model			
Block	23	159.91	$P < 0.0001$
Year	2	47.28	$P < 0.0001$
Parental effects	--	--	$P < 0.0001$
Year *x* Parental effects	--	--	$P < 0.0001$
Grey Cloud – fixed effects models for mean fitness			
2015 – Block	7	16.12	$P < 0.01$
2016 – Block	7	37.67	$P < 0.001$
2017 – Block	7	63.93	$P < 0.0001$
McCarthy Lake – *G x Year* model			
Block	23	159.91	$P < 0.001$
Year	2	47.28	$P < 0.001$
Parental effects	--	--	$P < 0.0001$
Year *x* Parental effects	--	--	$P < 0.0001$
McCarthy Lake – fixed effects models for mean fitness			
2015 – Block	7	4.26	$P < 0.05$
2016 – Block	7	63.32	$P < 0.0001$

2017 – Block	7	37.29	P < 0.0001
CERA – *G x Year* model			
Block	11	189.80	P < 0.0001
Year	2	58.95	P < 0.0001
Parental effects	--	--	P < 0.0001
Year *x* Parental effects	--	--	P < 0.0001
CERA – fixed effects models for mean fitness			
2015 – Block	3	50.09	P < 0.0001
2016 – Block	3	27.92	P < 0.0001
2017 – Block	3	46.16	P < 0.0001

Table 3. Additive genetic variance for fitness ($V_A(W)$), mean fitness ($\overline{W}$) and standard error, predicted change in mean fitness $\left(\frac{V_A(W)}{\overline{W}}\right)$, and predicted mean fitness of second generation determined for each of three sites and three years of *Chamaecrista fasciculata*. Block used to estimate mean fitness indicated in parentheses after year.

Site/Year (Block)	Number of Seeds Planted	$V_A(W)$ (SE)	$\overline{W}$ (SE)	$\frac{V_A(W)}{\overline{W}}$	Predicted Fitness of Gen. 2
Grey Cloud Dunes					
2015 (Block 6)	3658	1.858 (0.433)	1.11 (0.17)	1.67	2.78
2016 (Block 6)	3660	0.830 (0.211)	0.64 (0.13)	1.30	1.94
2017 (Block 7)	3660	6.491 (1.546)	1.06 (0.34)	6.11	7.17
McCarthy Lake					
2015 (Block 4)	3445	3.528 (1.143)	3.02 (1.23)	1.17	4.19
2016 (Block 5)	3480	1.572 (0.381)	1.88 (0.57)	0.83	2.72
2017 (Block 6)	3516	1.410 (0.331)	1.08 (0.18)	1.06	2.14
CERA					
2015 (Block 1)	1845	3.204 (1.068)	1.80 (0.24)	1.79	3.58
2016 (Block 2)	1750	0.856 (0.254)	0.73 (0.11)	1.18	1.91

| 2017 (Block 3) | 1750 | 1.573 | 1.24 | 1.27 | 2.51 |
| | | (0.659) | (0.253) | | |

611
612

Figure 1 (A) Graphical model used to estimate lifetime fitness. Each node represents a fitness component and

therefore response variable, and arrows represent conditional distributions. Probability of germination and

flowering (0 or 1; Bernulli distribution), and total number of fruits and seeds produced (Poisson distribution).

Additional node to account for subsampling of total fruit production (not shown) also followed a Poisson

distribution. (B) Map of three experimental populations. Seed photos from Anna Peschel and remaining photos

by MWK.

Figure 2. Additive genetic variation for lifetime fitness for three populations of *Chamaecrista fasciculata*, in

three consecutive years. Lines represent probability density distributions of estimated breeding values for

lifetime fitness (seeds set).

Figure 3. Paternal family-specific breeding values for lifetime fitness (seeds set) compared across three

consecutive years in three populations of *Chamaecrista fasciculata*. Pearson correlation coefficients do not

account for error associated with estimates of breeding values and are therefore presented as a coarse

description of the interannual relationship between breeding values. Please note differences in scale across

panels.

Figure 4. Estimated mean fitness of parental (circles with standard errors) and progeny generations (triangles with standard errors) for three populations of *Chamaecrista fasciculata* in three consecutive years. Horizontal dashed lines represent mean fitness of 1, indicating individual replacement and population stability.