Native molecule sequencing by nano-ID reveals synthesis and stability of RNA isoforms

Kerstin C. Maier[†1], Saskia Gressel[1], Patrick Cramer[1*], and Björn Schwalb[1†*]

[1]Max-Planck-Institute for Biophysical Chemistry, Department of Molecular Biology, Am
Faßberg 11, 37077 Göttingen, Germany.

[†]These authors contributed equally.

[*]Correspondence to: Patrick Cramer, (patrick.cramer@mpibpc.mpg.de), and Björn Schwalb,
(bjoern.schwalb@mpibpc.mpg.de).

Abstract

Eukaryotic genes often generate a variety of RNA isoforms that can lead to functionally distinct
protein variants. The synthesis and stability of RNA isoforms is however poorly characterized.
The reason for this is that current methods to quantify RNA metabolism use 'short-read'
sequencing that cannot detect RNA isoforms. Here we present nanopore sequencing-based
Isoform Dynamics (nano-ID), a method that detects newly synthesized RNA isoforms and
monitors isoform metabolism. nano-ID combines metabolic RNA labeling, 'long-read' nanopore
sequencing of native RNA molecules and machine learning. Application of nano-ID to the heat
shock response in human cells reveals that many RNA isoforms change their synthesis rate,
stability, and splicing pattern. nano-ID also shows that the metabolism of individual RNA
isoforms differs strongly from that estimated for the combined RNA signal at a specific gene
locus. And although combined RNA stability correlates with poly(A)-tail length, individual RNA
isoforms can deviate significantly. nano-ID enables studies of RNA metabolism on the level of
single RNA molecules and isoforms in different cell states and conditions.

Main

In metazoan cells, a single gene locus can give rise to a variety of different RNA molecules that are generally referred to as isoforms. These RNA isoforms can differ in their 5'- and 3'-ends that arise from the use of different transcription start sites and polyadenylation sites, respectively [1-4]. In addition, alternative splicing results in RNA isoforms that differ in the composition of their RNA body [5,6]. Different mRNA isoforms can result in functionally different proteins. Vulnerabilities in splicing can lead to non-functional protein products. Diseases have been linked to alternative splicing, which can generate malignant RNA isoforms [7]. Duchenne muscular dystrophy (DMD), for example, can be pinpointed to a single gene encoding the protein dystrophin. The underlying malignant RNA isoform exhibits a different splicing pattern and leads to a non-functional protein, which disrupts muscular cell integrity [8]. Likewise, the three most common types of breast tumors are linked to exon skipping and intron retention [9].

RNA isoforms can also differ in their stability. The untranslated region of an RNA isoform can differ in length and contains regulatory elements [10]. The length of the poly(A)-tail at the 3'-end of RNA isoforms can also differ and influence RNA stability [11,12], and this is relevant for disease as well [13]. Finally, introns may be retained in RNAs and can influence stability [14].

Little is known however about the synthesis and stability of single RNA isoforms in cells. This is because the systematic characterization of RNA isoforms and their metabolism is technically difficult. In particular, the detection, quantification and estimation of the stability of RNA isoforms is essentially impossible with 'short-read' RNA sequencing methods because reads generally cannot be assigned to RNA isoforms. Also, alternative splicing patterns can be manifold and are difficult to identify using 'short-read' sequencing approaches [15]. Finally, although the length of poly(A)-tails of RNAs can be measured genome-wide [16,17], they can currently not be obtained at the level of individual RNA isoforms.

The architecture of RNA isoforms has been addressed so far by 'short-read' RNA sequencing approaches such as DARTS [18], VastDB [19] and MPE-seq [20] to study alternative splicing or TIF-seq [1,3] to elucidate combinations of paired 5'- and 3'-ends of individual RNAs. More recent approaches include 'long-read' sequencing approaches on the PacBio SMRT Sequencing platform [6] or Oxford Nanopore Technologies nanopore sequencing platform [5,21,22].

These methods however are not able to study the metabolism of individual RNA isoforms
because they lack the ability to assign age to single reads.

Methods to measure the synthesis and stability of combined RNA for entire gene loci are
available [23-25]. Transient transcriptome sequencing (TT-seq) is a protocol that allows to
distinguish newly synthesized from pre-existing RNA in human cells [26]. TT-seq involves a brief
exposure of cells to the nucleoside analogue 4-thiouridine (4sU). 4sU is incorporated into RNA
during transcription, and the resulting 4sU-labeled RNA can be purified and sequenced to
provide a snapshot of immediate transcription activity. This then enables to computationally
infer RNA synthesis and stability at the level of the combined RNA signal from a gene locus.

Recent methods to assess RNA stability include SLAM-seq [27] and TimeLapse-seq [28].
Like TT-seq, SLAM-seq and TimeLapse-seq involve an exposure of cells to 4sU for labeling of
newly synthesized RNA. A chemical modification of the incorporated 4sU then allows for the
identification of labeled RNA *in silico* without the need for purification. All of these methods,
however, have limitations. First, sequencing reads can normally only be assigned to entire gene
loci and not to RNA isoforms and thus only allow a combined RNA stability assessment.
Second, they require template amplification, which can lead to an imbalance in measured
sequences and information loss, e.g. modified RNA bases [29]. Third, labeled RNA purification
(TT-seq) and cDNA library preparation (TT-seq, SLAM-seq & TimeLapse-seq) can also
introduce biases.

Therefore, monitoring RNA metabolism at the level of RNA isoforms requires a method
that can detect individual RNA molecules. Recent advances in 'long-read' nanopore sequencing
indeed enable the sequencing of single, full-length RNA molecules [5]. Nanopore technology can
directly sequence the original native RNA molecule with its modifications, may they be natural
or acquired by metabolic RNA labeling. Moreover, the availability of the entire RNA and coding
sequence (CDS) within a single read allows to unambiguously and directly determine exon usage
[30]. Direct RNA 'long-read' nanopore sequencing also has the potential to detect the position and
length of the poly(A)-tail along with each single isoform.

Here we developed nanopore sequencing-based Isoform Dynamics (nano-ID), which
combines metabolic RNA labeling with native RNA 'long-read' nanopore sequencing for RNA
isoform detection. In combination with computational modeling and machine learning this

allows for a full characterization of RNA isoforms dynamics. nano-ID can identify and quantify

RNA isoforms along with their synthesis rate, stability and poly(A)-tail length in the human

myelogenous leukemia cell line K562. We show that this is possible with nano-ID in a

quantitative manner in steady state and also during the transcriptional response to heat shock.

nano-ID is able to resolve the dynamic metabolism of RNA isoforms upon heat shock and

demonstrates the need for individual RNA isoform assessment. Taken together, nano-ID can be

used to elucidate a largely unexplored complex layer of gene regulation at the level of single

native RNA isoforms and their metabolism.

Results

Experimental design

To monitor the metabolism of RNAs at the level of single isoforms, we sought to combine

metabolic RNA labeling with direct, single-molecule RNA nanopore sequencing (**Figure 1a**). By

culturing cells in the presence of a nucleoside analogue, cells will take up and incorporate the

analogue in nascent RNA during transcription, allowing to distinguish newly synthesized RNA

isoforms from pre-existing RNA isoforms *in silico* based on the quantification of analogue-

containing subpopulations. This will allow to infer the synthesis rate and stability of single RNA

isoforms. In order to dynamically characterize functional and fully processed RNA transcripts,

we decided to measure poly-adenylated RNA species. The library preparation kit offered by

Oxford Nanopore Technologies for direct RNA sequencing (SQK-RNA001) is specifically

optimized for this purpose. A 3' poly(A)-tail specific adapter is ligated to the transcript in a first

step. Then a second sequencing adapter equipped with a motor protein is ligated to the first

adapter. The preparation of RNA libraries from biological samples for direct RNA nanopore

sequencing is established and can be carried out within 2h [31]. Major challenges that we faced

were however the search of a suited nucleoside analogue for RNA labeling and the detection of

labeled RNA isoforms, provided that the labeling efficiency is known to be limited to about 2-

3%, i.e. only two or three out of 100 natural nucleotides are replaced by the analogue [32].

5-Ethynyluridine ([5E]U) can be detected in RNA by nanopore sequencing

To investigate if nucleoside analogues incorporated into RNA are detectable in the nanopore, we
used synthetic RNAs derived from the ERCC RNA spike-in mix (Life Technologies). These
synthetic RNAs of an approximate length of 1,000 nucleotides were chosen with similar U
content (**Supplementary Table 3**). RNAs were transcribed *in vitro* using either the standard
bases A, U, C, G as a control, or with one of the natural bases exchanged for a nucleoside
analogue (**Figure 1b**, **Methods**). Subsequently, we subjected these synthetic RNAs to direct
RNA nanopore sequencing (**Supplementary Figure 1a-b**). We compared the nucleoside
analogues 5-Ethynyluridine (^{5E}U), 5-bromouridine (^{5Br}U), 5-iodouridine (^{5I}U), 4-thiouridine (^{4s}U)
and 6-thioguanine (^{6s}G). To this end we used the base-called and mapped direct RNA sequencing
results to calculate how probable the identification would be on the level of single nucleotides. In
particular, we compared the error rate in single nucleotide base-calls of nucleoside analogues to
that of natural U or G (**Figure 1c, Methods**).

The thiol-based analogues, ^{4s}U and ^{6s}G, showed lower incorporation efficiencies during
in vitro transcription (IVT) and led to blockages during nanopore sequencing. ^{5E}U and ^{5I}U could
be detected to a similar extent by nanopore sequencing, whereas ^{5Br}U was less easily recognized
(**Figure 1c**). Since ^{5E}U is not toxic to cells [32-34], we used ^{5E}U for a more detailed analysis.
Approximately 50% of all U positions in ^{5E}U-containing synthetic RNAs are consistently
miscalled by the standard base-calling algorithm and can thus be discerned from U (**Figure 1d,**
Supplementary Figure 1b). This is clearly visible in the raw data. Aberrations caused by
stretches of RNA containing ^{5E}U are distinguishable from stretches of RNA containing the
naturally occurring U in the nanopore (**Figure 1d**). Taken together, ^{5E}U-based RNA labeling is
well suited for nanopore sequencing.

Detection and sequencing of newly synthesized RNA isoforms

We next investigated whether it is possible to use metabolic RNA labeling with ^{5E}U in human
cells to detect single RNA molecules by nanopore sequencing. Calculations on the direct RNA
nanopore sequencing results of the ^{5E}U-containing synthetic RNAs showed that RNAs are
recognizable as ^{5E}U containing with a probability of 0.9 when a minimum length of 500
nucleotides is reached (**Supplementary Figure 1c-d**). This covers the vast majority (93%) of all
mature RNAs in the human organism (UCSC RefSeq GRCh38).

We then established direct RNA nanopore sequencing in the human myelogenous

leukemia cell line K562. We cultured K562 cells in the presence of [5E]U for 60 minutes ([5E]U 60

min) in 4 biological replicates (**Methods**). For comparison, we created 3 biological replicates

exposed to [5E]U labeling for 24 h ([5E]U 24 h) and 3 biological replicates that were not labeled

(Control). After standard base-calling, we could map reads to support 13,110 RefSeq annotated

transcription units (RefSeq-TUs, **Methods**), 8,098 of these were supported in all conditions and

1,726 were supported in all samples.

All combined samples were then used to perform a full-length alternative RNA isoform

analysis by means of the FLAIR algorithm [22]. This allows defining instances of unique exon-

intron architecture with unique start and end sites in human K562 cells. Raw human direct RNA

nanopore reads were corrected with the use of short-read sequencing data (RNA-seq) to increase

splice site accuracy. We could detect 33,199 distinct RNA isoforms with an average of 3

isoforms per gene. This shows that direct RNA nanopore sequencing uncovers individual RNA

isoforms in human K562 cells (**Figure 2**) with high reproducibility (**Supplementary Figure 2**).

A neural network identifies newly synthesized RNA isoforms

The next step was to derive a computational method that could classify each sequenced RNA

molecule into one of two groups, newly synthesized ([5E]U-labeled) or pre-existing (unlabeled)

RNA. To this end, the nucleoside analogue [5E]U had to be detected in RNA molecules. This

would allow the quantification of RNA isoforms generated during the [5E]U labeling pulse. Due to

the high error rate of nanopore sequencing, a single [5E]U base-call is inappropriate as an indicator.

We rather used the raw signal of the entire RNA nanopore read, including the base-calls and the

alignment, to discriminate labeled from unlabeled RNAs. This discrimination was implemented

as a classifying neural network. We developed a custom multi-layered data collection scheme to

train a neural network for the classification of human RNA isoforms under the assumption that

the [5E]U 24 h samples solely contain labeled reads and the fact that the Control samples solely

contain unlabeled reads (**Figure 3a, Methods**).

We then trained a neural network (**Methods**) on the [5E]U 24 h versus Control samples

with an accuracy of 0.87 and a false discovery rate (FDR) of 0.025 (5-fold cross-validated). A

ROC analysis (1 − specificity versus sensitivity) for all reads of the test set showed an area under

the curve (AUC) of 0.94. For reads with an alignment length larger than 500 nt and 1,000 nt the

AUC improved to 0.96 (**Figure 3b, Supplementary Figure 3a, b**). Subsequently we used the

trained neural network to classify reads of the [5E]U 60 min samples into [5E]U-labeled and

unlabeled. Taken together, [5E]U containing RNA isoforms are computationally detectable with

high accuracy (**Figure 3c**). For validation purposes, we used another machine learning approach.

We trained a random forest on the same data, which yielded similar results (**Supplementary**

Figure 3c, d). Thus, we were able to determine for each single RNA molecule if it has been

produced during [5E]U labeling or before, with a low false discovery rate (**Figure 3c**).

nano-ID provides the stability and poly(A) tail length of RNA isoforms

The ability to distinguish newly synthesized and pre-existing RNA molecules allowed us to

derive estimates for the stability of RNA isoforms. For each single direct RNA nanopore read we

were able to assign the RNA isoform it reflects. Additionally, we were able to assess the stability

of RNA for single RNA isoforms by applying a first-order kinetic model (**Methods,**

Supplementary Figure 3e-f) to derive estimates for RNA isoform specific synthesis and

stability. This can be done based on the number of reads classified as [5E]U-labeled and unlabeled

by the neural network. Taken together, nano-ID has the capability to infer synthesis and stability

of individual RNA isoforms in different cell states and conditions, and thus to monitor their

dynamic metabolism.

Moreover, we developed an algorithm to determine poly(A)-tail lengths for each RNA

isoform (**Figure 4**). This is possible by estimating the dwell time of the poly(A)-tail in the

nanopore by factoring in the measurement frequency in kHz and the speed of RNA translocation

through the nanopore (**Methods**). Sequencing adaptor ligation in the direct RNA nanopore

sequencing library preparation guarantees full-length poly(A)-tails because ligation of the

adapter would not be successful otherwise. The resulting poly(A)-tail length distribution is in

line with the current literature [16] (**Figure 4a**) and reveals a pattern that likely corresponds to the

26 nucleotide footprint of the poly(A) binding protein (**Supplementary Figure 4a**) [35]. The direct

RNA nanopore sequencing kit contains the so-called RNA calibration strand (RCS). The RCS is

a synthetic RNA with a poly(A)-tail of exactly 30 adenines. Using the RCS of the direct RNA

nanopore sequencing kit, we could assess the accuracy of the poly(A)-tail length estimates

(coefficient of variation 0.63). Our algorithm derives this length for the added RCS

subpopulation (**Figure 4b**). Taken together, nano-ID reveals the synthesis, stability, and poly(A)

tail length for individual RNA isoforms in human cells.

nano-ID monitors RNA isoform dynamics during heat shock

To demonstrate the advantages of nano-ID, we subjected human K562 cells to heat shock (42

°C) for 60 min in the presence of ^{5E}U (^{5E}U 60 min HS) (**Figure 5a**). The heat shock response

provides a well-established model system [36-41] (**Supplementary Figure 5**). We first asked

whether RNA isoforms do retain more introns after heat shock as this was shown in the mouse

system [42]. Indeed, we observed widespread intron retention which significantly increased upon

heat shock (**Figure 5b**). Although intron retention generally influences the stability of an RNA, it

does not explain changes in RNA isoform stability upon heat shock (**Figure 5c**). This finding is

consistent with the idea that specific RNA elements occurring only in specific RNA isoforms

influence RNA stability.

We next asked if RNA isoform synthesis is altered by heat shock and observed

significant differential RNA isoform synthesis for 285 isoforms (fold change > 1.25 and p-value

< 0.1). 187 RNA isoforms were significantly upregulated, while 98 were downregulated (**Figure**

5d). RNA isoforms that changed their synthesis during heat shock were also observed to alter

their stability (**Figure 5e-f**). In particular, RNA isoforms that were upregulated in their synthesis

during heat shock also showed a lower stability, and the other way around, resembling typical

stress response behavior [24]. The destabilization of upregulated RNA isoforms is likely to ensure

their rapid removal toward the end of the stress response. Similarly, downregulated RNA

isoforms are stabilized, perhaps to preserve them for translation at later stages.

nano-ID reveals the biogenesis of RNA isoforms

Although standard native RNA isoform sequencing can reveal isoforms present in a sample after

perturbation, it cannot distinguish whether these isoforms were derived by synthesis, stability,

splicing, or any combination of these. nano-ID however is able to disentangle these parameters.

For example, although we observe a general increase in intron retention upon heat shock, we find

exceptions at the level of RNA isoforms. This can be clearly seen at the human C1orf63 gene

locus (**Supplementary Figure 5g**). Here, the majority of reads, that retain the entire 3[rd] intron,

were newly synthesized in the control samples. It is however unclear if this intron will be

retained throughout the existence of these RNA molecules. Investigation of the same gene locus

upon heat shock showed that the vast majority of reads were pre-existing RNAs. This indicates

that this RNA is not transcribed anymore upon heat shock and allows for the conclusion that

intron retention is not altered, rather, less introns are seen retained when only old RNA is

detected. Taken together, this shows that nano-ID is able to resolve the dynamic behavior of

RNA isoforms upon stimuli that could not be seen otherwise. It demonstrates the need for

individual RNA isoform detection and classification into newly synthesized and pre-existing

molecules. By providing information on the age of RNA molecules, nano-ID enables an analysis

of the biogenesis of RNA isoforms.

The metabolism of individual RNA isoforms differs from combined RNA estimates

To demonstrate the importance of individual RNA isoform assessment, we first derived

estimates for the half-lives of combined RNAs that stem from entire gene loci under steady state

conditions (**Methods, Supplementary Figure 3e-f**). We found a robust correlation of combined

RNA stability with poly(A)-tail length (Spearman's rank correlation coefficient 0.48) (**Figure

5g**). We now asked whether changes in RNA stability would also be reflected in changes in

poly(A)-tail length upon heat shock, and this was not the case (**Figure 5h**). Instead, we found

genes that showed the opposite behavior to the overall correlation as demonstrated for the human

HSPB1 locus (**Figure 6a-b**). Here, destabilization of combined RNAs is accompanied by

lengthening of the poly(A)-tail. This view changes dramatically when considering individual

RNA isoforms (**Figure 6c**). For those three RNA isoforms at the human HSPB1 gene locus for

which stability estimates were supported by all 3 biological replicates (**Methods**) we found that

poly(A)-tails were generally longer. RNA stability however was decreased for 2 out of the 3

RNA isoforms and increased for the third. This clearly indicates the need for detailed individual

RNA isoform assessment as individual RNA isoforms can lead to functionally distinct protein

variants. Thus, it is crucial to also study the behavior of individual RNA isoforms instead of

breaking it down to the combined view of the entire gene locus.

As a second example, we picked RNA isoforms at the human TAGLN2 gene locus
(**Figure 6d**). We could identify 7 different RNA isoforms and reliably calculate RNA stability
for 6 RNA isoforms. Two of them were stabilized upon heat shock, 4 of them were destabilized.
All 4 destabilized RNA isoforms include the second to last exon, which might cause this change
in stability. RNA isoform 7 is an exception to this observation as it is stabilized upon heat shock.
It, however, also contains a 3' UTR that is 42 bases shorter than all the other RNA isoforms. We
asked whether there is differential behavior of individual RNA isoforms genome-wide or if RNA
isoforms generally reflect the changes in stability of the combined RNA from their respective
gene loci. To that end, we compared RNA stability estimates of individual RNA isoforms to
those from combined RNAs and found that the dynamics of individual RNA isoform during heat
shock varies globally (**Figure 6e**, **Supplementary Figure 6**). Taken together, this shows that
conclusions can be misleading when combined RNAs are used and how much can be learned on
the level of single RNA isoforms by using nano-ID.

Discussion
Here we develop nano-ID, a method that allows for dynamic characterization of functional and
fully processed RNA isoforms on the level of single native RNA molecules. nano-ID combines
metabolic RNA labeling with native RNA nanopore sequencing to enable RNA isoform
identification, estimation of its stability, and a measurement of its poly(A)-tail length from a
single sample. nano-ID is able to visualize changes in RNA isoform synthesis and stability and
reveals a hidden layer of gene regulation. nano-ID thus allows to study transcriptional regulation
in unprecedented detail and can prevent misleading conclusions that would be obtained when
only combined RNAs from an entire gene locus are considered, as is done by RNA-seq, 4sU-seq
or TT-seq.
nano-ID has many advantages over other sequencing-based transcriptomic strategies as it
allows to sequence the original native RNA molecule. In particular, there is no need for
fragmentation of RNA prior to sequencing and hence no ambiguity in assigning reads to RNA
isoforms. nano-ID also does not require template amplification and thus omits copying errors and
sequence-dependent biases. It comes without a lengthy library protocol and eliminates
sequencing by synthesis and therefore prevents loss of information on epigenetic modifications
and artificially introduced RNA base analogues. It is PCR-free and shows neither sequence bias

nor read duplication events. Taken together, it overcomes drawbacks and limitations of state-of-

the-art approaches and increases the gathered information vastly.

295   Generally, nanopore sequencing has still limitations in throughput and accuracy. These

drawbacks, however, are outweighed by the information obtained on the sequencing substrates.

The longer the sequenced molecules are, the less problematic is the lack in accuracy in

identifying their origin or classifying it into newly-synthesized or pre-existing. On top of that,

there are strategies to improve splice site calling with already existing high accuracy 'short-read'

sequencing data to reduce sequencing errors or to assess the likelihood of real nucleotide

variants. We can however show that our algorithmic strategies are already sufficient to address

metabolic rate estimation in a reliable manner. Technical improvements in nanopore sequencing

or their computational processing will strongly improve the accuracy of individual read

sequences and thus detectability of ^{5E}U. The task at hand will be the development of a novel

base-calling algorithm for direct RNA nanopore sequencing with extended base alphabet (A, C,

G, U & ^{5E}U). Furthermore, increased throughput will foster statistical precision of metabolic rate

estimation and will also allow to elucidate low abundant or transient processes.

308   Nanopore-based transcriptomic studies will allow us to monitor the formation of

transcripts, post-transcriptional processing, export and translation at the level of single RNA

isoforms. nano-ID is in principle also transferable to single cell methodologies, to catch

heterogeneity of the RNA population in any state of the cell. This however requires sequencing

library preparation with lower input amounts. The use of ^{5E}U is widely established for *in vivo*

applications in the field such as fluorescence microscopy. We thus envision that nano-ID is in

principle applicable to many types of organisms, cells and conditions.

Methods

Labeling and direct RNA nanopore sequencing of synthetic RNAs. Labeled synthetic RNAs

and synthetic control RNAs are derived from selected RNAs of the ERCC RNA Spike-in Mix

(Ambion) as described in [26]. Characteristics of selected RNAs of the ERCC RNA Spike-in Mix

are listed in (**Supplementary Table 3**). Briefly, selected spike-in sequences were cloned into a

pUC19 cloning vector and verified by Sanger sequencing. For IVT template generation, the

plasmid (3 µg) was linearized using EcoRV-HF (blunt end cut) digestion mix containing

CutSmart buffer and EcoRV-HF enzyme. The digestion mix was incubated at 37 °C for 1 h and

the reaction was terminated adding 1/20 volume of 0.5 M EDTA. Subsequently, DNA was

precipitated in 1/10 volume of 3 M sodium acetate pH 5.2, and 2 volumes of 100 % ethanol at -

20 °C for 15 min. DNA was collected by centrifugation at 4 °C and 16,000 x g for 15 min. The

pellet was washed twice using 75 % ethanol. DNA was air-dried and resuspended in 5 µL of

H_2O at a concentration of 0.1-1.0 µg/µL (quantified by NanoDrop). Synthetic RNAs were *in*

vitro transcribed using the MEGAscript T7 kit (Ambion). *In vitro* transcription (IVT) of

synthetic control RNAs was performed following the manufacturer's instruction. For IVT of

labeled synthetic RNAs, 100 % of UTP (resp. GTP) was substituted with either 5-ethynyl-UTP

(^{5E}U, Jena Bioscience), 5-bromo-UTP (^{5Br}U, Sigma), 5-iodo-UTP (^{5I}U, TriLink BioTechnologies

LLC), 4-thio-UTP (^{4S}U, Jena Bioscience) or 6-thio-GTP (^{6S}G, Sigma). Note that, for performing

a successful IVT with 4-thio-UTP and 6-thio-GTP, only a reduction to 80% substitution gave

successful yield. IVT reactions were incubated at 37 °C. After 4 h, reaction volume was filled up

with H_2O to 40 µL, then 2 µL of TURBO DNase was added and incubated at 37 °C for

additional 15 min. Synthetic RNAs were purified with RNAClean XP beads (Beckman Coulter)

following the manufacturer's instructions. The final synthetic RNA pool contained equal mass of

all respective synthetic RNAs in a given library (**Supplementary Table 1**). RNA was quantified

using Qubit (Invitrogen). RNA quality was assessed with the TapeStation System (Agilent)

Synthetic RNA pools were poly(A)-tailed using the *E. coli* Poly(A) Polymerase (NEB). The

reaction was incubated for 5 min and stopped with 0.1 M EDTA. Spike-ins were then purified

with phenol:chloroform:isoamyl alcohol and precipitated. Poly(A)-tailed synthetic RNA pools

were subsequently subjected to direct RNA nanopore sequencing library preparation (SQK-

RNA001, Oxford Nanopore Technologies) following manufacturer's protocol. All libraries were

sequenced on a MinION Mk1B (MIN-101B) for 20 h, unless reads sequenced per second
stagnated dramatically.

Culturing of human K562 cells. Human K562 erythroleukemia cells were obtained from
DSMZ (Cat. # ACC-10). K562 cells were cultured antibiotic-free in accordance with the DSMZ
Cell Culture standards in RPMI 1640 medium (Thermo Fisher Scientific) containing 10 % heat
inactivated fetal bovine serum (FBS) (Thermo Fisher Scientific), and 1x GlutaMAX supplement
(Thermo Fisher Scientific) at 37 °C in a humidified 5 % CO_2 incubator. Cells used in this study
display the phenotypic properties, including morphology and proliferation rate, that have been
described in literature. Cells were verified to be free of mycoplasma contamination using Plasmo
Test Mycoplasma Detection Kit (InvivoGen). Biological replicates were cultured independently.

[5E]U labeling and direct RNA nanopore sequencing of human K562 cells. K562 cells were
kept at low passage numbers (<6) and at optimal densities (3x10^5 - 8x10^5) during all
experimental setups. Per biological replicate, K562 cells were diluted 24 h before the experiment
was performed (**Supplementary Table 1**). Per [5E]U 60 min sample (4 replicates), cells were
incubated at 37 °C, 5 % CO_2 for 1 h after a final concentration of 500 µM 5-Ethynyluridine ([5E]U,
Jena Bioscience) was added. Per [5E]U 24 h sample (3 replicates), cells were incubated at 37 °C,
5% CO_2 for 24 h. [5E]U was added 3 times during the 24h incubation, i.e. every 8 hours (0h, 8h,
16h) at a final concentration of 500 µM. Control samples were not labeled (3 replicates). Per [5E]U
60 min HS (heat shock) sample (3 replicates), cells were incubated at 42 °C for 5 min (until cell
suspension reached 42 °C), and then [5E]U was added at a final concentration of 500 µM. Further,
heat shock treatments were performed in a water bath (LAUDA, Aqualine AL12) at 42 °C. for 1
h. Temperature was monitored by thermometer. To avoid transcriptional changes by freshly
added growth medium, fresh growth medium was added ~24 h prior to heat shock treatments [43].
Exactly after the labeling duration, cells were centrifuged at 37 °C and 1,500 x g for 2 min. Total
RNA was extracted from K562 cells using QIAzol (Quiagen) according to manufacturer's
instructions. Poly(A) RNA was purified from 1 mg of total RNA using the µMACS mRNA
Isolation Kit (Milteny Biotec) following the manufacturer's protocol. The quality of poly(A)
RNA selection was assessed using the TapeStation System (Agilent). Poly(A) selected RNAs

were subsequently subjected to direct RNA nanopore sequencing library preparation (SQK-

RNA001, Oxford Nanopore Technologies) following manufacturer's protocol with 1000 ng

input. All libraries were sequenced on a MinION Mk1B (MIN-101B) for 48 h, unless reads

sequenced per second stagnated dramatically.

RNA-seq. Two biological replicates of K562 cells were diluted 24 h before the experiment was

performed. Per replicate, 3.6×10^7 cells in growth medium were labeled at a final concentration

of 500 µM 4-thio-uracil (4sU, Sigma-Aldrich), and incubated at 37 °C, 5 % CO_2 for 5 min.

Exactly after 5 min of labeling, cells were harvested at 37 °C and 1,500 x g for 2 min. Total

RNA was extracted from K562 cells using QIAzol according to manufacturer's instructions

except for the addition of 150 ng RNA spike-in mix [26] together with QIAzol. To isolate polyA

RNA from 75 µg of total RNA, two subsequent rounds of purification by Dynabeads

Oligo (dT)$_{25}$ (invitrogen) were performed. Purification based on manufacturer's instructions was

performed twice, using 1 mg of Dynabeads Oligo (dT)$_{25}$ beads for the first round and 0.5 mg for

the second round of purification. The quality of polyadenylated RNA selection was assessed

using RNA ScreenTape on a TapeStation (Agilent). Sequencing libraries were prepared using the

NuGEN Ovation Universal RNA-seq kit according to manufacturer's instructions. Fragments

were amplified by 10 cycles of PCR, and sequenced on an Illumina NextSeq 550 in paired-end

mode with 75 bp read length.

Direct RNA nanopore sequencing data preprocessing of synthetic RNAs. Direct RNA

nanopore sequencing reads were obtained for each of the samples (**Supplementary Table 1**).

FAST5 files were base-called using Albacore 2.3.1 (Oxford Nanopore Technologies) with the

following parameters: read_fast5_basecaller.py -f FLO-MIN106 -k SQK-RNA001. Direct RNA

nanopore sequencing reads were mapped with GraphMap 0.5.2 [44] to the synthetic RNA reference

sequence with the following parameters: graphmap align --evalue 1e-10. Further data processing

was carried out using the R/Bioconductor environment.

Direct RNA nanopore sequencing data preprocessing of human K562 cells. Direct RNA

nanopore sequencing reads were obtained for each of the samples (**Supplementary Table 1**).

FAST5 files were base-called using Albacore 2.3.1 (Oxford Nanopore Technologies) with the

following parameters: read_fast5_basecaller.py -f FLO-MIN106 -k SQK-RNA001. Direct RNA

nanopore sequencing reads were mapped with Minimap2 2.10 [45] to the hg20/hg38 (GRCh38)

genome assembly (Human Genome Reference Consortium) with the following parameters:

minimap2 -ax splice -k14 --secondary=no. Samtools [46] was used to quality filter SAM files,

whereby alignments with MAPQ smaller than 20 (-q 20) were skipped. Further data processing

was carried out using the R/Bioconductor environment and custom python scripts.

Probability of [5E]U-labeled RNA isoform identification based on synthetic RNAs. The

following parameters were collected on the direct RNA nanopore sequencing data of synthetic

RNAs and used to calculate the probability of identification of a [5E]U-labeled RNA isoform as

labeled. Detectability d - the number of [5E]U caused mismatches in the [5E]U-labeled sample.

Background b - the number of U caused mismatches in the unlabeled control sample. Given

these parameters, the probability of identification p can be calculated as the probability of a U-

based mismatch being caused by a [5E]U in the transcript as

$$p = 0.25 \cdot 0.028 \cdot \left(d \cdot (1 - b) \right)$$

with *0.25* - the empirical probability of U content, and labeling efficiency *0.028* - the empirical

probability of a U being replaced by a [5E]U in the labeling process [32]. This then allows to calculate

the probability of labeled RNA identification p^{RNA} as

$$p^{RNA} = 1 - (1 - p)^{\#bases}$$

, the probability, that an RNA contains at least 1 detectable [5E]U.

Definition of transcription units based on the UCSC RefSeq genome assembly GRCh38

(RefSeq-TUs). For each annotated gene, transcription units were defined as the union of all

existing inherent transcript isoforms (UCSC RefSeq GRCh38).

Definition of isoform-independent exonic and intronic regions (constitutive exons and

introns). Isoform-independent exonic and intronic regions were determined using a model for

constitutive exons [47] and constitutive introns respectively based on UCSC RefSeq annotation

(GRCh38).

Isoform determination for human K562 cells. The FLAIR (Full-Length Alternative Isoform

analysis of RNA) algorithm [22] was used for the correction and isoform definition of raw human

K562 direct RNA nanopore reads. Corrected and collapsed isoforms were obtained by adding

short-read data (RNA-seq) to help increase splice site accuracy of the nanopore read splice

junctions (https://github.com/BrooksLabUCSC/FLAIR).

Parameter collection for neural network training and classification. For each read in each

human K562 sample ([5E]U 60 min, Control, [5E]U 24 h & [5E]U 60 min HS) we obtained ~1500

parameters from three different layers: Raw signal (ionic current), base-call event probabilities

and alignment derived mismatch properties. As raw signal, 1193 parameters were gathered

consisting of the raw ionic current measurements gathered for each possible 5-mer of nucleotides

as well as the raw ionic current measurements gathered for each possible 3-mer centered in a 5-

mer. The latter parameters were collected for U-containing and non-U-containing instances. In

addition to that, raw ionic current measurements were gathered for 5-mers with all possible

nucleotides in their center position also for U-containing and non-U-containing instances, as well

as 5-mers exclusively leading or lagging U content. All collected raw signal parameters were z-

score normalized on all non-U-containing instances given the mean values of the pore model on

which the original base-calling algorithm is based provided by Oxford Nanopore Technologies.

As base-call event probabilities, 120 parameters were gathered including 'model state', 'move',

'weights', 'p model state', the probability that 'model state' gave rise to the observation of the

event, the most probable 'model state', the probability that 'p model state' gave rise to the

observation of the event and the probabilities that events may be associated with the certain base

from the event probabilities table provided by the base-calling algorithm. As alignment derived

mismatch properties, 135 parameters were gathered including length of the reads, nucleotide

occurrences, number of nucleotide transitions, number of inserts and deletions on a single

nucleotide basis as well as on a 5-mer basis for U-containing and non-U-containing instances.

Neural network training, validation and classification of human RNA isoforms into [5E]U-

labeled and unlabeled. Neural network was trained on the [5E]U 24 h versus Control samples

under the assumption that [5E]U 24 h sample solely contains labeled reads and the fact that the

Control sample solely contains unlabeled reads. The trained neural network consists of a batch

normalization layer and 3 dense layers with decreasing output shape (**Supplementary Figure**

3a). 2 dropout layers (with 25% dropout) in between regularize the attempted classification.

Training was conducted on 404.201 reads, validation was performed on 173.240 reads in 40

epochs with the R interface to Keras on a TensorFlow backend [48], as

475                    $model <- keras\_model\_sequential()$

476                              $model \%>\%$

477            $layer\_batch\_normalization(input\_shape = 1448) \%>\%$

478        $layer\_dense(units = 64, activation = "relu", input\_shape = 1448) \%>\%$

479                    $layer\_dropout(rate = 0.25) \%>\%$

480              $layer\_dense(units = 8, activation = "relu") \%>\%$

481                    $layer\_dropout(rate = 0.25) \%>\%$

482              $layer\_dense(units = 1, activation = "sigmoid")$

484                          $model \%>\% compile($

485                    $optimizer = optimizer\_rmsprop(),$

486                    $loss = 'binary\_crossentropy')$

The neural network was 5-fold cross-validated with an accuracy of 0.87 and a false discovery

rate (FDR) of 0.025 and used to classify reads of the [5E]U 60 min and [5E]U 60 min HS samples

into [5E]U-labeled and unlabeled. A ROC analysis (1 − specificity vs sensitivity) for all reads of the

test set showed an area under the curve (AUC) of 0.94. For reads with an alignment length larger

than 500 nt and 1000 nt the AUC improved to 0.96. Note that, limiting the neural network

classification to reads produced in the first few hours of sequencing, i.e. reads with a generally

higher accuracy, improves the AUC to 0.98.

Random forest training, validation and classification of human RNA isoforms into [5E]U-

labeled and unlabeled. For validation purposes, a random forest [49] was trained on the [5E]U 24 h

versus Control samples on the same data as the neural network above. The random forest was 5-

fold cross-validated with an accuracy of 0.85 and a false discovery rate (FDR) of 0.32 and used

to classify reads of the [5E]U 60 min sample into [5E]U-labeled and unlabeled.

Poly(A)-tail length determination. Poly(A)-tail length is estimated by identifying the dwell

time of the poly(A)-tail in the nanopore. For each direct RNA nanopore sequencing read, the raw

signal readout of the nanopore in pico-Ampere [pA] was extracted from the FAST5 file. Every

data point above the 99.99% quantile or below the 0.001% quantile was set to the respective cut-

off value for reasons of robustness (**Supplementary Figure 5c, upper panel**). Subsequently

kmeans clustering was used to define two trend lines at 1/3 and 2/3 the distance between the two

cluster centers. The two trend lines were then used to squish the raw data by taking the parallel

minimum or maximum (**Supplementary Figure 5c, lower panel**). A loss score of a piecewise

linear function of two consecutive segments of the trend lines is then used to identify segments

along the squished data points (**Supplementary Figure 5c, middle panel**). The length of the

third identified segment r_j is used to calculate the length of the poly(A)-tail l_j of read j in sample

i as

$$l_j = \underset{j}{\mathrm{median}}(s_j) \cdot \frac{r_j}{hertz_i} + 5$$

with the sequencing read speed s_j of read j in [nt/s] and the frequency $hertz_i$ in [Hz] used in

measuring sample i and 5 additional adenines that are concealed in the flanking 5-mers.

Intron retention ratio. For each RefSeq-TU (UCSC RefSeq GRCh38) the intron retention ratio

for the [5E]U 60 min and [5E]U 60 min HS samples were calculated using the above defined model of

constitutive exons and introns by calculating the ratio of length normalized coverages of the

maximum value for all respective introns and the average of all respective exons. This yielded

358 gene loci with at least 5% intron retention in either of the samples.

RNA stability (degradation rate λ_{ij}, half-life hl_{ij}) and synthesis rate μ_{ij} estimation of

human RNA isoforms. Each neural network classified direct RNA nanopore sequencing read of

the [5E]U 60 min and [5E]U 60 min HS samples was assigned to a FLAIR defined human isoform (or

RefSeq-TU) either as [5E]U-labeled L_{ij} and unlabeled $T_{ij} - L_{ij}$. The resulting counts were

subsequently converted into synthesis rates μ_{ij} and degradation rates λ_{ij} for isoform i in sample

j assuming first-order kinetics as in [24] using the following equations:

$$\lambda_{ij} = -\alpha_j - \frac{1}{t} \cdot log\left(1 - L_{ij}/T_{ij}\right)$$

$$\mu_{ij} = T_{ij}\left(\alpha_j + \lambda_{ij}\right)$$

532   where t is the labeling duration in minutes and α is the growth rate (dilution rate, i.e. the

533   reduction of concentration due to the increase of cell volume during growth) defined as

$$\alpha_j = \frac{log\,(2)}{CCL_j}$$

535   with cell cycle length CCL_j [min]. The half-life hl_{ij} for isoform i in sample j can thus be

536   calculated as

$$hl_{ij} = \frac{log\,(2)}{\lambda_{ij}}$$

in minutes [min].

RNA-seq data preprocessing and antisense bias correction. Paired-end 75 base reads with

additional 6 base reads of barcodes were obtained for each of the samples (**Supplementary**

Table 1).   Reads were demultiplexed and mapped with STAR 2.3.0 [50] to the hg20/hg38

(GRCh38) genome assembly (Human Genome Reference Consortium). Samtools [46] was used to

quality filter SAM files, whereby alignments with MAPQ smaller than 7 (-q 7) were skipped and

only proper pairs (-f2) were selected. Further data processing was carried out using the

R/Bioconductor environment. We used a spike-in (RNAs) normalization strategy essentially as

described [26] to allow observation of antisense bias ratio c_j (ratio of spurious reads originating

from the opposite strand introduced by the reverse transcription reaction). Antisense bias ratios

were calculated for each sample j according to

550
$$c_j = \underset{i}{\mathrm{median}}\left(\frac{k_{ij}^{antisense}}{k_{ij}^{sense}}\right)$$

for all available spike-ins i. Read counts (k_{ij}) for spike-ins were calculated using HTSeq [51]. The

number of transcribed bases (tb_j) for all samples was calculated as the sum of the coverage of

evident (sequenced) fragment parts (read pairs only) for all fragments in addition to the sum of

the coverage of non-evident fragment parts for fragments with an inner mate interval not entirely

overlapping a Refseq annotated intron (UCSC RefSeq GRCh38). The number of transcribed

bases (tb_j) or read counts (k_j) for all features (RefSeq-TUs) were corrected for antisense bias c_j as

follows using the parameter calculated as described above. The real number of read counts or

coverage s_{ij} for transcribed unit i in sample j was calculated as

559
$$s_{ij} = \frac{S_{ij} - c_j A_{ij}}{1 - c_j^2}$$

where S_{ij} and A_{ij} are the observed numbers of read counts or coverage on the sense and antisense

strand. RPKs were calculated upon antisense bias corrected read counts (k_j) falling into the

region of a RefSeq-TU divided by its length in kilobases. Coverages were calculated upon

antisense bias corrected number of transcribed bases (tb_j) falling into the region of a RefSeq-TU

divided by its length in bases.

References

1   Chen, Y. *et al.* Principles for RNA metabolism and alternative transcription initiation
within closely spaced promoters. *Nat Genet* **48**, 984-994, doi:10.1038/ng.3616 (2016).

2   Core, L. J. *et al.* Analysis of nascent RNA identifies a unified architecture of initiation
regions at mammalian promoters and enhancers. *Nat Genet* **46**, 1311-1320,
doi:10.1038/ng.3142 (2014).

3   Pelechano, V., Wei, W. & Steinmetz, L. M. Extensive transcriptional heterogeneity
revealed by isoform profiling. *Nature* **497**, 127-131, doi:10.1038/nature12121 (2013).

4   Turner, R. E., Pattison, A. D. & Beilharz, T. H. Alternative polyadenylation in the
regulation and dysregulation of gene expression. *Semin Cell Dev Biol* **75**, 61-69,
doi:10.1016/j.semcdb.2017.08.056 (2018).

5   Garalde, D. R. *et al.* Highly parallel direct RNA sequencing on an array of nanopores.
Nat Methods, doi:10.1038/nmeth.4577 (2018).

6   Tilgner, H. *et al.* Comprehensive transcriptome analysis using synthetic long-read
sequencing reveals molecular co-association of distant splicing events. *Nat Biotechnol*
33, 736-742, doi:10.1038/nbt.3242 (2015).

7   Li, Y. I. *et al.* RNA splicing is a primary link between genetic variation and disease.
Science **352**, 600-604, doi:10.1126/science.aad9417 (2016).

8   Long, C. *et al.* Correction of diverse muscular dystrophy mutations in human engineered
heart muscle by single-site genome editing. *Sci Adv* **4**, eaap9004,
doi:10.1126/sciadv.aap9004 (2018).

9   Eswaran, J. *et al.* RNA sequencing of cancer reveals novel splicing alterations. *Sci Rep* **3**,
1689, doi:10.1038/srep01689 (2013).

10   Mayr, C. Regulation by 3'-Untranslated Regions. *Annu Rev Genet* **51**, 171-194,
doi:10.1146/annurev-genet-120116-024704 (2017).

11   Falcone, C. & Mazzoni, C. RNA stability and metabolism in regulated cell death, aging
and diseases. *FEMS Yeast Res* **18**, doi:10.1093/femsyr/foy050 (2018).

12   Houseley, J. & Tollervey, D. The many pathways of RNA degradation. *Cell* **136**, 763-
776, doi:10.1016/j.cell.2009.01.019 (2009).

13   Yamaguchi, T. *et al.* The CCR4-NOT deadenylase complex controls Atg7-dependent cell
death and heart function. *Sci Signal* **11**, doi:10.1126/scisignal.aan3638 (2018).

14   Jacob, A. G. & Smith, C. W. J. Intron retention as a component of regulated gene
expression programs. *Hum Genet* **136**, 1043-1057, doi:10.1007/s00439-017-1791-x
(2017).
15   Steijger, T. *et al.* Assessment of transcript reconstruction methods for RNA-seq. *Nat*
Methods **10**, 1177-1184, doi:10.1038/nmeth.2714 (2013).
16   Chang, H., Lim, J., Ha, M. & Kim, V. N. TAIL-seq: genome-wide determination of
poly(A) tail length and 3' end modifications. *Mol Cell* **53**, 1044-1052,
doi:10.1016/j.molcel.2014.02.007 (2014).
17   Subtelny, A. O., Eichhorn, S. W., Chen, G. R., Sive, H. & Bartel, D. P. Poly(A)-tail
profiling reveals an embryonic switch in translational control. *Nature* **508**, 66-71,
doi:10.1038/nature13007 (2014).
18   Zhang, Z. *et al.* Deep-learning augmented RNA-seq analysis of transcript splicing. *Nat*
Methods **16**, 307-310, doi:10.1038/s41592-019-0351-9 (2019).
19   Tapial, J. *et al.* An atlas of alternative splicing profiles and functional associations reveals
new regulatory programs and genes that simultaneously express multiple major isoforms.
Genome Res **27**, 1759-1768, doi:10.1101/gr.220962.117 (2017).
20   Xu, H., Fair, B. J., Dwyer, Z. W., Gildea, M. & Pleiss, J. A. Detection of splice isoforms
and rare intermediates using multiplexed primer extension sequencing. *Nat Methods* **16**,
55-58, doi:10.1038/s41592-018-0258-x (2019).
21   Clark, M. *et al.* Long-read sequencing reveals the splicing profile of the calcium channel
gene CACNA1C in human brain. *bioRxiv*, 260562, doi:10.1101/260562 (2018).
22   Tang, A. D. *et al.* Full-length transcript characterization of SF3B1 mutation
in chronic lymphocytic leukemia reveals downregulation of retained introns. *bioRxiv*
(2018).
23   Dolken, L. *et al.* High-resolution gene expression profiling for simultaneous kinetic
parameter analysis of RNA synthesis and decay. *RNA* **14**, 1959-1972,
doi:10.1261/rna.1136108 (2008).
24   Miller, C. *et al.* Dynamic transcriptome analysis measures rates of mRNA synthesis and
decay in yeast. *Mol Syst Biol* **7**, 458, doi:10.1038/msb.2010.112 (2011).

25 Rabani, M. *et al.* Metabolic labeling of RNA uncovers principles of RNA production and
degradation dynamics in mammalian cells. *Nat Biotechnol* **29**, 436-442,
doi:10.1038/nbt.1861 (2011).

26 Schwalb, B. *et al.* TT-seq maps the human transient transcriptome. *Science* **352**, 1225-
1228, doi:10.1126/science.aad9841 (2016).

27 Herzog, V. A. *et al.* Thiol-linked alkylation of RNA to assess expression dynamics. *Nat*
Methods **14**, 1198-1204, doi:10.1038/nmeth.4435 (2017).

28 Schofield, J. A., Duffy, E. E., Kiefer, L., Sullivan, M. C. & Simon, M. D. TimeLapse-
seq: adding a temporal dimension to RNA sequencing through nucleoside recoding. *Nat*
Methods **15**, 221-225, doi:10.1038/nmeth.4582 (2018).

29 Shendure, J. *et al.* DNA sequencing at 40: past, present and future. *Nature* **550**, 345-353,
doi:10.1038/nature24286 (2017).

30 Clark, M. W., T.; Garcia-Bea, A.; Kleinman, J.; Hyde, T.; Weinberger, D.; Haerty, W.;
Tunbridge, E. Long-read sequencing reveals the splicing profile of the calcium channel
gene CACNA1C in human brain. *bioRxiv* **10.1101/260562**,
doi:https://doi.org/10.1101/260562 (2018).

31 Garalde, D. R. *et al.* Highly parallel direct RNA sequencing on an array of nanopores.
bioRxiv, doi:10.1101/068809 (2016).

32 Jao, C. Y. & Salic, A. Exploring RNA transcription and turnover in vivo by using click
chemistry. *Proc Natl Acad Sci U S A* **105**, 15779-15784, doi:10.1073/pnas.0808480105
(2008).

33 Abe, K. *et al.* Analysis of interferon-beta mRNA stability control after poly(I:C)
stimulation using RNA metabolic labeling by ethynyluridine. *Biochem Biophys Res*
Commun **428**, 44-49, doi:10.1016/j.bbrc.2012.09.144 (2012).

34 Bharmal, H., Clarke, S., Lemire, A., Agnew, B. & Kumar, K. Capture and Analysis of
Newly Synthesized RNA A Novel Enabling Technology to Study High Resolution Gene
Expression. *Journal of Biomolecular Techniques : JBT* **21**, S43-S43 (2010).

35 Rissland, O. S. The organization and regulation of mRNA-protein complexes. *Wiley*
Interdiscip Rev RNA **8**, doi:10.1002/wrna.1369 (2017).

36 Theodorakis, N. G., Zand, D. J., Kotzbauer, P. T., Williams, G. T. & Morimoto, R. I.
Hemin-induced transcriptional activation of the HSP70 gene during erythroid maturation

in K562 cells is due to a heat shock factor-mediated stress response. *Mol Cell Biol* **9**,

3166-3173 (1989).

37   Sistonen, L., Sarge, K. D., Phillips, B., Abravaya, K. & Morimoto, R. I. Activation of

heat shock factor 2 during hemin-induced differentiation of human erythroleukemia cells.

Mol Cell Biol **12**, 4104-4111 (1992).

38   Mathew, A., Mathur, S. K. & Morimoto, R. I. Heat shock response and protein

degradation: regulation of HSF2 by the ubiquitin-proteasome pathway. *Mol Cell Biol* **18**,

5091-5098 (1998).

39   Niskanen, E. A. *et al.* Global SUMOylation on active chromatin is an acute heat stress

response restricting transcription. *Genome Biol* **16**, 153, doi:10.1186/s13059-015-0717-y

(2015).

40   Vihervaara, A. *et al.* Transcriptional response to stress in the dynamic chromatin

environment of cycling and mitotic cells. *Proc Natl Acad Sci U S A* **110**, E3388-3397,

doi:10.1073/pnas.1305275110 (2013).

41   Vihervaara, A. *et al.* Transcriptional response to stress is pre-wired by promoter and

enhancer architecture. *Nat Commun* **8**, 255, doi:10.1038/s41467-017-00151-0 (2017).

42   Shalgi, R., Hurt, J. A., Lindquist, S. & Burge, C. B. Widespread inhibition of

posttranscriptional splicing shapes the cellular transcriptome following heat shock. *Cell*

Rep **7**, 1362-1370, doi:10.1016/j.celrep.2014.04.044 (2014).

43   Mahat, D. B. & Lis, J. T. Use of conditioned media is critical for studies of regulation in

response to rapid heat shock. *Cell Stress Chaperones* **22**, 155-162, doi:10.1007/s12192-

016-0737-x (2017).

44   Sovic, I. *et al.* Fast and sensitive mapping of nanopore sequencing reads with GraphMap.

Nat Commun **7**, 11307, doi:10.1038/ncomms11307 (2016).

45   Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094-

3100, doi:10.1093/bioinformatics/bty191 (2018).

46   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,

2078-2079, doi:10.1093/bioinformatics/btp352 (2009).

47   Bullard, J. H., Purdom, E., Hansen, K. D. & Dudoit, S. Evaluation of statistical methods

for normalization and differential expression in mRNA-Seq experiments. *BMC*

Bioinformatics **11**, 94, doi:10.1186/1471-2105-11-94 (2010).

48   keras: R Interface to 'Keras' (2018).

49   Classification and Regression by randomForest (2002).

50   Dobin, A. & Gingeras, T. R. Mapping RNA-seq Reads with STAR. *Curr Protoc*

Bioinformatics **51**, 11 14 11-19, doi:10.1002/0471250953.bi1114s51 (2015).

51   Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-

throughput sequencing data. *Bioinformatics* **31**, 166-169,

doi:10.1093/bioinformatics/btu638 (2015).

52   Robinson, J. T. *et al.* Integrative genomics viewer. *Nat Biotechnol* **29**, 24-26,

doi:10.1038/nbt.1754 (2011).

Acknowledgments

We would like to thank Johannes Söding, Christian Roth and Dmitry Tegunov (Max Planck Institute for Biophysical Chemistry) for advice on machine learning aspects. We would also like to thank Nikolaos Papadopoulos and Noah Wulff Mottelson (Max Planck Institute for Biophysical Chemistry) for help with python. In addition, we would like to thank Michael Lidschreiber (Max Planck Institute for Biophysical Chemistry) and Brian Lee (UCSC Genomics Institute) for help with the UCSC Genome Browser Track Hub. Moreover, we would like to thank Anna Sawicka and Kristina Žumer (Max Planck Institute for Biophysical Chemistry) for sharing the pUC19 spike-in plasmids. PC was supported by the Deutsche Forschungsgemeinschaft (SFB860, SPP1935), the European Research Council Advanced Investigator Grant TRANSREGULON (grant agreement No 693023), and the Volkswagen Foundation.

Competing interests

The authors declare that no competing interests exist.

Authors' contributions

KM, BS and SG carried out experiments. BS designed and carried out all bioinformatics analysis. BS conceptualized, designed and supervised research. BS and PC prepared the manuscript, with input from all authors.

Figures

Figure 1. Nanopore sequencing-based Isoform Dynamics (nano-ID) combines metabolic

RNA labeling with 'long-read' nanopore sequencing of native RNA molecules. (a)

Experimental schematic of 5EU-labeled RNA isoforms subjected to direct RNA 'long-read'

nanopore sequencing. Metabolic labeling of human K562 cells with the nucleoside analogue 5-

Ethynyluridine (5EU) *in vivo*. Newly-synthesized RNA isoforms will incorporate 5EU instead of

standard uridine (U) residues. This allows to distinguish the newly synthesized RNA isoforms

(Labeled) from pre-existing RNA isoforms (Unlabeled) *in silico* after sequencing the native full-

length molecules on an array of nanopores [5]. 5EU containing RNA isoforms are computationally

traceable and thus allow classification. Identification and quantification of RNA isoforms

subsequently enable assessment of RNA stability, exon usage, intron retention and polyA-tail

length. (b) Experimental schematic to derive synthetic RNAs for nucleoside analogue

benchmark. RNAs were *in vitro* transcribed using either the standard bases A, U, C, G as a

control, or one of the natural bases was exchanged for a nucleoside analogue (shown for ^{5E}U). (c)

Barplot showing the probability of nucleoside analogue identification compared to natural

UTP/GTP based on base-miscalls (**Methods**) of all tested nucleoside analogues (^{5E}U, 5-

bromouridine (^{5Br}U), 5-iodouridine (^{5I}U), 4-thiouridine (^{4S}U) and 6-thioguanine (^{6S}G)). (d) Upper

panel: Base miscalls (colored vertical bars) of the standard base-calling algorithm for synthetic

RNAs containing ^{5E}U instead of U (-^{5E}U-, 3.563 molecules) and synthetic control RNAs (-U-,

15.840 molecules) in comparison to the original sequence (Reference) of an exemplary region on

synthetic RNA 'Spike-in 8' (**Methods, Supplementary Table 3**). Middle panel: Synthetic RNA

sequences with (-^{5E}U-) and without ^{5E}U (-U-) depicted above the reference sequence (Reference).

Lower panel: Alignment of the raw signal readout of the nanopore in pico-Ampere [pA] to the

reference sequence. Synthetic control RNAs (-U-) are shown in black. ^{5E}U containing synthetic

RNAs are shown in red (-^{5E}U-). ^{5E}U containing synthetic RNAs show a clear deviation from the

expected signal level in blue. Blue boxes indicate mean and standard deviation of the pore model

on which the original base-calling algorithm is based.

Figure 2. Direct RNA 'long-read' nanopore sequencing of [5E]U-labeled RNA isoforms in human K562 cells. Upper panel: Illustration of the experimental set-up. Human K562 cells were cultured in the presence of the nucleoside analogue [5E]U for 60 minutes ([5E]U 60 min, 4 replicates) and 24 h ([5E]U 24 h, 3 replicates). Control samples were not labeled (Control, 3 replicates). Lower panel: Genome browser view of direct RNA 'long-read' nanopore sequencing results of the human GAPDH gene locus on chromosome 12 (~8 kbp, chr12: 6,532,405-6,540,375) visualized with the Integrative Genomics Viewer (IGV, version 2.4.10; human hg38) [52]. From top to bottom: raw nanopore sequencing reads (light grey, shown are typical aligned raw reads below the accumulated coverage of all measured reads), corrected and collapsed isoforms (dark grey) determined with the FLAIR algorithm [22] based on raw reads and RefSeq GRCh38 annotation (blue).

Figure 3. Neural network based classification of human RNA isoforms into [5E]U-labeled and unlabeled. (a) Multi-layered data collection scheme. Parameter collection of human K562 samples ([5E]U 60 min, Control & [5E]U 24 h) was realized on three different layers: Raw signal (ionic current), base-call event probabilities and alignment derived U based mismatch properties (**Methods**). Neural network was trained on the [5E]U 24 h versus Control samples with an accuracy of 0.87 and a false discovery rate (FDR) of 0.025 and used to classify reads of the [5E]U 60 min samples into [5E]U-labeled and unlabeled. (b) ROC analysis of 5-fold cross-validated neural network training. Plot shows ROC curves (1 – specificity versus sensitivity) for all reads of the test set (black, alignment length >=0 nt, AUC = 0.94), for reads with an alignment length

larger than 500 nt (grey, alignment length >=500 nt, AUC = 0.96) and for reads with an

alignment length larger than 1000 nt (dashed grey, alignment length >=1000 nt, AUC = 0.96). (c)

Genome browser view of classified direct RNA 'long-read' nanopore sequencing reads of the

human GAPDH gene locus on chromosome 12 (~8 kbp, chr12: 6,532,405-6,540,375) visualized

with the Integrative Genomics Viewer (IGV, version 2.4.10; human hg38) [52]. Unlabeled reads

are shown in grey, [5E]U-labeled reads are shown in red.

Figure 4. Poly(A)-tail length determination of human RNA isoforms. (a) Histogram of

poly(A)-tail length estimates of 714,536 RNA isoforms (mean: 81 nt, median: 73 nt). (b)

Cumulative distribution function of poly(A)-tail length estimates of the RNA calibration strand

(RCS, yeast derived spike-in RNAs that are equipped with a poly(A)-tail of exactly 30 adenines

(ONT, SQK-RNA001)). Vertical solid black line indicates optimal result of 30 nt (median: 30.6,

coefficient of variation: 0.62). Vertical dashed black lines indicate 2-fold in either direction. (c)

Genome browser view of classified direct RNA 'long-read' nanopore sequencing reads with

poly(A)-tail (green) of the human GAPDH gene locus on chromosome 12 (~8 kbp, chr12:

6,532,405-6,540,375) visualized with the Integrative Genomics Viewer (IGV, version 2.4.10;

human hg38) [52].

Figure 5. nano-ID monitors RNA isoform dynamics during heat shock. (a) Experimental set-

up of the heat shock treatment (60 min at 42 °C) in human K562 cells. (b) Boxplot shows intron

retention rate (**Methods**, min 5% in either condition) of 358 gene loci comparing heat shock ([5E]U

60 min HS) against control ([5E]U 60 min). (c) Bar plot shows correlation (Spearman's rank

correlation coefficient) of RNA half-lives and intron retention ratios before and after heat shock

(1,027 loci). The third bar shows the correlation of their respective folds. (d) Boxplot shows

upregulated (red) and downregulated (blue) RNA isoforms upon 60 min of heat shock (42 °C). A

minimum fold change of 1.25 and a maximum p-value of 0.1 was set for calling a significant

expression change. (e) Boxplot shows half-lives of significantly upregulated RNA isoforms

comparing heat shock ([5E]U 60 min HS) against the control ([5E]U 60 min). (f) As in (e) for

significantly downregulated RNA isoforms. (g) Scatter plot with color-coded density of RNA

half-lives and RNA poly(A)-tail lengths in both conditions. Shown are 1,230 highly expressed

RefSeq GRCh38 annotated genes. Correlation is calculated as Spearman's rank correlation

coefficient (0.48) rounded to the second decimal. (h) As in (c) using the RNA poly(A)-tail

lengths (1,230 loci).

Figure 6. nano-ID resolves the characteristics of individual RNA isoforms. (a) Boxplot

shows half-life estimates of RNAs from the human HSPB1 gene locus (chr6:31,813,514-

31,819,942) comparing heat shock (HS, ^{5E}U 60 min HS) against control (Ctrl, ^{5E}U 60 min).

Standard deviation is shown as error bars. Red points depict half-life estimate of merged

replicates in each condition. (b) Boxplot shows the poly(A)-tail length distributions of RNAs

from the human HSPB1 gene locus. 437 RNAs from heat shocked samples (HS, [5E]U 60 min HS)

are compared to 341 RNAs in the respective control sample (Ctrl, [5E]U 60 min). (c) Schematic

shows direct RNA nanopore sequencing derived RNA isoforms at the human HSPB1 gene locus

above annotated transcription start sites (TSSs) from published GRO-cap data generated in K562

cells [2] and RefSeq GRCh38 annotation. Bar plots show RNA isoform half-life fold changes,

poly(A)-tail length fold changes and their respective significance as standard deviation (error

bars) or $-\log_{10}$(p-value). Red lines indicate no fold change or $-\log_{10}$(p-value) with p-value 0.01.

(d) As in (c) for RNA isoforms at the human TAGLN2 gene locus (chr1:159,916,107-

159,927,542). (e) Half-life fold change (y-axis) depicted for RNAs encoded by 306 high

confident gene loci (x-axis). All estimates are supported across biological replicates (n≥3) and

conditions (n=2). Half-life estimates for RNA encoded by the entire gene loci (combined) are

depicted as a black line (sorted in decreasing order). Blue dots represent individual RNA isoform

half-life estimates at respective gene loci (1,169 isoforms in total). Perpendicular blue and black

lines represent standard deviations of individual estimates. For individual RNA isoform half-life

estimates, standard deviations are only shown if not overlapping with the standard deviation of

the respective combined half-life estimates (black).

832