Proteome analysis of xylose metabolism in *Rhodotorula toruloides* during lipid production

Ievgeniia A. Tiukova [1,2], Jule Brandenburg [2], Johanna Blomqvist [2,3], Sabine Sampels [2], Nils Mikkelsen [2], Morten Skaugen [4], Magnus Øverlie Arntzen [4], Jens Nielsen [1], Mats Sandgren [2], Eduard J. Kerkhoven [1]*

[1]Systems and Synthetic Biology, Department of Biology and Biological Engineering, Chalmers University of Technology, Göteborg, Sweden

[2]Department of Molecular Sciences, Swedish University of Agricultural Sciences, Uppsala, Sweden

[3]Faculty of Science and Technology, Norwegian University of Life Sciences, Ås, Norway

[4]Department of Chemistry, Biotechnology and Food Science, Norwegian University of Life Sciences, Ås, Norway

Running title: proteome of *Rhodotorula toruloides*

Keywords: *Rhodotorula toruloides*, proteome, xylose, lipid production

∗Corresponding author: Eduard J Kerkhoven

E-mail: eduardk@chalmers.se

Abstract

Background

Rhodotorula toruloides is a promising platform organism for production of lipids from

lignocellulosic substrates. Little is known about the metabolic aspects of lipid production from

the lignocellolosic sugar xylose by oleaginous yeasts in general and *R. toruloides* in particular.

This study presents the first proteome analysis of the metabolism of *R. toruloides* during

conversion of xylose to lipids.

Results

R. toruloides cultivated on either glucose or xylose was subjected to comparative analysis of

its growth dynamics, lipid composition, fatty acid profiles and proteome. The maximum growth

and sugar uptake rate of glucose-grown *R. toruloides* cells were almost twice that of xylose-

grown cells. Cultivation on xylose medium resulted in a lower final biomass yield although

final cellular lipid content was similar between glucose- and xylose-grown cells. Analysis of

lipid classes revealed the presence of monoacylglycerol in the early exponential growth phase

as well as a high proportion of free fatty acids. Carbon source-specific changes in lipid profiles

were only observed at early exponential growth phase, where C18 fatty acids were more

saturated in xylose-grown cells. Proteins involved in sugar transport, initial steps of xylose

assimilation and NADPH regeneration were among the proteins whose levels increased the

most in xylose-grown cells across all time points. The levels of enzymes involved in the

mevalonate pathway, phospholipid biosynthesis and amino acids biosynthesis differed in

response to carbon source. In addition, xylose-grown cells contained higher levels of enzymes

involved in peroxisomal beta-oxidation and oxidative stress response compared to cells

cultivated on glucose.

Conclusions

The results obtained in the present study suggest that sugar import is the limiting step during

xylose conversion by *R. toruloides* into lipids. NADPH appeared to be regenerated primarily

through PPP although it may also involve malic enzyme as well as alcohol and aldehyde

dehydrogenases. Increases in enzyme levels of both fatty acid biosynthesis and beta-oxidation

in xylose-grown cells was predicted to result in a futile cycle. The results presented here are

valuable for the development of lipid production processes employing *R. toruloides* on xylose-

containing substrates.

Background

Successful replacement of fossil diesel fuels with biological oils currently faces a number of
challenges such as competing technologies, including electric vehicles, as well as unsustainable
biological oils production practices such as clearing rainforests for palm oil cultivation (1, 2).
There is also an increasing competition for lipids from the food sector due to a growing
population, increasing standards of human nutrition, and sustainability concerns. Lipids are
required in animal feeds, for instance oils from wild pelagic fish are used to feed fish in
aquaculture. So-called "single cell oils" (SCOs) are lipids derived from microorganisms, which
represent a potentially more sustainable source of lipids to partially replace fish-derived oils
(3). Some microorganisms can produce lipids from lignocellulose-based substrates. The five-
carbon (C5) monosaccharide xylose is the second most abundant sugar in plant biomass and
its utilization by microorganisms is regarded as a challenge to reach a sustainable valorization
of plant biomass and commercialization of lignocellulosic fuels and chemicals (4). Some
oleaginous yeasts can convert xylose to lipids up to more than 20% of its dry cell weight: 22%
Rhodotorula glutinis (5), 58% *Trichosporon fermentans* (6), 37% *Cryptococcus curvatus* (7)
33% *Cryptococcus albidus* (8), 42% *Rhodotorula toruloides* (9-11). Cultivation on xylose in
combination with glucose was also investigated using *Lipomyces starkeyi* with 61% lipids of
dry cell weight (DCW) produced (12), while engineering of *Yarrowia lipolytica* strain enabled
it to assimilate xylose (13, 14). Microbial production of lipids from xylose was furthermore
investigated in filamentous fungi: *Cunninghamella echinulata* (lipids constituted 58% DCW),
Mortierella isabellina (lipids constituted 66% DCW) (15).

Establishment of sustainable lucrative microbial lipid production in industrial scale includes:
i) use of complex substrate (generated from plant biomass and containing a mixture of C6 and
C5- sugars and lignocellulose-derived inhibitors); ii) energy-consuming aerated cultivation; iii)
production of oil with high value compositional fatty acid (FA) profile including n-3 long chain
polyunsaturated FA; iv) production of value-added byproducts. The basidiomycete yeast *R.*
toruloides is relatively fast growing, can assimilate pentoses (11), is resistant to lignocellulosic
inhibitors (16), has a beneficial FA composition (9) including n-3 linolenic acid, and produces
carotenoids (17). *R. toruloides* is one of the most promising species for lipid production with
reported lipid yields of up to 65.5% DCW under high cell density cultivation with low-nitrogen
feeding (18).

R. toruloides was originally applied in 1980 as a production organism for industrial-scale
production of oil substitutes to replace cocoa butter (19). The mono-unsaturated 18-carbon FA
oleic acid (C18:1 n-9) is the major FA and is accumulated to ~55% of total FA in *R. toruloides*
Y4. Less than half of this amount, ~20 % composes palmitic acid (C16:0, saturated). Stearic

(C18:0) and linoleic (C18:2 n-6) acid can each reach up to 10% of total FA (11, 18, 20). A

number of *R. toruloides* strains have been subjected to genome sequencing (21-25), genetic

engineering protocols are established (26-30) and the yeast is recognized as a novel platform

organism for production of oleochemicals (31, 32). A number of cultivation conditions of *R.*

toruloides have therefore been examined, e.g. growth on mixed substrates (33) or different

cultivation modes on lignocellulosic substrates (34, 35) and lipid production at sulfate (36) or

phosphate-limitation (37). Improved genetically modified strains of *R. toruloides* for

conversion of biomass have been generated (38).

The molecular physiology of *R. toruloides* during lipid production from different sugar

substrates have been investigated using both proteomics (39, 40) and transcriptomics (25, 41,

42) approaches. Meanwhile, the molecular physiology during conversion of xylose into lipids

in *R. toruloides* and in oleaginous yeasts in general has been poorly investigated. Xylose

assimilation involves the enzymes NAD(P)H-dependent xylose reductase (XR) and NADH-

dependent xylitol dehydrogenase (XDH). XR reduces xylose to xylitol, which is then oxidized

by XDH to xylulose. Xylulose is subsequently phosphorylated by xylulose kinase before

entering the pentose-phosphate pathway (PPP). *R. toruloides* also possesses the enzyme

phosphoketolase, which cleaves xylulose-5-phosphate to acetyl phosphate and glyceraldehyde-

3-phosphate (43). Unlike glucose assimilation, xylose assimilation requires NADPH due to the

initial reduction of xylose into xylitol by XR. The cellular NADPH pool can be replenished by

a number of pathways including PPP, the malic enzyme (ME) transhydrogenase cycle or the

cytoplasmatic NADP$^+$-dependent isocitrate dehydrogenase. While xylose utilization by

oleaginous yeasts is poorly understood, a number of studies have investigated xylose

metabolism in ethanol-producing yeasts: e.g. in engineered *Saccharomyces cerevisiae* (44),

Schefferomyces stipitis (45), and *Ogataea polymorpha* (46). These studies reported that the

carbon source affected levels of proteins involved in e.g. glucose repression, sugar transport,

gluconeogenesis and amino acid catabolism. In the current study, comparative whole-proteome

analysis was employed to study lipid formation in *R. toruloides* cultivated on either glucose or

xylose as sole carbon source under conditions of nitrogen limitation.

Results

Batch cultivation of *Rhodotorula toruloides*

Growth, sugar consumption and lipid production of *R. toruloides* in medium containing either

glucose or xylose as a carbon source were monitored in batch cultivations (Fig. 1). The C/N

ratio was set to 75 in order to provide sufficient carbon surplus while averting inhibitory effect

of nitrogen limitation. Cell growth was monitored by OD and DCW measurements every eight

hours. Samples for lipid and protein analysis were taken at early exponential phase (8 or 16 h),

late exponential (16 or 40 hours) and lipid production phase (64 or 96 hours), with the later

sampling-points referring to xylose, as indicated in Fig. 1.

The maximum growth rate of *R. toruloides* on glucose was almost twice that of xylose, which

is also reflected in an uptake rate of glucose during exponential phase that is roughly double

the xylose uptake rate (Table 1). The exponential growth phase on glucose lasted

approximately 24 h compared to 40 h for xylose, after which the observed increase in DCW

and OD slowed down (Fig. 1). The xylose level was lower than the glucose level after 72 h,

even though the growth rate on xylose was lower. Possibly, nitrogen was depleted in xylose-

based medium later than in the medium containing glucose (47-49). Growth on xylose resulted

in lower final biomass yield and concentration as compared to glucose (Table 1, Fig. 1). This

could indicate a higher ATP maintenance requirement of xylose-grown cells during the longer

growth phase as compared to growth on glucose.

Significant lipid accumulation was observed at the end of the exponential growth phase, which

was attributed to nitrogen depletion from the culture medium as commonly observed for

cultivations of oleaginous yeasts (19). Although sugar levels decreased further after the late-

exponential phase, particularly in glucose (Fig. 1, cf. glucose at 64 and 96 h), this only resulted

in a slight increase in lipid content. The final lipid content under both growth conditions was

approximately 51%, while final lipid yields were slightly higher on glucose, following the same

trend as the biomass yield (Table 1). Interestingly, the higher lipid yield on glucose is

comparable to yields reported previously (11), although the previous study reported similar

biomass yields between glucose and xylose, which was not observed in the present study. In

addition, the previous study reported lower lipid content of xylose-grown cells compared to

glucose-grown cells. This disparity between studies could be explained by differences in the

C/N ratio used for cultivation (cf. 65 (11) and 75 (our study)) or elemental medium composition

(50). Meanwhile, *R. toruloides* CCT 0783 showed similar effects of glucose and xylose on the

biomass yield, while lipid yields remained unaffected (34).

Lipid classes

Samples were taken for lipid class profiling to investigate whether the distribution of lipid

classes changed during the cultivation (Fig. 1, Table 2). Monoacylglycerols (MAGs) were

detected in the early exponential phase. MAGs are not commonly reported in yeast lipidomes

(20, 51, 52), although it has been observed in cultivations of *R. toruloides* NP11 (53) and *Y.*

lipolytica (54). The presence of MAGs in *R. toruloides* could be connected with diacylglycerol

lipase activity, which cleaves one FA chain from diacylglycerols (DAGs) (55), as *Rhodotorula*

spp. are used in the food industry as sources of lipases (56). Furthermore, a high proportion of

free FA was observed in *R. toruloides* compared to other species where free FA are commonly

below 5 % of total lipids (57, 58). MAGs were only detected during early exponential phase of

the *R. toruloides* cultivations, while percentage of DAGs and free FA increased towards late

exponential phase. This indicates that the absence of MAGs in late stage is not necessarily due

to absence of lipase activity in *R. toruloides*.

Lipid accumulation in *R. toruloides* is mainly in the form of neutral lipids as triacylglycerols

(TAGs), irrespective of the carbon source (Table 2). The percentage of sterols was also

approximately two-fold higher by the time the cells entered stationary phase compared to early

exponential phase. In contrast, phospholipid proportions remained essentially unchanged,

which was reflected by a pronounced increase in TAGs/phospholipids ratio compared to *S.*

cerevisiae (51) or *Y. lipolytica* (57). The high TAGs/phospholipid ratio in *R. toruloides*

suggested that the accumulated neutral lipids are stored in giant lipid droplets rather than the

cell membrane. The *R. toruloides* lipid droplet proteome was shown to drive a formation of

giant lipid droplets in this organism (53). Even at the start of cultivation, the observed

TAGs/phospholipid ratio in *R. toruloides* was higher than that what has previously been

reported for *S. cerevisiae* (51, 59) and *Y. lipolytica* (57, 60, 61).

No major differences in the proportions of lipid classes in *R. toruloides* were observed between

cultivations containing either xylose and glucose. The high similarity in final relative content

of TAGs in cells cultivated in medium containing glucose and xylose fits well with absence of

differences in final total lipid content in examined cells.

Fatty acids profiles

Samples were taken at different *R. toruloides* growth phases for lipid analysis to determine

whether any changes in lipid classes and content correlates with changes in the FA profile (Fig.

1, Table 3). The dominant FA under all conditions was oleic acid. The final FA compositions

agreed somewhat with previous reports (11, 18). The content of linoleic acid and stearic acid

has previously been reported to be roughly 13 % for each of the total FA content in *R. toruloides*

strains CBS 14 and Y4 (11, 18). However, in the present study we found that the amount of

linoleic acid was almost 1.7 times lower than stearic acid. This difference can be due to

differences in media composition and cultivation conditions: *R. toruloides* cultivations were

carried out at 25°C in the present study compared to 30°C in previous reports (11, 18). Higher

proportions of polyunsaturated FA have previously been reported to accumulate in *R. glutinis*

when this yeast was cultivated at 25°C as compared to 30°C (62). Small amounts of

heptadecenoic (C17:1) and heptadecanoic (C17:0) acid could also be detected in some samples

(Table 3). Production of heptadecenoic acid has previously been reported in the two closely

related species *Rhodotorula babjevae* and *Rhodotorula glutinis* (63, 64).

The second major FA detected was either palmitic or linoleic acid, and this was growth phase

dependent even while the average chain length remained relatively unchanged. Some

differences in the FA chain lengths were observed compared to previous reports of the Y4

strain of *R. toruloides* (18), although the dominance of oleic acid, the average chain length and

saturation levels were in agreement with published data.

Carbon source-specific changes in lipid profiles of *R. toruloides* were observed only at early

exponential phase, where C18 FA were more saturated in xylose compared to glucose. Wiebe

et al. observed a higher desaturation of C18 FA and an increase in palmitic acid proportion in

response to xylose in *R. toruloides* CBS 14 (11), which was not observed in this study.

Combined, it shows that medium composition and cultivation conditions (especially

temperature) more significantly affect FA composition than carbon source and growth phase.

Effect of growth phases on proteome composition of *R. toruloides*

As glucose and xylose assimilation proceed through distinct catabolic pathways, we anticipated

that alternative carbon sources result in adjustments of the *R. toruloides* proteome. To

investigate this, we examined relative proteome differences at different growth phases (Fig. 1).

A total of 2995 proteins (corresponding to 36.7 % of the genome) could consistently be

identified in at least two replicates of each tested condition (Additional file 2). Early and late

exponential and stationary (lipid production) phases were compared separately for glucose and

xylose cultures, in order to evaluate the effect of growth phase.

Of the 2995 identified proteins, 795 and 902 were differentially expressed when comparing

early exponential and with stationary phase for xylose and glucose cultures, respectively

(Additional file 1: Fig. S1, Additional file 3, 4). Gene set analysis (GSA) performed to identify

systemic changes revealed that genes involved in ribosome biogenesis and function (Additional

file 1: Fig. S1) were downregulated during the cultivation on both sugars. These observed

changes are more likely associated with the growth phase rather than lipogenesis, as

transcriptional downregulation of ribosomes is connected to inactivation of the TOR signalling

pathway upon nitrogen depletion (65). A similar response has been reported for the *R.*

toruloides strain NP11 as well as for the oleaginous yeast *Y. lipolytica* (66).

Proteins showing the largest differential expression during lipid accumulation phase on both

glucose and xylose were identified. Proteins increasing 5- to 8-fold in abundance included urea

transporter, NCS2 allantoate transporter, major facilitator superfamily (MFS) transporter of

unknown specificity, siderochrome-iron transporter, amine oxidase, perilipin-like protein and

aspartic proteinase (Additional file 1: Fig. S1, Additional file 3, 4). The observed upregulation

of proteins involved in the assimilation of nitrogen agreed well with the expected cessation of

the nitrogen catabolite repression (NCR) under conditions of nitrogen limitation (25).

Regarding genes that are downregulated at the later growth phase, little consensus could be

found between the two carbon sources (Additional file 1: Fig. S1, Additional file 3, 4).

Carbon source dependent proteome changes

Comparison between carbon source for each time point resulted in the identification of 457,

230 and 181 proteins for early exponential, late exponential and lipid accumulation stage

respectively (Additional file 1: Fig. S2, Additional file 5, 6, 7). Only limited conclusions could

be drawn from gene-set analysis of these results (Fig. 2): proteins involved in ribosomal

function were among the top upregulated proteins in early exponential phase on glucose, which

is consistent with the observed higher growth rate on this carbon sources (Fig. 2, Table 1).

Proteins involved in transport (MFS transporters) and metabolism of C5 sugar (xylitol

dehydrogenase, L-iditol 2-dehydrogenase), NADPH generation (NADPH-dependent alcohol

dehydrogenase; were among the top upregulated proteins on xylose at all time points (3- to 11-

fold increase), while there were little similarities among the top downregulated proteins.

Mitochondrial ME was among the top downregulated proteins in xylose grown yeast at early

stationary phase.

As the untargeted GSA did not elucidate systemic transcriptional responses, we proceeded with

targeted analysis of the largest observed transcriptional changes complemented with metabolic

pathways and genes suspected to be affected by different carbon sources.

Sugar transport. The expression levels of various sugar transporters show a carbon source

specific regulation. Thirteen putative sugar transporters were identified in the proteome dataset,

of which eight (M7WUK2, M7X9W8, M7WRB9, M7WT66, M7X183, M7WLS1, M7X6W5,

M7WXP0) had significantly elevated levels (5.6- to 338-fold) for xylose cultivations as

compared to glucose, suggesting their involvement in xylose transport (Fig. 3 and Additional

file 8). Conversely, three transporters (M7WGD5, M7WWM0, M7WTU3) with 16-fold

elevated levels in cells fermenting glucose are likely involved in glucose transport. The

relatively high level of the transporter M7X183 irrespective of carbon source during

exponential phase and M7WTU3 and M7WLS1 at stationary phase might indicate that these

proteins function as unspecific sugar transporters. The transporter M7WLS1 might have a high

affinity to both sugars, while M7WTU3 to xylose only. Lower levels of sugar transporters

induced by glucose rather than xylose suggest high efficiency of expressed transporters.

Xylose assimilation. Proteins predicted to be involved in xylose assimilation were highly upregulated in the presence of xylose, in particular xylitol dehydrogenase (Fig. 4 and Additional file 8). Four NADPH-dependent XRs where strongly upregulated (6.8-68.5 fold) at all three time points during xylose cultivations. Besides xylose reductase M7X8C7, also aldo/keto reductases M7WPA0, M7X3P9, M7XQJ0 are therefore likely involved in xylose assimilation. Three of these reductases also showed a more moderate (3.7- to 5.6-fold) increase during lipid accumulation on glucose, indicating that these enzymes are under lower glucose repression than xylose reductase M7X8C7.

Apart from induction of proteins involved in xylose assimilation, the absence of glucose from the xylose-containing medium also resulted in a strong upregulation (64-1000 fold) of enzymes involved in assimilation of alternative carbon sources (beta-glucosidase M7WLX6, mandelate racemase/muconate lactonizing enzyme M7WKJ1) (Additional file 5, 6, 7 and Fig. 4).

Glycolytic proteins and TCA cycle. Differences in expression levels of glycolytic and TCA cycle proteins were less pronounced than those of sugar transporters, although this might be expected due to dominance of post-translational regulation of glycolytic enzymes in yeast (67, 68). Apart from upregulation of phosphoglycerate mutase during growth on xylose, no significant differences were observed in the levels of enzymes involved in glycolysis, PPP and phosphoketolase metabolic routes (Additional file 8, Fig. 4). From these data it was therefore not possible to infer whether xylulose-5-phosphate is metabolized via phosphoketolase or the non-oxidative steps of PPP and further glycolysis.

Oleaginous yeasts under nitrogen limitation are known to transfer citrate from the mitochondrion into the cytoplasm thereby reducing the flow of metabolites through the TCA cycle. In contrast to previous work (25) where higher C/N ratios were used, only aconitate hydrotase and succinate dehydrogenase showed moderate decreased levels during lipid accumulation in the present study.

NADPH generation. Each addition of a two-carbon unit during fatty acid elongation requires one molar equivalent of NADPH. The observed upregulation of PPP enzymes during lipid accumulation on both sugar substrates would suggest that NADPH generation in *R. toruloides* occurs mainly through PPP (Fig. 4 and Additional file 8). Such a scenario was further supported by the lack of significant differences in the levels of NAD(P)-dependent isocitrate dehydrogenase or cytosolic ME. Meanwhile, the upregulation of two other enzymes of the transhydrogenation cycle, i.e. pyruvate carboxylase and L-malate dehydrogenase (2.1-fold), suggests that this path has some contribution into regeneration of NADPH during lipogenesis. In addition, we identified five aldehyde dehydrogenases (M7WKL7, M7X6W7, M7WN91,

M7X0N9, M7WMR7) that might use NAD(P) as cofactor. A number of these dehydrogenases

increased their expression during lipid accumulation (Additional file 8), while also some

response to carbon source could be observed. Protein M7WTN9 (potentially mitochondrial),

annotated as alcohol dehydrogenase (NADPH), was among the top upregulated proteins (832

times) on xylose at all time points.

Oleaginicity. The levels of ATP-citrate lyase (M7WHC9), acetyl-CoA carboxylase

(M7XLR4) as well as both subunits of FA synthase (FAS) (M7XM89 and M7WSW5) were

upregulated in *R. toruloides* during conversion of both xylose and glucose into lipids

(Additional file 8). Apart from cytoplasmic ME our results agree with earlier studies showing

upregulation of key enzymes of lipid production machinery during lipogenesis (25, 39).

Levels of pyruvate decarboxylase decreased 2-fold during lipid accumulation on both carbon

sources while levels of alcohol dehydrogenase (ADH; M7XN51) decreased sharply by 588 and

1024-fold on glucose and xylose, respectively. This suggests rechannelling carbon towards the

pyruvate dehydrogenase—citrate synthase—ATP-citrate lyase path rather than to pyruvate

decarboxylase—acetaldehyde dehydrogenase—acetyl CoA synthetase path that provides more

energy efficient carbon utilization.

Beta-oxidation. The *R. toruloides* proteomic data suggested that peroxisomal beta-oxidation

triggered by nitrogen starvation was more pronounced in cells cultivated on xylose-based

substrate than on glucose. The levels of enzymes involved in peroxisomal oxidation

(M7WHA9, M7XII7, M7WMF3, M7XHL1, M7WZ16, M7X3J1) was 8- to 16-fold higher in

xylose than in glucose-grown cells already during the exponential growth phase (Additional

file 8). The lipid production phase coincided with a decrease in levels of some enzymes of

peroxisomal oxidation (M7WZ16 and M7X774) in xylose-grown cells while levels of others

(M7XA05, M7WEV1) increased in glucose-grown cells. These differences might indicate that

TOR signaling is affected by carbon source in line with (69).

Similar trends were observed for mitochondrial beta oxidation: 8-128 times upregulation of the

enzymes (M7X3A8, M7XIH3, M7WUJ6, M7XPW4) happened in xylose grown cells already

in the beginning of cultivation, however in glucose grown cells 2.3-8 times upregulation

(M7X3A8, M7WL19 and M7XPW4) occurred only at lipid accumulation phase.

Simultaneous upregulation of the two inverse pathways beta-oxidation and FA biosynthesis in

cells grown on xylose during induction of lipogenesis is expected to result in a futile cycle,

which may partially explain the lower final biomass yields on xylose as compared to glucose.

TAG synthesis. The majority of enzymes involved in TAG synthesis did not change in protein

levels during *R. toruloides* cultivation with the exception of a 4.1-fold increase in the level of

glycerol-3-phosphate dehydrogenase (M7WSY9) on both carbon sources as well as a 2-fold increase in the level of the phosphatidic acid biosynthetic 1-acyl-sn-glycerol-3-phosphate acyltransferase (M7XIE8) in glucose-grown cells (Additional file 8). An increase in TAG lipase (M7Y014) level was observed during the lipid accumulation phase on xylose, that is probably counteracted by increase of perilipins, which protect lipids in droplets from degradation. The observed increase of TAG coincided with elevated absolute levels of DAGs and free FAs during stationary phase. However, MAGs were only detected during early exponential phase.

Phospholipids. Observed differences in levels of phospholipid biosynthetic enzymes suggest that phospholipid biosynthesis is affected by the choice of carbon source. Xylose-grown cells displayed a 2.5-fold increase in phosphatidate cytidylyltransferase (M7XN88) levels (converting phosphatidic acid to CDP-DAG), 2-fold upregulation of phosphatidyl-*N*-methylethanolamine *N*-methyltransferase M7WQ32 (forming phosphatidyl-*N*-dimethylethanolamine) and 2.3-fold downregulation of CDP-diacylglycerol-inositol 3-phosphatidyltransferase (M7X4H7, synthesising phosphatidyl-1D-myo-inositol) during lipid production phase (Additional file 8). In glucose-treated cells the level of phosphatidylserine decarboxylase M7WRH3 forming phosphatidylethanolamine was upregulated 16 times. Meanwhile, analysis of lipid classes did not reveal any difference in proportion of phospholipids in *R. toruloides* cells in response to carbon source.

Mevalonate pathway. The mevalonate pathway is involved in the synthesis of dolichol, carotenoids and sterols from for acetyl-CoA and ATP and is therefore in competition with FA biosynthesis. Some enzymes of this pathway were regulated differentially in *R. toruloides* in response to carbon source. For instance, during the lipid production phase in glucose-treated cells, the level of acetyl-CoA C-acetyltransferase M7WL18 decreased 3.5-fold. Xylose-grown cells displayed a 2-fold increase in the enzyme geranylgeranyl diphosphate synthase (M7WR53, Additional file 8). Phytoene dehydrogenase (M7XKL7) was upregulated in stationary phase cells irrespective of carbon source.

Amino acids biosynthesis. While we did not anticipate that carbon source would affect other metabolic pathways, we did observe changes in e.g. amino acid biosynthesis. A 2.5-fold decrease in the levels of leucine biosynthetic enzymes was observed in both xylose- and glycose grown cells upon the initiation of the lipid production phase (Additional file 8), while the downregulation of enzymes involved in the biosynthesis of proline (proline synthetase associated protein M7X2S8) and leucine (branched-chain-amino-acid aminotransferase M7WE11 and M7WQ90, acetolactate synthase M7XNS5 and acetolactate synthase small

subunit M7X8E9) was detected in xylose- as compared to glucose-grown cells at the exponential phase. This might be related to the decrease of protein biosynthesis in xylose-treated cells.

The downregulation of enzymes involved in the synthesis of leucine (3-isopropylmalate dehydratase M7X0Z1, branched-chain-amino-acid aminotransferase M7WE11, acetolactate synthase M7XNS5, ketol-acid reductoisomerase M7XVW7) and the interconversion of methionine and cysteine (cystathionine beta-lyase M7WT61) was observed during lipid production in glucose-grown *R. toruloides* cells compared to earlier growth phase. Lipid accumulation in *Y. lipolytica* has previously been shown to be accompanied by downregulation of leucine biosynthesis (66).

Oxidative stress response. Levels of proteins involved in glutathione metabolism increased in *R. toruloides* cells cultivated on xylose but also during lipid production, possibly as a response to oxidative stress induced under these conditions. Levels of lactoylglutathione lyase (M7X2H6), glutathione peroxidase (M7WYV4), glutathione *S*-transferases (M7X890, M7WZE5) and NAD(P)H-dependent *S*-(hydroxymethyl) glutathione dehydrogenase (M7XJI6) increased during exponential phase in xylose-grown cells compared to glucose-grown cells (Additional file 8). Levels of lactoylglutathione lyase (M7X2H6), glutathione *S*-transferases (M7WZE5, M7X890) and glutathione peroxidase (M7WYV4) also increased during the lipid production phase in xylose-grown cells relative to exponential growth phase. The levels of glutathione peroxidase (M7WYV4) and glutathione *S*-transferase (M7X890) also increased during the lipid accumulation phase in glucose-grown *R. toruloides* cells. Formate dehydrogenase (M7XUW4) was detected among top upregulated enzymes during cultivation in xylose-based medium. Formate dehydrogenase is involved in the oxidation of formic acid to carbon dioxide as the final step of glutathione-dependent formaldehyde detoxification (70).

Nitrogen starvation response. A number of parallel changes were observed in both glucose- and xylose-grown *R. toruloides* cells in response to nitrogen starvation. The quicker depletion of nitrogen from glucose-containing medium meant that differences in levels of proteins involved in the nitrogen starvation response were first observed in glucose-grown cells during exponential phase. Therefore the levels of amino acid transmembrane transporters (M7WMG2, M7X7B2, M7WSQ5) and proteins involved in amino acid degradation (branched-chain amino acid aminotransferase, M7WE11; tyrosine aminotransferase, M7WV12; aromatic amino acid aminotransferase I, M7X1I9; L-asparaginase, M7WYU3) appeared lower in xylose-grown *R. toruloides* cells compared to glucose-grown cells during early exponential phase as the xylose-grown cells had not yet fully exhausted their nitrogen source and hence not activated their

nitrogen starvation response (Additional file 8). Further depletion of nitrogen from the medium

during the cultivation in xylose eventually triggered a nitrogen starvation response that was

consistent between the two carbon sources, such as increased levels of glutamate

dehydrogenase, allantoicase, autophagy related proteins and urea and amino acid transporters.

Discussion

A number of previous proteomic and transcriptomic studies have investigated metabolism of

xylose in ethanol-producing yeasts (44-46), while this study of *R. toruloides* represents the first

proteomic investigation of an oleaginous yeast during conversion of xylose to lipids. As the

major constituent of hemicellulose, xylose is the second most abundant monosaccharide in

nature after glucose (4).

Many notable differences were observed between the proteomes of cells cultivated in medium

containing either glucose or xylose. Glucose- and xylose-grown cells had similar lipid content

and profiles, but xylose-grown cells displayed lower biomass yields, in agreement with

previous observations (34). The growth rate of xylose-grown *R. toruloides* cells in exponential

phase was roughly half of the growth rate on glucose. The observed decrease in the levels of

ribosomal proteins and other proteins associated with translation of mRNA in xylose-grown

cells was the most prominent change in proteome between the two sugar substrates and is likely

rather connected with differences in the growth rates.

The lower biomass yield and growth rate are possibly an effect of higher ATP demands by *R.*

toruloides when grown on xylose. In the presence of excess sugar, and in particular during lipid

accumulation, beta-oxidation is a futile cycle, and as proteins involved in beta-oxidation were

upregulated during growth on xylose it is feasible that this futile cycle is underlying the higher

ATP requirements. This is potentially an effect of absence of glucose, which would otherwise

down-regulate beta-oxidation. Abolition of this pathway is therefore expected to have a

significant beneficial impact on not only lipid formation but also on growth in excess of sugar

with more pronounced effect on xylose as compared to glucose. Disruption of peroxisome

biogenesis in *Y. lipolytica* has previously been reported to impair β-oxidation, which resulted

in increased lipid accumulation from glucose (71). However, another study showed that

disruption of peroxisome biogenesis in *R. toruloides* had the opposite effect, namely a

decreased lipid production in glucose medium (72).

There were indications of that sugar transport is the underlying reason for the observed

differences in physiology between glucose- and xylose-grown *R. toruloides* cells. A number of

candidate glucose- and xylose-specific transporters were identified from the *R. toruloides*

proteome datasets. As assimilation on xylose required strongly increased levels of some transport proteins, while only a reduced growth rate could be obtained, it is suggested that the xylose transporters are less efficient than those responsible for glucose transport. It is possible that the applied protein extraction methods have a bias against membrane proteins; however, improving xylose transport could be a valuable strategy for further optimization of *R. toruloides* for use of lignocellulosic biomass.

We have detected expression of *S. cerevisiae* homologs of members of glucose catabolite repression pathway in *R. toruloides*. However, apart from the monosaccharide transporter, M7WTU3 (homolog of *SNF3*), we did not find any homologs to *S. cerevisiae* proteins-members of glucose induction pathway in genome of *R. toruloides*, which in the absence of glucose represses proteins required for glucose assimilation, such as glucose transporters (73). Our results showed that expression of some of glucose transporters was lower in xylose-grown cells. It remains unclear which mechanisms in *R. toruloides* provide function of glucose induction pathway. Possibly, release from strong inhibition of glucose-induced proteins can provide high fitness of microorganism in nutrient poor environment, such as soil and phyllosphere - natural habitats of *R. toruloides* (74).

Analysis of metabolic routes that can be used by *R. toruloides* to assimilate sugars did not elucidate a dominant pathway. As the main catabolic flux of glucose is funneled through glycolysis, this results in a lower flux through xylulose-5-P compared to catabolism of xylose. From this reduced availability of xylulose-5-P one might expect a downregulation of phosphoketolase during growth on glucose, however, its protein level showed no significant change as response to the carbon source. One possibility is that once cells pre-cultured on glucose were transferred to xylose medium, the xylose flux into cells was low enough that PPP enzymes levels did not have to be increased further to ensure efficient xylose catabolism.

The phosphoketolase reaction normally enables higher efficiency of carbon metabolism since bypasses the wasteful decarboxylation pyruvate to acetyl-CoA, which causes a loss of one third of the carbon substrate, while phosphoketolase can produce an acetyl residue directly from a C5-substrate. However, under conditions of excess of carbon and simultaneous nitrogen limitation, the carbon saving phosphoketolase reaction may become unnecessary, which could result in xylulose-5-P being redirected towards PPP. Such a scenario would also explain the decrease in phosphoketolase levels during the lipid production phase on both substrates since the NADPH-regenerating reactions of PPP occur upstream of xylulose-5-P and therefore phosphoketolase is not expected to be regulated by NADPH demand. It was speculated in a previous study that assimilation of xylose via phosphoketolase would enable slightly higher

lipid yields when grown on xylose (0.34 g/g sugar) as compared to growth on glucose (0.32
g/g) (75). It should be noted that partial assimilation of xylose via PPP would still be required
for synthesis of essential nucleic acid precursors as well as regeneration of NADPH for
lipogenesis.

NADPH generation for lipid synthesis was assumed to occur through PPP although the
observed increase of the levels of transhydrogenase pathway enzymes indicates that ME could
contribute to $NADP^+$ reduction during stationary phase. Previous studies showed that in other
oleaginous yeasts, *Y. lipolytica* and *L. starkeyi* PPP provides major contribution to NADPH
pool during lipogenesis (76, 77). It has been previously demonstrated, that overexpression of
genes encoding native ME slightly increased lipid accumulation in *R. toruloides* (72).
Interestingly, an increase in levels of a putative $NADP^+$-dependent alcohol dehydrogenase was
observed, which suggests that oxidation of one or more alcohols into the corresponding
aldehydes could contribute to supply of NADPH pool. Aldehyde dehydrogenase could also
potentially generate NADPH, but the substrate specificity of this enzyme must first be
determined. Lipid synthesis is an NADPH demanding process, but the absence of significant
changes in the levels of NADPH regenerating enzymes in the present study would suggest that
NADPH availability is not limiting lipid accumulation in xylose-grown *R. toruloides* cells.

Additional metabolic processes displaying significant changes in protein levels between
glucose- and xylose-grown *R. toruloides* cells included biosynthesis of amino acids and
phospholipids as well as oxidative stress response. Similar effect of xylose on ethanol
producing yeasts has been reported previously (44-46). Some of these effects, such as
downregulation of leucine biosynthesis and upregulation of xylose transporters during growth
on xylose were reminiscent of proteomic changes observed in ethanol producing yeasts during
cultivation on either xylose or glucose. The increased levels of two of the three enzymes
involved in the glutathione-dependent formaldehyde detoxification pathway during cultivation
in xylose-based medium was curious, as there was no obvious source of formaldehyde
formation equivalent to what occurs during assimilation of methylated substrates such as
methanol or methylated amines.

As the difference in *R. toruloides* growth rate were likely underlying many of the detected
protein levels, it would be beneficial to perform continuous cultivations on both carbon sources
to provide additional insight on the response on protein levels. In addition, multicomponent
lignocellulose-based medium could provide valuable information for further optimization of
hemicellulose fermentation by *R. toruloides*. While proteomics measurements are instrumental
to inform how transcriptional regulation affects metabolism, further augmenting with

transcriptomics analysis could provide a higher depth of analysis due to its larger coverage of the total genome.

Conclusions

This study is the first investigation of proteome of oleaginous yeast *R. toruloides* during conversion of xylose into lipids. Our study showed two times lower maximum growth rate and sugar uptake rate of *R. toruloides* cells in xylose as compared to glucose- based substrate. Proteome analysis revealed significantly lower levels of ribosomal proteins and translation associated factors in xylose-grown cells as compared to glucose-grown cells. In addition, xylose-grown cells contained higher levels of enzymes involved in peroxisomal beta-oxidation and oxidative stress response and lower levels of enzymes involved in leucine biosynthesis compared to cells cultivated on glucose. The levels of enzymes involved in sugar transport and phospholipid biosynthesis differed in response to carbon source. In addition, the proteomic data reported here identifies several potential target sugar transporters (M7WUK2, M7X9W8, M7WRB9, M7WT66, M7X183, M7WLS1) and enzymes involved in peroxisomal beta-oxidation (M7WHA9, M7XII7, M7WMF3, M7XHL1, M7WZ16, M7X3J1) for genetic engineering in order to optimize xylose metabolism.

Methods

Strain and Media

R. toruloides CBS 14 was obtained from the Westerdijk Fungal Biodiversity Institute (Utrecht, the Netherlands) and stored at $-80°C$ in 50% v/v glycerol. For cultivation experiments, frozen *R. toruloides* stocks were revived on solid YPD medium (20 g/l glucose, 20 g/l peptone and 10 g/l yeast extract, 16 g/l agar) and incubated for 3 days at 25°C. *R. toruloides* was precultured in 200 ml YPD liquid medium in 500 ml Erlenmeyer flasks and incubated for 3 days at 25°C on an orbital shaker at 150 rpm. The chemically defined cultivation medium used for growth experiments contained: 1.173 g/l $(NH_4)_2SO_4$; 1.7 g/l yeast nitrogen base without amino acids or ammonium sulfate; 3 g/l KH_2PO_4; 0.5 g/l $MgSO_4$-$7H_2O$; 40 g/l carbon source (either xylose or glucose). The carbon/nitrogen (C/N) ratio of the medium was set to 75. 0.07% (v/v) polypropylene glycol was used to prevent adhesion and foaming of yeast cells to glass surfaces within the bioreactors.

Yeast cultivation

Batch cultivations were performed in 0.7-l bioreactors (Multifors 2, Infors-HTBottmingen, Switzerland). Cultivation experiments for each carbon source (glucose and xylose) for analysis

of lipid accumulation were performed in triplicate. *R. toruloides* precultures were washed twice

with a saline solution (9 g/l NaCl) and then used to inoculate the bioreactors to a final optical

density at 600 nm (OD_{600}) of approx. 1.0 in a total volume of 0.5 l chemically defined

cultivation medium containing either glucose or xylose as carbon source. Batch cultivations

were performed at pH 5.5 and 25 °C. The dissolved oxygen tension (DOT) was kept above 20

by regulating a stirring speed between 200 and 600 rpm at an aeration rate of 1 l/min. Samples

for HPLC analysis as well as OD_{600} and DCW measurements were taken every 8 hours.

Samples for lipid analysis were taken at early exponential phase, late exponential phase and

early stationary phase for each cultivation experiment.

Analytical methods

Cell OD_{600} was measured using an Ultrospec 1100 pro spectrophotometer (GE Healthcare,

Chicago IL). Yeast cell biomass was determined by cell dry weight measurement from 2 ml of

culture samples. The samples were centrifuged at $17,000 \times g$, washed twice with deionized

water, incubated in an oven at 105°C for 72 h and then weighed. Each DCW measurement was

done in triplicate. HPLC measurements of ethanol, acetate and glycerol were performed as

described previously (78). Yeast lipids were extracted using the method described by Folch et

al. (79) with some modifications (80). Briefly, 100 mg of freeze-dried yeast cells were

suspended in 1 ml 1 M HCl and soaked for 15 min followed by 1 h incubation at 75 °C. 5 ml

of chloroform:methanol (2:1, v/v) was added to the sample tube, which was then vortexed for

15 min at maximum speed. The sample tube was centrifuged at $5,000 \times g$ for 3 min at room

temperature followed by the transfer of the lower lipid layer to a tube of known weight. The

extraction was repeated with remaining upper layer of the sample. The lipid phases from the

first and second extractions were pooled followed by evaporation of the solvent under $N_{2\ gas}$

flow. The total amount of lipids was determined gravimetrically. Total lipid content was

determined at the start of fermentation instead of at early exponential phase due to a lack of

material.

Dried lipid samples were dissolved in 0.5 ml hexane and methylated with BF_3 according to

Appelqvist (81). Methyl esters were analysed by gas chromatography (GC) as described

previously (80, 82). The standard mixture 68A (Nu-Check, Elysian, MN) and retention times

were used to identify FA. The double binding or unsaturation index (UI) was calculated as UI

$[\%] = [\%16{:}1 + \%17{:}1 + \%18{:}1 + (\%18{:}2)\cdot2 + (\%18{:}3)\cdot3]/100$.

Lipid class composition was determined according to Olsen & Henderson (1989) (83) with

slight modifications. Samples were diluted to a final concentration of 1 g/l in hexane, and 5 µl

of each sample was then applied with a CAMAG TLC Sampler ATS4 (Camag, Switzerland) 2

cm from the base edge of the TLC plates (pre-coated with silica gel TLC plates (20 × 10 cm;

Silicagel 60; 0.20 mm layer, Merck, Darmstadt, Germany) in 2 mm bands with an application

speed of 250 nl/s. N_2 was used as spray gas. All samples were applied in duplicate, and the

distance between tracks was 9.8 mm. Separation of the lipid classes was executed with a

CAMAG Automatic Developing Chamber 2 (ADC 2) (Camag, Switzerland). Lipid classes

were separated using a hexane:diethyl ether:acetic acid (85:15:2; v/v/v) mobile phase.

Afterwards, separation procedure plates were submerged in a solution of 3 % (w/v) cupric

acetate in 8 % (v/v) phosphoric acid and then charred for 20 min at 140 °C. Quantitative

analysis of the separated lipid classes was done by scanning the plates with a CAMAG TLC

Scanner 3 (Camag, Switzerland). The scanning was performed at a speed of 20 mm/s and a

data resolution of 100 μm/step, with a slit dimension of 6.00 × 0.45 mm at a wavelength of 350

nm. Lipid classes were identified by comparison to external standards (TLC 18-4A, Nu-Chek

Prep, Elysian, USA; Ergosterol, PHR1512, Sigma-Aldrich, Sweden). The mode Savitsky-

Golay 7 was used for data filtering. Manual baseline and peak correction were used when

necessary.

Preparation of samples for proteome analysis

Cell pellets were thawed and resuspended in 500 μl of a solution of 4 % (w/v) SDS, 10 mM

DTT and 100 mM Tris-HCl (pH 8). Resuspended cell pellets were then transferred to 2 ml

FastPrep tubes containing ca 500 μl of a 3:1 (w/w) glass bead mixture of bead sizes 106 μm

(Sigma Aldrich, product no. G4649) and 425-600 μm (Sigma Aldrich, product no. G8772),

respectively. Cells were disrupted using a FastPrep homogenizer (MP Biomedicals, CA) with

three 45-s cycles at full speed interspersed with 5 min cooling on ice between runs. The tubes

were then centrifuged for 15 min at 20,000 × g, and the supernatant was transferred to fresh

tubes. An upper red layer of the supernatant could be observed at this point and was carefully

avoided. The bottom layer was used to generate tryptic peptides according to the Filter Aided

Sample Prep (FASP) protocol (84). Briefly, 50 μl of each FastPrep extract was incubated for

10 min at 95 °C, and thereafter processed according to the FASP protocol (in Microcon YM-

30). Tryptic peptides were desalted using STAGE spin tips, and redissolved in 30 μl loading

solution consisting of 2 % (v/v) acetonitrile and 0.05 % (w/v) trifluoroacetic acid in water.

Approximate peptide concentrations were determined using a NanoDrop 2000 instrument

(ThermoFisher Scientific, MA) and the A205 protocol (85).

LC-MS/MS analysis

Approximately 1 μg of tryptic peptides was loaded onto a trap column (Acclaim PepMap100,

C18, 5 μm, 100 Å, 300 μm i.d. × 5 mm) and then backflushed onto a 50 cm × 75 μm analytical

column (Acclaim PepMap RSLC C18, 2 μm, 100 Å, 75 μm i.d. × 50 cm, nanoViper). A 120

minutes gradient from 4 to 40 % solution B (80 % [v/v] acetonitrile, 0.1% [v/v] formic acid)

was used for separation of the peptides, at a flow rate of 300 nl/min. The Q-Exactive mass

spectrometer was set up as follows (Top12 method): a full scan (300-1600 m/z) at R=70.000

was followed by (up to) 12 MS2 scans at R=35000, using an NCE setting of 28 eV. Singly

charged precursors and precursors with charge>5 were excluded for MS/MS. Dynamic

exclusion was set to 20 seconds.

Bioinformatics

The raw MS data were analyzed using MaxQuant (86) version 1.5.3.40 and proteins were

identified and quantified using the MaxLFQ algorithm (87). The data were searched against

the UniProt *R. toruleoides* proteome (UP000016926; 8,138 sequences) supplemented with

common contaminants such as human keratin and bovine serum albumin. In addition, reversed

sequences of all protein entries were concatenated to the database to allow for estimation of

false discovery rates. The tolerance levels used for matching to the database were 4.5 ppm for

MS and 20 ppm for MS/MS. Trypsin/P was used as digestion enzyme and 2 miscleavages were

allowed. Carbamidomethylation of cysteine was set as fixed modification and protein N-

terminal acetylation, oxidation of methionines and deamidation of asparagines and glutamines

were allowed as variable modifications. All identifications were filtered in order to achieve a

protein false discovery rate (FDR) of 1%. Perseus version 1.5.2.6 (88) were used for data

analysis, and the quantitative values were $\log_2$-transformed, and grouped according to carbon

source. Proteins were only considered detected if they were present in at least two replicates

on one of the two carbon sources. Missing values were imputed by drawing numbers from a

normal distribution centred around the level of detection of the mass spectrometer. Student's

t-tests including permutation-based FDR ($p<0.05$) were used to detect significant differences

between groups and volcano plots were constructed by plotting the negative logarithm of the

p-values against the fold difference between two groups in a scatter plot, as shown in Additional

file 1: Fig. S1. Differential expression between groups was assigned if proteins had a significant

t-test and showed $\geq$ 2-fold difference. Due to increased variation in growth dynamics at early

stationary phase, we used the p values ($p\leq0.05$) instead of permutation-based FDR for t-test

truncation to detect differences in cells cultivated with glucose or xylose as sole carbon source

at the late stage of cultivation. Individual proteins were referenced by their UniProt accession

numbers as described for the *R. toruloides* NP11 genome annotation (25). Gene set analysis

was performed as described earlier (57) using available functional genome annotation (25).

Declarations

Authors' contributions

IAT provided major contributions to study design and performed a major part of laboratory

work, data analysis and writing the manuscript. JBr, JB and SS performed chemical analysis,

data evaluation and was involved in manuscript writing. NM performed parts of the laboratory

work and data analysis. MSk and MOA performed major parts of laboratory work. JN provided

a major contribution for study design, data analysis and final manuscript writing. MS and EK

coordinated the project and provided major contributions to study design, data analysis and

final manuscript writing. All authors reviewed the final manuscript.

Acknowledgements

We thank Dr. Tomas Linder for proofreading the manuscript.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

All the data analyzed in this study are included in this manuscript and its additional files. The

proteomics   data   has   been   deposited   to   the   ProteomeXchange   consortium

(http://proteomecentral.proteomexchange.org) via the PRIDE partner repository (89) with the

dataset identifier PXD012332.

Funding

Financial support was provided by a strategic grant of The Swedish Research Council for

Environment, Agricultural Sciences and Spatial Planning (Formas), grant number 213-2013-

80 and Formas Mobility grant 2016-00767.

References

1. McDonald BC, de Gouw JA, Gilman JB, Jathar SH, Akherati A, Cappa CD, et al.

Volatile chemical products emerging as largest petrochemical source of urban organic

emissions. Science. 2018;359(6377):760-4.

2. Wallington TJ, Anderson JE, Kurtz EM, Tennison PJ. Biofuels, vehicle emissions, and

urban air quality. Faraday Discuss. 2016;189:121-36.

3. Blomqvist J, Pickova J, Tilami SK, Sampels S, Mikkelsen N, Brandenburg J, et al.

Oleaginous yeast as a component in fish feed. Sci Rep. 2018;8(1):15945.

4. Sanchez Nogue V, Karhumaa K. Xylose fermentation as a challenge for
commercialization of lignocellulosic fuels and chemicals. Biotechnol Lett.
2015;37(4):761-72.
5. Easterling ER, French WT, Hernandez R, Licha M. The effect of glycerol as a sole and
secondary substrate on the growth and fatty acid composition of *Rhodotorula glutinis*.
Bioresour Technol. 2009;100(1):356-61.
6. Zhu LY, Zong MH, Wu H. Efficient lipid production with *Trichosporon fermentans*
and its use for biodiesel preparation. Bioresour Technol. 2008;99(16):7881-5.
7. Evans CT, Scragg AH, Ratledge C. Regulation of citrate efflux from mitochondria of
oleaginous and non-oleaginous yeasts by long-chain fatty acyl-CoA esters. Eur J
Biochem. 1983;132(3):617-22.
8. Hansson L, Dostálek M. Influence of cultivation conditions on lipid production by
Cryptococcus albidus. Appl Microbiol Biotechnol. 1986;24(1):12–8.
9. Jagtap SS, Rao CV. Production of D-arabitol from D-xylose by the oleaginous yeast
Rhodosporidium toruloides IFO0880. Appl Microbiol Biotechnol. 2018;102(1):143-
51.
10. RS M. Physiology of lipid accumulation yeast. In: RS M, editor. Single cell oil.
London: Longman; 1988. p. 1-32.
11. Wiebe MG, Koivuranta K, Penttila M, Ruohonen L. Lipid production in batch and fed-
batch cultures of *Rhodosporidium toruloides* from 5 and 6 carbon carbohydrates. BMC
Biotechnol. 2012;12:26.
12. Zhao X, Kong X, Hua Y, Feng B, Zhao Z. Medium optimization for lipid production
through co- fermentation of glucose and xylose by the oleaginous yeast *Lipomyces*
starkeyi. Eur J Lipid Sci Technol. 2008;110(5):405-12.
13. Ledesma-Amaro R, Nicaud JM. Metabolic Engineering for Expanding the Substrate
Range of *Yarrowia lipolytica*. Trends Biotechnol. 2016;34(10):798-809.
14. Li H, Alper HS. Enabling xylose utilization in *Yarrowia lipolytica* for lipid production.
Biotechnol J. 2016;11(9):1230-40.
15. Fakas S, Papanikolaou S, Batsos A, Galiotou-Panayotou M, Mallouchos A, Aggelis G.
Evaluating renewable carbon sources as substrates for single cell oil production by
Cunninghamella echinulata and *Mortierella isabellina*. Biomass Bioenerg.
2009;33(4):573–80.

16. Hu C, Zhao X, Zhao J, Wu S, Zhao ZK. Effects of biomass hydrolysis by-products on oleaginous yeast *Rhodosporidium toruloides*. Bioresour Technol. 2009;100(20):4843-7.

17. Lee JJ, Chen L, Shi J, Trzcinski A, Chen WN. Metabolomic profiling of *Rhodosporidium toruloides* grown on glycerol for carotenoid production during different growth phases. J Agric Food Chem. 2014;62(41):10203-9.

18. Li Y, Zhao Z, Bai F. High-density cultivation of oleaginous yeast *Rhodosporidium toruloides* Y4 in fed-batch culture. Enzyme and Microbial Technology. 2007;41(3):312-7

19. Ratledge C, Wynn J. The biochemistry and molecular biology of lipid accumulation in oleaginous microorganisms. Adv Appl Microbiol. 2002;51:1–52.

20. Wei Y, Siewers V, Nielsen J. Cocoa butter-like lipid production ability of non-oleaginous and oleaginous yeasts under nitrogen-limited culture conditions. Appl Microbiol Biotechnol. 2017;101(9):3577-85.

21. Hu J, Ji L. Draft Genome Sequences of *Rhodosporidium toruloides* Strains ATCC 10788 and ATCC 10657 with Compatible Mating Types. Genome Announc. 2016;4(2).

22. Kumar S, Kushwaha H, Bachhawat AK, Raghava GP, Ganesan K. Genome sequence of the oleaginous red yeast *Rhodosporidium toruloides* MTCC 457. Eukaryot Cell. 2012;11(8):1083-4.

23. Morin N, Calcas X, Devillers H, Durrens P, Sherman DJ, Nicaud JM, et al. Draft Genome Sequence of *Rhodosporidium toruloides* CECT1137, an Oleaginous Yeast of Biotechnological Interest. Genome Announc. 2014;2(4).

24. Sambles C, Middelhaufe S, Soanes D, Kolak D, Lux T, Moore K, et al. Genome sequence of the oleaginous yeast *Rhodotorula toruloides* strain CGMCC 2.1609. Genom Data. 2017;13:1-2.

25. Zhu Z, Zhang S, Liu H, Shen H, Lin X, Yang F, et al. A multi-omic map of the lipid-producing yeast *Rhodosporidium toruloides*. Nat Commun. 2012;3:1112.

26. Coradetti ST, Pinel D, Geiselman GM, Ito M, Mondo SJ, Reilly MC, et al. Functional genomics of lipid metabolism in the oleaginous yeast *Rhodosporidium toruloides*. Elife. 2018;7.

27. Koh CM, Liu Y, Moehninsi, Du M, Ji L. Molecular characterization of KU70 and KU80 homologues and exploitation of a KU70-deficient mutant for improving gene deletion frequency in *Rhodosporidium toruloides*. BMC Microbiol. 2014;14:50.

28. Lin X, Wang Y, Zhang S, Zhu Z, Zhou YJ, Yang F, et al. Functional integration of
multiple genes into the genome of the oleaginous yeast *Rhodosporidium toruloides*.
FEMS Yeast Res. 2014;14(4):547-55.

29. Liu Y, Koh CM, Sun L, Hlaing MM, Du M, Peng N, et al. Characterization of
glyceraldehyde-3-phosphate dehydrogenase gene RtGPD1 and development of genetic
transformation method by dominant selection in oleaginous yeast *Rhodosporidium*
toruloides. Appl Microbiol Biotechnol. 2013;97(2):719-29.

30. Zhang S, Skerker JM, Rutter CD, Maurer MJ, Arkin AP, Rao CV. Engineering
Rhodosporidium toruloides for increased lipid production. Biotechnol Bioeng.
2016;113(5):1056-66.

31. Park YK, Nicaud JM, Ledesma-Amaro R. The Engineering Potential of
Rhodosporidium toruloides as a Workhorse for Biotechnological Applications. Trends
Biotechnol. 2018;36(3):304-17.

32. Yaegashi J, Kirby J, Ito M, Sun J, Dutta T, Mirsiaghi M, et al. *Rhodosporidium*
toruloides: a new platform organism for conversion of lignocellulose into terpene
biofuels and bioproducts. Biotechnol Biofuels. 2017;10:241.

33. Bommareddy RR, Sabra W, Maheshwari G, Zeng AP. Metabolic network analysis and
experimental study of lipid production in *Rhodosporidium toruloides* grown on single
and mixed substrates. Microb Cell Fact. 2015;14:36.

34. Bonturi N, Matsakas L, Nilsson R, Christakopoulos P, Miranda E, Berglund K, et al.
Single cell oil producing yeasts *Lipomyces starkeyi* and *Rhodosporidium toruloides*:
selection of extraction strategies and biodiesel property prediction. Energies.
2015;8:5040–52.

35. Fei Q, O'Brien M, Nelson R, Chen X, Lowell A, Dowe N. Enhanced lipid production
by *Rhodosporidium toruloides* using different fed-batch feeding strategies with
lignocellulosic hydrolysate as the sole carbon source. Biotechnol Biofuels. 2016;9:130.

36. Wu S, Zhao X, Shen H, Wang Q, Zhao ZK. Microbial lipid production by
Rhodosporidium toruloides under sulfate-limited conditions. Bioresour Technol.
2011;102(2):1803-7.

37. Wu S, Hu C, Jin G, Zhao X, Zhao ZK. Phosphate-limitation mediated lipid production
by *Rhodosporidium toruloides*. Bioresour Technol. 2010;101(15):6124-9.

38. Diaz T, Fillet S, Campoy S, Vazquez R, Vina J, Murillo J, et al. Combining
evolutionary and metabolic engineering in *Rhodosporidium toruloides* for lipid

production with non-detoxified wheat straw hydrolysates. Appl Microbiol Biotechnol. 2018;102(7):3287-300.

39. Liu H, Zhao X, Wang F, Li Y, Jiang X, Ye M, et al. Comparative proteomic analysis of *Rhodosporidium toruloides* during lipid accumulation. Yeast. 2009;26(10):553-66.

40. Shi J, Feng H, Lee J, Ning Chen W. Comparative proteomics profile of lipid-cumulating oleaginous yeast: an iTRAQ-coupled 2-D LC-MS/MS analysis. PLoS One. 2013;8(12):e85532.

41. Qi F, Zhao X, Kitahara Y, Li T, Ou X, Du W, et al. Integrative transcriptomic and proteomic analysis of the mutant lignocellulosic hydrolyzate-tolerant *Rhodosporidium toruloides*. Eng Life Sci. 2017;17:249-61.

42. Wang Y, Zhang S, Zhu Z, Shen H, Lin X, Jin X, et al. Systems analysis of phosphate-limitation-induced lipid accumulation by the oleaginous yeast *Rhodosporidium toruloides*. Biotechnol Biofuels. 2018;11:148.

43. Evans CT, Ratledge C. Induction of xylulose-5-phosphate phosphoketolase in a variety of yeasts grown ond-xylose: the key to efficient xylose metabolism. C Arch Microbiol. 1984;139(48).

44. Salusjarvi L, Kankainen M, Soliymani R, Pitkanen JP, Penttila M, Ruohonen L. Regulation of xylose metabolism in recombinant *Saccharomyces cerevisiae*. Microb Cell Fact. 2008;7:18.

45. Yuan T, Ren Y, Meng K, Feng Y, Yang P, Wang S, et al. RNA-Seq of the xylose-fermenting yeast *Scheffersomyces stipitis* cultivated in glucose or xylose. Appl Microbiol Biotechnol. 2011;92(6):1237-49.

46. Kim OC, Suwannarangsee S, Oh D, Kim S, Seo J, Kim C, et al. Transcriptome analysis of xylose metabolism in the thermotolerant methylotrophic yeast *Hansenula polymorpha*. Bioprocess Biosyst Eng. 2013;36:1509.

47. Evans CT, Ratledge C. Influence of Nitrogen Metabolism on Lipid Accumulation by *Rhodosporidium toruloides* CBS 14. Journal of General Microbiology. 1984;130:1705-10.

48. Shen H, Jin G, Hu C, Gong Z, Bai F, Zhao ZK. Effects of dilution rate and carbon-to-nitrogen ratio on lipid accumulation by *Rhodosporidium toruloides* under chemostat conditions. Sheng Wu Gong Cheng Xue Bao. 2012;28(1):56-64.

49. Turcotte G, Kosaric N. The effect of C/N ratio on lipid production by *Rhodosporidium toruloides* ATCC 10788. Biotechnol Lett 1989; 11(9):637.

50. Ismail KS, Sakamoto T, Hasunuma T, Zhao XQ, Kondo A. Zinc, magnesium, and
calcium ion supplementation confers tolerance to acetic acid stress in industrial
Saccharomyces cerevisiae utilizing xylose. Biotechnol J. 2014;9(12):1519-25.

51. Ejsing CS, Sampaio JL, Surendranath V, Duchoslav E, Ekroos K, Klemm RW, et al.
Global analysis of the yeast lipidome by quantitative shotgun mass spectrometry. Proc
Natl Acad Sci U S A. 2009;106(7):2136-41.

52. Klose C, Surma MA, Gerl MJ, Meyenhofer F, Shevchenko A, Simons K. Flexibility of
a eukaryotic lipidome - insights from yeast lipidomics. PLoS One. 2012;7(4):e35063.

53. Zhu Z, Ding Y, Gong Z, Yang L, Zhang S, Zhang C, et al. Dynamics of the lipid droplet
proteome of the Oleaginous yeast *Rhodosporidium toruloides*. Eukaryot Cell.
2015;14(3):252-64.

54. Beopoulos A, Verbeke J, Bordes F, Guicherd M, Bressy M, Marty A, et al. Metabolic
engineering for ricinoleic acid production in the oleaginous yeast *Yarrowia lipolytica*.
Appl Microbiol Biotechnol. 2014;98(1):251-62.

55. Van Heusden GP, Nebohacova M, Overbeeke TL, Steensma HY. The *Saccharomyces*
cerevisiae TGL2 gene encodes a protein with lipolytic activity and can complement an
Escherichia coli diacylglycerol kinase disruptant. Yeast. 1998;14(3):225-32.

56. Strehaiano P, Ramon-Portugal F, Taillandier P. Yeasts as biocatalysts. . In: Querol A,
Fleet G, editors. Yeasts in food and beverages,. Berlin: Springer; 2006.

57. Kerkhoven EJ, Pomraning KR, Baker SE, Nielsen J. Regulation of amino-acid
metabolism controls flux to lipid accumulation in *Yarrowia lipolytica*. NPJ Syst Biol
Appl. 2016;2:16005.

58. Rattray JB, Schibeci A, Kidby DK. Lipids of yeasts. Bacteriol Rev. 1975;39(3):197-
231.

59. Casanovas A, Sprenger RR, Tarasov K, Ruckerbauer DE, Hannibal-Bach HK,
Zanghellini J, et al. Quantitative analysis of proteome and lipidome dynamics reveals
functional regulation of global lipid metabolism. Chem Biol. 2015;22(3):412-25.

60. Beopoulos A, Mrozova Z, Thevenieau F, Le Dall MT, Hapala I, Papanikolaou S, et al.
Control of lipid accumulation in the yeast *Yarrowia lipolytica*. Appl Environ Microbiol.
2008;74(24):7779-89.

61. Hein EM, Hayen H. Comparative Lipidomic Profiling of *Saccharomyces cerevisiae* and
Four Other Hemiascomycetous Yeasts. Metabolites. 2012;2(1):254-67.

62. Granger LM, Perlot P, Goma G, Pareilleux A. Kinetics of growth and fatty acid
production of *Rhodotorula glutinis*. Appl Microbiol Biotechnol. 1992;37(1):13-7.

63. Brandenburg J, Poppele I, Blomqvist J, Puke M, Pickova J, Sandgren M, et al.
Bioethanol and lipid production from the enzymatic hydrolysate of wheat straw after
furfural extraction. Appl Microbiol Biotechnol. 2018;102(14):6269-77.

64. Rezanka T, Kolouchova I, Sigler K. Precursor directed biosynthesis of odd-numbered
fatty acids by different yeasts. Folia Microbiol (Praha). 2015;60(5):457-64.

65. Hardwick JS, Kuruvilla FG, Tong JK, Shamji AF, Schreiber SL. Rapamycin-modulated
transcription defines the subset of nutrient-sensitive signaling pathways directly
controlled by the Tor proteins. Proc Natl Acad Sci U S A. 1999;96(26):14866-70.

66. Kerkhoven EJ, Kim YM, Wei S, Nicora CD, Fillmore TL, Purvine SO, et al. Leucine
Biosynthesis Is Involved in Regulating High Lipid Accumulation in *Yarrowia*
lipolytica. MBio. 2017;8(3).

67. Daran-Lapujade P, Rossell S, van Gulik WM, Luttik MA, de Groot MJ, Slijper M, et
al. The fluxes through glycolytic enzymes in *Saccharomyces cerevisiae* are
predominantly regulated at posttranscriptional levels. Proc Natl Acad Sci U S A.
2007;104(40):15753-8.

68. van den Brink J, Canelas AB, van Gulik WM, Pronk JT, Heijnen JJ, de Winde JH, et
al. Dynamics of glycolytic regulation during adaptation of *Saccharomyces cerevisiae*
to fermentative metabolism. Appl Environ Microbiol. 2008;74(18):5710-23.

69. Zhang J, Vaga S, Chumnanpuen P, Kumar R, Vemuri GN, Aebersold R, et al. Mapping
the interaction of Snf1 with TORC1 in *Saccharomyces cerevisiae*. Mol Syst Biol.
2011;7:545.

70. Popov VO, Lamzin VS. NAD(+)-dependent formate dehydrogenase. Biochem J.
1994;301 (Pt 3):625-43.

71. Xue Z, Sharpe PL, Hong SP, Yadav NS, Xie D, Short DR, et al. Production of omega-
3 eicosapentaenoic acid by metabolic engineering of *Yarrowia lipolytica*. Nat
Biotechnol. 2013;31(8):734-40.

72. Zhang S, Ito M, Skerker JM, Arkin AP, Rao CV. Metabolic engineering of the
oleaginous yeast *Rhodosporidium toruloides* IFO0880 for lipid overproduction during
high-density fermentation. Appl Microbiol Biotechnol. 2016;100(21):9393-405.

73. Kaniak A, Xue Z, Macool D, Kim JH, Johnston M. Regulatory network connecting two
glucose signal transduction pathways in *Saccharomyces cerevisiae*. Eukaryot Cell.
2004;3(1):221-31.

74. Yurkov AM. Yeasts of the soil - obscure but precious. Yeast. 2018;35(5):369-78.

75. Papanikolaou S, Aggelis G. Lipids of oleaginous yeasts. Part I; Biochemistry of single cell oil production. Eur J Lipid Sci Technol. 2011;113:1031–51.

76. Beopoulos A, Nicaud JM, Gaillardin C. An overview of lipid metabolism in yeasts and its impact on biotechnological processes. Appl Microbiol Biotechnol. 2011;90(4):1193-206.

77. Vorapreeda T, Thammarongtham C, Cheevadhanarak S, Laoteng K. Repertoire of malic enzymes in yeast and fungi: insight into their evolutionary functional and structural significance. Microbiology. 2013;159(Pt 12):2548-57.

78. Fredlund E, Blank LM, Schnurer J, Sauer U, Passoth V. Oxygen- and glucose-dependent regulation of central carbon metabolism in *Pichia anomala*. Appl Environ Microbiol. 2004;70(10):5905-11.

79. Folch J, Lees M, Sloane Stanley GH. A simple method for the isolation and purification of total lipides from animal tissues. J Biol Chem. 1957;226(1):497-509.

80. Brandenburg J, Blomqvist J, Pickova J, Bonturi N, Sandgren M, Passoth V. Lipid production from hemicellulose with *Lipomyces starkeyi* in a pH regulated fed-batch cultivation. Yeast. 2016;33(8):451-62.

81. Appelqvist L. Rapid methods of lipid extraction and fatty acid methyl ester preparation for seed and leaf tissue with special remarks on preventing accumulation of lipid contaminants. Chem Sci. 1968;28:551–70

82. Olstorpe M, Pickova J, Kiessling A, Passoth V. Strain- and temperature-dependent changes of fatty acid composition in *Wickerhamomyces anomalus* and *Blastobotrys adeninivorans*. Biotechnol Appl Biochem. 2014;61(1):45-50.

83. Olsen RE, J. HR. The rapid analysis of neutral and polar marine lipids using double-development HPTLC and scanning densitometry. Journal of Experimental Marine Biology and Ecology. 1989;129(2):189-97.

84. Wisniewski JR. Filter-Aided Sample Preparation: The Versatile and Efficient Method for Proteomic Analysis. Methods Enzymol. 2017;585:15-27.

85. Simonian MH. Spectrophotometric determination of protein concentration. Curr Protoc Toxicol. 2004;Appendix 3:A 3G 1-7.

86. Tyanova S, Temu T, Cox J. The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. Nat Protoc. 2016;11(12):2301-19.

87. Cox J, Hein MY, Luber CA, Paron I, Nagaraj N, Mann M. Accurate proteome-wide label-free quantification by delayed normalization and maximal peptide ratio extraction, termed MaxLFQ. Mol Cell Proteomics. 2014;13(9):2513-26.

88. Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, et al. The Perseus
computational platform for comprehensive analysis of (prote)omics data. Nat Methods.
2016;13(9):731-40.

89. Vizcaino JA, Cote RG, Csordas A, Dianes JA, Fabregat A, Foster JM, et al. The
PRoteomics IDEntifications (PRIDE) database and associated tools: status in 2013.
Nucleic Acids Res. 2013;41(Database issue):D1063-9

Table 1. Physiological parameters for batch cultivation of *R. toruloides* CBS 14 in minimal
medium containing either glucose or xylose.

	Glucose	Xylose
Lag phase (h)	8	16
Expontential phase (h)	54	80
μ_{max} (h-1)	0.14	0.07
Final DCW (g/l)	13.8 ± 0.13	11.1 ± 0.1
Starting sugar (g/l)	40.8 ± 0.3	39.4 ± 0.25
Residual sugar (g/l)	3.19 ± 0.81	1.1 ± 1
Final $Y_{X/S}$ (gDW/g glucose^{-1})	0.36 ± 0.004	0.28 ± 0.003
Lipid content at t=0 (%) *	7.59 ± 1.54	
Lipid content at t=2 (%) *	10.58 ± 1.00	13.99 ± 1.5
Lipid content at t=3 (%) *	47.51 ± 1.81	47.05 ± 1.4
Final $Y_{L/S}$ (g/g)	$0.19 \pm 0,01$	$0.15 \pm 0,009$

* T=0 represents the beginning of cultivation, t=2 - late exponential growth phase, t=3 - lipid
accumulation phase.

Table 2. Lipid class composition of *R. toruloides* presented as percentage of total identified
lipids. Samples were collected during early exponential phase (t1) and lipid production phase
(t3). Total lipids were not determined at t1, but instead the values reported here are the mean
total lipid contents between the first two total lipid measurements (Fig. 1), assuming a linear
increase over time. Absolute levels of lipid classes were determined from the mean percentages
of lipid classes and mean value of total lipid content.

	t1				t3			
	Glucose % of lipid classes	mg/gDCW	Xylose % of lipid classes	mg/gDCW	Glucose % of lipid classes	mg/gDCW	Xylose % of lipid classes	mg/gDCW
Phospholipids	15.09 ± 0.34	13.71	16.27 ± 1.5	17.56	4.11 ± 0.34	19.53	3.9 ± 0.21	18.37
Monoglycerols	1.02 ± 0.33	0.93	1.35 ± 0.5	1.46	nd	nd	nd	nd
Diglycerols	8.35 ± 0.7	7.59	7.88 ± 0.26	8.50	4.9 ± 0.3	23.28	4.02 ± 0.36	18.93
Sterols	14.11 ± 0.34	12.82	14.61 ± 0.79	15.76	6.49 ± 0.3	30.83	5.67 ± 0.15	26.71
Free fatty acids	38.15 ± 0.9	34.66	40.77 ± 3.1	43.99	11.9 ± 0.3	56.54	10.80 ± 0.25	50.87
Triacylglycerols	23.82 ± 1.7	21.64	19.39 ± 4.5	20.92	72.64 ± 0.76	345.11	75.61 ± 0.63	356.12
TAG/PL	1.58		1.19		17.66		19.37	
Total lipid content		9.085±1		10.79±1		47.51±1.8		47.1±1.4

Abbreviations: DCW: dry cell weight, TAG: Triacylglycerols, PL: Phospholipids

Table 3. Fatty acid profiles of *R. toruloides* during early exponential phase and lipid
accumulation phase following batch cultivation in medium containing either glucose or xylose
(presented as percentage of total identified, average ± standard deviation). Samples were
collected at the beginning of cultivation (t0) as well as during early exponential phase (t1), late
exponential phase (t2) and lipid production phase (t3). Polyunsaturated fatty acids were
counted for the number of unsaturations according to (18).

	t0	t1		t2		t3	
	Glc = Xyl	Xylose	Glucose	Xylose	Glucose	Xylose	Glucose
C14:0	0.98 ± 0.25	1.03 ± 0.26	0.91 ± 0.03	1.23 ± 0.09	1.25 ± 0.05	1.28 ± 0.04	1.1 ± 0.07
C16:0	14.35 ± 1.9	17.91 ± 2.5	17.16 ± 0.01	26.63 ± 0.22	25.76 ± 0.24	26.73 ± 0.26	25.55 ± 0.3
C16:1	1.39 ± 0.28	0.96 ± 0.1	0.51 ± 0.02	0.98 ± 0.05	0.53 ± 0.01	1.00 ± 0.04	0.57 ± 0.05
C17:0		0.76 ± 0.13					
C17:1	0.75 ± 0.18						
C18:0	3.84 ± 0.44	5.55 ± 0.57	7.63 ± 0.5	8.11 ± 0.28	9.31 ± 0.19	7.73 ± 0.22	7.83 ± 0.56
C18:1(n-7)	66.66 ± 2.28	46.27 ± 3.05	36.99 ± 1.3	45.26 ± 0.5	47.23 ± 0.61	45.07 ± 0.7	47.25 ± 1.06
C18:2(n-6)	6.62 ± 0.33	20.36 ± 4.16	23.56 ± 0.71	13.51 ± 0.69	11.58 ± 0.5	13.83 ± 0.57	13.06 ± 0.68
C18:3(n-3)	3.38 ± 0.86	5.74 ± 1.9	11.25 ± 0.98	2.78 ± 0.15	2.94 ± 0.22	2.99 ± 0.19	3.01 ± 0.25
C24:0	0.9 ± 0.01	1.15 ± 0.13	0.66 ± 0.05	0.73 ± 0.05	0.76 ± 0.02	0.67 ± 0.05	0.91 ± 0.13
Avg length	17.4891	17.5942	17.4104	17.3038	17.3546	17.3084	17.3586
Avg desaturation	0.9218	1.0517	1.1837	0.816	0.7974	0.827	0.8297

Fig. 1. Growth and lipid production kinetics of *R. toruloides* during batch cultivation in
medium containing either glucose (A) or xylose (B). Sample points for fatty acids profile (F),
lipid classes (L) and proteome analysis (P) at early exponential (t1), late exponential (t2) and
lipid production (t3) phase are all indicated. Mean values of three independent cultivation
experiments are shown with error bars indicating one standard deviation.

Fig. 2. GO term enrichment analysis of differentially expressed proteins in *R. toruloides* grown
on glucose or xylose during early exponential growth phase (A), late exponential growth phase
(B), lipid accumulation phase (C). For each GO term showing significant change (rank score
of ≥ 5), the direction and significance of the relative changes in protein levels are shown,
together with the total number of proteins within each GO term.

Fig. 3. Heatmap showing relative abundance levels of proteins involved in sugar transport
during batch cultivation of *R. toruloides* in minimal medium containing either glucose or
xylose.

Fig. 4. The figure shows a map of the central carbon metabolism of *R. toruloides* augmented
with $\log_2$-transformed fold change of protein levels relative to the level at glucose early
exponential phase. For most proteins, color scale ranges from -4 (blue) to +4 (red). For proteins
with very large changes (*), the scale ranges from -4 (blue) to + 10 (red). More gene and protein
names, annotations and expression abundance data are included in Additional file 8.

Additional files

Additional file 1: Fig. 1. GO term enrichment analysis and volcano plots of differentially expressed proteins in *R. toruloides* during either exponential growth phase or lipid production phase in medium containing either glucose (A) or xylose (B). For each GO term showing significant change (rank score of ≥ 5), the direction and significance of the relative changes in protein levels are shown, together with the total number of proteins within each GO term.

Additional file 1: Fig. 2. Volcano plots of differentially expressed proteins in *R. toruloides* cultivated on either glucose or xylose during early exponential growth phase (A), late exponential growth phase (B), or lipid accumulation phase (C)

Additional file 2. Complete list of proteins detected during batch cultivation of *R. toruloides* in minimal medium containing either glucose or xylose and their relative expression values

Additional file 3. Relative levels of differentially expressed proteins in glucose-grown *R. toruloides* during either exponential cultivated phase or lipid production phase.

Additional file 4. Relative levels of differentially expressed proteins in xylose-grown *R. toruloides* during either exponential cultivated phase or lipid production phase.

Additional file 5. Relative levels of differentially expressed proteins in *R. toruloides* cultivated on either glucose or xylose during early exponential growth phase

Additional file 6. Relative levels of differentially expressed proteins in *R. toruloides* cultivated on either glucose or xylose during late exponential growth phase

Additional file 7. Relative levels of differentially expressed proteins in *R. toruloides* grown on either glucose or xylose during lipid accumulation phase

Additional file 8. Relative levels of proteins involved in central carbon metabolism of *R. toruloides.*

1 - 2	hexokinase
3	glucose-6-phosphate isomerase
4	6-phosphofructokinase
5	fructose-bisphosphate aldolase
6	triosephosphate isomerase
7	glyceraldehyde 3-phosphate dehydrogenase
8	phosphoglycerate kinase
9 - 11	phosphoglycerate mutase
12	enolase
13	pyruvate kinase
14	pyruvate carboxylase
15 - 16	malate dehydrogenase
17	malic enzyme
18	pyruvate decarboxylase
19 - 22	aldehyde dehydrogenase
23	acetyl-CoA synthetase
24 - 29	alcohol dehydrogenase
30 - 32	pyruvate dehydrogenase
33 - 34	citrate synthase
35 - 36	aconitate hydrotase
37 - 39	isocitrate dehydrogenase
40 - 41	α-ketoglutarate dehydrogenase
42 - 43	succinyl-CoA synthetase
44 - 47	succinate dehydrogenase
48	fumarate hydratase
49 - 50	malate dehydrogenase
51	ATP citrate synthase
52 - 53	fatty acid synthase
54 - 55	glycerol-3-phosphate dehydrogenase
56	glucose-6-phosphate dehydrogenase
57	6-phosphogluconolactonase
58	6-phosphogluconate dehydrogenase
59	ribose 5-phosphate isomerase
60	ribulose-phosphate 3-epimerase
61	transketolase
62	transaldolase
63 - 64	xylose reductase
65 - 69	xylitol dehydrogenase
70	phosphoketolase

Xylose Early exponential-Ref
Xylose Late lipid accumulation-Ref
Xylose Lipid accumulation-Ref
Ref=Glucose Early exponential
Glucose Late exponential-Ref
Glucose Lipid accumulation-Ref

* -4 ▬ 10
-4 ▬ 4
Log₂FC