Integrative taxonomy confirms the species status of the Himalayan langurs,

Semnopithecus schistaceus **Hodgson 1840**

Kunal Arekar[a,b,*] S. Sathyakumar[b], K. Praveen Karanth[a]

[a]Centre for Ecological Sciences, Indian Institute of Science, Bangalore 560012, India

[b]Wildlife Institute of India, Dehradun 248001, India

*Corresponding author Email – kunal.cyan@gmail.com

Abstract

Taxonomy is replete with groups where the species identity and classification remain
unresolved. One such group is the widely distributed Hanuman langur (Colobinae:
Semnopithecus). For most part of the last century, Hanuman langur was considered to be a
single species with multiple subspecies. Nevertheless, recent studies using an integrative
taxonomy approach suggested that this taxon is a complex, with at least three species.
However, these studies did not include the Himalayan population of the Hanuman langur
whose taxonomic status remains unresolved. The Himalayan population of Hanuman langurs
have been classified as a distinct species with multiple subspecies or have been subsumed
into other species. These classification schemes are wholly based on morphological
characters and which are sometimes insufficient to delimit different species. Here, we have
integrated data from multiple sources viz. morphology, DNA, and ecology to resolve the
taxonomy of the Himalayan langur and to understand its distribution limit. Our results with
three lines of evidence corresponding to three different species concepts show that Himalayan
langur is a distinct species from *S. entellus* of the plains. Additionally, these results did not
show any support for splitting of the Himalayan langur into multiple subspecies. Our study
supports the classification proposed by Hill (1939) and we recommend *Semnopithecus*
schistaceus, Hodgson 1840 as species name for the Himalayan langur and subsume all the
known subspecies into it.

Introduction:

Species is the fundamental unit of study in many fields of biology such as systematics, ecology, evolution, behaviour and many more (de Queiroz, 2007). Thus, the first step in these studies is accurate and unambiguous identification of species for which it is imperative to have a well resolved taxonomy of the group of interest. Nevertheless, taxonomy is replete with "problematic groups" wherein species identity and classification remain unresolved. One such group are the colobine monkeys broadly referred to as Hanuman langur (Genus: *Semnopithecus*, Subfamily: Colobinae). Hanuman langur is a widely distributed primate in the Indian subcontinent (Bishop, 1977; Newton, 1988) which exhibits extensive morphological variation across its range. A multitude of classification schemes have been proposed mostly during early-mid 20th century to resolve the taxonomic status of Hanuman langurs (Brandon-jones, 2004; Ellerman, J. R. and Morrison-Scott, 1966; Groves & C, 2001; Hill, 1939; Napier & Napier, 1967; Pocock, 1928, 1939; Roonwal, 1984; Roonwal & Mohnot, 1977).

Hanuman langur is broadly divided into two categories; the northern type (NT) which is characterised by forward looping tail towards the head and southern type (ST) with the tail looping backwards away from the head (Roonwal, 1979, 1984). The Tapti-Godavari rivers in central India form the borderline with NT distributed to the north and ST distributed to the south of these rivers (Roonwal, 1979, 1984). Recent studies support the splitting of ST Hanuman langur into two species, namely *S. priam* and *S. hypoleucos*, based on an integrative approach wherein multiple lines of evidence from molecular, morphological and ecological data were used (Ashalakshmi, Nag, & Karanth, 2014; Nag, Karanth, & Vasudeva, 2014; Nag, Pramod, & Karanth, 2011). Similarly, genetic and morphological data suggest that the plains population of NT Hanuman langur is a separate species, *S. entellus* (Karanth, Singh, & Stewart, 2010). However, taxonomy of the Himalayan population (hereafter

Himalayan langur) is still unresolved. In order to understand the exact number of species of

Hanuman langur in the Indian subcontinent, it is important to resolve the taxonomy of all the

populations of Hanuman langur.

Himalayan langur is the northernmost population of Hanuman langur (Sugiyama,

1976) distributed in the Himalayan region of India, Nepal and parts of Pakistan and Bhutan

(Blanford, 1888; Pocock, 1939). Altitudinally, they are distributed from the Himalayan

foothills up to 4270m asl (above sea level) (Bishop, 1977) and is one of the few colobine

monkey living in temperate climate (Nijman, 2013), while the rest are distributed

predominantly in tropical regions (Bishop, 1979). Himalayan langurs are classified in the NT

category based on the tail loop character (Roonwal, 1979, 1984). Behavioural characters

suggests that the Himalayan langurs might be distinct from *S. entellus* of the northern plains.

(Bishop, 1975; Bishop, 1979; Curtin, 1982; Sugiyama, 1976), Morphologically, the

Himalayan langur can be distinguished from the plains' population (*S. entellus*) by the tail

carriage pattern (discussed in methods section) and by their pelage – the langurs from the

Himalayas have a bushy white head which is very distinct from the darker grey-brown body

(Bishop, 1979).

A variety of classification schemes have been proposed to resolve the taxonomy of

the Himalayan langurs (Table 1). One of the earliest comprehensive classification of Indian

colobines was by Pocock (1928). He assigned the Himalayan langurs to five subspecies

schistaceus, *ajax*, *achilles*, *lanius* and *hector* under the species *Pithecus entellus*. Later

Pocock (1939) renamed *Pithecus* as *Semnopithecus* with three subspecies under it;

schistaceus, *ajax* and *achilles*. The subspecies *lanius* and *hector* from Pocock's (1928) earlier

classification were not included here. Hill (1939) considered the Himalayan langur to be a

single species *S. schistaceus* with five subspecies *hector*, *lanius*, *achilles*, *schistaceus* and

ajax.

Subsequent classification schemes (Ellerman and Morrison-Scott, 1966; Napier &

Napier, 1967; Roonwal, 1984; Roonwal & Mohnot, 1977) synonymised *Semnopithecus* with

Presbytis and subsumed all the Himalayan species into a single species *P. entellus* along with

the subspecies *entellus* from the northern plains. The subspecies *lanius* (Hill, 1939) was

changed to *lania* and the subspecies *hector* (Hill, 1939) was removed.

Later, Groves (2001) reverted back to using *Semnopithecus* for Hanuman langurs and

recognized three species of Himalayan langur *S. schistaceus*, *S. ajax* and *S. hector*. He

elevated the three subspecies from previous classification schemes to species level. Lastly,

Brandon-Jones (2004) included all the Himalayan species as subspecies of *S. entellus*, except

for *achilles* and *lanius* (Hill, 1939) which he did not include in the classification.

Thus, the Himalayan langur has had a convoluted taxonomic history which fall into

three broad groups of taxonomic schemes (TS). TS1 – Various populations of Himalayan

langurs are placed in multiple subspecies under *Pi. entellus*, *S. entellus* or *P. entellus*

(Pocock, 1928, 1939; Ellerman and Morrison-Scott, 1966; Napier and Napier, 1967; Roonwal

and Mohnot, 1977; Roonwal, 1984; Brandon-Jones et al., 2004). TS2 – Himalayan langurs is

considered a distinct species with multiple subspecies (Hill, 1939). TS3 – Himalayan langurs

is split into multiple species (Groves, 2001).

With the advent of molecular techniques many recent studies have used genetic data

to resolve taxonomic ambiguities in primates of the Indian subcontinent (Arekar, Parigi, &

Karanth, 2019; Ashalakshmi et al., 2014; Chakraborty, Ramakrishnan, Panor, Mishra, &

Sinha, 2007; Karanth, Singh, Collura, & Stewart, 2008; Karanth et al., 2010; Osterholz,

Walter, & Roos, 2008; Wangchuk, Inouye, & Hare, 2008). However, the use of molecular

data does not guarantee a robust description and identification (Will, Mishler, & Wheeler,

2005). Molecular data is often considered as another singular data type, such as the

morphological data, which can be used as a line of evidence to characterise and describe

species. In order to achieve a robust delineation of species, we need to integrate methods

from different discipline (Dayrat, 2005)..

Given this background, we have used data from multiple sources viz. morphology,

DNA and ecology to address the following questions 1) Is Himalayan langur a distinct

species from *S. entellus*? 2) Does Himalayan langur occupy different niche than *S.* entellus?

3) Does Himalayan langur population consist of multiple species/subspecies? And 4) What is

the potential distribution range of the Himalayan langurs.

Materials and Methods

Molecular data

Data collection –

We conducted field work in four Himalayan states of India – Jammu and Kashmir (J&K),

Himachal Pradesh (HP), Uttarakhand and Sikkim. These states were chosen for field work

based on distribution records from past studies (Pocock, 1939; Hill, 1939; Sugiyama, 1975;

Bishop, 1979: Choudhury, 2001) as well as from IUCN website (www.iucnredlist.org).

Furthermore, we included our field data and published sequences from Nepal.

Molecular data was based on 175 faecal samples collected from 46 locations (Table

S1) from across the distribution range of Himalayan langur in India and which included

multiple samples collected from each location. Additionally, five faecal samples of *S. entellus*

were collected from one location in the northern plains (22.88220N, 88.39970E). Fresh faecal

samples were collected by following the troops in the morning and the evening hours.

Samples were collected by two different methods – First, as described in Kawamoto et al.

(2013), a sterile cotton swab was rolled multiple times over the surface of the faeces and

thoroughly rinsed in the lysis buffer (White & Densmore, 1992). Second method involved

collection of the whole faeces which was then stored in absolute alcohol. These samples were

stored at -20 °C in the laboratory until DNA extraction. Samples stored in lysis buffer were

first treated with starch to remove potential PCR inhibitors like bilirubin and bile salts

(Kawamoto et al., 2013; Zhang, Li, Ma, & Wei, 2006) and then DNA was extracted by using

Wizard® SV Gel and PCR Clean-Up System, Promega, Madison, WI, USA and stored in

pure nuclease free water at -20 °C until further use. DNA from whole faecal samples was

extracted using the commercially available QIAamp DNA stool mini kit (QIAGEN Inc.),

following the manufacturer's protocol with slight modifications as mentioned in Mondol et

al. (2009), however, we did not add the carrier RNA (Poly A) (Kishore, Reef Hardy,

Anderson, Sanchez, & Buoncristiani, 2006;). Each extraction had a negative control to

monitor contamination. The quantity of extracted DNA was measured using a NanoDrop

2000 Spectrophotometer (Thermo Fisher Scientific Inc).

PCR amplification and sequencing –

A 700 bp region of mitochondrial cytochrome *b* (Cyt-*b*) gene was PCR amplified using the

primer pair Cytb_278F (5' – GCCTATTTCTACACGTAGGCCG – 3') and Cytb_1052R

(5'– CCAATTGCAATGAAGGGTTGGT – 3'). A 25 µl of reaction was set with standard 1X

reaction buffer premixed with 1.5 mM $MgCl_2$ (New England BioLabs® Inc.), 0.25mM of

dNTPs (Bangalore Genei, Bangalore), 0.3 µM of each primer (Sigma-Aldrich Inc.), 1.5U Taq

polymerase (New England BioLabs® Inc.) and 2 µl of template DNA (DNA concentration

between samples varied from 20 ng/µl to 80 ng/µl). We also added 2 µl of BSA (Bovine

Serum Albumin, Fisher Scientific) to augment the PCR reaction. Further, the template DNA

was diluted to 1:5 (DNA extract: water) ratio and used for the reaction in order to reduce the

amount of PCR inhibitors. The PCR cycling conditions were carried out with initial

denaturation at 94 °C for 5 mins followed by 50 cycles with denaturation at 94 °C for 40 sec,

annealing at 57.5 °C for 30 sec and elongation at 72 °C for 30 sec and final extension at 72

°C for 10 mins. The PCR products were outsourced for purification and sequencing to

Medauxin, Bangalore.

Phylogenetic analysis –

The sequence files obtained were viewed and edited manually in ChromasLite v2.01

(Technelysium Pty Ltd). Sequences Himalayan langur as well as *S. entellus* were also

downloaded from a previous molecular study (Supplementary material, Table S5)

(Ashalakshmi et al., 2014; Khanal, Chalise, Wan, & Jiang, 2018). The sequences were

aligned using MUSCLE algorithm (Edgar, 2004) incorporated in MEGA v7 (Kumar, Stecher,

& Tamura, 2016).

We used jModelTest 2.1.3 (Darriba, Taboada, Doallo, & Posada, 2012) to pick the

best model of sequence evolution. Phylogenetic reconstruction was performed using

Maximum Likelihood (ML) and Bayesian methods. ML analysis was performed in

RAxML7.4.2 incorporated in raxmlGUI v1.3 (Stamatakis, 2006). We used GTR+G model in

RAxML as there is no provision to use other models. 1000 replicates were performed to

assess support for different nodes. We used MrBayes 3.2.2 (Ronquist et al., 2012) to perform

the Bayesian analysis with HKY+G nucleotide substitution model. Two parallel runs with

four chains each were run for 10 million generations with sampling frequency every 1000

generations. Convergence between the two runs was determined based on standard deviation

of split frequencies. The program Tracer v1.6 (Rambaut, A., Suchard, M.A., Xie, D.,

Drummond, 2013) was used to determine stationarity, an effective sample size (ESS) value of

>200 for each parameter was used as a cut-off for run length. The first 25% of trees were

discarded as burn-in.

Hypothesis testing –

We compared the Bayesian tree (Fig. 1), with an *a priori* hypothesis where the phylogeny

was constrained to be consistent with Groves' (2001) three species of Himalayan langurs; *S.*

ajax, *S. hector* and *S. schistaceus*. RAxML7.4.2 incorporated in raxmlGUI v1.3 (Stamatakis,

2006) was used to generate a constraint tree. The likelihood of the constraint tree was then

compared with the best tree using the SH test (Shimodaira & Hasegawa, 1999) and AU test

(Shimodaira, 2002) in PAUP* Version 4.0a (build 164) (Swofford, 2001). For both the SH

and AU test, we performed 10000 bootstrap replicates and the nonparametric bootstrap with

reestimated log likelihoods (RELL) approximation (Kishino, Miyata, & Hasegawa, 1990)

was used for resampling the loglikelihoods.

Morphological data

Data collection –

Morphological data was collected by direct observations during field survey as well as from

photographs. Based on past studies (Nag et al., 2011) and our field observations, six color-

independent morphological characters were used to differentiate between *S. entellus* and the

Himalayan langur. These included the four characters described by Nag et al. (2011) and two

characters unique to Himalayan langurs. Characters specific to Himalayan langur included

tail carriage (Roonwal, 1979, 1984) and demarcation between the head and the body (HBC)

(Bishop, 1979; Groves, 2001; Hill, 1939; Pocock, 1939).Within the NT langurs, two forms of

tail carriage are observed, in *S. entellus* from the northern plains the tail loops over the back

and the tip of the tail hangs perpendicular to the ground, here onwards TC3 (Fig. 1b),

whereas, in the Himalayan populations the tail loops well behind the back and the tip ends

above the base of the tail, here onwards TC4 (Fig. 1a). We coded them TC3 and TC4 to be

consistent with Nag et al.'s (2011) coding system. The tail carriage pattern was recorded

when the individual was walking on a flat surface and not while climbing up or down the hill

and nor while it was running or standing (as per Roonwal, 1984). The Himalayan langur has a

distinct demarcation between the head and the body, the head is bushy and white coloured

distinct from the grey-brown body (Fig. 2a) (Bishop, 1979; Groves, 2001; Hill, 1939; Pocock,

1939). On the other hand, the langurs from the plains (*S. entellus*) have a uniform colour

without any distinction between the head and the body (Fig. 2b). We call this character Head-

Body contrast (HBC) character. Apart from these two characters we also used the four

characters described in Nag et al. (2011), these characters are not seen in Himalayan langurs

and therefore were coded '0'(absent). We recorded these characters by direct observations

using 10 X 50 binoculars (Olympus) and through photographs taken from a digital camera

(Canon PowerShot SX20 IS). These characters were scored for multiple adult individuals per

location.

Analysis –

We typed 65 adult individuals from 29 locations (Supplementary material, Table S6). We

also included one of the southern species from this complex, *S. hypoleucos* (St1 and St2

morphotypes in Nag et al., 2011) in the analysis. The characters were coded as described in

Table 2. All the characters were coded in a way that they are consistent with Nag et al.'s

(2011) coding system. The codes for *S. hypoleucos* were similar to ones used in Nag et al.

(2011).

219   We prepared a character matrix with the terminal taxa in the rows and the six

morphological characters in the columns. Using this matrix, first the mean character

difference was calculated between the individuals in PAUP* Version 4.0a (build 164)

(Swofford, 2001). Then a Neighbour Joining (NJ) tree was built using these distances, with

the default settings. Mid-point rooting was used to root the tree.

Ecological data

Data collection –

For ecological data, we obtained 192 occurrence records of the Himalayan langurs (Available

with authors on request). Out of these, 79 records were from the field surveys conducted for

this study, 58 occurrence records were obtained from previous studies (Khanal et al., 2018;

R.A. Minhas et al., 2018; Minhas et al., 2012) and 55 occurrence records were downloaded

from GBIF (Global Biodiversity Information Facility) database (www.gbif.org). For *S.*

entellus, 69 occurrence records were used, out of these, 21 records were obtained from Nag et

al. (2014) and 48 records were downloaded from GBIF database (www.gbif.org). For the

occurrence records downloaded from GBIF database, we plotted these occurrence records on

the map and included only those records which were falling within the known distribution

zones of the respective taxa. Further, we used 22 environmental layers, 19 were bioclimatic

layers downloaded from www.worldclim.org, one altitude layer (USGS website), two more

layers – slope and aspect were derived from the altitude layer in ArcGIS 10.2.1 (Table S2).

All the layers were of 30 arcsec resolution, projected in WGS84 projection. These bioclimatic

variables were clipped to the region from 68 °E to 97.4 °E and from 6.7 °N to 37 °N using

ArcGIS 10.2.1. These clipped layers were then exported to ASCII format using QGIS

2.18.12. The 22 environmental layers were tested for multicollinearity by calculating

Pearson's correlation coefficient (r). The layers with $r \leq 0.75$ were selected for further

analysis (Supplementary material, Table S3).

Ecological Niche Modelling analysis –

Model selection:

We first performed model selection using the maximum entropy algorithm available in

Maxent v3.4.1 (Phillips, Anderson, & Schapire, 2006). We tested 48 models, each for

Himalayan langur and *S*. *entellus*, by employing different combinations of features and

regularisation multiplier (RM) (Supplementary material, Table S4) in MaxEnt v3.4.1

(Phillips et al., 2006).

MaxEnt algorithm:

Separate Maxent analyses for the Himalayan langur and for *S. entellus* were implemented in

MaxEnt v3.4.1. Maxent was run with the following modifications. Random test percentage

was set to 30%, maximum number of background points was set to 10000 and the replicates

were set to 10 with replicated run type changed to Subsample. 5000 iterations were

performed with the convergence threshold set to 1×10^{-5}. Jackknife test was used to estimate

the contribution of each environmental variable. The feature type and RM value were based

on the best model selected (Table S4). To overcome the sampling bias, a bias file was created

in ArcGIS 10.2.1 by applying Gaussian kernel density function to 10000 background points

(Elith, Kearney, & Phillips, 2010). The output format was chosen as Cloglog (Phillips,

Anderson, Dudík, Schapire, & Blair, 2017). AUC values were examined to check for the

predictive ability of the model.

Schoener's D (Schoener, 1968) was calculated as a measure of niche overlap between

the two distribution models (Warren & Seifert, 2011) implemented in ENMTools (Warren,

Glor, & Turelli, 2010)

Results

Phylogenetic analysis

Out of the total 182 samples subjected to DNA extraction we obtained sequences from 76

samples. We also performed DNA extraction and sequencing for one *S. hypoleucos* sample, it

was used as an outgroup in this study. These 76 sequences contained multiple identical

sequences, so for building the phylogenetic tree, we trimmed the dataset to one unique

sequence per location. Therefore, our final dataset contained 746 bp of Cyt-*b* sequence from

57 samples (Table S5), which included sequences from previous studies.

The Bayesian and ML analysis recovered two major clades, the *S. entellus* clade –

containing the sequences from the northern plains; and the Himalayan clade – containing

sequences from the Himalayan region (Fig 3). Within the Himalayan clade was a well-

supported subclade consisting of haplotypes from the western Himalayas, whereas sequences

from eastern Himalayas did not form a separate cluster. Both, the Bayesian and ML trees,

showed similar topology where in all the major clades were retrieved. Furthermore, in the

Bayesian tree (Fig. 1), two samples i.e.128_Nepal and 129_Nepal were placed within the

clade containing samples from the Indian Himalayan Region (IHR), whereas in the ML tree

(Supplementary material, Fig. S1), these two samples were sister to the above mentioned

clade.

Hypothesis testing

The likelihood score of the best tree was significantly higher than that of the constraint tree

for both SH test and AU tests. Therefore, these trees based on the molecular data did not

support the splitting of Himalayan langurs into three species (Table 3).

Morphological data

Our final NJ tree (Fig. 4) was based on morphological data collected from 64 adult

individuals from 60 localities. The NJ dendrogram retrieved a distinct cluster consisting of

Himalayan samples that was sister to *S. entellus*. The UPGMA method also generated a

similar topology.

Ecological niche modelling

Model prediction –

The best model selected for the Himalayan langurs had features 'LQPTH' and a RM value of

2.5; whereas for *S. entellus*, the best model had features 'Auto' with a RM value of 3 (Table

S4). Based on these models and the environmental variables selected, we obtained different

distribution maps for *S. entellus* and the Himalayan langurs (Fig 5). In the distribution maps,

the warmer colour indicates suitable area whereas the cooler colours indicate unsuitable

areas. The AUC values for the training and test data for the Himalayan langur dataset were

0.9663 and 0.9621, respectively. And the AUC values for the training and test data for *S.*

entellus were 0.889 and 0.87, respectively. These AUC values indicate that the potential

distribution of these species fits well with our data.

Variable selection and their importance –

For the Himalayan langur, precipitation of driest quarter (Bio17) was the highest ranked

variable (Table 5). Jackknife test also illustrates the importance of Bio17 (Supplementary

material, Fig. S4). The response curve (Supplementary material, Fig. S2) shows that the

habitat suitability increases with the precipitation of the driest quarter (Bio17) but eventually

attains stationarity. Annual mean temperature (Bio1) and mean diurnal range (Bio2) were the

next two contributing variables (Table 5).

Precipitation seasonality (Bio15) was the most contributing variable towards the

distribution of *S. entellus* (Table 5). The response curve for Bio15 (Supplementary material,

Fig S3) shows that the values for habitat suitability were high towards higher values of

precipitation seasonality (Bio15), suggesting that *S. entellus* prefers areas with high variation

in rainfall. Mean temperature of warmest quarter (Bio10) and annual mean temperature

(Bio1) were the next two highly contributing variables (Table 5).

Niche overlap between *S. entellus* and Himalayan langur was 17 percent (Schoener's

D value = 0.17). However, when we compare the areas with high probability of distribution

(> 0.75), there was no overlap.

Discussion

The so-called Hanuman langur has been known to be a species complex since a long time.

Nevertheless, recent studies have brought some clarity to their confused taxonomy. These

studies suggest that the so-called Hanuman langur consists of at least three species: *S.*

entellus, distributed in the plains of North India (Karanth et al., 2010); *S. priam* and *S.*

hypoleucos distributed in peninsular India (Nag et al., 2011, Ashalakshmi et al., 2014, Nag et

al., 2014). However, the taxonomic status of the Himalayan population of this complex

remained unresolved. Here we address this issue by applying multiple lines of evidence to

resolve the taxonomic status of the Himalayan langur.

Our molecular analysis based on mitochondrial Cyt-*b* gene retrieved a monophyletic

Himalayan langur distinct from *S. entellus* of the plains. The Himalayan langur is also

morphologically distinct and occupies a specific niche in the Himalayas. Thus, three lines of

evidences, corresponding to different species concepts, suggest that the Himalayan langurs is

separately evolving metapopulation lineage (de Queiroz, 1998). Phylogenetic species concept

(PSC) II (de Queiroz, 1998; Donoghue, 1985; Mishler, 1985) identifies species as

monophyletic groups on the basis of shared derived characters. Our genetic data shows that

Himalayan langur and *S. entellus* are reciprocally monophyletic. Results from the

morphological analysis suggests that the Himalayan langur is a separate species as per the

Phenetic Species Concept (Sokal & Crovello, 1970). Our niche modelling analysis too shows

that these two lineages occupy distinct ecological regions and thus show separation along the

ecological axis (Van Valen, 1976).

Among the plethora of classification schemes only Hill (1939) placed the Himalayan

langurs in a separate species, *S. schistaceus*, with multiple subspecies – *ajax*, *achilles*, *lanius*

and *hector* (Table 1). Our study supports this classification, however there is no support for

further splitting this species into multiple subspecies/species (see result under hypotheses

testing). Therefore, we recommend *Semnopithecus schistaceus* Hodgson, 1840 as a species

name for all the Himalayan langurs and subsume all the earlier described species and
subspecies within it until further detailed taxonomic studies address this issue.

Two of the characters used in this study – HBC and forms of tail carriage (TC3 and
TC4), can be used as field identification characters for differentiating S. *schistaceus* and *S.*
entellus. We do not recommend the use any of the external morphological characters listed in
the earlier classification schemes (Table 1) for distinguishing between different subspecies of
Himalayan langurs. The morphological characters used in earlier classification schemes to
describe these subspecies, are highly plastic, variable, and subjective in nature (Nag et al.,
2011). For instance, in the identification key Pocock (1939) describes *schistaceus* as follows
"General colour paler, salty or greyish-buff; coat shorter and less woolly" and for *achilles* he
writes, "General colour dark earthy brown; coat thick and woolly". Hill (1939) describes
schistaceus as "A slatey-grey race, with shorter, less woolly coat than those found at higher
altitudes". These characters tend to differ based on what month of the year the langur is being
observed and what is the altitude at that location. Roonwal (1981) describes four types of tail
carriage in the NT langurs, however we recorded only two tail carriage types that are used in
this study, the other two tail carriage types have not been observed.

We used ecological niche modelling (ENM) to determine the distribution range of *S.*
schistaceus. It predicted the distribution range of the *S. schistaceus* (Fig. 5a) with high
accuracy (AUC = 0.96). The distribution was mainly governed by precipitation of the driest
quarter (Bio17). Response curves (Supplementary material, Fig. S2) for the top three
contributing variables (Table 4) suggests that *S. schistaceus* prefers areas with high
precipitation but with moderate temperatures. Interestingly, slope and aspect did not
contribute to the model prediction. A recent study (Khanal et al., 2018) also showed that
precipitation plays a major role in the distribution of *S. schistaceus* in Nepal. Langurs in the
Himalaya inhabit broad leaves subtropical forest at lower altitudes and temperate broadleaf

forest at higher altitudes (Bishop, 1979; Curtin, 1982; Minhas et al., 2013; Sayers &

Norconk, 2008; Sugiyama, 1976). Occasionally, these langurs are found in temperate

coniferous forests above 4000m altitude (Bishop, 1977). These forests receive high rainfall

during the monsoon as well as precipitation in the form of snowfall in winter (Bhattarai &

Vetaas, 2003; J. S. Singh & Singh, 1987; P. Singh, Ramasastri, & Kumar, 1995).

We also used ENM to determine if the ecological niches of *S. schistaceus* and *S.*

entellus are separate. The ENM distinctly demarcated ecological niches of these two species.

The AUC value for both the distribution models was significant, implying that the results

greatly differ from the random predictions. Precipitation was the most important variable in

demarcating these two species, with *S. entellus* requiring less precipitation and the Himalayan

langurs need higher precipitation. Bishop (1979) also pointed out that climate is the factor

that governs the distribution of the *S. schistaceus* and *S. entellus*. We also checked for niche

overlap between these two species to determine if they are divergent in their ecological axis.

The niche overlap between these two taxa was not significant suggesting that their ecological

niches are separate.

This study provides a comprehensive evidence for elevating the Himalayan langur to

a species that is distinct from *S. entellus* of the northern plains. Based on our results, we

recommend that this species be assigned to *S. schistaceus* as per Hill (1939).

Acknowledgments

We would like to thank the Department of Biotechnology, Government of India for providing

funds for laboratory work. We would like to thank Dr. Chetan Nag and Ms. Mehreen Khaleel

for providing some of the samples used in this study. Ms. Himani Nautiyal provided

photographs from Bhutan for the morphological analysis. Dr. Bilal Habib for help during

sample collection and Ms. Neha Tiwari for help in lab work. We would also like to thank the

forest departments of Uttarakhand, Himachal Pradesh and Sikkim for providing necessary

collection permits. Additionally, thanks to Jammu and Kashmir forest department for permits

provided to Mehreen Khaleel. KA would like to thank the Rufford foundation and Primate

Conservation Inc. (PCI) for providing funding for field work. Dr. Aravind Madhyastha and

Mr. Bipin Charles for guidance in the niche modelling analysis.

References

Arekar, K., Parigi, A., & Karanth, P. K. (2019). The convoluted evolutionary history of the

capped-golden langur lineage (Cercopithecidae: Colobinae) – concatenation versus

coalescent analyses. *BioRxiv*. https://doi.org/http://dx.doi.org/10.1101/508929

Ashalakshmi, N. C., Nag, K. S. C., & Karanth, K. P. (2014). Molecules support morphology:

species status of South Indian populations of the widely distributed Hanuman langur.

Conservation Genetics, *16*(1939), 43–58. https://doi.org/10.1007/s10592-014-0638-4

Bhattarai, K. R., & Vetaas, O. R. (2003). Variation in plant species richness of different life

forms along a subtropical elevation gradients in the Himalayas, East Nepal. *Global

Ecology   and   Biogeography*,   *12*(4),   327–340.   https://doi.org/10.1046/j.1466-

822X.2003.00044.x

Bishop, N. H. (1975). *Vocal behavior of adult male langurs (presbytis entellus) in a high

altitude environment*. PhD Thesis, University of California, Berkeley.

Bishop, N. H. (1977). Langurs living at high altitudes. *Journal of The Bombay Natural

History Society*, *74*(3), 518–520.

Bishop, N. H. (1979). Himalayan langurs: Temperate colobines. *Journal of Human

Evolution*. https://doi.org/10.1016/0047-2484(79)90092-7

Blanford, W. T. (1888). *Mammalia,* in *The Fauna of British India*. London: Taylor &

Francis.

Brandon-jones, D. (2004). a Taxonomic Revision of the Langurs and Leaf Monkeys

(Primates: Colobinae) of South Asia. *Zoo's Print Journal*, *19*, 1552–1594.

Chakraborty, D., Ramakrishnan, U., Panor, J., Mishra, C., & Sinha, A. (2007). Phylogenetic

relationships and morphometric affinities of the Arunachal macaque *Macaca munzala*, a

newly described primate from Arunachal Pradesh, northeastern India. *Molecular

Phylogenetics   and   Evolution*,   *44*(2),   838–849.   https://doi.org/10.1016/j.ympev.

2007.04.007

Choudhury, A. (2001). Primates in Northeast India: An Overview of their Distribution and Conservation Status. *Envis Bulletin: Wildlife and Protected Areas*, *1*, 92–101.

Curtin, R. . (1982). Range Use of Gray Langurs in Highland Nepal. *Folia Primatologica*, *38*, 1–18.

Darriba, D., Taboada, G. L., Doallo, R., & Posada, D. (2012). jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods*, *9*(8), 772–772. https://doi.org/10.1038/nmeth.2109

Dayrat, B. (2005). Towards integrative taxonomy. *Biological Journal of the Linnean Society*, *85*(3), 407–415. https://doi.org/10.1111/j.1095-8312.2005.00503.x

de Queiroz, K. (1998). The general lineage concept of species, and the process of speciation: A conceptual unification and terminological recommendations. In *Endless forms: Species and speciation* (pp. 57–75). Oxford University Press, New York.

de Queiroz, K. (2007). Species concepts and species delimitation. *Systematic Biology*, *56*(6), 879–886. https://doi.org/10.1080/10635150701701083

Donoghue, M. J. (1985). A Critique of the Biological Species Concept and Recommendations for a Phylogenetic Alternative. *The Bryologist*, *88*(3), 172–181. https://doi.org/10.2307/3243026

Edgar, R. C. (2004). MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, *32*(5), 1792–1797. https://doi.org/10.1093/nar/gkh340

Elith, J., Kearney, M., & Phillips, S. (2010). The art of modelling range-shifting species. *Methods in Ecology and Evolution*, *1*(4), 330–342. https://doi.org/10.1111/j.2041-210X.2010.00036.x

Ellerman, J. R. and Morrison-Scott, T. C. S. (1966). *Checklist of palaearctic and Indian mammals, 1758– 1946* (2nd ed.). London: Trustees of the British Museum (Natural History).

Groves, C. P., & C, G. (2001). *Primate Taxonomy*. Smithsonian Institution Press.

Hill, W. C . (1939). An Annotated Systematic List of the Leaf-monkeys. *Ceylon Journal of Science*, *XXI*.

Karanth, K. P., Singh, L., Collura, R. V, & Stewart, C.-B. (2008). Molecular phylogeny and biogeography of langurs and leaf monkeys of South Asia (Primates: Colobinae). *Molecular Phylogenetics and Evolution*, *46*, 683-694.

Karanth, K. P., Singh, L., & Stewart, C.-B. (2010). Mitochondrial and nuclear markers

suggest Hanuman langur (Primates: Colobinae) polyphyly: Implications for their species status. *Molecular Phylogenetics and Evolution*, *54*(2), 627–633. https://doi.org/10.1016/j.ympev.2009.10.034

Kawamoto, Y., Takemoto, H., Higuchi, S., Sakamaki, T., Hart, J. a., Hart, T. B., … Furuichi, T. (2013). Genetic Structure of Wild Bonobo Populations: Diversity of Mitochondrial DNA and Geographical Distribution. *PLoS ONE*, *8*(3), 1–9. https://doi.org/10.1371/journal.pone.0059660

Khanal, L., Chalise, M. K., Wan, T., & Jiang, X. (2018). Riverine barrier effects on population genetic structure of the Hanuman langur (*Semnopithecus entellus*) in the Nepal Himalaya. *BMC Evolutionary Biology*, *18*(159), 1–16.

Kishino, H., Miyata, T., & Hasegawa, M. (1990). Maximum likelihood inference of protein phylogeny and the origin of chloroplasts. *Journal of Molecular Evolution*, *31*(2), 151–160. https://doi.org/10.1007/BF02109483

Kishore, R., Reef Hardy, W., Anderson, V. J., Sanchez, N. A., & Buoncristiani, M. R. (2006). Optimization of DNA extraction from low-yield and degraded samples using the BioRobot EZ1 and BioRobot M48. *Journal of Forensic Sciences*, *51*(5), 1055–1061. https://doi.org/10.1111/j.1556-4029.2006.00204.x

Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Molecular Biology and Evolution*, *33*(7), 1870–1874. https://doi.org/10.1093/molbev/msw054

Minhas, R. A., Ahmed, K. B., Awan, M. S., Zaman, Q., Dar, N. I., & Ali, H. (2012). Distribution patterns and population status of the Himalayan grey langur (*Semnopithecus ajax*) in Machiara National Park, Azad Jammu and Kashmir, Pakistan. *Pakistan Journal of Zoology*, *44*(3), 869–877.

Minhas, R. A., Ali, U., Awan, M. S., Ahmed, K. B., Khan, M. N., Dar, N. I., … Tsuji, Y. (2013). Ranging and foraging of Himalayan grey langurs (*Semnopithecus ajax*) in Machiara National Park, Pakistan. *Primates*, *54*(2), 147–152. https://doi.org/10.1007/s10329-013-0345-7

Minhas, R. A., Khan, M. N., Awan, M. S., Ahmad, B., Bibi, S. S., Hanif, M., & Mian, A. (2018). RAPD based Genetic Diversity of Endangered Himalayan Gray Langur (*Semnopithecus ajax*) Populations of Pakistan. *Pakistan Journal of Zoology*, *50*(6), 2059–2071. https://doi.org/10.17582/journal.pjz/2018.50.6.2059.2071

Mishler, B. D. (1985). The Morphological , Developmental, and Phylogenetic Basis of Species Concepts in Bryophytes. *The Bryologist*, *88*(3), 207–214.

Mondol, S., Ullas Karanth, K., Samba Kumar, N., Gopalaswamy, A. M., Andheria, A., &
Ramakrishnan, U. (2009). Evaluation of non-invasive genetic sampling methods for
estimating tiger population size. *Biological Conservation*, *142*(10), 2350–2360.
https://doi.org/10.1016/j.biocon.2009.05.014

Nag, C., Karanth, K. P., & Gururaja, K. V. (2014). Delineating Ecological Boundaries of
Hanuman Langur Species Complex in Peninsular India Using MaxEnt Modeling
Approach. *PLoS ONE*, *9*(2), e87804. https://doi.org/10.1371/journal.pone.0087804

Nag, K. S. C., Pramod, P., & Karanth, K. P. (2011). Taxonomic Implications of a Field Study
of Morphotypes of Hanuman Langurs (*Semnopithecus entellus*) in Peninsular India.
International Journal of Primatology, *32*(4), 830–848. https://doi.org/10.1007/s10764-
011-9504-0

Napier, J. R., & Napier, P. H. (1967). *A handbook of living primates: morphology, ecology*
and behaviour of nonhuman primates. Academic Press.

Newton, P. N. (1988). The Variable Social Organization of Hanuman Langurs (*Presbytis*
entellus), Infanticide, and the Monopolization of Females. *International Journal of*
Primatology, *9*(1), 59-77.

Nijman, V. (2013). Distribution and Ecology of the Most Tropical of the High-Elevation
Montane Colobines: The Ebony Langur of Java. In N. B. et al. Grow (Ed.), *High*
Altitude Primates (pp. 115–132). Springer. https://doi.org/10.1007/978-1-4614-8175-1

Osterholz, M., Walter, L., & Roos, C. (2008). Phylogenetic position of the langur genera
Semnopithecus and *Trachypithecus* among Asian colobines , and genus affiliations of
their species groups. *BMC Evolutionary Biology*, *12*, 1–12.
https://doi.org/10.1186/1471-2148-8-58

Phillips, S. J., Anderson, R. P., Dudík, M., Schapire, R. E., & Blair, M. E. (2017). Opening
the black box: an open-source release of Maxent, (March), 887–893.
https://doi.org/10.1111/ecog.03049

Phillips, S. J., Anderson, R. P., & Schapire, R. E. (2006). Maximum entropy modeling of
species geographic distributions. *Ecological Modelling*, *190*, 231–259.
https://doi.org/10.1016/j.ecolmodel.2005.03.026

Pocock, R. I. (1928). The Langurs, Or Leaf Monkeys, of British India. *Journal of The*
Bombay Natural History Society, *32*, 472–504.

Pocock, R. I. (1939). *Primates and Carnivora (in part)* (Vol. 1). Taylor & Francis. Retrieved
from http://f.letmeprint.ru/14757241-7a331b5b/fragment_311854.pdf

Rambaut, A., Suchard, M.A., Xie, D., Drummond, A. J. (2013). Tracer v1.6. Available from

http://beast.bio.ed.ac.uk/Tracer.

Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D. L., Darling, A., Höhna, S., …

Huelsenbeck, J. P. (2012). Mrbayes 3.2: Efficient bayesian phylogenetic inference and

model choice across a large model space. *Systematic Biology*, *61*(3), 539–542.

https://doi.org/10.1093/sysbio/sys029

Roonwal, M. L. (1979). Field observations on distribution and tail carriage in the central

Himalayan langur, *Presbytis entellus schistaceous* (Primates). *Proceedings of the Indian*

National Science Academy, *45*(B1), 43-55.

Roonwal, M. L. (1984). Tail form and carriage in Asian and other primates, and their

behavioral and evolutionary significance. In N. S. Roonwal, M. L., Mohnot, S. M. and

Rathore (Ed.), *Current primate research* (pp. 93–151). Jodhpur University Press, India.

Roonwal, M. L., & Mohnot, S. M. (1977). *Primates of South Asia: Ecology, Sociobiology,*

and Behavior. Harvard University Press.

Sayers, K., & Norconk, M. A. (2008). Himalayan Semnopithecus entellus at Langtang

National Park, Nepal: Diet, activity patterns, and resources. *International Journal of*

Primatology, *29*(2), 509–530. https://doi.org/10.1007/s10764-008-9245-x

Schoener, T. W. (1968). The Anolis lizards of Bimini: Resource partitioning in a complex

fauna. *Ecology*, *49*(4), 704–726.

Shimodaira, H. (2002). An approximately unbiased test of phylogenetic tree selection.

Systematic Biology, *51*(3), 492–508. https://doi.org/10.1080/10635150290069913

Shimodaira, H., & Hasegawa, M. (1999). Multiple Comparisons of Log-Likelihoods with

Applications to Phylogenetic Inference. *Molecular Biology and Evolution*, *16*(8), 1114-

1116. https://doi.org/10.1093/oxfordjournals.molbev.a026201

Singh, J. S., & Singh, S. P. (1987). Forest Vegetation of the Himalaya. *The Botanical Review*,

53(1), 80–192.

Singh, P., Ramasastri, K. S., & Kumar, N. (1995). Topographical Influence on Precipitation

Distribution in Different Ranges of Western Himalayas. *Nordic Hydrology*, (26), 259–

284.

Sokal, R. R., & Crovello, T. J. (1970). The Biological Species Concept: A Critical

Evaluation. *The American Naturalist*, *104*(936), 127–153.

Stamatakis, A. (2006). RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses

with thousands of taxa and mixed models. *Bioinformatics*, *22*(21), 2688–2690.

https://doi.org/10.1093/bioinformatics/btl446

Sugiyama, Y. (1976). Characteristics of the Ecology of the Himalayan Langurs. *Journal of*

Human Evolution, *5*, 249–277.

Swofford, D. (2001). PAUP*—Phylogenetic Analysis Using Parsimony and Other Methods,
Version 4. Sunderland, MA: Sinauer Associates.

Van Valen, L. (1976). Ecological Species, Multispecies, and Oaks. *Taxon*, *25*(2/3), 233–239.

Wangchuk, T., Inouye, D. W., & Hare, M. P. (2008). The Emergence of an Endangered
Species: Evolution and Phylogeny of the *Trachypithecus geei* of Bhutan. *International*
Journal of Primatology, *29*, 565‑582.

Warren, D. L., Glor, R. E., & Turelli, M. (2010). ENMTools⬚: a toolbox for comparative
studies   of   environmental   niche   models.   *Ecography*,   *33*,   607-611.
https://doi.org/10.1111/j.1600-0587.2009.06142.x

Warren, D. L., & Seifert, S. N. (2011). Ecological niche modeling in Maxent⬚: the
importance of model complexity and the performance of model selection criteria.
Ecological Applications, *21*(2), 335–342.

White, P. S., & Densmore III, L. D. (1992). Mitochondrial DNA isolation. In A. R. Hoelzel
(Ed.), *Molecular genetic analysis of populations: a practical approach* (pp. 29–58).
Oxford: IRL press, Oxford University Press.

Will, K. W., Mishler, B. D., & Wheeler, Q. D. (2005). The Perils of DNA Barcoding and the
Need   for   Integrative   Taxonomy.   *Systematic   Biology*,   *54*(5),   844–851.
https://doi.org/10.1080/10635150500354878

Zhang, B.-W., Li, M., Ma, L.-C., & Wei, F.-W. (2006). A Widely Applicable Protocol for
DNA Isolation from Fecal Samples. *Biochemical Genetics*, *44*(11–12), 494–503.
https://doi.org/10.1007/s10528-006-9050-1

Tables

Table 1: Different Taxonomic Schemes (TS) for Himalayan langur proposed by various

TS 1					TS 2	TS 3
Pocock (1928)	Pocock (1939)	Roonwal and Mohnot (1977)	Roonwal (1984)*	Brandon-Jones (2004)	Hill (1939)	Groves (2001)
Pi. e. schistaceus	*S. e. schistaceus*	*P. e. schistaceus*	*P. e. schistaceus*	*S. e. schistaceus*	*S. schistaceus*	*S. schistaceus*
Pi. e. ajax	*S. e. ajax*	*P. e. ajax*	*P. e. ajax*	*S. e. ajax*	*S. s. ajax*	*S. ajax*
Pi. e. achilles	*S. e. achilles*	*P. e. achilles*	*P. e. achilles*	—	*S. s. achilles*	—
Pi. e. lanius	—	*P. e. lania*	*P. e. lania*	—	*S. s. lanius*	—
Pi. e. hector	—	—	—	*S. e. hector*	*S. s. hector*	*S. hector*

592   authors

593   *Also includes Ellerman and Morison-Scott (1966) and Napier and Napier (1967)
594   *Pi = Pithecus*; *S = Semnopithecus*; *e. = entellus*
595

Table 2: Coding system used for the 6 morphological characters used in this study. The coding system is designed to be consistent with Nag et al. (2011), unless mentioned otherwise.

	Characters					
	Crest	Streak	EOB	Tail loop (TL)	Tail carriage (TC)	HBC
Himalayan langur	0	0	0	1	3	1
*Semnopithecus entellus**	0	0	3	1	$4^{\#}$	0
Semnopithecus hypoleucos	0	1	$3/4^{\dagger}$	2	1	0
*Semnopithecus priam***	1	0	1	2	2	0

**S. entellus* = Nt from Nag et al. (2011).*

[#]Nag et al. (2011) coded Nt as 0, here we have coded it 4.

†*S. hypoleucos* contains two morphotypes (St1/St2; Nag et al., 2011).

** We did not use this species in the analysis.

605 **Table 3: Topology test results**

Tree	-ln L	Diff -ln L	SH	AU	p-value
Best tree	2107.65046	(best)			
Constrained tree	2371.94530	264.29484	0.0000*	~0*	*P < 0.05

607   **Table 4:** Maxent results showing the most important variables ranked on the basis of the
608   amount of variation they explain in predicted distribution of *Semnopithecus entellus* and the
609   Himalayan langur along with the permutation importance for the variables.

Semnopithecus entellus			Himalayan langur		
Variable	**Percent contribution**	**Permutation importance**	**Variable**	**Percent contribution**	**Permutation importance**
Bio 15	35.5	39.6	Bio 17	42.1	5.9
Bio 10	26.5	7.2	Bio 1	20.6	48.2
Bio 1	20.7	19	Bio 2	15.5	16.5
Bio 12	11.7	19.9	Bio 4	13.7	23.7
Bio 18	2.5	5.7	Bio 12	8.1	5.7
Bio 3	1	5.4	Slope	0.1	0
Bio 8	0.8	0	Aspect	0	0
Slope	0.3	0.1			
Bio 9	0.3	1.7			
Aspect	0.3	0.7			
Bio 14	0.2	0.6			
Bio 19	0.1	0			
Bio 4	0.1	0			

Figure Captions

Fig. 1: Tail carriage pattern in the Himalayan langur (a) and *Semnopithecus entellus* (b).

Fig. 2: Contrast between the head and the dorsal region of the body in the Himalayan langur (a) and *Semnopithecus entellus* (b).

Fig. 3: Bayesian phylogeny of Himalayan langur for mitochondrial cytochrome *b* (Cyt-b) gene. Node support is indicated by black circles and black squares. Black circle indicates high Bayesian posterior probability support (> 0.75) as well as a high maximum likelihood bootstrap support (>75). Black square indicates only a high Bayesian posterior probability support (> 0.75). HP = Himachal Pradesh; JnK = Jammu & Kashmir; Uk = Uttarakhand. [#]Sequenced in this study.

Fig. 4: Neighbour-Joining tree based on 6 colour independent characters. *S. hypoleucos* is one of the two species of Hanuman langur from peninsular India

Fig. 5: Species distribution maps of Himalayan langur (a) and *S. entellus* (b).

a   b