Development of specific molecular markers to distinguish and quantify broomrape species in a soil sample from infected field

Radi Aly[1]* , Vinay K. Bari[1] , Avishai Londner[1], Kocherman Marina[1], Jackline Abu Nassar[1], Leena Taha-Salaime[1], Eizenberg Hanan[1] and Ran Lati[1]

[1]Department of Plant Pathology and Weed Research, Newe Ya'ar Research Center, Agricultural Research Organization (ARO), Israel.

*Corresponding Author: Radi Aly

radi@volcani.agri.gov.il

    https://orcid.org/0000-0002-5670-3798
    https://orcid.org/0000-0002-6038-9586

Keywords: Plant parasite, *Phelipanche aegyptiaca, Orobanche* spp., ribulose-1,5-bisphosphate carboxylase (*rbcL1*), Internal transcribed spacer (*ITS*).

Conflict of interest: The authors have declared no conflict of interest

Abstract

Broomrapes (*Orobanche* and *Phelipanche*) are obligate holoparasites that cause heavy

damage to numerous crops, reducing the yield and its quality. The parasite develops in the

soil and exerts the greatest damage prior to its emergence; therefore the majority of field loss

may occur before diagnosis of infection. Because of the parasite tiny seed size (200 to 300

μm) and dormancy for several decades in the field, it is very difficult to diagnose the parasite

by conventional methods. Therefore, to restrict the parasite seeds spread and contamination to

other commercial fields, development of DNA-based molecular markers to identify and

quantify broomrape species in a soil sample is much needed. In this study, we developed a

specific molecular marker (RbcL-M) based on *rbcL1* (large subunit of the ribulose-

bisphosphate carboxylase) gene from *Orobanche crenata* to differentiate between *Orobanche*

crenata and *Orobanche cumana.* Likewise, a specific marker (ITS100) based upon unique

sequences in the internal transcribed spacer (ITS) regions of the nuclear ribosomal DNA of

Phelipanche aegyptiaca to quantify three species of the parasite (*P.aegyptiaca*, *O.crenata*

and *O.cumana*) in a soil sample was developed. Genomic DNA was extracted from soil

samples artificially infested with broomrape seeds or tissue of *P.aegyptiaca*, *O.cumana* and

O.crenata and subjected to PCR analysis. RbcL-M marker successfully amplified a PCR

product (1300bp) when *O.crenata* seeds or tissues (collected from several locations in Israel)

were added to the soil samples. The same marker amplified a PCR product (1000bp) when *O.*

cumana seeds or tissues were added to the soil samples. RbcL-M marker did not amplify soil

samples with seeds or tissues of *P.aegyptiaca* or any soil-borne DNA. Furthermore, using

ITS-100 marker and Real-Time PCR analysis, allowed quantitative diagnostic of the parasite

in a soil sample from infected sunflower field. As expected the universal internal control

primer (UCP-555) amplified a PCR product (555bp) when genomic DNA extracted from soil

samples with or without broomrape tissues. The development of an efficient, simple and

robust molecular marker to detect and distinguish between broomrape species, has a

significant insights on assessment the level of infestation and planning eradication program of

the parasite in a field crop.

INTRODUCTION

Broomrapes (*Orobanche* and *Phelipanche* spp.) are member of Orobanchaceae family

(Parker C., 1993), is an underground root parasite mostly affecting agricultural crop plants

(McNeal et al., 2013). These parasitic plants are unable to synthesize their photosynthetic

assimilate due to lack of chlorophyll; and totally depend on host plants for the germination

and growth by feeding through haustorium (Westwood et al., 2010a). The Orobanchaceae

family contains the largest number of parasitic species that form haustoria in roots, for

example *Orobanche, Phelipanche* species (Yoshida et al., 2016). Haustoria are special organ

of parasitic plant which invades host tissue and serve as connecting bridge that allows the

parasite to obtain water, mineral and nutrient from host (Joel and Losnergoshen, 1994;

Westwood et al., 2010b).The natural habitats of broomrapes are the warm and temperate

regions of the northern hemisphere, including the Mediterranean region (Parker C., 1993),

however branched broomrape *O.ramosa* native to central and south-western Europe, and

considered as a major threat to agricultural crops. They cause extensive damage by reducing

the yield of all economically important crops belongs to Solanaceae, Fabaceae, Brasicaceae,

Asteraceae and Apiaceae family (Joel, 2006; Parker C., 1993; Westwood et al., 2010b). In

Israel the significance of infested fields has increased dramatically during the last decade,

causing heavy crop damage or even total yield losses.

In general, parasitized crops suffer from reductions in total biomass at the greatest expense to

their productive tissue (Joel, 2006). Broomrapes cause many problems for agriculture farmers

because of their tiny seed size (200-300µm) which makes them difficult to detect in harvested

crop seeds and in agriculture soil (Joel, 1987). Moreover, small broomrape germinates and

grows only in the presence of a susceptible host. In addition, small broomrape seeds can

survive more than a decade in the soil (Habimana et al., 2014). Therefore, identification and

quantification of broomrape seeds in a soil sample urgently required and will be helpful in

prevention of crop damages by these parasites (Joel et al., 1996). The conventional methods

of detection and differentiation such as seed coat morphological feature (Joel, 1987), Random

amplified polymorphic DNA technique (RAPD) (Katzir et al., 1996; Paran et al., 1997),

Intersimple sequence repeat (ISSR) (Benharrat et al., 2002) and high resolution melting

analysis (Rolland et al., 2016) has some limitation to distinguish and quantify broomrapes

exist in the soil. Recent progress in sequencing broomrape plastid genome, which was made

available, providing new insight to develop molecular marker for specific broomrape species

identification (Cusimano and Wicke, 2016; Wicke et al., 2013). Genetic or DNA based

molecular marker such as internal transcribe spacer (ITS), plastidial ribulose bisphosphate

carboxylase (Rubisco) large subunit (*rbcL1*) and maturase K (*matK*) are among the most

widely used molecular markers in evolutionary, taxonomy, phylogenetic and diversity

identification of many organisms including *Orobanche*, have been established in recent years

(Agarwal et al., 2008; Park et al., 2008; Schneeweiss et al., 2004).

In order to develop a complementary methodology for mapping parasitic plants in the field, a

procedure includes: a known geographical information systems (GIS) for soil sampling that

will characterize the spatial variation in the field (Eizenberg et al., 2012), and at the same

time, use of molecular markers to identifying broomrapes in a soil sample. Those molecular

markers would assist detection and diagnosis of broomrapes species and population level in

the soil. In this study we developed a simple and easy method based on *rbcL1* gene to

differentiate between three broomrape species *O.crenata, O.cumana* and *P.aegyptiaca*, the

most common and destructive weed in Israel. We also developed a specific molecular marker

to quantify the parasite seeds in soil sample in an infected sunflower field.

RESULTS

Design of *rbcL1* specific marker to differentiate between *O. cumana* and *O. crenata*

In our previous study, we reported amplification of ITS region using ITS350 primer; which

specifically detect broomrape in soil sample (Aly et al., 2012). However, ITS350 marker does

not differentiate between broomrape species because of the high sequence similarity among

ITS regions. Therefore, we compared and analysed the *rbcL1* gene sequences from these

species using Clustal Omega and designed new molecular marker to differentiate broomrape

species such as *O.cumana* and *O.crenata* (Figure S1). To distinguish between the above

broomrape species, probes differing at least by five-base mismatches or more were designed.

Based on sequence alignment, a molecular marker (RbcL-M) was designed using *rbcL1* gene

from *O.cumana* (GenBank accession number: AF090349.1) Soil samples were artificially

contaminated with seeds or shoots of *P. aegyptiaca*, *O.crenata* and *O.cumana* and subjected

to PCR analysis. By PCR amplification using RbcL-M, we were able to differentiate between

O.crenata and *O.cumana* according to the length of the PCR products. For *O.crenata* the

PCR product size was 1300bp compared to 1000bp for *O. cumana*. However, RbcL-M

marker failed to amplify any PCR product while soil samples were artificially contaminated

with *P. aegyptiaca* seeds or shoots (Figure 1A). As expected, the universal internal control

primers amplified a PCR product (555bp) from soil samples with or without broomrape

shoots or seeds (Figure 1B).

Development of specific ITS100 marker to detect seeds of three broomrape species in a

soil sample

To analyse the number of the broomrape seeds present in a soil sample from infected field,

we designed ITS100 primers from the highly conserved region of ITS350 region (Aly et al.,

2012). To determine the specificity of the marker to each broomrape specie, we first extracted

genomic DNA from soil samples (250mg) containing (0, 1, 10, 25, 50 100 and 250 seeds) of

P.aegyptiaca, *O.crenata* or *O.cumana* and subjected to PCR analysis. ITS100 primers

successfully amplified a PCR product (100bp) (Figure 2A) when *P.aegyptiaca*, *O.crenata* or

O.cumana were mixed with soil sample, except soil sample without *Orobanche* seed.

However, we have got PCR product in all the sample including soil without *Orobanche* seeds

using universal primer UCP-555 (Figure 2B).

In addition, the specificity of the markers generated in this study were evaluated by testing

DNA extracted from sunflower, carrot and tomato leaves and roots corresponding to possible

respond to the host plants. Fortunately, the qPCR assay always yielded negative results for

these plant species (data not shown). Next to quantify *Orobanche*, we generated standard

curve using *O.cumana* genomic DNA and ITS100 marker utilizing qPCR procedure. *O.

cumana* genomic DNA was purified and serial dilutions (1:10 fold) ranging from 0.001 to 10

ng/µl were used for qPCR assay. The standard curve for *O.cumana* exhibited a slope of -1.28

and (R^2) of 0.9994 (Figure 2C). The same type of assay was performed with *O.cumana*

genomic DNA extracted from soil sample (250 mg) mixed with the parasite seeds (0.1, 1, 10

and 100 mg seeds) and subjected to qPCR. The standard curve for *O.cumana* seeds mixed

with soil sample, exhibited a slope of -1.332 and (R^2) of 0.987 (Figure 2D).

Prediction and quantification of *O. cumana* seeds in sunflower field

Previously, we have developed a protocol that allows extraction of genomic DNA from few

tiny seeds of *Orobanche* spp. in a soil sample and we were able to subject the sample DNA to

a rapid PCR-assay (Aly et al., 2012). Here, we extended the protocol, allowing for the

identification and distinguishing between soil-borne and parasite seed species that are

collected from a field soil sample. Following confirmation and validation of the ITS100

marker specificity to *O.cumana* seeds, geo-statistics model for soil sampling that

characterized the spatial variation in the field, was performed according to (Eizenberg et al.,

2012) in a sunflower field located at Havat Eden, Emic Betshan– Israel. This field was

infected with *O. cumana* (Figure 3A). Five soil samples were collected from a plot of 1 acre

according to the illustration (Figure 3B). Genomic DNA was extracted from each soil sample

and subjected to qPCR using ITS100 primers to predict and quantify *O.cumana* DNA in

naturally infected field. Then, genomic DNA from soil samples was extrapolated with Ct

values to quantify the number of seeds in each sample according to the generated Ct values

(Figure 2C and D). The results showed variable quantity of *O.cumana* seeds in the soil

sample ranged from 2 to 22 parasite seeds (Figure 3C) in 250mg soil sample processed

originally from 1kg soil sample. Sample no. 6 showed the highest density of *O.cumana* seeds

(22 seeds) in the sunflower infected field as compared to the other samples. Sample no. 6 was

collected from a highly infected area as was shown by the geo-statistics model proposed by

(Eizenberg et al., 2012). Concomitantly, we monitored and counted the parasite

inflorescences in adjacent to the position were the soil samples were taken. Our results

revealed that average density of 0.6 inflorescences /m^2, in a total area of 1acre was recorded.

Inflorescences counts in the selected area in the field were also variable ranged from 2 to 15

and were less than the seed numbers found in the soil sample when qPCR was applied

(Figure 3C).

Evaluation viability of the parasite seeds in sunflower infected field

To determine viability of broomrape seeds in a sunflower infected field, a tetrazolium test

(TZ) (Lopez-Granados and Garcia-Torres, 1999; Thorogood et al., 2009) was conducted.

According to TZ test, viable parasite seeds tends to have orange to brown colour as a result of

its metabolite activity while non-viable tends to have yellow to faint colour (Figure 4A). To

evaluate the parasite seed viability, two treatments were assayed: soil samples (organic matter

only) collected from sunflower field located at Havat Eden, Emic Betshan, naturally infested

with *O. cumana* and pure *O. cumana* seeds (1 mg) prepared from a seed stock. Our results

indicate that the highest percentage (% of total) of viable *O. cumana* seeds were found in soil

samples collected from the sunflower field (83%) as compared to 25% found in a seed stock

(Figure 4B).

DISCUSSION

Parasitic weeds are often exert their greatest damage prior to their emergence; therefore the

majority of field loss may occur before diagnosis of infection. Early detection of infection by

visual inspection is not possible. This hampers early control strategy efforts and the detection

methods used by crop scientists are laborious and time consuming. Most of the methods

developed to detect broomrape seeds in soil samples, need several steps such as seed

purification (flotation, filtration and binocular microscope observation (Kroschel, 2001).

Despite of the great advances in genome sequencing approaches that has already been

achieved, our understanding towards genomic evolution of Orobancheaceae is still

incomplete due to aberrant evolution of plastid genome of holoparasite (Park et al., 2007).

Available DNA sequence data are insufficient to differentiate closely related species, such as

the weedy parasites *O.cumana, O.crenata* and *P.aegyptiaca*.

Genetic or DNA based marker techniques such as ITS or *rbcL1* are routinely being used in

evolutionary, taxonomical, phylogenetic and diversity studies (Agarwal et al., 2008; Park et

al., 2008; Schneeweiss et al., 2004). Diagnosis and early identification of the parasite species

in the field by soil sampling is of great importance to farmers and it is particularly needed due

to the host-parasite specific interaction, the difficulty to detect the tiny parasite seeds by

conventional methods in contaminated fields and because seeds will only germinate and grow

in the presence of a susceptible host. The gene encoding the large subunit of ribulose-1,5-

bisphosphate carboxylase/oxygenase (Rubisco; *rbcL1*) is retained as a pseudogene in

Orobanche and *Phelipanche* (Wicke et al., 2013). In the present work, we designed sets of

primers targeting the plastid *rbcL1* gene sequence from *O.cumana* to differentiate between

three broomrape species (*O.cumana, O.crenata* and *P.aegyptiaca*) the most common and

harmful species in field crops in Israel. Another sets of primers were designed to target ITS

sequences (Schneeweiss et al., 2004) to quantify the parasite seeds potentially present in a

soil sample to help for seed bank assessment in the field. We found that RbcL-M primers

(marker) were useful to distinguish between DNA from *O.cumana* and *O. crenata* producing

different amplicon sizes (a specific PCR product about 1300bp for *O. crenata* and 1000bp for

O.cumana). Subsequent BLASTn with previously published broomrape genomic data

(Cusimano and Wicke, 2016) reconfirmed the differences between the two species. In

contrast, DNA from *P.aegyptiaca* failed to produce PCR amplicon suggesting possibly that

Phelipanche species completely lost *rbcL1* gene (Delavault et al., 1995; Leebens-Mack and

de Pamphilis, 2002).To allow differentiation between *P.aegyptiaca* from other species, a

specific molecular marker was recently designed (Aly et al., 2019), (GenBank accession

numbers MK637618-637624).

To diagnose and quantify broomrape species in a seed stock (Dongo et al., 2012), different

types of nuclear and plastid DNA markers have been proposed. (Schneeweiss et al., 2004)

were the first to present molecular phylogenetic analysis using nuclear ITS sequences. ITS-

based markers were also used to detect *P.aegyptiaca* seeds in a soil sample (Aly et al., 2012)

and quantify contamination of *O.ramosa* and *O. cumana* in crop seed lots (Dongo et al.,

2012). Here we developed ITS100 marker that was based on primers consisting of unique

sequences in the internal transcribed spacer (ITS) regions of the nuclear ribosomal DNA

(nrDNA) of *O. crenata*. ITS-100 marker was used with qPCR assay to quantify *O.cumana*

seeds in a soil sample from sunflower field located at Havat Eden, Emic Betshan – Israel.

We were able to detect 0.1 mg *O.cumana* seeds in 250 mg soil sample (wt ∕ wt). A detection

threshold of 0.1 mg broomrape seeds in 20 g seed samples was previously reported (Dongo et

al., 2012). For detection, mapping and quantifying *O. cumana* seeds in the field, we used

geographical information systems (GIS) for soil sampling and other advanced technologies

for parasitic weed mapping and field history data storage (Eizenberg et al., 2012) followed by

qPCR assay. Soil samples (500gm) were collected from the sunflower field, organic material

was extracted from each sample ending with 250mg then, genomic DNA was extracted and

subjected to qPCR using ITS100 primers. This method allowed specifically detecting and

quantifying the DNA of *O.cumana* in a total DNA extract from sunflower soil sample.

Accordingly, the results of this assay can be also expressed as the number of parasite seeds

per kilogram of soil following extrapolation with the standard curve was prepared. Our

results indicate that samples collected from highly infected area (Fig. 4A) according to the

geo-statistics model proposed by Eizenberg et al. (2012), were with agreement with our

qPCR assay (sample no. 6 showed the highest density of *O.cumana* seeds (22 seeds) in the

sunflower infected field as compared to the other samples. However, no correlation was

found between qPCR assay (parasite seed number) compared to number of the parasite

inflorescences collected from the same location in the sunflower infected field. An

explanation for that may be related to the viability of *O.cumana* seeds, genomic DNA from a

soil sample will contain viable and non-viable parasite seeds, add to that, we cannot exclude

the presence of some related parasite seeds in the same sample therefore, the we counted

more parasite seeds using the qPCR assay. Specificity of the qPCR assay was tested against

several possible contaminants of soil-borne pathogens by using universal internal control

primer UCP-555 (White, 1990) or harvested crop seeds like sunflower and tomato using

ITS100. No amplification was observed, confirming the specificity of the marker.

The powerful of broomrape-infected field to distribute and contaminate the neighbouring

non-infested fields depends on the soil seed-bank and viability of the parasite seeds. Viability

of broomrape seeds in the sunflower infected field was determined by tetrazolium test. Our

results showed the highest percentage of viable broomrape seeds were found in soil samples

collected from the sunflower field (83%) as compared to 25% found in a seed stock. We

assume that the highest count recorded in the sunflower field was due to the release and

distribution of the fresh parasite seeds by the newly parasite inflorescences showed up

through the crop growth. We have to take in consideration that we used an old seed stock (10

years old) that doesn't represent newly harvested seed stocks. Our experience with seed

germination of *O. cumana* with germination stimulant (GR) could reach more than 90%.

Additionally, we cannot exclude the presence of some related parasite seeds in the same

sample from previous growth seasons.

In this study, we provide a simple, fast and non-expensive approach to distinguish and

quantify broomrape seeds exist in a soil sample from a crop field. Molecular markers would

assist accurate detection and population level of *Phelipanche* and *Orobanche* spp. in a soil

sample and offer numerous advantages over conventional phenotype based alternatives, as

they are stable and detectable in all tissues regardless of growth, differentiation or

development. These methods could be helpful in precision agriculture, in which they provide

answers routinely questioned by the farmers: are there parasite seeds in my crop field? What

species? And how much seeds are exist in a soil sample?

EXPERIMENTAL PROCEDURE

Plant material

Seeds, shoots and inflorescences of three broomrape species (*P.aegyptiaca, O.cumana and O.*

crenata) were collected from different Israeli locations (Golan heights, Havat Eden- Emic

Betshan and Jordan valley respectively). Seeds were harvested from freshly inflorescences

and stored at 4°C while shoots and inflorescences were dried and stored at −80°C until used.

Two hundred milligrams of parasite plant seeds/tissue were ground manually in presence of

liquid nitrogen with the help of pestle and mortar. Total genomic DNA was extracted using

the GeneGET Plant genomic DNA extraction kit according to the manufacturer instructions

(Thermo Scientific).

ITS and *rbcL1* sequences and primers design

The internal transcribed spacer (ITS) regions of *Phelipanche aegyptiaca* (GenBank-

Accession No. AY209327) was used to design ITS100 marker. The PCR product yielded a

100bp fragment in length. UCP-555 used as universal internal control primer, which

amplifies a region of the small subunit of nrDNA (555 bp) from a wide variety of

microorganism such as protists, fungi, and plants. The length of the *rbcL1* sequences was

1211bp and 1290bp for *O.cumana* (GenBank: AF090349.1) and *O. crenata* (GenBank:

AY582191.1) respectively. Sequence similarities between broomrape species were done

using Clustal Omega software. Design of primers for the qPCR detection of broomrape

species was based on the alignment of the ITS of nuclear ribosomal DNA regions. All

primers were designed with Gene Runner software. PCR amplifications of *rbcL1* sequences

were performed with the RbcL-M-F and RbcL-M-R primers using PCR Ready Mix

(BioLabs, Israel), and 50ng DNA-template. The following PCR conditions were used:

denaturation for 5 min at 95°C; 32 cycles with 30 second at 95°C, 30 second at 55°C, 1.30

min at 72°C; and final elongation for 10 min at 72°C.

Soil sampling and extraction of Genomic DNA

Design and mapping the soil samples in the sunflower tested field were performed according

to (Eizenberg et al., 2012). Soil samples were collected from the infected field on May and

counting the inflorescences was performed on July. Each sample includes 500gm of a soil

sample that was introduced into one-litter container, mixed thoroughly with 1.5L of (13.3 M

$CaCl_2$) solution and kept at room temperature for overnight. The next day, the upper phase

including the organic materials and broomrape seeds were collected, dried on Whatmann

paper and the parasite seeds were obtained by filtration through 50 and 100 mesh. Genomic

DNA was extracted by PowerSoil kit (MO BIO Lab. Inc., Loker Ave West, Carlsbad, CA).

according to the manufacturer's instructions (Sagova-Mareckova et al., 2008).

Quantitative real time PCR of soil sample from the sunflower fields

The quantitative real time PCR was performed using PerfeCTa® SYBR® Green Fast Mix®,

ROX™ (Quanta biosciences) and ABI-Prism 7000 Real-Time PCR Detection System

(Applied Biosystems) according to the manufacturer's protocol. The qPCR was performed in

quantitative reaction with final volume of 10 µl including: 100ng DNA, 5 µl of SYBR Green

Fast Mix- ROX and 500nM of each ITS-100 primers. *O.cumana* actin was used as

endogenous control and the relative gene expression level was calculated using the $2^{-\Delta\Delta Ct}$

method (Livak and Schmittgen, 2001). Two standard curves were prepared: The first curve

included standard points from *O.cumana* seeds ranging from 0.001 to 10 ng DNA per tube,

were made using 1:10-fold serial dilutions of broomrape DNA and were used for qPCR

assay. The second curve was generated by mixing different amounts of genomic DNA from

O.cumana (0.1 to 100 ng) with total genomic DNA (10 μg) extracted from 250 mg soil

sample. Each point on the standard curve was assayed in triplicate.

Viability evaluation of the parasite seeds by Tetrazolium test

Tetrazolium test was performed as described earlier (Lopez-Granados and Garcia-Torres,

1999; Thorogood et al., 2009). Briefly, 50-100 seeds were hydrated between paper towel

sheets for 2-3 days. After pre-conditioning, they were placed in flasks, covered with a 1%

solution of 2,3, 5 triphenyl tetrazolium chloride (TZ) and incubated at 40°C for 2h. After that,

the TZ solution was discarded; and the seeds were washed thoroughly with sterile water.

Seeds were directly evaluated for viability under the microscope using magnification (40X

and 100X) and classified in two categories: viable seeds – orange to brown light coloured or

non-viable seeds –yellow to white colour.

ACKNOWLEDGMENTS

Results of this research were supported by the Chief Scientist of the Ministry of Agriculture

and Rural Development - Israel, grant No.132-1499-10 (MEZAMALEKET). VKB is grateful

to the ARO-Volcani Center, Agricultural Ministry of Israel for providing the Postdoctoral

fellowship.

REFERENCES

Agarwal, M., Shrivastava, N., and Padh, H. (2008). Advances in molecular marker techniques and

their applications in plant sciences. Plant Cell Rep *27*, 617-631.

Aly, R., Eizenberg, H., Kocherman, M., Abu-Nassar, J., Taha, L., and Saadi, I. (2012). Use of ITS

nuclear sequences from Phelipanche aegyptiaca as a direct tool to detect single seeds of broomrape

species in the soil. Eur J Plant Pathol *133*, 523-526.

Aly, R., Lati, R., Abu-Nassar, J., Ziadna, H., Achdari, G., Münchow, C.S.v., Wicke, S., Bari, V.K.,

and Eizenberg, H. (2019). The weedy parasite Phelipanche aegyptiaca attacks Brassica rapa var. rapa

L. for the first time in Israel. Plant Disease Note.

Benharrat, H., Veronesi, C., Theodet, C., and Thalouarn, P. (2002). Orobanche species and population

discrimination using intersimple sequence repeat (ISSR). Weed Res *42*, 470-475.

Cusimano, N., and Wicke, S. (2016). Massive intracellular gene transfer during plastid genome

reduction in nongreen Orobanchaceae. New Phytol *210*, 680-693.

Delavault, P., Sakanyan, V., and Thalouarn, P. (1995). Divergent evolution of two plastid genes, rbcL

and atpB, in a non-photosynthetic parasitic plant. Plant Mol Biol *29*, 1071-1079.

Dongo, A., Leflon, M., Simier, P., and Delavault, P. (2012). Development of a high-throughput real-

time quantitative PCR method to detect and quantify contaminating seeds of Phelipanche ramosa and

Orobanche cumana in crop seed lots. Weed Res *52*, 34-41.

Eizenberg, H., Aly, R., and Cohen, Y. (2012). Technologies for Smart Chemical Control of

Broomrape (Orobanche spp. and Phelipanche spp.). Weed Sci *60*, 316-323.

Habimana, S., Nduwumuremyi, A., and Chinama, R.J.D. (2014). Management of orobanche in field

crops- A review. J Soil Sci Plant Nut *14*, 43-62.

Joel, D.M. (1987). Detection and Identification of Orobanche Seeds Using Fluorescence Microscopy.

Seed Sci Technol *15*, 119-124.

Joel, D.M., Hershenhorn, Y., Eizenberg, H., Aly, R., Ejeta, G., Rich, P.J., Ransom, J.K., Sauerborn, J.

and Rubiales, D. (2006). Biology and Management of Weedy Root Parasites. In Horticultural

Reviews, New Jersey: Wiley, 267–349.

Joel, D.M., and Losnergoshen, D. (1994). The Attachment Organ of the Parasitic Angiosperms
Orobanche Cumana and O-Aegyptiaca and Its Development. Can J Bot *72*, 564-574.

Joel, D.M., Portnoy, V., and Katzir, N. (1996). Identification of single tiny seeds of Orobanche using
RAPD analysis. Plant Mol Biol Rep *14*, 243-248.

Katzir, N., Portnoy, V., Tzuri, G., Joel, D.M., and Castejon-Munoz, M. (1996). Use of random
amplified polymorphic DNA (RAPD) markers in the study of the parasitic weed Orobanche. Theor
Appl Genet *93*, 367-372.

Kroschel (2001). A Technical Manual for Parasitic Weed Research and Extension. Springer,
Dordrecht.

Leebens-Mack, J., and de Pamphilis, C. (2002). Power analysis of tests for loss of selective constraint
in cave crayfish and nonphotosynthetic plant lineages. Mol Biol Evol *19*, 1292-1302.

Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods *25*, 402-408.

Lopez-Granados, F., and Garcia-Torres, L. (1999). Longevity of crenate broomrape (Orobanche
crenata) seed under soil and laboratory conditions. Weed Sci *47*, 161-166.

McNeal, J.R., Bennett, J.R., Wolfe, A.D., and Mathews, S. (2013). Phylogeny and origins of
holoparasitism in Orobanchaceae. Am J Bot *100*, 971-983.

Paran, I., Gidoni, D., and Jacobsohn, R. (1997). Variation between and within broomrape
(Orobanche) species revealed by RAPD markers. Heredity *78*, 68-74.

Park, J.M., Manen, J.F., Colwell, A.E., and Schneeweiss, G.M. (2008). A plastid gene phylogeny of
the non-photosynthetic parasitic Orobanche (Orobanchaceae) and related genera. J Plant Res *121*,
365-376.

Park, J.M., Manen, J.F., and Schneeweiss, G.M. (2007). Horizontal gene transfer of a plastid gene in
the non-photosynthetic flowering plants Orobanche and Phelipanche (Orobanchaceae). Mol
Phylogenet Evol *43*, 974-985.

Parker C., R.C.R. (1993). Parasitic weeds of the world : biology and control (Wallingford, Oxon :
CAB International).

Rolland, M., Dupuy, A., Pelleray, A., and Delavault, P. (2016). Molecular Identification of

Broomrape Species from a Single Seed by High Resolution Melting Analysis. Front Plant Sci *7*.

Sagova-Mareckova, M., Cermak, L., Novotna, J., Plhackova, K., Forstova, J., and Kopecky, J. (2008).

Innovative methods for soil DNA purification tested in soils with widely differing characteristics.

Appl Environ Microbiol *74*, 2902-2907.

Schneeweiss, G.M., Colwell, A., Park, J.M., Jang, C.G., and Stuessy, T.F. (2004). Phylogeny of

holoparasitic Orobanche (Orobanchaceae) inferred from nuclear ITS sequences. Mol Phylogenet Evol

30, 465-478.

Thorogood, C.J., Rumsey, F.J., and Hiscock, S.J. (2009). Seed viability determination in parasitic

broomrapes (Orobanche and Phelipanche) using fluorescein diacetate staining. Weed Res *49*, 461-

468.

Westwood, J.H., Yoder, J.I., Timko, M.P., and dePamphilis, C.W. (2010a). The evolution of

parasitism in plants. Trends in Plant Science *15*, 227-235.

Westwood, J.H., Yoder, J.I., Timko, M.P., and dePamphilis, C.W. (2010b). The evolution of

parasitism in plants. Trends Plant Sci *15*, 227-235.

White, T.J., T. D. Bruns, S. B. Lee, and J. W. Taylor (1990). Amplification and direct sequencing of

fungal ribosomal RNA Genes for phylogenetics.

Wicke, S., Muller, K.F., de Pamphilis, C.W., Quandt, D., Wickett, N.J., Zhang, Y., Renner, S.S., and

Schneeweiss, G.M. (2013). Mechanisms of Functional and Physical Genome Reduction in

Photosynthetic and Nonphotosynthetic Parasitic Plants of the Broomrape Family. Plant Cell *25*, 3711-

3725.

Yoshida, S., Cui, S.K., Ichihashi, Y., and Shirasu, K. (2016). The Haustorium, a Specialized Invasive

Organ in Parasitic Plants. Annu Rev Plant Biol *67*, 643-667.

Figure legends

Figure 1. (A) Specificity of RbcL-M marker and PCR amplification patterns in three broomrape species (*P.aegyptiaca, O.cumana and O.crenata*). Soil samples (250 mg) were artificially contaminated with broomrape seeds (100) or 10 mg shoots of *P.aegyptiaca, O.crenata and O.cumana* and subjected to PCR analysis using RbcL-M primers. Genomic DNA was extracted and 100 ng were used for PCR amplification. A PCR product of 1300bp was obtained from *O.crenata* seeds and shoots, A PCR product of 1000bp was obtained from *O.cumana* seeds and shoots. However, RbcL-M marker failed to amplify PCR product with soil sample alone or with soil contaminated with *P.aegyptiaca* seeds or shoots. **(B)** The same samples were subjected to PCR detection using universal internal control primers (UCP-555). Arrows indicates the PCR product sizes.

Figure 2. PCR amplification patterns of ITS100 primers in *O.cumana* seeds **(A)** gDNA was extracted from soil samples (250 mg) containing the parasite seeds (1, 10, 25, 50, 100 and 250 seeds). For each sample, 100 ng gDNA were used for PCR amplification. The same samples were subjected to PCR detection using universal internal control primers (UCP-555). Arrows indicates the PCR product sizes **(B)** Standard curves generated for qPCR to quantify *O.cumana* seeds in a soil sample. standard curve generated by plotting the value of threshold cycle value (Ct) against log of the amount of template DNA (ng/μl) from *O.cumana.***(C)** Detection and quantification of *O. cumana* template DNA (ng) from known amount of the parasite seeds mixed with extracted soil sample DNA (10 μg). Each point on the standard curve was assayed in triplicate.

Figure 3. Quantification of *O.cumana* seeds in an infected sunflower field by qPCR and manually counting the inflorescences adjacent to the soil sample. **(A)** represnts a section of the sunflower infected field. Red arrows show *O.cumana* inflorescences. **(B)** Representing a

(GIS) model (Eizenberg et al., 2012) for soil sampling and characterize the spatial variation

in sunflower field infected with *O.cumana* in Havat Eden. Five soil samples (2, 4, 6, 8, 10)

of 500gm were collected from the infected field on May. Each sample was collected from a

depth of 0 to 20 cm in a total area of 1x1 meter. The blue spots indicate location of the

sample in the infected field. Red color in the plot represents high density of the parasite

inflorescences. Monitoring infectivity by counting the inflorescences was performed on July

and restricted to the total area 1x1 meter of the five soil samples. **(C)** quantification of *O.*

cumana seeds in a soil sample by qPCR (black bars) and monitoring the parasite

inflorescences counts in fields (Grey bars). The data are the means of three soil sample or

biological replicates. Vertical lines indicate SD of three independent measurements.

Figure 4. Evaluation the viability of *O.cumana* seeds by Tetrazolium test. **(A).** Tetrazolium

test was used to visually differentiate between metabolically active tissue from (cleaned

viable seed (a), viable seed with organic matter from an infected sunflower field (c), and

inactive tissues from (cleaned non-viable seed (b), non-viable seed with organic mater from

an infected sunflower field (d). Images were acquired using fluorescence microscope using

40X and 100X magnification. **(B)** The same test was also performed to evaluate percentage

of viability of *O.cumana* seeds in soil samples from the infected field (83%) as compared to

viability of the seeds from cleaned seed bank (25%). The data are the means of three separate

experiments.

A

B

Figure 1_Radi Aly et al., 2019

A

+Soil

← **100bp**

← **555bp**

B

y = -1.282ln(x) + 16.807
R² = 0.9994

Ct Value

gDNA from *O.cumana* seeds (ng/µl)

C

y = -1.332ln(x) + 20.566
R² = 0.9878

Ct Value

gDNA from *O.cumana* seeds, ng in
soil sample DNA (10µg)

Figure 2_Radi Aly et al., 2019

A

B

C

Figure 3_Radi Aly et al., 2019

A

B

Figure 4_Radi Aly et al., 2019

Development of specific molecular markers to distinguish and quantify broomrape species in agricultural field soil sample

Radi Aly[1]* (iD), Vinay K. Bari[1] (iD), Avishai Londner[1], Kocherman Marina1, Jackline Abu Nassar[1], Leena Taha-Salaime[1], Eizenberg Hanan[1] and Ran Lati[1]

[1]Department of Plant Pathology and Weed Research, Newe Ya'ar Research Center, Agricultural Research Organization (ARO), Israel.

Figure S1. Multiple sequence alignment of *O.cumana* and *O.crenata rbcL1* gene using online

tool CLUSTAL Omega

```
                                    RbcL-M-F
O.cumana    ------------------------------------------------------------ 0
O.crenata   ATGTCACCACAAACAGAACTAAAGCAAGTGTTGGATTCAAAGCGGGTGTTAAAGATAAA 60

O.cumana    ------------------------------------------------------------ 0
O.crenata   GAGTACAAATTGACTTATTATATTCCTGAATACAAAACTAAAGATACTGATATTTTAGCA 120

O.cumana    ------------------------------------------------------------ 0
O.crenata   GCATTTCGAGTAACTCCTCAACCTGAAGTTCCGCCTGAACAGGGGCTGCAGTAGCTGCCG 180

O.cumana    ------------------------------------------------------------ 0
O.crenata   AATCTTCTATTGGTACATGGACAACTGTGTGGACTGATGGACCCTTGATCGTTACAAAGG 240

O.cumana    ---------------------------TATCGAGTA----GACCTTGTGTTGTTATC 26
O.crenata   TCGATACTACCACATCACATCGAGCCCAGTTCTGGAGAAACAGATCAATATATTTGTTAT 300
                                     *  *  *** *       *   *  * **   *
                                                     RbcL-M-F
O.cumana    AGAATTTTTAATTCATAAGTTTTAGGGAGAAATTTATGTCACC--------ACAAACAGA 78
O.crenata   ATAGCTTACCCTTTAGAC-TTTTTTGAGGAAGGTTCTGTTACTAACATGTTTACTTCCAT 359
            * *  **   ** * *  **** *   *** **  ** *** **                *
O.cumana    AACTAAAGCAAGTGTTGGATTCAAAGTCCTGCGTGATCTATGTCTGGAATATCTGCGAAT 138
O.crenata   TATAGGAAATATATTTGGATTTAAAGCCCTGCGTGCTCTCACGTCTGGAAGATCTACGAAT 419
             *    *   * ******* **** ******** **** ***** ********* **** *****
O.cumana    CCCTCCTGCTTATATTAAAA---TTTTCCAAGGCCCGCTCCATGGTATTCAAGTTTAAAG 195
O.crenata   TCCTCCTGCTTATATTAAAATTATTTTTCCAAAGCCCGCCCCATGAGATTAAAGTTGAAAT 479
             ****************** *   ******** ****** ***** *** ***** ***
O.cumana    AGATAAATTAAACAAGTATGGTCGTCCCCTGTTGGGATGTATTATTAAACCTAAA----T 251
O.crenata   AGATAAATTGAACAAGTATGGTCGTCCCCTGTTGGGATGTACTATTAAATAAACCTAAAT 539
            ********* ***************************** *******   *       *
O.cumana    TGGGGTTATCCACTAAAAACTATGGGGGAACAGTTTATGAATGTCTTCGCGGTGGACTTG 311
O.crenata   TAGGATTATCCGC----------------------------------------TAAAAACC 560
            * ** ****** *                                         *   *
O.cumana    ATTGTACCAAAGATGATGAGAACGTAAATTCCCAGCCATTTATGCGCTGGAGAGATCGTT 371
O.crenata   ATGGGAGAACAGTTGATGAGAACGTAAACTCCCAGCCATTTATGCGCTGGAGAAATCGTT 620
            ** *  *   * ** ************* ********************** ******
O.cumana    TTTTATTTTGTGCTGAAGCAATTTATAAATCACAGGCTGAAACAGGCGAAATTAAAGGCC 431
O.crenata   TCTTATTTTGTGCCGAAGCAATTTATAAAACACAGGCTGAAACAGGCGAAATCAAAGGTC 680
            * *********** *************** ****************** ***** *
O.cumana    ATTACTTGAATACTACTGCAGGTACATGCGAGGAAATGATCAAAAGAGCTGTCTTTGCTA 491
O.crenata   ATTACTTGAATACTACTGCAGGTACATGTGAGGAAATGATCAAAAGAGCTGTTTTTGCCA 740
            **************************** ********************** ***** *
O.cumana    GAGAATTGGGAGTTCCTATTATAA--ATGTACGACTACTTAACAGTAGGATTCACTGCAA 549
O.crenata   GAGAATTGGGAGTTCCTATTATAATAATGCACGACTACTTAACAGGAGGATTCACTTCAA 800
            ************************   *** *************** ********* ***
O.cumana    ATACTAGCTTGGCTCATTATTGCCGTAATAATGTCC-----TACTTCTCACCGTGCAATG 604
O.crenata   ATACTAGCTTGGCTCATTATTGCCATAATAATGGCCTGGCCTACTTCTTCACCTGCAATG 860
            ************************ ******** **       *******   * *******
O.cumana    CATGTAGTTATTGATAGACAGAAGAACCATAGTATACATTTCCGTTGTACTAGCTAAAGC 664
O.crenata   CATGCAGTTATTTATAGACAGAAGAACCATGGTATACATTTCC-GTGTACTAGCTAAAGC 919
            **** ******* ***************** ************  **************
O.cumana    GTTACGTATGTCTGGTGGAGATCATATTCACTCTAGGACCGTAGTAGGTAAACTTGAAGG 724
O.crenata   ATTACGTATGTCTGGTGGAGATCATATTCACTTTGGGACTGCAGTAGGTAAACTTGAAGG 979
             ******************************** * **** * ***** * ****************
```

```
O.cumana    AGAAAGAGACATTACTTTGGGCTTTGTTGATTTATTGCGTGATGATTTTATTAA-AAAGA  783
O.crenata   AGAAAGAGACATTACTTTTGACTTTGTTGATTTATTGCGTGATGATTTTATTGAAAAAGA  1039
            *************** * *************************** * *****

O.cumana    TCGAAGTCACGGTATTTATTTTACTCAAGATTGGGTTTCTTTACCAGGTGTTTTTACTGT  843
O.crenata   TCGAAGTCGCGATATTTATTTCACCCAAGATTGGGTTTCTCTACCAGGTGTTATTCCTGT  1099
            ******** ** ********* ** ************** ********** ** ****

O.cumana    GGCTTCAGGGGGGTATTCACGTTTGGCATATGCCTTCATCT-GACGGAGATCTTTGGGGAT  902
O.crenata   GGCTTCAGGGGGGTATTCACGTTTGGCATATGCCTTCCCTGACCGCGAGATCTTTGGGGAT  1159
            ************************************        ***************

O.cumana    GATTCCGTACTACAGTTTGACGGAGGAACTTTAGGACATCCTTAGCGTAATGCACCAGGC  962
O.crenata   GATTCCATACTACAGTTTGGCGGAGGAACTTTAGGACATCCTTGGGGTAATGCACCAGGT  1219
            ****** *********** ********************** * *************

O.cumana    GCTCTGTAGCTAATCGAGTAGCTATAGAAGCATGTGTACAAGCTCGTAATGAAGGATGTA  1022
O.crenata   GC--TGTAGCTAATCGAGTAGCTATATAAGCATATGTACAAGCTCGTAATGAAGGACGTG  1277
            **  ********************* ****** ******************** **

                                                       RbcL-M-R
O.cumana    ATCTTGCTACTGAGGGGAATGCAATTATACGCGAGGCTAGGAAATGGAGCCCTGAACTAG  1082
O.crenata   ATCTTGCTGCTGAGGGTAATTATA-----CGTGAGGCTAGCAAAC<GGAGTCCTGAACTAG  1332
            ******** ******* ***     *   ** ******* *** **** **********
                                                         RbcL-M-R

O.cumana    TTGCTGTTTGTGAGGTATGTAAAGATATCAAATTTGAGTTTAAAGCAGTCGATACTTTGG  1142
O.crenata   TTGCTG------------------------------------------------------  1338
            ******

O.cumana    GTAAGTGTAAGTAAGATAACATTACTCTTCATTCTCTTAATTGAATTTCAATTAAATTCG  1202
O.crenata   ------------------------------------------------------------  1338

O.cumana    GCTTAATC   1210
O.crenata   --------   1338
```

Table S1. List of oligo's used in this study

Primer's name	Sequence 5 ⟶ 3'	Use in the study
UCP555-F	GTAGTCATATGCTTGTCTC	Universal internal control
UCP555-R	GGCTGCTGGCACCAGACTTGC	
RbcL-M-F	ATGTCACCACAAACAGAAAC	Molecular marker to identify *O.cumana* and *O.crenata*
RbcL-M-R	CAGCAACTAGTTCAGGCTCC	
ITS100-F	CAACGGATATCTCGGCTCTC	qRT-PCR to quantify broomrape
ITS100-R	TTGCGTTCAAAGACTCGATG	
ACT1-F	ATGGGCCAGAAAGATGCATATGTT	Housekeeping gene used for qRT-PCR expression normalization
ACT1-R	GTGTGATGCCAAATTTTCTCCATGT	