Development of specific molecular markers to distinguish and quantify broomrape species in a

soil sample from infected field

Radi Aly[1]* , Vinay K. Bari[1] , Avishai Londner[1], Jackline Abu Nassar[1], Leena Taha-Salaime[1],

Eizenberg Hanan[1] and Ran Lati[1]

[1]Department of Plant Pathology and Weed Research, Newe Ya'ar Research Center, Agricultural

Research Organization (ARO), Israel.

*Corresponding Author: Radi Aly

radi@volcani.agri.gov.il

https://orcid.org/0000-0002-5670-3798

https://orcid.org/0000-0002-6038-9586

Keywords: Plant parasite, *Phelipanche aegyptiaca, Orobanche* spp., ribulose-1,5-

bisphosphate carboxylase (*rbcL*), Internal transcribed spacer (*ITS*).

Conflict of interest: The authors have declared no conflict of interest

Abstract

Broomrapes (*Orobanche* and *Phelipanche*) are obligate holoparasites that cause heavy damage to

numerous crops, reducing the yield and its quality. The parasite develops in the soil and exerts the

greatest damage prior to its emergence; therefore the majority of field loss may occur before diagnosis

of infection. Because of the parasite tiny seed size (200 to 300 μm) and dormancy for several decades

in the field, it is very difficult to diagnose the parasite by conventional methods. Therefore, to restrict

the parasite seeds spread and contamination to other commercial fields, development of DNA-based

molecular markers to identify and quantify broomrape species in a soil sample is much needed. In this

study, we developed a specific molecular marker (RbcL-M) based on *rbcL* (large subunit of the

ribulose-bisphosphate carboxylase) gene from *Orobanche crenata* to differentiate between *Orobanche*

crenata and *Orobanche cumana.* Likewise, a specific marker (ITS100) based upon unique sequences

in the internal transcribed spacer (ITS) regions of the nuclear ribosomal DNA of *Phelipanche*

aegyptiaca to quantify three species of the parasite (*P. aegyptiaca*, *O. crenata* and *O. cumana*) in a

soil sample was developed. Genomic DNA was extracted from soil samples artificially infested with

broomrape seeds or tissue of *P. aegyptiaca*, *O. cumana* and *O. crenata* and subjected to PCR analysis.

RbcL-M marker successfully amplified a PCR product (1300bp) when *O. crenata* seeds or tissues

(collected from several locations in Israel) were added to the soil samples. The same marker amplified

a PCR product (1000bp) when *O. cumana* seeds or tissues were added to the soil samples. RbcL-M

marker did not amplify soil samples with seeds or tissues of *P. aegyptiaca* or any soil-borne DNA.

Furthermore, using ITS-100 marker and Real-Time PCR analysis, allowed quantitative diagnostic of

the parasite in a soil sample from infected sunflower field. As expected the universal internal control

primer (UCP-555) amplified a PCR product (555bp) when genomic DNA extracted from soil samples

with or without broomrape tissues. The development of an efficient, simple and robust molecular

marker to detect and distinguish between broomrape species, has a significant insights on assessment

the level of infestation and planning eradication program to the parasite in a field crop.

INTRODUCTION

Broomrapes (*Orobanche* and *Phelipanche* spp.) are member of Orobanchaceae family (C. Parker, 1993), is an underground root parasite mostly affecting agricultural crop plants (McNeal et al., 2013). These parasitic plants are unable to synthesize their photosynthetic assimilate due to lack of chlorophyll; and totally depend on host plants for the germination and growth by feeding through haustorium (Westwood et al., 2010). The Orobanchaceae family contains the largest number of parasitic species that form haustoria in roots, for example *Orobanche, Phelipanche* species (Yoshida et al., 2016). Haustoria are special organ of parasitic plant which invades host tissue and serve as connecting bridge that allows the parasite to obtain water, mineral and nutrient from host (Joel and Losnergoshen, 1994; Westwood et al., 2010).The natural habitats of broomrapes are the warm and temperate regions of the northern hemisphere, including the Mediterranean region (C. Parker, 1993), however branched broomrape *O. ramosa* is native to central and south-western Europe, and considered as a major threat to agricultural crops. They cause extensive damage by reducing the yield of all economically important crops belongs to Solanaceae, Fabaceae, Brasicaceae, Asteraceae and Apiaceae family (Aly, 2007; C. Parker, 1993; Joel, 2006). In Israel the significance of infested fields has increased dramatically during the last decade, causing heavy crop damage or even total yield losses.

In general, parasitized crops suffer from reductions in total biomass at the greatest expense to their productive tissue (Joel, 2006). Broomrapes cause many problems for agriculture farmers because of their tiny seed size (200-300µm) which makes them difficult to detect in harvested crop seeds and in agriculture soil (Joel, 1987). Moreover, small broomrape germinates and grows only in the presence of a susceptible host. In addition, small broomrape seeds can survive more than a decade in the soil (Habimana et al., 2014). Therefore, identification and quantification of broomrape seeds in a soil sample urgently required and will be helpful in prevention of crop damages by these parasites (Joel et al., 1996). The conventional methods of detection and differentiation such as seed coat morphological feature (Joel, 1987), Random amplified polymorphic DNA technique (RAPD) (Katzir et al., 1996; Paran et al., 1997), Intersimple sequence repeat (ISSR) (Benharrat et al., 2002) and high resolution

melting analysis (Rolland et al., 2016) has some limitation to distinguish and quantify broomrapes

exist in the soil. Recent progress in sequencing broomrape plastid genome, which was made available,

providing new insight to develop molecular marker for specific broomrape species identification

(Cusimano and Wicke, 2016; Wicke et al., 2013). Genetic or DNA based molecular marker such as

internal transcribe spacer (ITS), plastidial ribulose bisphosphate carboxylase (Rubisco) large subunit

(*rbcL*) and maturase K (*matK*) are among the most widely used molecular markers in evolutionary,

taxonomy, phylogenetic and diversity identification of many organisms including *Orobanche*, have

been established in recent years (Agarwal et al., 2008; Park et al., 2008; Schneeweiss et al., 2004).

In order to develop a complementary methodology for mapping parasitic plants in the field, a

procedure includes: a known geographical information systems (GIS) for soil sampling that will

characterize the spatial variation in the field (Eizenberg et al., 2012), and at the same time, use of

molecular markers to identifying broomrapes in a soil sample. Those molecular markers would assist

detection and diagnosis of broomrapes species and population level in the soil. In this study we

developed a simple and easy method based on *rbcL* gene to differentiate between three broomrape

species *O. crenata, O. cumana* and *P. aegyptiaca*, the most common and destructive weed in Israel.

We also developed a specific molecular marker to quantify the parasite seeds in soil sample in an

infected sunflower field.

MATERIAL AND METHODS

Plant material

Seeds, shoots and inflorescences of three broomrape species (*P. aegyptiaca, O. cumana and O.*

crenata) were collected from different Israeli locations (Golan heights, Havat Eden- Emic Betshan

and Jordan valley respectively). Seeds were harvested from freshly inflorescences and stored at 4°C

while shoots and inflorescences were dried and stored at −80°C until used. Two hundred milligrams

of parasite plant seeds/tissue were ground manually in presence of liquid nitrogen with the help of

pestle and mortar. Total genomic DNA was extracted using the GeneGET Plant genomic DNA

extraction kit according to the manufacturer instructions (Thermo Scientific).

ITS and *rbcL* sequences and primers design

The internal transcribed spacer (ITS) regions of *Phelipanche aegyptiaca* (GenBank-Accession No. AY209327) was used to design ITS100 marker. The PCR product yielded a 100bp fragment in length. UCP-555 was used as universal internal control primer, which amplifies a region of the small subunit of nrDNA (555 bp) from a wide variety of microorganism such as protists, fungi, and plants. The length of the *rbcL* sequences was 1211bp and 1290bp for *O. cumana* (GenBank: AF090349.1) and *O. crenata* (GenBank: AY582191.1) respectively. Sequence similarities between broomrape species were done using Clustal Omega software. Design of primers for the qPCR detection of broomrape species was based on the alignment of the ITS of nuclear ribosomal DNA regions. All primers were designed with Gene Runner software. PCR amplifications of *rbcL* sequences were performed with the RbcL-M-F and RbcL-M-R primers using PCR Ready Mix (BioLabs, Israel), and 50ng DNA-template. The following PCR conditions were used: denaturation for 5 min at 95°C; 32 cycles with 30 second at 95°C, 30 second at 55°C, 1.30 min at 72°C; and final elongation for 10 min at 72°C.

Soil sampling and extraction of Genomic DNA

Design and mapping the soil samples in the sunflower tested field were performed according to (Eizenberg et al., 2012). Soil samples were collected from the infected field on May and counting the inflorescences was performed on July. Each sample includes 500gm of a soil sample that was introduced into one-litter container, mixed thoroughly with 1.5L of (13.3 M $CaCl_2$) solution and kept at room temperature for overnight. The next day, the upper phase including the organic materials and broomrape seeds were collected, dried on Whatmann paper and the parasite seeds were obtained by filtration through 50 and 100 mesh. Genomic DNA was extracted by PowerSoil kit (MO BIO Lab. Inc., Loker Ave West, Carlsbad, CA). according to the manufacturer's instructions (Sagova-Mareckova et al., 2008).

Quantitative real time PCR of soil sample from the sunflower fields

The quantitative real time PCR was performed using PerfeCTa® SYBR® Green Fast Mix®, ROX™

(Quanta biosciences) and ABI-Prism 7000 Real-Time PCR Detection System (Applied Biosystems)

according to the manufacturer's protocol. The qPCR was performed in quantitative reaction with final

volume of 10 µl including: 100ng DNA, 5 µl of SYBR Green Fast Mix- ROX and 500nM of each

ITS-100 primers. *O. cumana* actin was used as endogenous control and the relative gene expression

level was calculated using the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001). Two standard curves were

prepared: The first curve included standard points from *O. cumana* seeds ranging from 0.001 to 10 ng

DNA per tube, were made using 1:10-fold serial dilutions of broomrape DNA and were used for

qPCR assay. The second curve was generated by mixing different amounts of genomic DNA from *O.*

cumana (0.1 to 100 ng) with total genomic DNA (10 µg) extracted from 250 mg soil sample. Each

point on the standard curve was assayed in triplicate.

Viability evaluation of the parasite seeds by Tetrazolium test

Tetrazolium test was performed as described earlier (Lopez-Granados and Garcia-Torres, 1999;

Thorogood et al., 2009). Briefly, 50-100 seeds were hydrated between paper towel sheets for 2-3

days. After pre-conditioning, they were placed in flasks, covered with a 1% solution of 2,3, 5

triphenyl tetrazolium chloride (TZ) and incubated at 40°C for 2h. After that, the TZ solution was

discarded; and the seeds were washed thoroughly with sterile water. Seeds were directly evaluated for

viability under the microscope using magnification (40X and 100X) and classified in two categories:

viable seeds – orange to brown light coloured or non-viable seeds –yellow to white colour.

RESULTS

Design of *rbcL* specific marker to differentiate between *O. cumana* and *O. crenata*

In our previous study, we reported amplification of ITS region using ITS350 primer; which

specifically detect broomrape in soil sample (Aly et al., 2012). However, ITS350 marker does not

differentiate between broomrape species because of the high sequence similarity among ITS regions.

Therefore, we compared and analysed the *rbcL* gene sequences from these species using Clustal

Omega and designed new molecular marker to differentiate broomrape species such as *O. cumana* and

O. crenata (Fig. S1). To distinguish between the above broomrape species, probes differing at least by

five-base mismatches or more were designed. Based on sequence alignment, a molecular marker

(RbcL-M) was designed using *rbcL* gene from *O. cumana* (GenBank accession number: AF090349.1)

Soil samples were artificially contaminated with seeds or shoots of *P. aegyptiaca*, *O. crenata* and *O.*

cumana and subjected to PCR analysis. By PCR amplification using RbcL-M, we were able to

differentiate between *O. crenata* and *O .cumana* according to the length of the PCR products. For *O.*

crenata the PCR product size was 1300bp compared to 1000bp for *O. cumana*. However, RbcL-M

marker failed to amplify any PCR product while soil samples were artificially contaminated with *P.*

aegyptiaca seeds or shoots (Fig. 1a). As expected, the universal internal control primers amplified a

PCR product (555bp) from soil samples with or without broomrape shoots or seeds (Fig. 1b).

Development of specific ITS100 marker to detect seeds of three broomrape species in a soil

sample

To analyse the number of the broomrape seeds present in a soil sample from infected field, we

designed ITS100 primers from the highly conserved region of ITS350 region (Aly et al., 2012). To

determine the specificity of the marker to each broomrape specie, we first extracted genomic DNA

from soil samples (250mg) containing (0, 1, 10, 25, 50 100 and 250 seeds) of *P. aegyptiaca*, *O.*

crenata or *O. cumana* and subjected to PCR analysis. ITS100 primers successfully amplified a PCR

product (100bp) when *P. aegyptiaca*, *O. crenata* or *O. cumana* were mixed with soil sample, except

soil sample without *Orobanche* seeds. However, a PCR product (555bp) was amplified in all samples

including soil with or without broomrape seeds by using universal control primer UCP-555 (Fig. 2a).

In addition, the specificity of the markers generated in this study were evaluated by testing DNA

extracted from sunflower, carrot and tomato leaves and roots corresponding to possible respond to the

host plants. Fortunately, the qPCR assay always yielded negative results for these plant species (data

not shown). Next to quantify *Orobanche*, we generated standard curve using *O. cumana* genomic

DNA and ITS100 marker utilizing qPCR procedure. *O. cumana* genomic DNA was purified and serial

dilutions (1:10 fold) ranging from 0.001 to 10 ng/µl were used for qPCR assay. The standard curve

for *O. cumana* exhibited a slope of -1.28 and (R^2) of 0.9994 (Fig. 2b). The same type of assay was

performed with *O. cumana* genomic DNA extracted from soil sample (250 mg) mixed with the

parasite seeds (0.1, 1, 10 and 100 mg seeds) and subjected to qPCR. The standard curve for *O.*

cumana seeds mixed with soil sample, exhibited a slope of -1.332 and (R^2) of 0.987 (Fig. 2c).

Prediction and quantification of *O. cumana* seeds in sunflower field

Previously, we have developed a protocol that allows extraction of genomic DNA from few tiny seeds

of *Orobanche* spp. in a soil sample and we were able to subject the sample DNA to a rapid PCR-assay

(Aly et al., 2012). Here, we extended the protocol, allowing for the identification and distinguishing

between soil-borne and parasite seed species that are collected from a field soil sample. Following

confirmation and validation of the ITS100 marker specificity to *O. cumana* seeds, geo-statistics model

for soil sampling that characterized the spatial variation in the field, was performed according to

(Eizenberg et al., 2012) in a sunflower field located at Havat Eden, Emic Betshan– Israel, which was

infected with *O. cumana* (Fig. 3a and b). Five soil samples (spot number 2,4,6,8 and 10) were

collected from a plot of 1 acre according to the illustration (Fig. 3c). Genomic DNA was extracted

from each soil sample and subjected to qPCR using ITS100 primers to predict and quantify *O.*

cumana DNA in naturally infected field. Then, genomic DNA from soil samples was extrapolated

with Ct values to quantify the number of seeds in each sample according to the generated Ct values

(Fig. 2b and c). The results showed variable quantity of *O. cumana* seeds in the soil sample ranged

from 2 to 22 parasite seeds (Fig. 3d) in 250mg soil sample processed originally from 1kg soil sample

from the infected field. Sample no. 6 showed the highest density of *O. cumana* seeds (22 seeds) in the

sunflower infected field as compared to the other samples. Sample no. 6 was collected from a highly

infected area as was shown by the geo-statistics model proposed by (Eizenberg et al., 2012).

Concomitantly, we monitored and counted the parasite inflorescences in adjacent to the position were

the soil samples were taken. Our results revealed that average density of 0.6 inflorescences /m^2, in a

total area of 1acre was recorded. Inflorescences counts in the selected area in the field were also

variable ranged from 2 to 15 and were less than the seed numbers found in the soil sample when

qPCR was applied (Fig. 3d).

Evaluation viability of the parasite seeds in sunflower infected field

To determine viability of broomrape seeds in a sunflower infected field, a tetrazolium test (TZ) (Lopez-Granados and Garcia-Torres, 1999; Thorogood et al., 2009) was conducted. According to TZ test, viable parasite seeds tends to have orange to brown colour because of its metabolite activity, while non-viable tends to have yellow to faint colour (Fig. 4a). To evaluate the parasite seed viability, two treatments were assayed: soil samples (organic matter only) collected from sunflower field located at Havat Eden, Emic Betshan, naturally infested with *O. cumana* and pure *O. cumana* seeds (1 mg) prepared from an old seed stock. Our results indicate that the highest percentage (% of total) of viable *O. cumana* seeds were found in soil samples collected from the sunflower field (83%) as compared to 25% found in the old seed stock (Fig. 4b).

DISCUSSION

Parasitic weeds are often exert their greatest damage prior to their emergence; therefore, the majority of field loss may occur before diagnosis of infection. Early detection of infection by visual inspection is not possible. This hampers early control strategy efforts and the detection methods used by crop scientists are laborious and time consuming. Most of the methods developed to detect broomrape seeds in soil samples, need several steps such as seed purification (flotation, filtration and binocular microscope observation (Kroschel, 2001). Despite of the great advances in genome sequencing approaches that has already been achieved, our understanding towards genomic evolution of Orobancheaceae is still incomplete due to aberrant evolution of plastid genome of holoparasite (Park et al., 2007). Available DNA sequence data are insufficient to differentiate closely related species, such as the weedy parasites *O. cumana, O. crenata* and *P. aegyptiaca*.

Genetic or DNA based marker techniques such as ITS or *rbcL* are routinely being used in evolutionary, taxonomical, phylogenetic and diversity studies (Agarwal et al., 2008; Park et al., 2008; Schneeweiss et al., 2004). Diagnosis and early identification of the parasite species in the field by soil sampling is of great importance to farmers and it is particularly needed due to the host-parasite specific interaction, the difficulty to detect the tiny parasite seeds by conventional methods in

contaminated fields and because seeds will only germinate and grow in the presence of a susceptible

host. The gene encoding the large subunit of ribulose-1,5-bisphosphate carboxylase/oxygenase

(Rubisco; *rbcL*) is retained as a pseudogene in *Orobanche* and *Phelipanche* (Wicke et al., 2013). In

the present work, we designed sets of primers targeting the plastid *rbcL* gene sequence from *O.*

cumana to differentiate between three broomrape species (*O. cumana, O. crenata* and *P. aegyptiaca*)

the most common and harmful species in field crops in Israel. Another sets of primers were designed

to target ITS sequences (Schneeweiss et al., 2004) to quantify the parasite seeds potentially present in

a soil sample to help for seed bank assessment in the field. We found that RbcL-M primers (marker)

were useful to distinguish between DNA from *O. cumana* and *O. crenata* producing different

amplicon sizes (a specific PCR product about 1300bp for *O. crenata* and 1000bp for *O. cumana*).

Subsequent BLASTn with previously published broomrape genomic data (Cusimano and Wicke,

2016) reconfirmed the differences between the two species. In contrast, DNA from *P. aegyptiaca*

failed to produce PCR amplicon suggesting possibly that *Phelipanche* species completely lost *rbcL*

gene (Delavault et al., 1995; Leebens-Mack and de Pamphilis, 2002).To allow differentiation between

P. aegyptiaca from other species, a specific molecular marker was recently designed (Aly et al.,

2019), (GenBank accession numbers MK637618-637624).

To diagnose and quantify broomrape species in a seed stock (Dongo et al., 2012), different types of

nuclear and plastid DNA markers have been proposed. (Schneeweiss et al., 2004) were the first to

present molecular phylogenetic analysis using nuclear ITS sequences. ITS-based markers were also

used to detect *P. aegyptiaca* seeds in a soil sample (Aly et al., 2012) and quantify contamination of

O. ramosa and *O. cumana* in crop seed lots (Dongo et al., 2012). Here we developed ITS100 marker

that was based on primers consisting of unique sequences in the internal transcribed spacer (ITS)

regions of the nuclear ribosomal DNA (nrDNA) of *O.crenata*. ITS-100 marker was used with qPCR

assay to quantify *O. cumana* seeds in a soil sample from sunflower field located at Havat Eden, Emic

Betshan – Israel.

We were able to detect 0.1 mg *O. cumana* seeds in 250 mg soil sample (wt ⁄ wt). A detection threshold

of 0.1 mg broomrape seeds in 20g seed samples was previously reported (Dongo et al., 2012). For

detection, mapping and quantifying *O. cumana* seeds in the field, we used geographical information

systems (GIS) for soil sampling and other advanced technologies for parasitic weed mapping and field

history data storage (Eizenberg et al., 2012) followed by qPCR assay. Soil samples (500gr) were

collected from the sunflower field, organic material was extracted from each sample ending with

250mg then, genomic DNA was extracted and subjected to qPCR using ITS100 primers. This method

allowed specifically detecting and quantifying the DNA of *O. cumana* in a total DNA extract from

sunflower soil sample. Accordingly, the results of this assay can be also expressed as the number of

parasite seeds per kilogram of soil following extrapolation with the standard curve was prepared. Our

results indicate that samples collected from highly infected area (Fig. 3c) according to the geo-

statistics model proposed by Eizenberg et al. (2012), were with agreement with our qPCR assay

(sample no. 6 showed the highest density of *O. cumana* seeds (22 seeds) in the sunflower infected

field as compared to the other samples. However, no correlation was found between qPCR assay

(parasite seed number) compared to number of the parasite inflorescences collected from the same

location in the sunflower infected field. An explanation for that may be related to the viability of *O.*

cumana seeds, genomic DNA from a soil sample will contain viable and non-viable parasite seeds,

add to that, we cannot exclude the presence of some related parasite seeds in the same sample

therefore, we counted more parasite seeds using the qPCR assay. Specificity of the qPCR assay was

tested against several possible contaminants of soil-borne pathogens by using universal internal

control primer UCP-555 (White, 1990) or harvested crop seeds like sunflower and tomato using

ITS100. No amplification was observed, confirming the specificity of the marker.

The powerful of broomrape-infected field to distribute and contaminate the neighbouring non-infested

fields depends on the soil seed-bank and viability of the parasite seeds. Viability of broomrape seeds

in the sunflower infected field was determined by tetrazolium test. Our results showed the highest

percentage of viable broomrape seeds were found in soil samples collected from the sunflower field

(83%) as compared to 25% found in an old seed stock. We assume that the highest count recorded in

the sunflower field was due to the release and distribution of the fresh parasite seeds by the newly

parasite inflorescences showed up through the crop growth. We have to take in consideration that we

used an old seed stock (10 years old) that doesn't represent newly harvested seed stocks. Our

experience with seed germination of *O. cumana* with germination stimulant (GR) could reach more

than 90%. Additionally, we cannot exclude the presence of some related parasite seeds in the same sample from previous growth seasons.

In this study, we provide a simple, fast and non-expensive approach to distinguish and quantify broomrape seeds exist in a soil sample from a crop field. Molecular markers would assist accurate detection and population level of *Phelipanche* and *Orobanche* spp. in a soil sample and offer numerous advantages over conventional phenotype based alternatives, as they are stable and detectable in all tissues regardless of growth, differentiation or development. These methods could be helpful in precision agriculture, in which they provide answers routinely questioned by the farmers: are there parasite seeds in my crop field? What species? and how much seeds are exist in a soil sample?

ACKNOWLEDGMENTS

Results of this research were supported by the Chief Scientist of the Ministry of Agriculture and Rural Development - Israel, grant No.132-1499-10 (MEZAMALEKET). VKB is grateful to the ARO-Volcani Center, Agricultural Ministry of Israel for providing the Postdoctoral fellowship.

REFERENCES

Agarwal, M., Shrivastava, N., and Padh, H. (2008). Advances in molecular marker techniques and

their applications in plant sciences. Plant Cell Rep *27*, 617-631.

Aly, R. (2007). Conventional and biotechnological approaches for control of parasitic weeds. In Vitro

Cell Dev-Pl *43*, 304-317.

Aly, R., Eizenberg, H., Kocherman, M., Abu-Nassar, J., Taha, L., and Saadi, I. (2012). Use of ITS

nuclear sequences from Phelipanche aegyptiaca as a direct tool to detect single seeds of broomrape

species in the soil. Eur J Plant Pathol *133*, 523-526.

Aly, R., Lati, R., Abu-Nassar, J., Ziadna, H., Achdari, G., Münchow, C.S.v., Wicke, S., Bari, V.K.,

and Eizenberg, H. (2019). The weedy parasite Phelipanche aegyptiaca attacks Brassica rapa var. rapa

L. for the first time in Israel. Plant Disease Note.

Benharrat, H., Veronesi, C., Theodet, C., and Thalouarn, P. (2002). Orobanche species and population

discrimination using intersimple sequence repeat (ISSR). Weed Res *42*, 470-475.

C. Parker, C.R.R. (1993). Parasitic weeds of the world : biology and control (Wallingford, Oxon :

CAB International).

Cusimano, N., and Wicke, S. (2016). Massive intracellular gene transfer during plastid genome

reduction in nongreen Orobanchaceae. New Phytol *210*, 680-693.

Delavault, P., Sakanyan, V., and Thalouarn, P. (1995). Divergent evolution of two plastid genes, rbcL

and atpB, in a non-photosynthetic parasitic plant. Plant Mol Biol *29*, 1071-1079.

Dongo, A., Leflon, M., Simier, P., and Delavault, P. (2012). Development of a high-throughput real-

time quantitative PCR method to detect and quantify contaminating seeds of Phelipanche ramosa and

Orobanche cumana in crop seed lots. Weed Res *52*, 34-41.

Eizenberg, H., Aly, R., and Cohen, Y. (2012). Technologies for Smart Chemical Control of

Broomrape (Orobanche spp. and Phelipanche spp.). Weed Sci *60*, 316-323.

Habimana, S., Nduwumuremyi, A., and Chinama, R.J.D. (2014). Management of orobanche in field

crops- A review. J Soil Sci Plant Nut *14*, 43-62.

Joel, D.M. (1987). Detection and Identification of Orobanche Seeds Using Fluorescence Microscopy.

Seed Sci Technol *15*, 119-124.

Joel, D.M., Hershenhorn, Y., Eizenberg, H., Aly, R., Ejeta, G., Rich, P.J., Ransom, J.K., Sauerborn, J.

and Rubiales, D. (2006). Biology and Management of Weedy Root Parasites. In Horticultural

Reviews, New Jersey: Wiley, 267–349.

Joel, D.M., and Losnergoshen, D. (1994). The Attachment Organ of the Parasitic Angiosperms

Orobanche Cumana and O-Aegyptiaca and Its Development. Can J Bot *72*, 564-574.

Joel, D.M., Portnoy, V., and Katzir, N. (1996). Identification of single tiny seeds of Orobanche using

RAPD analysis. Plant Mol Biol Rep *14*, 243-248.

Katzir, N., Portnoy, V., Tzuri, G., Joel, D.M., and Castejon-Munoz, M. (1996). Use of random

amplified polymorphic DNA (RAPD) markers in the study of the parasitic weed Orobanche. Theor

Appl Genet *93*, 367-372.

Kroschel (2001). A Technical Manual for Parasitic Weed Research and Extension. Springer,

Dordrecht.

Leebens-Mack, J., and de Pamphilis, C. (2002). Power analysis of tests for loss of selective constraint

in cave crayfish and nonphotosynthetic plant lineages. Mol Biol Evol *19*, 1292-1302.

Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time

quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods *25*, 402-408.

Lopez-Granados, F., and Garcia-Torres, L. (1999). Longevity of crenate broomrape (Orobanche

crenata) seed under soil and laboratory conditions. Weed Sci *47*, 161-166.

McNeal, J.R., Bennett, J.R., Wolfe, A.D., and Mathews, S. (2013). Phylogeny and origins of

holoparasitism in Orobanchaceae. Am J Bot *100*, 971-983.

Paran, I., Gidoni, D., and Jacobsohn, R. (1997). Variation between and within broomrape

(Orobanche) species revealed by RAPD markers. Heredity *78*, 68-74.

Park, J.M., Manen, J.F., Colwell, A.E., and Schneeweiss, G.M. (2008). A plastid gene phylogeny of

the non-photosynthetic parasitic Orobanche (Orobanchaceae) and related genera. J Plant Res *121*,

365-376.

Park, J.M., Manen, J.F., and Schneeweiss, G.M. (2007). Horizontal gene transfer of a plastid gene in

the non-photosynthetic flowering plants Orobanche and Phelipanche (Orobanchaceae). Mol

Phylogenet Evol *43*, 974-985.

Rolland, M., Dupuy, A., Pelleray, A., and Delavault, P. (2016). Molecular Identification of

Broomrape Species from a Single Seed by High Resolution Melting Analysis. Front Plant Sci *7*.

Sagova-Mareckova, M., Cermak, L., Novotna, J., Plhackova, K., Forstova, J., and Kopecky, J. (2008).

Innovative methods for soil DNA purification tested in soils with widely differing characteristics.

Appl Environ Microbiol *74*, 2902-2907.

Schneeweiss, G.M., Colwell, A., Park, J.M., Jang, C.G., and Stuessy, T.F. (2004). Phylogeny of

holoparasitic Orobanche (Orobanchaceae) inferred from nuclear ITS sequences. Mol Phylogenet Evol

30, 465-478.

Thorogood, C.J., Rumsey, F.J., and Hiscock, S.J. (2009). Seed viability determination in parasitic

broomrapes (Orobanche and Phelipanche) using fluorescein diacetate staining. Weed Res *49*, 461-

468.

Westwood, J.H., Yoder, J.I., Timko, M.P., and dePamphilis, C.W. (2010). The evolution of parasitism

in plants. Trends in Plant Science *15*, 227-235.

White, T.J., T. D. Bruns, S. B. Lee, and J. W. Taylor (1990). Amplification and direct sequencing of

fungal ribosomal RNA Genes for phylogenetics.

Wicke, S., Muller, K.F., de Pamphilis, C.W., Quandt, D., Wickett, N.J., Zhang, Y., Renner, S.S., and

Schneeweiss, G.M. (2013). Mechanisms of Functional and Physical Genome Reduction in

Photosynthetic and Nonphotosynthetic Parasitic Plants of the Broomrape Family. Plant Cell *25*, 3711-

3725.

Yoshida, S., Cui, S.K., Ichihashi, Y., and Shirasu, K. (2016). The Haustorium, a Specialized Invasive

Organ in Parasitic Plants. Annu Rev Plant Biol *67*, 643-667.

 Figure legends

Fig. 1. (a) Specificity of RbcL-M marker and PCR amplification patterns in three broomrape species

(*P.aegyptiaca, O.cumana and O.crenata*). Soil samples (250 mg) were artificially contaminated with

broomrape seeds (100) or 10 mg shoots of *P.aegyptiaca, O.crenata and O.cumana* and subjected to

PCR analysis using RbcL-M primers. Genomic DNA was extracted and 100 ng were used for PCR

amplification. A PCR product of 1300bp was obtained from *O.crenata* seeds and shoots, A PCR

product of 1000bp was obtained from *O.cumana* seeds and shoots. However, RbcL-M marker failed

to amplify PCR product with soil sample alone or with soil contaminated with *P.aegyptiaca* seeds or

shoots. **(b)** The same samples were subjected to PCR detection using universal internal control

primers (UCP-555). Arrows indicates the PCR product sizes.

Fig. 2. PCR amplification patterns of ITS100 primers in *O.cumana* seeds **(a)** gDNA was extracted

from soil samples (250 mg) containing the parasite seeds (1, 10, 25, 50, 100 and 250 seeds). For each

sample, 100 ng gDNA were used for PCR amplification. The same samples were subjected to PCR

detection using universal internal control primers (UCP-555). Arrows indicates the PCR product sizes

(b) Standard curves generated for qPCR to quantify *O.cumana* seeds in a soil sample. standard curve

generated by plotting the value of threshold cycle value (Ct) against log of the amount of template

DNA (ng/µl) from *O.cumana*.**(c)** Detection and quantification of *O. cumana* template DNA (ng) from

known amount of the parasite seeds mixed with extracted soil sample DNA (10 µg). Each point on the

standard curve was assayed in triplicate.

Fig. 3. Quantification of *O.cumana* seeds in an infected sunflower field by qPCR and manually

counting the inflorescences adjacent to the soil sample. **(a and b)** represents a section of the sunflower

infected field. Red arrows show *O.cumana* inflorescences. **(c)** Representing a (GIS) model (Eizenberg

et al., 2012) for soil sampling and characterize the spatial variation in sunflower field infected with

O.cumana in Havat Eden. Five soil samples (2, 4, 6, 8, 10) of 500gm were collected from the infected

field on May. Each sample was collected from a depth of 0 to 20 cm in a total area of 1x1 meter. The

blue spots indicate location of the sample in the infected field. Red color in the plot represents high

density of the parasite inflorescences. Infectivity was also monitored by counting the inflorescences

on July and was restricted to the total area 1x1 meter of the five selected soil samples. **(d)**

quantification of *O. cumana* seeds in a soil sample by qPCR (black bars) and monitoring the parasite

inflorescences counts in fields (Grey bars). The data are the means of three soil samples or biological

replicates. Vertical lines indicate SD of three independent measurements.

Fig. 4. Evaluation the viability of *O.cumana* seeds by Tetrazolium test. **(a)** Tetrazolium test was used

to visually differentiate between metabolically active tissue from (cleaned non-viable seed (a-1), non-

viable seed with organic matter from an infected sunflower field (a-3), and active tissues from

(cleaned viable seed (a-2), viable seed with organic mater from an infected sunflower field (a-4).

Images were acquired using fluorescence microscope using 40X and 100X magnification. **(b)** The

same test was also performed to evaluate percentage of viability of *O.cumana* seeds in soil samples

from the infected field (83%) as compared to viability of the seeds from cleaned old seed bank (25%).

The data are the means of three separate experiments with vertical lines indicating SD.

Fig. 1

Fig. 2

a

b

y = -1.282ln(x) + 16.807
R² = 0.9994

gDNA from *O.cumana* seeds (ng/µl)

c

y = -1.332ln(x) + 20.566
R² = 0.9878

gDNA from *O.cumana* seeds, ng in
soil sample DNA (10µg)

Fig. 3

a   b   c

d

Fig. 4

100X   40X

a

b

Development of specific molecular markers to distinguish and quantify broomrape species in a soil sample from infected field

Radi Aly[1]* iD **,** Vinay K. Bari[1] iD **,** Avishai Londner[1]**,** Jackline Abu Nassar[1], Leena Taha-Salaime[1], Eizenberg Hanan[1] and Ran Lati[1]

[1]Department of Plant Pathology and Weed Research, Newe Ya'ar Research Center, Agricultural Research Organization (ARO), Israel.

Corresponding author: Radi Aly

radi@volcani.agri.co.il

Fig. S1. Multiple sequence alignment of *O.cumana* and *O.crenata rbcL* gene using online

tool CLUSTAL Omega

```
                              RbcL-M-F
O.cumana    ------------------------------------------------------------ 0
O.crenata   ATGTCACCACAAACAGAAACTAAAGCAAGTGTTGGATTCAAAGCGGGTGTTAAAGATAAA 60

O.cumana    ------------------------------------------------------------ 0
O.crenata   GAGTACAAATTGACTTATTATATTCCTGAATACAAAACTAAAGATACTGATATTTTAGCA 120

O.cumana    ------------------------------------------------------------ 0
O.crenata   GCATTTCGAGTAACTCCTCAACCTGAAGTTCCGCCTGAACAGGGGCTGCAGTAGCTGCCG 180

O.cumana    ------------------------------------------------------------ 0
O.crenata   AATCTTCTATTGGTACATGGACAACTGTGTGGACTGATGGACCCTTGATCGTTACAAAGG 240

O.cumana    -----------------------------TATCGAGTA----GACCTTGTGTTGTTATC 26
O.crenata   TCGATACTACCACATCACATCGAGCCCAGTTCTGGAGAAACAGATCAATATATTTGTTAT 300
                                        *  *  *** *      *  * * **  *
                                                           RbcL-M-F
O.cumana    AGAATTTTTAATTCATAAGTTTTAGGGAGAAATTTATGTCACC--------ACAAACAGA 78
O.crenata   ATAGCTTACCCTTTAGAC-TTTTTTGAGGAAGGTTCTGTTACTAACATGTTTACTTCCAT 359
            *  *   **     ** * * *    ****   *  ***   ** *** **        *
O.cumana    AACTAAAGCAAGTGTTGGATTCAAAGTCCTGCGTGATCTATGTCTGGAATATCTGCGAAT 138
O.crenata   TATAGGAAATATATTTGGATTTAAAGCCCTGCGTGCTCTACGTCTGGAAGATCTACGAAT 419
             *      *    ******* **** ******* **** **** ******* **** *****
O.cumana    CCCTCCTGCTTATATTAAAA---TTTTCCAAGGCCCGCTCCATGGTATTCAAGTTTAAAG 195
O.crenata   TCCTCCTGCTTATATTAAAATTATTTTCCAAAGCCCGCCCCATGAGATTAAAGTTGAAAT 479
             ****************      *******  ****** *****    *** ***** ***
O.cumana    AGATAAATTAAACAAGTATGGTCGTCCCCTGTTGGGATGTATTATTAAACCTAAA----T 251
O.crenata   AGATAAATTGAACAAGTATGGTCGTCCCCTGTTGGGATGTACTATTAAATAAACCTAAAT 539
            ********* ************************************* *******   *       *
O.cumana    TGGGGTTATCCACTAAAAACTATGGGGGAACAGTTTATGAATGTCTTCGCGGTGGACTTG 311
O.crenata   TAGGATTATCCGC----------------------------------------TAAAAACC 560
            *  ** ****** *                                            *  *
O.cumana    ATTGTACCAAAGATGATGAGAACGTAAATTCCCAGCCATTTATGCGCTGGAGAGATCGTT 371
O.crenata   ATGGGAGAACAGTTGATGAGAACGTAAACTCCCAGCCATTTATGCGCTGGAGAAATCGTT 620
            ** * *   * ** ************** ************************* ******
O.cumana    TTTTATTTTGTGCTGAAGCAATTTATAAATCACAGGCTGAAACAGGCGAAATTAAAGGCC 431
O.crenata   TCTTATTTTGTGCCGAAGCAATTTATAAAACACAGGCTGAAACAGGCGAAATCAAAGGTC 680
            *  *********** ***************** ******************** ***** *
O.cumana    ATTACTTGAATACTACTGCAGGTACATGCGAGGAAATGATCAAAAGAGCTGTCTTTGCTA 491
O.crenata   ATTACTTGAATACTACTGCAGGTACATGTGAGGAAATGATCAAAAGAGCTGTTTTTGCCA 740
            **************************** ************************ ***** *
O.cumana    GAGAATTGGGGAGTTCCTATTATAA--ATGTACGACTACTTAACAGTAGGATTCACTGCAA 549
O.crenata   GAGAATTGGGGAGTTCCTATTATAATAATGCACGACTACTTAACAGGAGGATTCACTTCAA 800
            ************************* *** ************** ******* *** ***
O.cumana    ATACTAGCTTGGCTCATTATTGCCGTAATAATGTCC-----TACTTCTCACCGTGCAATG 604
O.crenata   ATACTAGCTTGGCTCATTATTGCCATAATAATGGCCTGGCCTACTTCTTCACCTGCAATG 860
            ************************ ******* **      ******  * *  *******
O.cumana    CATGTAGTTATTGATAGACAGAAGAACCATAGTATACATTTCCGTTGTACTAGCTAAAGC 664
O.crenata   CATGCAGTTATTTATAGACAGAAGAACCATGGTATACATTTCC-GTGTACTAGCTAAAGC 919
            **** ******* ***************** ************ **************
O.cumana    GTTACGTATGTCTGGTGGAGATCATATTCACTCTAGGACCGTAGTAGGTAAACTTGAAGG 724
O.crenata   ATTACGTATGTCTGGTGGAGATCATATTCACTTTGGGACTGCAGTAGGTAAACTTGAAGG 979
            *********************************  * ***  * ****************

O.cumana    AGAAAGAGACATTACTTTGGGCTTTGTTGATTTATTGCGTGATGATTTTATTAA-AAAGA 783
O.crenata   AGAAAGAGACATTACTTTTGACTTTGTTGATTTATTGCGTGATGATTTTATTGAAAAAGA 1039
```

```
                  ***************** * ****************************** * *****
O.cumana   TCGAAGTCACGGTATTTATTTTACTCAAGATTGGGTTTCTTTACCAGGTGTTTTTACTGT 843
O.crenata  TCGAAGTCGCGATATTTATTTCACCCAAGATTGGGTTTCTCTACCAGGTGTTATTCCTGT 1099
           ******** ** ********* ** ************** ********** ** ****

O.cumana   GGCTTCAGGGGGTATTCACGTTTGGCATATGCCTTCATCT-GACGGAGATCTTTGGGGAT 902
O.crenata  GGCTTCAGGGGGTATTCACGTTTGGCATATGCCTTCCCTGACCGCGAGATCTTTGGGGAT 1159
           ************************************        ***************

O.cumana   GATTCCGTACTACAGTTTGACGGAGGAACTTTAGGACATCCTTAGCGTAATGCACCAGGC 962
O.crenata  GATTCCATACTACAGTTTGGCGGAGGAACTTTAGGACATCCTTGGGGTAATGCACCAGGT 1219
           ****** *********** ********************** * *************

O.cumana   GCTCTGTAGCTAATCGAGTAGCTATAGAAGCATGTGTACAAGCTCGTAATGAAGGATGTA 1022
O.crenata  GC--TGTAGCTAATCGAGTAGCTATATAAGCATATGTACAAGCTCGTAATGAAGGACGTG 1277
           **   ******************** ****** ******************** **

                                                          RbcL-M-R
O.cumana   ATCTTGCTACTGAGGGGAATGCAATTATACGCGAGGCTAGGAAATGGAGCCCTGAACTAG 1082
O.crenata  ATCTTGCTGCTGAGGGTAATTATA-----CGTGAGGCTAGCAAAG GGAGTCCTGAACTAG 1332
           ******** ****** *** *      ** ******* *** * *** **********
                                                          RbcL-M-R

O.cumana   TTGCTGTTTGTGAGGTATGTAAAGATATCAAATTTGAGTTTAAAGCAGTCGATACTTTGG 1142
O.crenata  TTGCTG------------------------------------------------------ 1338
           ******

O.cumana   GTAAGTGTAAGTAAGATAACATTACTCTTCATTCTCTTAATTGAATTTCAATTAAATTCG 1202
O.crenata  ------------------------------------------------------------ 1338

O.cumana   GCTTAATC   1210
O.crenata  --------   1338
```

Table S1. List of oligo's used in this study

Primer's name	Sequence 5 ⟶ 3'	Use in the study
UCP555-F	GTAGTCATATGCTTGTCTC	Universal internal control
UCP555-R	GGCTGCTGGCACCAGACTTGC	
RbcL-M-F	ATGTCACCACAAACAGAAAC	Molecular marker to identify *O.cumana* and *O.crenata*
RbcL-M-R	CAGCAACTAGTTCAGGCTCC	
ITS100-F	CAACGGATATCTCGGCTCTC	qRT-PCR to quantify broomrape
ITS100-R	TTGCGTTCAAAGACTCGATG	
ACT1-F	ATGGGCCAGAAAGATGCATATGTT	Housekeeping gene used for qRT-PCR expression normalization
ACT1-R	GTGTGATGCCAAATTTTCTCCATGT	