Using the basic reproduction number to assess the risk of

transmission of lumpy skin disease virus by biting insects

Simon Gubbins*

The Pirbright Institute, Ash Road, Pirbright, Surrey GU24 0NF, U.K.

* Corresponding author (email: simon.gubbins@pirbright.ac.uk; Tel. +44 1483 232441; Fax

+44 1483 232448); ORCID: 0000-0003-0538-4173

Short running title: R_0 for LSDV

Summary

In recent years, lumpy skin disease virus (LSDV) has emerged as a major threat to cattle

outside Africa, where it is endemic. Although evidence suggests that LSDV is transmitted by

the bites of blood sucking arthropods, few studies have assessed the risk of transmission

posed by particular vector species. Here this risk is assessed by calculating the basic

reproduction number (R_0) for transmission of LSDV by five species of biting insect: the

stable fly, *Stomoxys calcitrans*, the biting midge, *Culicoides nubeculosus*, and three mosquito

species, *Aedes aegypti*, *Anopheles stephensi* and *Culex quinquefasciatus*. Parameters relating

to mechanical transmission of LSDV were estimated using new analyses of previously-

published data from transmission experiments, while vector life history parameters were

derived from the published literature. Uncertainty and sensitivity analyses were used to

compute R_0 for each species and to identify those parameters which influence its magnitude.

Results suggest that *S. calcitrans* is likely to be the most efficient at transmitting LSDV, with

Ae. aegypti also an efficient vector. By contrast, *C. nubeculosus*, *An. stephensi*, and *Cx.*

quinquefasciatus are likely to be inefficient vectors of LSDV. However, there is considerable

uncertainty associated with the estimates of R_0, reflecting uncertainty in most of the

constituent parameters. Sensitivity analysis suggests that future experimental work should

focus on estimating the probability of transmission from insect to bovine and on the virus

inactivation rate in insects.

Keywords: cattle; epidemiology; transmission model; Bayesian methods; uncertainty

analysis; sensitivity analysis

1   Introduction

Lumpy skin disease (LSD) is an important transboundary disease of cattle and is caused by lumpy skin disease virus (LSDV). Historically, LSD outbreaks have largely been confined to Africa (EFSA Panel on Animal Health and Welfare, 2015), where the disease contributes to rural poverty and food insecurity (Tuppurainen & Oura, 2012; Molla, de Jong, Gari & Frankena, 2017). In recent years, however, LSDV has emerged as a major threat to cattle outside of Africa. In 2012 it spread to the Middle East, through Israel and the Lebanon, reaching Turkey in 2013 (Alkhamis & VanderWaal, 2016). In 2015, the first cases of LSD were reported in Greece (Tasioudi et al., 2016) and the virus subsequently spread to much of the Balkans (Mercier et al., 2018).

Evidence from both the field and from experiments indicates that transmission of LSDV by most direct and indirect routes is inefficient (Weiss, 1968; Carn & Kitching, 1995). The principal transmission route for LSDV is believed to be via the bites of blood sucking arthropods (Tuppurainen & Oura, 2012; EFSA Panel on Animal Health and Welfare, 2015). This has been shown experimentally for *Aedes aegypti* mosquitos (Chihota, Rennie, Kitching & Mellor, 2001) and for *Rhipicephalus appendiculatus* male ticks (Tuppurainen et al., 2013). The stable fly, *Stomoxys calcitrans*, has been incriminated as a potential vector in Israel by comparing its seasonal abundance with the seasonality of LSD cases (Kahana-Sutin, Klement, Lensky & Gottlieb, 2017) and because of its ability to transmit the closely related capripox virus between sheep and goats (Kitching & Mellor, 1986; Mellor, Kitching & Wilkinson, 1987). The potential for the biting midge, *Culicoides nubeculosus*, and two other mosquito species, *Culex quinquefasciatus* and *Anopheles stephensi*, to transmit LSDV has also been assessed, though for these three species attempts at transmission were unsuccessful (Chihota, Rennie, Kitching & Mellor, 2003).

In this study, the risk of transmission of LSDV posed by different insect species is explored by estimating the basic reproduction number for five species of biting insect (*S. calcitrans*, *C. nubeculosus*, *Ae. aegypti*, *An. stephensi* and *Cx. quinquefasciatus*). These species were selected as their potential role in LSDV transmission has been investigated previously (Chihota, Rennie, Kitching & Mellor, 2001, 2003) and they represent insect species (or at least genera) that are relevant to Europe. The basic reproduction number, denoted by R_0, is the "average number of secondary cases arising from the introduction of a single infected individual into an otherwise susceptible population" (Diekmann & Heesterbeek, 2000). An outbreak can occur only if $R_0 > 1$ and, consequently, R_0 provides a means to assess the risk posed by each vector species.

Using a transmission model, an expression for R_0 is derived that shows how it relates to the underlying transmission processes in insect vectors and cattle. These constituent parameters are then estimated using data from the published literature. In particular, data from transmission experiments involving the five putative vector species (Chihota, Rennie, Kitching & Mellor, 2001, 2003) are re-analysed using Bayesian methods to quantify the uncertainty in parameters relating to mechanical transmission. In addition, the latent, infectious and incubation periods for LSDV are estimated from the outcome of challenge experiments (Tuppurainen, Venter & Coetzer, 2005; Babiuk et al., 2008), again using Bayesian methods. Finally, uncertainty and sensitivity analyses are used to calculate R_0 and to determine which of the constituent parameters has the greatest influence on its magnitude.

2   Materials and methods

2.1   Basic reproduction number for LSDV

87   For an infection transmitted mechanically by biting insects the basic reproduction number can

88   be written as,

$$R_0 = \sqrt{\frac{ba}{(\gamma+\mu)} \times \frac{\beta ma}{r_I}}. \tag{1}$$

A full mathematical derivation of this expression for R_0, including the transmission model on

which it is based, is presented in the supporting information (Text S1). However, the

expression, (1), can be understood heuristically as follows. After it feeds on an infected

animal, an insect remains infected (and infectious) until the virus becomes inactivated or it

dies, a period which lasts on average $1/(\gamma+\mu)$ days, where γ is the virus inactivation rate and μ

is the vector mortality rate. During this time it will bite susceptible cattle a times per day

(where a is the reciprocal of the time interval between blood meals) and a proportion, b, of

these bites (i.e. the probability of transmission from insect to bovine) will result in a newly

infected host. Once infected, a bovine will remain infectious for the duration of its infectious

period, which lasts $1/r_I$ days on average. During this time the host will be bitten by

susceptible insects on average $m \times a$ times per day (here $m=N/H$ is the vector to host ratio and

N and H are the number of vectors and hosts, respectively), a proportion, β, of which will

result in a newly infected vector (i.e. the probability of transmission from bovine to insect).

2.2   Mechanical transmission of LSDV by insects

Data on mechanical transmission of lumpy skin disease virus to cattle by five species of

biting insect (*S. calcitrans*, *C. nubeculosus*, *Ae. aegypti*, *An. stephensi* and *Cx.*

quinquefasciatus) were extracted from the published literature (Chihota, Rennie, Kitching &

Mellor, 2001, 2003). This provided the number of positive insects (i.e. those for which viral

DNA was detected; virus isolation was only carried out for a small number of pooled

samples) and the number of insects tested after feeding on a LSDV-infected bovine (*S.*

calcitrans, *C. nubeculosus* or *Ae. aegypti*) or a blood-virus mix via a membrane (*An.*

stephensi and *Cx. quinquefasciatus*) at each day post feeding (Table S1). It also provided

information on whether or not transmission occurred when batches of insects that had

previously fed on an infected animal or received an infected blood meal were allowed to

refeed on a naïve bovine (Table S1).

These data were used to estimate the virus inactivation rate (γ), the probability of

transmission from bovine to insect (β) and the probability of transmission from insect to

bovine (b) for each species. Parameters were estimated in a Bayesian framework to facilitate

the incorporation of uncertainty in estimates of R_0. The likelihood for the data is given by,

121
$$L(\gamma, \beta, b) = \prod_i \binom{N_i}{Y_i} p_i^{Y_i} (1 - p_i)^{N_i - Y_i} \times$$
$$\prod_j q_j^{I_j} (1 - q_j)^{1 - I_j}, \tag{2}$$

122   where Y_i and N_i are the number of positive insects and number of insects tested at t_i days post

123   feeding, respectively, and I_j is a variable indicating whether ($I_j=1$) or not ($I_j=0$) transmission

124   occurred when insects were allowed to refeed on naïve animal j at t_j days post initial feed. In

125   equation (2),

126
$$p_i = \beta \exp(-\gamma t_i), \tag{3}$$

127   is the probability that an insect is positive at t_i days post feeding and,

128
$$q_j = 1 - (1 - b\beta \exp(-\gamma t_j))^{n_j}, \tag{4}$$

is the probability that transmission occurred from infected insects to bovine j at t_j days post

initial feed. The probability, (3), is the probability that an insect became infected (β)

multiplied by the probability that it was still infected when tested ($\exp(-\gamma t_i)$). The probability,

(4), is the probability that at least one insect (out of the n_j feeding) transmitted LSDV, where

the probability that an individual insect will transmit LSDV is the product of the probabilities

that it became infected (β), that it was still infected when refeeding occurred (exp($-\gamma t_j$)) and

that it subsequently transmitted LSDV to the animal during refeeding (*b*). Non-informative

priors were assumed for all three parameters: exponential with mean 100 for γ and uniform

with range (0,1) for *b* and β.

Samples from the joint posterior density were generated using an adaptive Metropolis scheme

(Haario, Saksman & Tamminen, 2001), modified so that the scaling factor was tuned during

burn-in to ensure an acceptance rate of between 20% and 40% for more efficient sampling of

the target distribution (Andrieu & Thoms, 2008). Two chains of 50,000 iterations were run,

with the preceding 10,000 iterations discarded to allow for burn-in of the chain. The chains

were then thinned (taking every fifth sample) to reduce autocorrelation amongst the samples.

The adaptive Metropolis scheme was implemented in Matlab (version R2018a; The

Mathworks Inc.). Convergence of the scheme was assessed visually and by examining the

Gelman-Rubin statistic provided in the coda package (Plummer, Best, Cowles & Vines,

2006) in R (R Core Team, 2018).

2.3   Duration of infectiousness in cattle

The mean infectious period ($1/r_I$) for LSDV in cattle was estimated using data from

experimental infections of cattle (Tuppurainen, Venter & Coetzer, 2005; Babiuk et al., 2008).

Three proxy measures of infectiousness were considered: detection of viral DNA in blood by

PCR; detection of virus in blood by virus isolation in cell culture; and detection of virus (by

transmission electron microscopy) or viral DNA (by PCR) in skin biopsies. For each animal,

the shortest and longest duration of infectiousness was calculated based on the times of the

last negative and first positive sample and the times of the last positive and first subsequent

negative sample for each measure (i.e. accounting for sampling frequency) (Table S2).

Although they are not needed for the calculation of R_0 (see equation (1)), the time to onset of

infectiousness (i.e. the latent period) and the time to onset of clinical signs (i.e. the incubation

period) were also estimated using data from experimental infections of cattle (Chihota,

Rennie, Kitching & Mellor, 2001; Tuppurainen, Venter & Coetzer, 2005; Babiuk et al.,

2008). In this case, the earliest and latest times of onset were determined based on the times

of the last negative and first positive sample for each measure (fever, appearance of skin

lesions, detection of viral DNA in blood and detection of virus in blood) (Table S2).

The infectious period was assumed to follow a gamma distribution with mean duration $1/r_I$

and shape parameter n_I, while the latent (or incubation) period was assumed to follow a

gamma distribution with mean duration $1/r_E$ and shape parameter n_E (see Text S1).

Parameters (i.e. mean and shape parameter) for each proxy measure were estimated using

Bayesian methods, with the likelihood given by,

$$L(r_i, n_i) = \prod_j \int_{t_{\min}^{(j)}}^{t_{\max}^{(j)}} f(\tau)\, \mathrm{d}\tau, \tag{5}$$

where $t_{\min}$ and $t_{\max}$ are the minimum and maximum duration of infectiousness ($i=I$) or earliest

and latest time of onset ($i=E$) for an animal, respectively, and f is the probability density

function for the gamma distribution. Non-informative priors (exponential with mean 100)

were assumed for both the mean and the shape parameter.

Samples from the joint posterior distribution were generated using a random walk

Metropolis-Hastings algorithm (Andrieu & Thoms, 2008), with each parameter updated in

turn. Two chains of 20,000 iterations were run, with the preceding 5,000 iterations discarded

to allow for burn-in of the chain. The chains were then thinned (taking every other sample) to

reduce autocorrelation amongst the samples. The Metropolis-Hastings scheme was

implemented in Matlab (version R2018a; The Mathworks Inc.). Convergence of the scheme

was assessed visually and by examining the Gelman-Rubin statistic provided in the coda

package (Plummer, Best, Cowles & Vines, 2006) in R (R Core Team, 2018).

2.4   Vector life history parameters

Life history parameters, specifically the reciprocal of the time interval between blood meals

(*a*), the vector to host ratio (*m*) and the vector mortality rate (*μ*), were estimated for *S.*

calcitrans, *C. nubeculosus*, *Ae. aegypti*, *An. stephensi* and *Cx. quinquefasciatus*. For each

parameter, plausible ranges were derived from the published literature (Table 1), so they

could be incorporated in the uncertainty and sensitivity analysis.

2.5   Uncertainty and sensitivity analyses

Replicated Latin hypercube sampling (LHS) was used to explore the parameters influencing

the basic reproduction number, R_0 for each insect species and proxy measure of

infectiousness (Blower & Dowlatabadi, 1994; Luz, Codeco, Massad & Struichner, 2003;

Gubbins et al., 2008). Parameters were sampled either from their marginal posterior

distributions (*b*, *β*, *γ* and $1/r_I$) or uniformly from plausible ranges (*a*, *m* and *μ*). The LHS

results were used to compute the median and 95% prediction interval for R_0. The sensitivity

of R_0 to changes in each parameter was assessed by calculating the partial rank correlation

coefficients (PRCCs). The uncertainty and sensitivity analyses were implemented in Matlab

(version R2018a; The Mathworks Inc.).

3   Results

3.1   Mechanical transmission of LSDV by insects

The model, (3), adequately captured the data for all five species of biting insect, with the

observed number of positive insects lying within the 95% credible intervals for the posterior

predictive distribution (Figure 1). Although there is uncertainty associated with all the

parameters related to mechanical transmission, there are still clear differences amongst the

species (Figure 2). The virus inactivation rate is lowest for *Ae. aegypti* (0.08 day^{-1}),

intermediate for *An. stephensi* (0.2 day^{-1}) and *Cx. quinquefasciatus* (0.3 day^{-1}) and highest for

S. calcitrans (1.6 day^{-1}) and *C. nubeculosus* (71.4 day^{-1}) (Table 2). The probability of

transmission from bovine to insect is highest for the three mosquito species, lower for *S.*

calcitrans and lowest for *C. nubeculosus* (Table 2; Figure 2).

There is considerable uncertainty in the estimates for the probability of transmission from

insect to bovine for all five insect species (Figure 2; Table 2). The posterior mode for this

probability was non-zero only for *Ae. aegypti*, though the probability could not be precisely

estimated for this species (Figure 2). The posterior median for the probability was low for *An.*

stephensi (0.03) and *Cx. quinquefasciatus* (0.04), though the upper 95% credible limit for

both species is an order of magnitude higher (Figure 2; Table 2). Finally, the probability of

transmission from insect to bovine could not be reliably estimated for either *S. calcitrans* or

C. nubeculosus (Figure 2; Figure S1). This is, in part, because the number of insects

refeeding when attempting transmission to cattle was not specified for these species (Chihota,

Rennie, Kitching & Mellor, 2003). For *C. nubeculosus*, the posterior distribution for the

probability was identical to the prior distribution (Figure S1), regardless of assumptions made

about the numbers of insects that refed (either 1, 5, 10, 20, 50 or 100). For *S. calcitrans* the

posterior mass for the probability was shifted towards zero if a larger number of insects was

assumed to refeed, though the 95% credible interval remained large (Figure S1). For both

species, estimates for the virus inactivation rate and probability of transmission from bovine

to insect were not affected by the number of insects assumed to refeed (Figure S1).

3.2   Duration of infectiousness in cattle

The mean infectiousness period depends on the proxy measure used to determine when an

animal is infectious (Table 3). It is shortest when based on virus isolation from blood (8.8

days), intermediate when based on detection of viral DNA in blood (16.3 days) and longest

when based on detection of virus or viral DNA in skin lesions (23.1 days). The mean time to

onset was estimated to be 5.4 days for fever, 7.3 days for skin lesions, 5.8 days for viraemia

(PCR) and 8.1 days for viraemia (virus isolation) (Table 3).

3.3   Uncertainty and sensitivity analyses

The basic reproduction number for LSDV was highest for transmission by *S. calcitrans*,

intermediate for *Ae. aegypti*, low for *C. nubeculosus* and *An. stephensi* and lowest *Cx.*

quinquefasciatus (Figure 3; Table 4). Indeed, the median R_0 for *Cx. quinquefasciatus* was

below the threshold for an outbreak to occur (i.e. $R_0=1$). The magnitude of R_0 depended on

the proxy measure of infectiousness used (through its influence on the duration of

infectiousness), with higher values when based on detection of virus or viral DNA in skin

biopsies, intermediate when based on detection of viral DNA in blood and lowest when based

on detection of virus in blood (Figure 3; Table 4).

The sensitivity of R_0 to changes in its constituent parameters differed amongst the insect

species, though some patterns emerge (Figure 4). For all species, the strongest correlations

were with the probability of transmission from insect to bovine (*b*; PRCC>0.66), the biting

rate (*a*; PRCC>0.75, except for *Ae. aegypti*), the vector to host ratio (*m*; PRCC>0.58) and the

virus inactivation rate (γ; PRCC<-0.47, except *An. stephensi* and *Cx. quinquefasciatus*). For

the three mosquito species, there was also correlation with the vector mortality rate (μ;

PRCC<-0.45), but this was not the case for the other dipteran species (PRCC$\approx$0). The

probability of transmission from bovine to insect (β) and the mean infectious period ($1/r_I$)

were also positively correlated with R_0 (PRCC=0.3-0.4 and PRCC=0.2-0.5, respectively).

These patterns are independent of the proxy measure of infectiousness used (Figure 4).

4   Discussion

The results of the uncertainty analysis for the basic reproduction number (Figure 3; Table 4)

suggest that *S. calcitrans* is likely to be the most efficient vector of LSDV (out of the five

species considered in the present study), with *Ae. aegypti* also an efficient vector of the virus.

By contrast, *C. nubeculosus*, *An. stephensi* and *Cx. quinquefasciatus* are likely to be

inefficient as vectors of LSDV.

These conclusions would potentially be different if the risk posed by each species was

assessed on the outcome of the transmission experiments alone (Table 2). While the

experimental outcome suggests *Ae. aegypti* might be an efficient vector, it could be

concluded that *S. calcitrans* is likely to be a poor vector given the very short duration of

infection (<1 day) and the comparatively low probabilities of transmission from bovine to

insect or insect to bovine for this species. By contrast, the results of the transmission

experiments suggest that *Cx. quinquefasciatus* could be a more efficient vector than it is

based on the estimates for R_0, because of the relatively long duration of infection and the high

probability of transmission from bovine to insect. This highlights the importance of

considering all factors associated with transmission, including vector life history, when

assessing the risk of transmission for each species. For both *S. calcitrans* and *Cx.*

quinquefasciatus the main reason for the difference in conclusions between transmission

experiments and R_0 is the time interval between blood meals, which is shorter than the

duration of infection for *S. calcitrans*, but longer than the duration of infection for *Cx.*

quinquefasciatus (Tables 1 & 2).

A certain amount of care should be taken in interpreting these results, however, because there

is substantial uncertainty associated with the estimates of R_0 for all five species, but

especially for *S. calcitrans* and *C. nubeculosus*. One of the major sources of uncertainty is the

probability of transmission from insect to bovine (Figure 2; Table 2). Although transmission

to naïve cattle was attempted (Chihota, Rennie, Kitching & Mellor, 2003), it was done at

times when *S. calcitrans* and *C. nubeculosus* were unlikely to still be infectious (see Figure 1;

refeeding was at 1-3 days post infection for *S. calcitrans* and 3-5 days post infection for *C.*

nubeculosus). Consequently, there is little information in the data to estimate this parameter,

which is reflected in the posterior distributions being almost the same as the prior

distributions, especially in the case of *C. nubeculosus* (Figure 2; Figure S1). Yet the

probability of transmission from insect to bovine was identified as one of the parameters to

which the basic reproduction number is most sensitive for all five vector species considered

(Figure 4).

The influence of the virus inactivation rate and mortality rate on R_0 differed amongst the

species (Figure 4). The duration of infection (i.e. the reciprocal of the virus inactivation rate)

for *C. nubeculosus* or *S. calcitrans* is much shorter than the lifespan of an insect and,

consequently, only the virus inactivation rate is important for determining the magnitude of

R_0. By contrast, the duration of infection and insect lifespan are comparable for *Ae. aegypti*,

An. stephensi and *Cx. quinquefasciatus*, so that both of these parameters have an influence on the basic reproduction number.

As with all uncertainty and sensitivity analyses, the conclusions drawn from them are valid only over the parameter ranges considered. For parameters related to mechanical transmission, these ranges were estimated from the available data on the outcome of transmission experiments (Chihota, Rennie, Kitching & Mellor, 2001, 2003) and so represent the best current estimates. Given the uncertainty in these parameters (which reflects the small numbers of animals and insects tested at each time point in the studies; Table S1), this suggests that a focus of future experimental work should be to more precisely measure parameters related to mechanical transmission and, in particular, the probability of transmission from insect to bovine and the virus inactivation rate.

All three vector life history parameters were identified as influencing the magnitude of R_0 (Figure 4). Plausible ranges for these parameters were drawn from the published literature (Table 1). Wherever possible, ranges were derived using data relating to the species themselves. If suitable data were not available, however, ranges were derived using data for related species. In addition, the vector life parameters were assumed to be constant, but they will depend on environmental factors, including temperature and rainfall. Consequently, the basic reproduction number, and so the risk of transmission, is unlikely to be constant over space or time, but will vary both geographically and seasonally.

This study considered the potential of five species of biting insects to transmit LSDV. This was primarily because their ability to transmit LSDV had been assessed experimentally for each of the species. However, the present analysis did not consider the effect of vector

feeding preferences on their potential role in transmission. The stable fly, *S. calcitrans*, is one

of the most damaging arthropod pests of cattle worldwide, both through its impact on

production (Taylor, Moon & Mark, 2012) and its ability to transmit a number of pathogens

(Baldacchino et al., 2013). Consequently, this species' feeding preferences are unlikely to

affect its efficiency as a vector of LSDV. Similarly, *C. nubeculosus* and other *Culicoides*

species for which *C. nubeculosus* can be considered a model are livestock-associated (Lassen,

Nielsen & Kristensen, 2012; Purse, Carpenter, Venter, Bellis & Mullens, 2015), suggesting

host preference will not reduce their efficiency as a vector of LSDV. By contrast, *Ae. aegypti*

feeds primarily on humans (Reiter, 2010), potentially limiting its role in the transmission of

LSDV, despite its apparent efficiency as a vector. However, there are other *Aedes* species (for

example, *Ae. cinereus* or *Ae. vexans*), which display a preference for feeding on cattle

(Hayes, Tempelis, Hess & Reeves, 1973; Magnarelli, 1977) and for which *Ae. aegypti* could

act as a model vector species. Many *Anopheles* species, including those in the Maculipennis

subgroup, which is of most relevance to Europe (Sinka et al., 2010; ECDC, 2019), are

described as zoophilic (Sinka et al., 2010). Accordingly, host preference is unlikely to limit

their ability to transmit LSDV. Finally, host preference in *Cx. quinquefasciatus* varies

according to environmental conditions and geographical area, though it does feed on

mammals (Hayes, Tempelis, Hess & Reeves, 1973; Molaei, Huang & Andreadis, 2012;

Janssen et al., 2015). However, any preference for non-bovine hosts in this species (or other

Culex species for which it is a suitable model) will only further reduce its importance in the

transmission of LSDV.

The magnitude of the basic reproduction number depends on the proxy measure used for

infectiousness (Figure 3), though the conclusions of the sensitivity analysis do not (Figure 4).

Viral titres in blood can be low and detection of virus intermittent (Tuppurainen, Venter &

Coetzer, 2005; Babiuk et al., 2008), suggesting this may not be the best proxy measure. By

contrast, skin nodules have high titres of virus (Babiuk et al., 2008) and insects which feed on

nodules become infected (Chihota, Rennie, Kitching & Mellor, 2001, 2003), indicating that

this may be a reasonable proxy measure. This could be tested experimentally by feeding

insects on areas with and without lesions of clinically-affected cattle and on infected, but

subclinical cattle and relating viral titres in skin and blood to the probability of insects

becoming infected.

The expression for the basic reproduction number presented in equation (1) assumes there is

negligible disease-associated mortality in cattle (see Text S1). Reported mortality for LSD

outbreaks in the Balkans was very low, with the median within-herd mortality being zero in

all affected countries (EFSA, 2017). In endemic settings, mortality is generally low (1-3%)

(Tuppurainen & Oura, 2012; Molla, Frankena & de Jong, 2017), but it may sometimes reach

40% (Tuppurainen & Oura, 2012). If disease-associated mortality (up to 40%) is included in

the computation of R_0, it does not greatly affect the conclusions of the present study regarding

either the magnitude of R_0 for each vector species or the sensitivity of R_0 to changes in its

constituent parameters (see Text S2).

Two previous studies have estimated the basic reproduction number for LSDV using

outbreak data. In the first, Magori-Cohen et al. (2012)[1] estimated R_0 to be around 4 for an

outbreak in a large dairy herd in Israel in 2006, where *S. calcitrans* was likely to be the

principal vector species (Kahana-Sutin, Klement, Lensky & Gottlieb, 2017). Because of the

control measures implemented on the farm (removal of cattle showing generalised disease on

the day of detection), this implicitly assumed an infectious period of around one day. If a

---

[1] Because of differences in model formulation, the value of R_0 presented by Magori-Cohen et al. (2012) is the square of R_0 in the current paper. Here the square root transformed values are presented.

mean infectious period of 23 days is used instead (i.e. the maximum estimated in the present

study; Table 3), the corresponding value for R_0 is around 19, which is similar to the median

estimate for transmission by *S. calcitrans* obtained in the present study (Figure 3; Table 4).

By contrast, the second study estimated R_0 to be around 1.1 for outbreaks in eight cattle herds

in Ethiopia in 2014-2015, using viraemia as a proxy for infectiousness (Molla, Frankena & de

Jong, 2017). This is substantially lower than the ranges estimated for either *S. calcitrans* or

Ae. aegypti, but is consistent with those for *C. nubeculosus*, *An. stephensi* or *Cx.*

quinquefasciatus (Figure 3; Table 4). However, the vector species involved in transmission of

LSDV in Ethiopia are not known (Molla, de Jong & Frankena, 2017).

Funding

This work was funded by the Biotechnology and Biological Sciences Research Council

(BBSRC) [grant code: BB/R002606/1]. SG was also partially supported by BBSRC-funded

Institute Strategic Programme grants BBS/E/I/00007033, BBS/E/I/00007036 and

BBS/E/I/00007037.

Conflict of interest

None to declare.

Acknowledgements

The author is grateful to Pip Beard, Beatriz Sanz-Bernardo and Christopher Sanders (all at

The Pirbright Institute) for helpful comments on the manuscript.

References

Alkhamis, M.A. & VanderWaal, K. (2016). Spatial and temporal epidemiology of lumpy skin disease in the Middle East, 2012-2015. *Frontiers in Veterinary Science*, *3*, 19. https://doi.org/10.3389/fvets.2016.00019

Andrieu, C. & Thoms, J. (2008). A tutorial on adaptive MCMC. *Statistics and Computing*, *18*, 343-373. https://doi.org/10.1007/s11222-008-9110-y

Babiuk, S., Bowden, T.R., Parkyn. G., Dalman, B., Manning, L., Neufeld, J., Embury-Hyatt, C., Copps, J. & Boyle, D.B. (2008). Quantification of lumpy skin disease virus following experimental infection in cattle. *Transboundary and Emerging Diseases*, *55*, 299-307. https://doi.org/10.1111/j.1865-1682.2008.01024.x

Baldacchino, F., Muenwom, V., Desquesnes, M., Desoli, F., Charoenviriyaphap, T. & Duvallet, G. (2013). Transmission of pathogens by *Stomoxys* flies (Diptera: Muscidae): a review. *Parasite*, *20*, 26. https://doi.org/10.1051/parasite/2013026

Birley, M.H. & Boorman, J.P.T. (1982). Estimating the survival and biting rates of hematophagous insects, with particular reference to the *Culicoides obsoletus* group (Diptera, Ceratopogonidae) in southern England. *Journal of Animal Ecology*, *51*, 135–148. https://doi.org/10.2307/4315

Blower, S.M. & Dowlatabadi, H. (1994). Sensitivity and uncertainty analysis of complex models of disease transmission: an HIV model as an example. *International Statistical Review*, *62*, 229-243. https://doi.org/10.2307/1403510

Brady, O.J., Johansson, M.A., Guerra, C.A., Bhatt, S., Golding, N., Pigott, D.M., Delatte, H., Grech, M.G., Leisnham, P.T., Maciel-de-Freitas, R., Styer, L.M., Smith, D.L., Scott, T.W., Gething, P.W. & Hay, S.I. (2013). Modelling adult *Aedes aegypti* and *Aedes albopictus* survival at different temperatures in laboratory and field settings. *Parasites & Vectors*, *6*, 351. https://doi.org/10.1186/1756-3305-6-351

Braverman, Y., Linley, J. R., Marcus, R. & Frish, K. (1985). Seasonal survival and

expectation of infectious life of *Culicoides* spp. (Diptera: Ceratopogonidae) in Israel, with

implications for bluetongue virus transmission and a comparison of the parous rate in *C.*

imicola from Israel and Zimbabwe. *Journal of Medical Entomology*, *22*, 476-484.

https://doi.org/10.1093/jmedent/22.5.476

Carn, V.M. & Kitching, R.P. (1995). An investigation of possible routes of transmission of

lumpy skin disease virus (Neethling). *Epidemiology and Infection*, *114*, 219-226.

https://doi.org/10.1017/S0950268800052067

Charlwood, J.D., Nenhep, S., Sovannaroth, S., Morgan, J.C., Hemingway, J., Chitnis, N. &

Briët, J.T. (2016). 'Nature or nurture': survival rate, oviposition interval and possible

gonotrophic discordance among South East Asian anophelines. *Malaria Journal*, *15*, 356.

https://doi.org/10.1186/s12936-016-1389-0

Chihota, C.M., Rennie, L.F., Kitching, R.P. & Mellor, P.S. (2001). Mechanical transmission

of lumpy skin disease virus by *Aedes aegypti* (Diptera: Culicidae). *Epidemiology and*

Infection, *126*, 317-321. https://doi.org/10.1017/S0950268801005179

Chihota, C.M., Rennie, L.F., Kitching, R.P. & Mellor, P.S. (2003). Attempted mechanical

transmission of lumpy skin disease virus by biting insects. *Medical and Veterinary*

Entomology, *17*, 294-300. https://doi.org/10.1046/j.1365-2915.2003.00445.x

Diekmann, O. & Heesterbeek, J.A.P. (2000). *Mathematical epidemiology of infectious*

diseases: model building, analysis and interpretation. Chichester: John Wiley & Sons.

ECDC (2019). European Centre for Disease Prevention and Control and European Food

Safety Authority: Mosquito maps. https://ecdc.europa.eu/en/disease-vectors/surveillance-and-

disease-data/mosquito-maps

EFSA (2017). Scientific report on lumpy skin disease: I. Data collection and analysis. *The*

EFSA Journal, *15*, 4773. https://doi.org/10.2903/j.efsa.2017.4773

EFSA Panel on Animal Health and Welfare (2015). Scientific opinion on lumpy skin disease. *The EFSA Journal*, *13*, 3986. https://doi.org/10.2903/j.efsa.2015.3986

Gachohi, J.M., Njenga, M.K., Kitala, P. & Bett, B. (2016). Modelling vaccinations strategies against Rift Valley fever virus in livestock in Kenya. *PLoS Neglected Tropical Diseases*, *10*, e0005049. https://doi.org/10.1371/journal.pntd.0005049

Gad, A.M., Feinsod, F.M., Soliman, B.A. & El Said, S. (1989). Survival estimates for adult *Culex pipiens* in the Nile Delta. *Acta Tropica*, *46*, 173-179. https://doi.org/10.1016/0001-706X(89)90034-X

Gerry, A.C., Mullens, B.A., MacLachlan, N.J. & Mecham, O.J. (2001). Seasonal transmission of bluetongue virus by *Culicoides sonorensis* (Diptera: Ceratopogonidae) at a southern California dairy and evaluation of vectorial capacity as a predictor of bluetongue virus transmission. *Journal of Medical Entomology*, *38*, 197-209. https://doi.org/10.1603/0022-2585-38.2.197

Griffith, J.S.R. & Turner, G.D. (1996). Culturing *Culex quinquefasciatus* mosquitoes with a blood substitute diet for the females. *Medical and Veterinary Entomology*, *10*, 265–268. https://doi.org/10.1111/j.1365-2915.1996.tb00741.x

Gubbins, S., Carpenter, S., Baylis, M., Wood, J.L.N. & Mellor, P.S. (2008). Assessing the risk of bluetongue to UK livestock: uncertainty and sensitivity analysis of a temperature-dependent model for the basic reproduction number. *Journal of the Royal Society Interface*, *5*, 363-371. https://doi.org/10.1098/rsif.2007.1110

Haario, H., Saksman, E. & Tamminen, J. (2001). An adaptive Metropolis algorithm. *Bernoulli*, *7*, 223-242. https://projecteuclid.org/euclid.bj/1080222083

Hayes, R.O., Tempelis, C.H., Hess, A.D. & Reeves, W.C. (1973). Mosquito preference studies in Hale County, Texas. *American Journal of Tropical Medicine and Hygiene*, *22*, 270-277. https://doi.org/10.4269/ajtmh.1973.22.270

Jacquiet, P., Rouet, D., Bouhsira, E., Salem, A., Lienard, E. & Franc, M. (2014). Population dynamics of *Stomoxys calcitrans* (L.) (Diptera: Muscidae) in southwestern France. *Revue de Médicine Vétérinaire*, *165*, 267-271. https://www.revmedvet.com/artdes-us.php?id=16015

Janssen, N., Fernandez-Salas, I., Díaz González, E.E., Gaytan-Burns, A., Medina-de la Garza, C.E., Sanchez-Casas, R.M., Börstler, J., Cadar, D., Schmidt-Chanasit, J. & Jöst, H. (2015). Mammalophilic feeding behaviour of *Culex quinquefasciatus* mosquitoes collected in the cities of Chetumal and Cancun, Yucatán Peninsula, Mexico. *Tropical Medicine and International Health*, *20*, 1488-1491. https://doi.org/10.1111/tmi.12587

Jones, C.E., Lounibos, P., Marra, P.P. & Kilpatrick, A.M. (2012). Rainfall influences survival of *Culex pipiens* (Diptera: Culicidae) in a residential neighbourhood in the mid-Atlantic United States. *Journal of Medical Entomology*, *49*, 467-473. https://doi.org/10.1603/ME11191

Kahana-Sutin, E., Klement, E., Lensky, I. & Gottlieb, Y. (2017). High relative abundance of the stable fly *Stomoxys calcitrans* is associated with lumpy skin disease outbreaks in Israeli dairy farms. *Medical and Veterinary Entomology*, *31*, 150-160. https://doi.org/10.1111/mve.12217

Killeen, G.F., McKenzie, F.E., Foy, B.D., Shieffelin, C., Billingsley, P.F. & Beier, J.C. (2000). A simplified model for predicting malaria entomologic inoculation rates based on entomologic and parasitologic parameters relevant to control. *American Journal of Tropical Medicine and Hygiene*, *62*, 535-544. https://doi.org/10.4269/ajtmh.2000.62.535

Kiszewski, A., Mellinger, A., Spielman, A., Malaney, P., Ehrlich Sachs, S. & Sachs, J. (2004). A global index representing the stability of malaria transmission. *American Journal of Tropical Medicine and Hygiene*, *70*, 486-498. https://doi.org/10.4269/ajtmh.2004.70.486

Kitching, R.P. & Mellor, P.S. (1986). Insect transmission of capripox virus. *Research in Veterinary Science*, *40*, 255-258. https://doi.org/10.1016/S0034-5288(18)30523-X

Kunz, S.E. & Monty, J. (1976). Biology and ecology of *Stomoxys nigra* Macquart and

Stomoxys calcitrans (L.) (Diptera, Muscidae) in Mauritius. *Bulletin of Entomological*

Research, *66*, 745-755. https://doi.org/10.1017/S0007485300010798

Lassen, S.B., Nielson, S.A. & Kristensen, M. (2012). Identity and diversity of blood meal

hosts of biting midges (Diptera: Ceratopogonidae: *Culicoides* Latreille) in Denmark.

Parasites & Vectors, *5*, 143. https://doi.org/10.1186/1756-3305-5-143

Liu-Helmersson, J., Stenlund, H., Wilder-Smith, A. & Rocklöv, J. (2014). Vectorial capacity

of *Aedes aegypti*: effects of temperature and implications for global dengue epidemic

potential. *PLoS ONE*, *9*, e89783. https://doi.org/10.1371/journal.pone.0089783

Luz, P.M., Codeco, C.T., Massad, E. & Struichner, C.J. (2003). Uncertainties regarding

dengue modelling in Rio de Janeiro, Brazil. *Memorias do Instituto Oswaldo Cruz*, *98*, 871-

878. https://doi.org/10.1590/S0074-02762003000700002

Magnarelli, L.A. (1977). Host feeding patterns of Connecticut mosquitoes (Diptera:

Culicidae). *American Journal of Tropical Medicine and Hygiene*, *26*, 547-552.

https://doi.org/10.4269/ajtmh.1977.26.547

Magori-Cohen, R., Louzoun, Y., Herziger, Y., Oron, E., Arazi, A., Tuppurainen, E., Shpigel,

N.Y. & Klement, E. (2012). Mathematical modelling and evaluation of the different routes of

transmission of lumpy skin disease virus. *Veterinary Research*, *43*, 1.

https://doi.org/10.1186/1297-9716-43-1

Mellor, P.S., Kitching, R.P. & Wilkinson, P.J. (1987). Mechanical transmission of capripox

virus and African swine fever virus by *Stomoxys calcitrans*. *Research in Veterinary Science*,

43, 109-112. https://doi.org/10.1016/S0034-5288(18)30753-7

Mercier, A., Arsevska, E., Bournez, L., Bronner, A., Calavas, D., Cauchard, J., Falala, S.,

Caufour, P., Tisseuil, C., Lefrançois, T. & Lancelot, R. (2018). Spread rate of lumpy skin

disease in the Balkans, 2015-2016. *Transboundary and Emerging Diseases*, *65*, 240-243.

https://doi.org/10.1111/tbed.12624

Molaei, G., Huang, S. & Andreadis, T.G. (2012). Vector-host interactions of *Culex pipiens*

complex in northeastern and southwestern USA. *Journal of the American Mosquito Control*

Association, *28(Suppl4)*, 127-136. https://doi.org/10.2987/8756-971X-28.4s.127

Molla, W., de Jong, M.C.M. & Frankena, K. (2017). Temporal and spatial distribution of

lumpy skin disease outbreaks in Ethiopia in the period 2000 to 2015. *BMC Veterinary*

Research, *13*, 310. https://doi.org/10.1186/s12917-017-1247-5

Molla, W., Frankena, K. & de Jong, M.C.M. (2017). Transmission dynamics of lumpy skin

disease in Ethiopia. *Epidemiology and Infection*, *145*, 2856-2863.

https://doi.org/10.1017/S0950268817001637

Molla, W., de Jong, M.C.M., Gari, G. & Frankena, K. (2017). Economic impact of lumpy

skin disease and cost effectiveness of vaccination for the control of outbreaks in Ethiopia.

Preventive Veterinary Medicine, *147*, 100-107.

https://doi.org/10.1016/j.prevetmed.2017.09.003

Mullens, B.A., Gerry, A.C., Lysyk, T.J. & Schmidtmann, E.T. (2004). Environmental effects

on vector competence and virogenesis of bluetongue virus in *Culicoides*: interpreting

laboratory data in a field context. *Veterinaria Italiana*, *40*, 160-166.

http://www.izs.it/vet_italiana/2004/40_3/32.pdf

Plummer, M., Best, N., Cowles, K. & Vines, K. (2006). CODA: Convergence Diagnosis and

Output Analysis for MCMC. *R News*, *6*, 7-11. https://cran.r-

project.org/doc/Rnews/Rnews_2006-1.pdf

Purse, B.V., Carpenter, S., Venter, G.J., Bellis, G. & Mullens, B.A. (2015). Bionomics of

temperate and tropical *Culicoides* midges: knowledge gaps and consequences for

transmission of Culicoides-borne viruses. *Annual Review of Entomology*, *60*, 373-392. https://doi.org/10.1146/annurev-ento-010814-020614

R Core Team 2018 R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org/

Reisen, W.K., Fang, Y. & Martinez, V.M. (2006). Effects of temperature on the transmission of West Nile virus by *Culex tarsalis* (Diptera: Culicidae). *Journal of Medical Entomology*, *43*, 309-317. https://doi.org/10.1093/jmedent/43.2.309

Reiter, P. (2010). Yellow fever and dengue: a threat to Europe? *EuroSurveillance*, *15*, 19509. https://www.eurosurveillance.org/content/10.2807/ese.15.10.19509-en

Ryan, S.J., Lippi, C.A., Boersch-Supan, P.H., Heydari, N., Silva, M., Adrian, J., Noblecilla, L.F., Ayala, E.B., Encalada, M.D., Larsen, D.A., Krisher, J.T., Krisher, L., Fregosi, L. & Stewart-Ibarra, A.M. (2017). Quantifying seasonal and diel variation in Anopheline and Culex human biting rates in Southern Ecuador. *Malaria Journal*, *16*, 479. https://doi.org/10.1186/s12936-017-2121-4

Salem, A., Franc, M., Jacquiet, P., Bouhsira, E. & Lienard, E. (2012). Feeding and breeding aspects of *Stomoxys calcitrans* (Diptera: Muscidae) under laboratory conditions. *Parasite*, *19*, 309-317. https://doi.org/10.1051/parasite/2012194309

Scott, T.W., Amerasinghe, P.H., Morrison, A.C., Lorenz, L.H., Clark, G.G., Strickman, D., Kittayapog, P. & Edman, J.D. (2000). Longitudinal studies of *Aedes aegypti* (Diptera: Culicidae) in Thailand and Puerto Rico: blood feeding frequency. *Journal of Medical Entomology*, *37*, 89-101. https://doi.org/10.1603/0022-2585-37.1.77

Sinka, M.E., Bangs, M.J., Manguin, S., Coetzee, M., Mbogo, C.M., Hemingway, J., Patil, A.P., Temperley, W.H., Gething, P.W., Kabaria, C.W., Okara, R.M., Van Boekel, T., Godfray, H.C.J., Harbach, R.E. & Hay, S.I. (2010). The dominant Anopheles vectors of

human malaria in Africa, Europe and the Middle East: occurrence data, distribution maps and

bionomic précis. *Parasites & Vectors*, *3*, 117. https://doi.org/10.1186/1756-3305-3-117

Skovgård, H. & Nachman, G. (2017). Modeling the temperature- and age-dependent survival,

development, and oviposition rates of stable flies (*Stomoxys calcitrans*) (Diptera: Muscidae).

Environmental Entomology, *46*, 1130-1142. https://doi.org/10.1093/ee/nvx118

Tasioudi, K.E., Antoniou, S.E., Iliadou, P., Sachpatzidis, A., Plevraki, E., Agianniotaki, E.I.,

Fouki, C., Mangana-Vougiouka, O., Chondrokouki, E. & Dile, C. (2016). Emergence of

lumpy skin disease in Greece, 2015. *Transboundary and Emerging Diseases*, *63*, 260-265.

https://doi.org/10.1111/tbed.12497

Taylor, D.B., Moon, R.D. & Mark, D.R. (2012). Economic impact of stable flies (Diptera:

Muscidae) on dairy and beef cattle production. *Journal of Medical Entomology*, *49*, 198-209.

https://doi.org/10.1603/ME10050

Tuppurainen, E.S.M., Venter, E.H. & Coetzer, J.A.W. (2005). The detection of lumpy skin

disease virus in samples of experimentally infected cattle using different diagnostic

techniques.   *Onderstepoort   Journal   of   Veterinary   Research*,   *72*,   153-164.

https://doi.org/10.4102/ojvr.v72i2.213

Tuppurainen, E.S. & Oura, C.A.L. (2012). Lumpy skin disease: an emerging threat to Europe,

the Middle East and Africa. *Transboundary and Emerging Diseases*, *59*, 40-48.

https://doi.org/10.1111/j.1865-1682.2011.01242.x

Tuppurainen, E.S., Lubinga, J.C., Stoltsz, W.H., Troskie, M., Carpenter, S.T., Venter, E.H. &

Oura, C.A. (2013). Mechanical transmission of lumpy skin disease virus by *Rhipicephalus*

appendiculatus   male   ticks.   *Epidemiology   and   Infection*,   *141*,   425–430.

https://doi.org/10.1017/S0950268812000805

Weiss, K.E. (1968). Lumpy skin disease. In: *Virology Monographs 3* (pp. 111-131). Vienna:

Springer-Verlag. https://doi.org/10.1007/978-3-662-39771-8_3

Table 1. Life history parameters for five putative insect vectors of lumpy skin disease virus.

parameter	symbol	range	comments	references
Stomoxys calcitrans				
biting rate (day^{-1})	a	0.33-6	time interval between blood meals (1/a) of 4-72 hours	Salem et al. (2013); Baldacchino et al. (2013)
vector to host ratio	m	30-145	-	Kunz & Monty (1976); Jacquiet et al. (2014)
mortality rate (day^{-1})	μ	0.04-0.11	median lifespan (1/μ) of 9-26 days	Salem et al. (2013); Skovgård & Nachman (2017)
Culicoides nubeculosus				
biting rate (day^{-1})	a	0-0.4	based on duration of gonotrophic cycle	Birley & Boorman (1982); Braverman et al. (1985); Mullens et al. (2004)
vector to host ratio	m	0-5000	based on a maximum host biting rate ($m \times a$) of 2500 bites per host per day	Gerry et al. (2001)
mortality rate (day^{-1})	μ	0.1-0.5	based on proportion of females surviving gonotrophic cycle	Birley & Boorman (1982); Braverman et al. (1985)
Aedes aegypti				
biting rate (day^{-1})	a	0.18-0.23	-	Scott et al. (2000); Liu-Helmersson et al. (2014)
vector to host ratio	m	0-80	modelled ratio of *Aedes* to cattle	Gachohi et al. (2016)
mortality rate (day^{-1})	μ	0.07-0.2	lifespan of 5-15 days estimated using a modelling analysis of 50 mark-release-recapture field studies	Brady et al. (2013)
Anopheles stephensi				
biting rate (day^{-1})	a	0.25-0.5	time interval between blood meals (1/a) of 2-4 days	Killeen et al. (2000); Charlwood et al. (2016)
vector to host ratio	m	0.6-60	based on a host biting rate ($m \times a$) of around 2 to 120 bites per host per day	Killeen et al. (2000); Ryan et al. (2017)
mortality rate (day^{-1})	μ	0.02-0.38	daily survival probability (exp(-μ)) of 0.68-0.98	Kiszewski et al. (2004)

parameter	symbol	range	comments	references
Culex quinquefasciatus				
biting rate (day^{-1})	*a*	0.08-0.25	time interval between blood meals (1/*a*) of 4-12 days	Griffith & Turner (1996); Reisen et al. (2006)
vector to host ratio	*m*	0-80	modelled ratio of *Culex* to cattle	Gachohi et al. (2016)
mortality rate (day^{-1})	*μ*	0.07-0.84	-	Gad et al. (1989); Jones et al. (2012)

Table 2. Parameters for mechanical transmission of lumpy skin disease virus for five species of biting insect.

species	virus inactivation rate (γ; day^{-1})		mean duration of infection ($1/\gamma$; days)		probability of transmission from bovine to insect (β)		probability of transmission from insect to bovine (b)	
	estimate†	95% CI‡	estimate	95% CI	estimate	95% CI	estimate	95% CI
S. calcitrans§	1.71	(0.80, 3.34)	0.58	(0.30, 1.26)	0.46	(0.23, 0.71)	0.07	(0.002, 0.64)
C. nubeculosus§	71.16	(4.65, 390.0)	0.01	(0.003, 0.22)	0.27	(0.09, 0.54)	0.50	(0.03, 0.98)
Ae. aegypti	0.09	(0.01, 0.19)	11.23	(5.19, 79.11)	0.90	(0.64, 0.99)	0.48	(0.04, 0.97)
An. stephensi	0.24	(0.10, 0.42)	4.08	(2.38, 9.64)	0.61	(0.36, 0.83)	0.03	(0.001, 0.16)
Cx. quinquefasciatus	0.34	(0.17, 0.55)	2.96	(1.81, 5.74)	0.72	(0.47, 0.91)	0.04	(0.002, 0.28)

† posterior median

‡ CI: credible interval

§ estimates are shown for the analysis assuming 100 insects refed when attempting transmission to cattle

Table 3. Parameters for the duration of latent, incubation and infectious periods for lumpy skin disease virus in cattle.

	shape parameter		mean (days)	
	estimate†	95% CI‡	estimate†	95% CI‡
time to onset (days)				
fever	151.5	(44.1, 442.6)	5.4	(5.1, 5.8)
skin lesions	25.4	(4.8, 82.7)	7.3	(5.9, 9.7)
viraemia (PCR)	3.4	(1.6, 6.4)	5.8	(4.4, 7.8)
viraemia (VI)	5.7	(1.8, 13.6)	8.1	(5.9, 12.1)
duration of infectiousness (days)				
skin biopsies	10.4	(2.1, 36.1)	23.1	(16.4, 36.2)
blood (PCR)	3.1	(1.1, 7.1)	16.3	(11.4, 25.4)
blood (virus isolation)	2.6	(0.7, 6.7)	8.8	(5.5, 17.8)

† posterior median

‡ CI: credible interval

Table 4. Basic reproduction number† (R_0) for lumpy skin disease virus (LSDV) for five species of biting insect.

species	proxy measure of infectiousness		
	skin biopsies	blood (PCR)	blood (virus isolation)
S. calcitrans‡	15.5 (1.4, 81.9)	13.1 (1.2, 70.8)	9.7 (0.9, 51.7)
C. nubeculosus‡	1.8 (0.06, 13.5)	1.5 (0.05, 11.3)	1.1 (0.04, 8.8)
Ae. aegypti	7.4 (1.3, 17.6)	6.3 (1.1, 14.5)	4.6 (0.8, 11.2)
An. stephensi	1.6 (0.2, 6.0)	1.4 (0.2, 5.0)	1.0 (0.1, 3.9)
Cx. quinquefasciatus	0.8 (0.09, 3.5)	0.7 (0.08, 2.8)	0.5 (0.06, 2.2)

† median (95% prediction interval) based on replicated Latin hypercube sampling

‡ estimates are shown for the analysis assuming 100 insects refed when attempting transmission to cattle

Figure 1. Proportion of biting insects positive for viral DNA after feeding on a bovine infected with lumpy skin disease virus. Results are shown for five species of biting insect: (*a*) *Stomoxys calcitrans*; (*b*) *Culicoides nubeculosus*; (*c*) *Aedes aegypti*; (*d*) *Anopheles stephensi*; and (*e*) *Culex quinquefasciatus*. Each plot shows the observed proportion of infected insects (open squares), the expected proportion of infected insects (black line: posterior median; shading: percentiles of the posterior distribution in 5% bands from 5% to 95%) and the posterior predictive distribution for the proportion of infected insects (grey circles: median; grey error bars: 95% prediction intervals).

Figure 2. Parameters for mechanical transmission of lumpy skin disease virus by five species of biting insect: (*a*) virus inactivation rate (γ; day^{-1}); (*b*) probability of transmission from bovine to insect (β; Pr(B to I)); and (*c*) probability of transmission from insect to bovine (*b*; Pr(I to B)). Each plot shows the median (black circle), 25th and 75th percentiles (black line) and density (shape) for the marginal posterior distribution. Results shown for *S. calcitrans* and *C. nubeculosus* assume 100 insects refed when attempting transmission to cattle.

Figure 3. Basic reproduction number (R_0) for lumpy skin disease virus (LSDV) when transmitted by *Stomoxys calcitrans*, *Culicoides nubeculosus*, *Aedes aegypti*, *Anopheles stephensi* or *Culex quinquefasciatus*. The estimated values for R_0 are shown for three proxy measures of infectiousness: detection of virus or viral DNA in skin biopsies (light grey); detection of viral DNA in blood (mid grey); or detection of virus in blood (dark grey). Box and whisker plots show the median (black horizontal line), interquartile range (coloured box) and 95% range (whiskers) based on replicated Latin hypercube sampling (100 replicates with the range for each parameter subdivided into 100 steps). Results shown for *S. calcitrans* and

C. nubeculosus used the posterior distributions assuming 100 insects refed when attempting transmission to cattle.

Figure 4. Sensitivity analysis of the basic reproduction number (R_0) for lumpy skin disease virus (LSDV) when transmitted by: (*a*) *Stomoxys calcitrans*; (*b*) *Culicoides nubeculosus*; (*c*) *Aedes aegypti*; (*d*) *Anopheles stephensi*; or (*e*) *Culex quinquefasciatus*. Plots show the partial rank correlation coefficients (PRCC) for each parameter when the estimated values for R_0 are calculated using three proxy measures of infectiousness: detection of virus or viral DNA in skin biopsies (light grey); detection of viral DNA in blood (mid grey); or detection of virus in blood (dark grey). Box and whisker plots show the median (black horizontal line), interquartile range (coloured box) and 95% range (whiskers) for the PRCCs based on replicated Latin hypercube sampling (100 replicates with the range for each parameter subdivided into 100 steps). Results shown for *S. calcitrans* and *C. nubeculosus* used the posterior distributions assuming 100 insects refed when attempting transmission to cattle.