Basal Contamination of Sequencing: Lessons from the GTEx dataset

Tim O. Nieuwenhuis[1,2], Stephanie Yang[2], Rohan X. Verma[1], Vamsee Pillalamarri[2], Dan E. Arking[2], Avi Z. Rosenberg[1], Matthew N. McCall[3], Marc K. Halushka[1*]

[1] Department of Pathology, Johns Hopkins University SOM, Baltimore, MD, USA

[2] McKusick-Nathans Institute, Department of Genetic Medicine, Johns Hopkins University SOM, Baltimore, MD, USA

[3] Department of Biostatistics and Computational Biology, University of Rochester Medical Center, Rochester, NY, USA

Email Addresses:

tnieuwe1@jhmi.edu

syang93@jhmi.edu

rverma6@jhmi.edu

vpillal1@jhmi.edu

arking@jhmi.edu

arosen34@jhmi.edu

matthew_mccall@urmc.rochester.edu

* Correspondence and address for reprints to:
Marc K. Halushka, M.D., Ph.D.
Johns Hopkins University School of Medicine
Ross Bldg. Rm 632B
720 Rutland Avenue
Baltimore, MD 21205
410-614-8138 (ph)
410-502-5862 (fax)
mhalush1@jhmi.edu

Abstract

One of the challenges of next generation sequencing (NGS) is read contamination. We used the Genotype-Tissue Expression (GTEx) project, a large, diverse, and robustly generated dataset, to understand the factors that contribute to contamination. We obtained GTEx datasets and technical metadata and validating RNA-Seq from other studies. Of 48 analyzed tissues in GTEx, 26 had variant co-expression clusters of four known highly expressed and pancreas-enriched genes (*PRSS1*, *PNLIP*, *CLPS*, and/or *CELA3A*). Fourteen additional highly expressed genes from other tissues also indicated contamination. Sample contamination by non-native genes was associated with a sample being sequenced on the same day as a tissue that natively expressed those genes. This was highly significant for pancreas and esophagus genes (linear model, p=9.5e-237 and p=5e-260 respectively). Nine SNPs in four genes shown to contaminate non-native tissues demonstrated allelic differences between DNA-based genotypes and contaminated sample RNA-based genotypes, validating the contamination. Low-level contamination affected 4,497 (39.6%) samples (defined as ≥10 *PRSS1* TPM). It also led to eQTL assignments in inappropriate tissues among these 18 genes. We note this type of contamination occurs widely, impacting bulk and single cell data set analysis. In conclusion, highly expressed, tissue-enriched genes basally contaminate GTEx and other datasets impacting analyses. Awareness of this process is necessary to avoid assigning inaccurate importance to low-level gene expression in inappropriate tissues and cells.

Key Words: GTEx, RNA-Seq, Contamination, scRNA-Seq, eQTL, PEER factors

Introduction

The rise of next generation sequencing has allowed for unparalleled data

generation for a variety of nucleic acid studies including RNA expression. As cost per

basepair decreases, more large-scale transcriptome projects can be performed that will

inform on tissue expression patterns in health and disease[1-4]. These data sources are

generally publicly-available and have been used by hundreds of researchers for

secondary analyses of high impact[5,6].

Limitations exist for all –omics technologies, including RNA sequencing (RNA-

Seq). Issues of hybridization biases, library preparation biases, and computational

biases such as positional fragment bias are known limitations of RNA-Seq experiments[7-

9]. Another challenge of high throughput RNA-Seq is contamination, leading to the

presence of sequence data within a dataset of one sample that originates from a

separate sample. This contamination can come from many different aspects of the

modern sequencing process, such as human error, machine or equipment

contamination, intrinsic preparation and sequencing errors, and computational errors,

including errors that can occur based on the multiplexing methods used[10-12]. For single

cell RNA-Seq (scRNA-Seq), doublets or multiplets (2+ cells partitioned together) can

cause cross-contamination resulting in expression hybrids[13]. Compared to RNA-Seq,

contamination has been better characterized for DNA sequencing projects[14-16].

The Genotype-Tissue Expression project (GTEx) aims to create a large publicly

available database of tissue-specific expression quantitative trait loci (eQTL) from over

40 tissues[1]. It is an ongoing project with over 700 individuals and 11,000 tissue

samples. GTEx identifies eQTLs by associating genotypes called from whole genome

sequencing with gene expression levels obtained from bulk RNA-Seq. GTEx has made

their RNA-Seq, phenotype, genotype, and technical data available for public access

with permission.

In an analysis of variation in the GTEx RNA-Seq data (V7), we detected

unexpected sources of variation that we hypothesized were likely contaminating

sequence reads found at low, but variable levels across different tissues. Herein we

describe how we identified the source of contamination and establish basal rates of

contamination in the GTEx bulk RNA-Seq data. We further demonstrate the universality

of highly-expressed genes contaminating other samples.

Results

Patterns of extreme tissue variation identified usual and unusual gene signatures

We embarked on a project to expand our initial description of the causes of lung

expression variation in GTEx to all tissue samples [17]. We used DESeq2 variance

stabilizing transformation to normalize read counts from 11,340 samples[17,18]. Then we

filtered genes in each tissue keeping those with a mean transformed count >5. The

median number of genes above the expression threshold was 17,729 with the highest

and lowest gene counts being 23,930 and 13,807 in the testis and whole blood

respectively. As previously described, we correlated and hierarchically clustered

variable genes (>4 variance across samples) for all tissues with >70 samples (N=48) in

the GTEx dataset V7 [17]. Our algorithm identified multiple gene clusters per tissue,

based on their Kendall's tau correlations (Fig. 1a). It additionally reported non-

clustering, highly variable genes. Most clusters were the result of biologic and

phenotypic features related to the tissues. For example, a cluster of Y chromosome

genes and *XIST* appeared in 42 of 43 non-sex specific tissues. However, there was one

consistent cluster of 3-4 genes (*PNLIP, PRSS1, CELA3A,* and/or *CLPS)* identified in 26

of the 48 tissues that did not have an intuitive biological explanation. These genes are

highly-expressed and pancreas acinar-cell specific[19]. To identify other highly expressed

tissue-enriched genes appearing variably in other samples, we cross-referenced a list of

tissue-enriched proteins generated by the Human Protein Atlas (HPA) to the GTEx

transcripts per million (TPM) data (Table 1)[20,21]. We noted 18 genes from seven tissues

including two esophagus genes *KRT13* and *KRT4* that are highly expressed in their

native tissue and identified as variable in five or more other unrelated tissues (Fig. 1a,

Supplementary Fig. 1).

Table 1 Eighteen highly expressed genes that appear to contaminate five or more

GTEx tissues

Gene	Times identified as variable in other tissues	Highest expressed GTEx tissue	GTEx TPM	2nd Highest expressed GTEx Tissue		
				Tissue	GTEx TPM	Independent Study TPM
PRSS1	43	Pancreas	99,100	Stomach	28.36	0.92
PNLIP	34	Pancreas	33,660	Ovary	3.67	0
CPA1	31	Pancreas	54,500	Testis	8.16	0
GP2	29	Pancreas	14,280	Prostate	18.34	13.5
CELA3A	24	Pancreas	27,130	Stomach	14.1	0.03
KRT13	20	Esophagus	33,960	Vagina	13,140	3,961
PGC	19	Stomach	36,720	Lung	83.84	12.42
KRT4	18	Esophagus	22,290	Vagina	1,375	15,069
PRL	17	Pituitary	54,500	Testis	5.15	0
LIPF	15	Stomach	29,380	Testis	4.33	0
CLPS	14	Pancreas	51,640	Stomach	5.35	0
CTRB2	9	Pancreas	20,760	Ovary	3.21	0
MYBPC1	6	Skeletal muscle	3,587	Prostate	54.5	21.3

MYH2	6	Skeletal muscle	1,064	Colon	0.52	0.02
ZG16B	6	Salivary gland	17,540	Prostate	48.1	187.1
FGA	5	Liver	5,717	Stomach	12.43	6.02
HP	5	Liver	12,710	Adipose	140.1	2.7
CKM	5	Skeletal muscle	11,138	Heart	2,987	2,339
Independent study samples were taken from RNA-seq experiments of the 2[nd] highest GTEx tissue and were not co-sequenced with the highest expressed GTEx tissue.						

Fig. 1 Identification and explanation of sequencing contamination **a** A correlation

heatmap of highly variable subcutaneous adipose tissue genes across 442 subjects.

Red shows a positive correlation. The genes within the contamination cluster and the

sex cluster are given. The meaning of cluster A is unknown. Cluster B may relate to the

percentage of smooth muscle cells and cluster C includes acute phase reactants **b**

Contamination normalized score values for non-pancreas tissue samples (N=11,366)

colored relative to being sequenced on the same day as a pancreas tissue. **c** Violin plot

of the same data showing a strong, but not complete correlation of sequencing on a

pancreas day. **d** Ranked order of all samples either sequenced on the same day as a

pancreas sample (black) or on a non-pancreas sequencing day (colors) for *PRSS1* read

counts in log10. Among samples not sequenced on a pancreas day, 91% of samples

with >100 reads were sequenced within 4 days of a known sequenced pancreas. The

dashed line represents 100 reads. **e** *KRT4* contaminating reads in GTEX1's fibroblast

RNA-Seq appear to have originated from GTEX2 esophagus mucosa tissue. By DNA

and RNA of the appropriate tissue source of *KRT4*, sample GTEX1 is homozygous for

the C allele at rs7956809. The fibroblast sample is 87% G reads, primarily matching

sample GTEX2. The read count depth at the SNP in the GTEX1 esophagus was 85,803

and 204 for the GTEX1 fibroblast.

As both abundant and tissue-enriched genes were unlikely to be randomly and

lowly expressed in a range of other tissues, we performed analyses to determine the

source of the contamination.

We first questioned if the contamination occurred during tissue harvesting,

hypothesizing that occasionally small fragments of a tissue could contaminate a

140   separate sample from shared dissection tools or surfaces. A pancreas gene

141   contamination cluster was found in transformed fibroblasts which were grown over

142   multiple passages and would not retain other cell types over that time period,

143   challenging this as a source of contamination (Supplementary Fig. 1).

Linear mixed models reveal sequencing date as the source of contamination

We then queried if other technical sources of contamination may be related.

Other GTEx metadata available included nucleic acid isolation date and sequencing

date, which were both interrogated for associations with this contamination. The

normalized score of four pancreas-enriched genes (*PRSS1*, *CLPS*, *PNLIP*, and

CELA3A) was significantly higher in non-pancreas samples if they were sequenced on

the same day as a pancreas sample (Wilcox Rank Sum Test, p< 5e-324, Fig 1b,c). We

then performed linear mixed model analyses across non-contaminating tissues to

understand expression levels of genes known to be highly-enriched and highly-

expressed in the pancreas or esophagus, but were found lowly and variably in other

tissues (ex. coronary artery, liver, tibial nerve). Because the majority of samples were

sequenced on the same day as a pancreas or esophagus – mucosa sample, for model

robustness, we limited our models to only tissues with >40 samples not sequenced on

the same data as a respective contaminating tissue (Supplementary Data 1 and 2).

After adjusting for tissue type, the pancreas contamination score (average of

normalized expression values for *PRSS1*, *CLPS*, *PNLIP*, and *CELA3A*) was on average

0.175 higher if the tissue was isolated on the same day a pancreas sample (p=3.9e-12),

but on average 0.863 higher if the tissue was sequenced on the same day as a

pancreas sample (p=9.5e-237) (Supplementary Data 3).

A similar linear mixed model was used to evaluate the esophagus contamination

score (average of normalized expression values for two highly-expressed esophagus-

enriched genes *KRT4* and *KRT13*). The same model as above showed only sequencing

a tissue on the same day as a esophagus sample to be significant (p = 5e-260)

(Supplementary Data 4).

Despite the significance of with sequencing date, some high scores came from

samples that were not sequenced on the same days as pancreata. Focusing on just one

these genes, *PRSS1*, it was clear that all of these samples were sequenced within a

few days of a pancreas (Fig. 1d). This additionally implicated the library preparation

process (for which date information is lacking in GTEx) as a potential source for

contamination, as it is temporally related, but not identical, to sequencing date.

With this understanding of the temporal importance of sequencing date to gene

expression, we then revisited the four genes making up the pancreas normalized score

gene cluster. For all tissues, median normalized scores were higher on pancreas

sequencing dates, and the division between the 26 tissues with clustering vs. the 21

without was a result of the high threshold for TPM we set at the VST step of our pipeline

(Supplementary Figure 2).

Genetic polymorphisms confirm contamination is derived from other samples

To prove that pancreas/esophagus transcripts were contaminating from other

(non-self) samples we identified incongruencies between a person's genotype (from

DNA data) and the genotype in matching loci in the pancreas/esophagus contaminated

RNA-Seq samples. We required both the individuals' DNA genotype and RNA-Seq files

from contaminating tissues, in order to account for both RNA editing and preferential

allele expression. Based on these sample requirements and limited by available raw

sequencing files, we identified 11 contaminated tissues to evaluate. For each, we

obtained and processed their raw RNA-Seq FASTQ sequences to identify nucleotide

variants in both their contaminated tissues and their matched pancreas or esophagus

tissue (depending on the gene source of contamination). Additionally, we used the

GTEx filtered VCF file from the individual's sequenced DNA to further establish their

SNP allele patterns. Across all tissues, 533 SNPs, rare variants, and private variants,

were investigated in pancreas associated gene coding sequences (*PNLIP*, *CLPS*, and

CELA3A) and 190 in esophagus associated gene coding sequences (*KRT13*, *KRT4*).

As a comparison group, 287 variants were investigated in two control gene coding

sequences (*GAPDH*, and *RAB7A*) that have near ubiquitous expression across all

tissues. Of 1,010 variants obtained from the combined VCF files, 11 had some degree

of allelic heterogeneity (Table 2). No incongruencies were found in the 287 variants of

the two control genes.

Table 2 Allelic inconsistencies found in contaminated samples

Individual	Gene	SNP	Major/ Minor	Enriched Tissue		Contaminated		
				Reads	Major Allele %	Tissue Type	Reads	Major Allele %
GTEX-1	*KRT13*	rs903	C/A	101,908	0%	Fibroblast Cells	252	50%
GTEX-1	*KRT4*	rs7959052	T/C	74,468	100%	Fibroblast Cells	203	12%
GTEX-1	*KRT4*	rs7956809	C/G	85,803	100%	Fibroblast Cells	204	13%
GTEX-1	*KRT4*	rs2035879	T/C	72,978	51%	Fibroblast Cells	164	7%
GTEX-1	*KRT4*	rs17119475	G/A	71,592	49%	Fibroblast Cells	226	98%
GTEX-8	*CELA3A*	rs9187	C/T	162,318	73%	Tibial Nerve	1,155	100%
GTEX-8	*CELA3A*	rs12908	G/A	169,394	74%	Tibial Nerve	1,215	100%
GTEX-9	*CELA3A*	rs3820285	C/G	98,896	1%	Adipose	5,178	48%
GTEX-9	*CELA3A*	rs9187	C/T	105,462	75%	Adipose	6,082	97%
GTEX-9	*CELA3A*	rs12908	G/A	108,681	75%	Adipose	6,313	98%

| GTEX-10 | *CLPS* | rs3748050 | T/C | 80,019 | 47% C | Artery | 1,117 | 99% |

One SNP site, rs7956809, was particularly informative. SNP rs7956809 (C/G),

located in *KRT4*, had a relatively low allelic variation, with only 5 individuals in the entire

GTEx cohort homozygous for the alternative allele (G). One individual (arbitrarily

GTEX1) was homozygous C at rs7956809 in both its DNA (VCF file) and matched

esophagus (RNA-Seq FASTQ data) (Fig. 1e). However, the rs7956809 SNP in the

GTEX1 fibroblast sample was 87% G and 13% C. Six esophagus samples were

sequenced on the same day as the GTEX1 fibroblast sample. No other esophagus

samples were sequenced within 4 days. One of those six samples, GTEX2, was

homozygous G at rs7956809. The five other samples were homozygous C. This

strongly implicates the GTEX2 esophagus sample as the dominant contaminant of the

GTEX1 fibroblast sample.

We further investigated the relationship between the GTEX1 fibroblast sample

and the GTEX2 esophagus sample finding no clear connection. The two samples were

sequenced on different machines and in different flow cells. Of some interest, the

sequencing sample adapters (molecular indexes) were similar (Supplementary Data 5).

The extent of highly expressed, tissue-enriched gene contamination in GTEx

After establishing that contamination exists in GTEx by identification of a temporal

association and validation through polymorphisms, we then attempted to address the

extent of contamination in the GTEx dataset. To characterize this, we investigated the

various levels of pancreas enhanced gene expression in non-pancreatic tissue (Fig. 2).

In the 11,366 non-pancreas samples investigated, roughly 25% had 0 reads for each of

4 pancreas-enriched genes and another ~50% of samples had TPM <10.

Fig. 2 A density plot of 11,366 non-pancreas RNA-seq samples and their cumulative

TPM expression of four pancreas genes.

PEER factor normalization does not fully correct for contamination

The GTEx analysis pipeline uses probabilistic estimation of expression residuals

(PEER) factor to correct for possible confounders[22,23]. This method identifies hidden

factors that explain much of the expression variability and can be used to normalize

RNA expression data. We focused on just one tissue, lung, and followed the GTEx

analysis pipeline to determine the extent to which PEER factor normalization can

identify and correct for this contamination. Sixty PEER factors were identified, with the

top two capturing a difference between "in hospital" (short postmortem interval) and

"outside of hospital" (longer postmortem interval) deaths (Fig. 3a). This relationship is

consistent with our prior report of variation in lung[17]. Similar to the global findings of Fig.

1, *PNLIP* expression was increased in lung samples sequenced on the same day as a

pancreas. Despite correcting for 35 or even 60 PEER factors, this difference was not

fully accounted for (Fig. 3b). Indeed, of five genes evaluated, only one gene (*KRT4*)

was fully corrected for by PEER factors (Table 3). We then explored if this lack of full

correction impacted eQTL analysis in the GTEx program.

Table 3. Significance of same-day sequencing of lung with contaminating tissues

on gene expression

Gene	P. value before PEER correction	P. value after correcting for 35 PEER factors	P. value after correcting for 60 PEER factors	Beta estimate after correction
PNLIP	4.34e-14	1.39e-07	1.94e-05	0.49
PRSS1	6.29e-14	5.83e-07	2.63e-05	0.48
CELA3A	5.91e-14	3.36e-07	2.41e-05	0.49
KRT4	0.0034	0.17	0.15	0.18
KRT13	8.18e-17	0.00048	0.0036	0.37
P. values are shown before and after PEER correction.				

Fig. 3 Impact of PEER factors on contamination. **a** The top two PEER factors separated in hospital from out of hospital deaths (N=427). **b** With no PEER factor correction there is a significant increase in *PNLIP* expression normalized scores in lung samples if they were sequenced on the same day as a pancreas (No = 96, Yes = 331; linear regression, p= 4.34e-14). After 35 (p= 1.38e-11) or 60 (p= 3.03e-06) PEER factor corrections, the difference remained. **c** *PRL* read counts in pituitary (high), placenta (medium), and uterus (low), where *PRL* is known to be expressed across GTEx, HPA, and the RNA Atlas. The numbers in colored boxes indicate sample sizes and the color

indicates respective study. **d** *PRL* contamination reads across 6 tissues from 3 studies

that correlate with levels of likely contamination based on the other sequenced organs.

e *INS* contamination across three scRNA-Seq data sets. Only in the pancreas dataset

(GSE84133), where beta cells were also sequenced, does *INS* appear to be lowly

expressed in endothelial and mesenchymal cells. Cells with expression above the

dotted line at 1,000 TPM/100 are likely doublets or multiplets.

Contamination affects GTEx eQTL reporting

Using the GTEx eQTL browser, we identified 72 tissues reported as having significant

eQTLs for the 18 genes listed in Table 1. Seven tissues matched the known dominant

expression patterns of the genes. An additional 34 tissues were deemed possible based

on expression patterns noted by RNA and protein immunohistochemistry in which

expression (in TPM) was above the basal level of all tissues. However, 31 inappropriate

tissues were identified as harboring eQTLs even though these genes are not natively

expressed in these tissues, appearing only as a result of contamination (Table 4).

Table 4 Distribution of GTEx eQTLs by tissue type in contaminating genes

Genes	Appropriate Tissues	Possible tissues	Inappropriate tissues
PRSS1	--	Small intestine	Liver, coronary, skin, lung
PNLIP	--	--	--
CPA1	--	--	Coronary
GP2	--	--	Brain
CELA3A	Pancreas	Stomach	Liver
KRT13	Vagina	Lung	--
PGC	--	Lung, pancreas	Tibial artery
KRT4	Esophagus	Skin, lung	Colon, brain, thyroid
PRL	--	--	Gastroesophageal junction, skin, tibial artery
LIPF	Stomach	--	--
CLPS	Pancreas	--	--
CTRB2	Pancreas	--	Aorta, brain, lung, thyroid

FGA	Liver	Stomach	--
HP	--	Whole blood, adipose (2), artery (3), lung, tibial nerve, heart	Brain, esophagus mucosa
CKM	--	--	Aorta, whole blood
MYBPC1	--	Heart, testis, colon, prostate, brain (2)	Esophagus (2), lung, thyroid
MYH2	--	--	Colon, lung
ZG16B	--	Skin (2), adrenal, fibroblasts, lung, stomach, prostate, spleen, colon, testis, whole blood	Adipose, esophagus, pituitary

Non-GTEx bulk sequencing data sets confirm contamination

To determine if highly-expressed tissue-enriched contamination is a feature of sequencing in general, we identified two additional RNA-Seq datasets containing multiple organs/tissues[24,25,26]. Neither study sequenced the pituitary, which is the organ with the highest levels of prolactin (*PRL*) expression (Fig. 3c). Both studies performed multiplexed sequencing on Illumina 2000 or 3000 sequencers. These data sets demonstrate the extent of *PRL* expression contamination across six tissues is dependent on the amount of *PRL* expressed in the appropriate tissues (GTEx pituitary median 54,500 TPM and uterus median 4.01 TPM (Fig. 3d). To additionally characterize the breadth of contamination in the literature, we identified ten studies with two or more organs/cells sequenced together. All ten had cross-sample contamination of the type described herein (Supplementary Data 6). We also analyzed the METSIM study, containing bulk sequencing of adipose tissue in 434 subjects.[27] The data from that study demonstrates how, in the absence of sequencing at the same time as a contaminating tissue (ex. stomach, pancreas), contaminating reads are minimal do not correlate. This was opposite that observed in GTEx adipose tissue (Supplementary Fig. 3).

scRNA-Seq may be similarly impacted by this type of contamination.

We then investigated this type of highly-expressed gene contamination in

scRNA-Seq. Although doublets are well-characterized, and barcode swapping has been

reported, other forms of contamination are not well-described[13,28]. Also, due to the low

depth of sequencing in current scRNA-seq compared to bulk sequencing, the type of

contamination concerned herein would not be expected to generate large numbers of

contaminating reads. However, investigations across two cell types, endothelial cells

and mesenchymal cells (fibroblasts, activated stellate cells, cancer-associated

fibroblasts), suggested otherwise.

Three different scRNA-Seq data sets (GSE84133, GSE103322, GSE72056) from

normal pancreas, metastatic melanoma, and head and neck squamous cell carcinoma

were used[29-31]. From the pancreas sample, insulin (*INS*) represented a highly-

expressed, cell type-enriched gene that is exclusive to beta cells (TPM value of

167,144; transcripts per 10,000 cells = 1,671). *INS* expression was modestly elevated in

endothelial cells and mesenchymal cells from the same data set (GSE84133), but is

absent in endothelial and mesenchymal cells from the other data sets in which beta

cells were not sequenced (Fig. 3e). Contamination of pancreatic endothelial cells is

further supported by a recent snATAC-seq study that demonstrated closed chromatin at

the *INS* locus in pancreatic endothelial cells[32].

scRNA-seq contamination drives aberrant cell clustering

Tabula Muris is a single cell transcriptome atlas of 20 mouse tissues[33]. We

gathered endothelial cell data from 9 tissues and generated tSNE plots of this data

based on adjustment by either 6 or 10 principal components (PCs) (Supplementary Fig.

4a, b). With 10 PCs, a subset of pancreatic endothelial cells incorrectly clustered

separately as a result of contaminating pancreatic acinar genes that drive PCs 8 and

10. HPA staining validated the localization of these proteins to acinar cells

(Supplementary Fig. 4c).

Discussion

The GTEx dataset represents an ideal resource to study sequence

contamination. Its 11,000+ samples from 700+ individuals from a diverse set of tissues

with all library preparation and sequencing performed at one center is unique. During

our initial variation analysis of 48 tissues spanning 10,294 samples, we detected a

variable signal of pancreas genes in 26 of those tissues. From there we noticed genes

that were highly expressed in esophagus, stomach, pituitary and other tissues also

appeared in shared clusters across unrelated tissues. These highly expressed, tissue-

enriched genes were found at low, variable levels in other organs and represented

some of the most frequent causes of variation between samples of the same tissue

type.

We found that contamination is best linked to the date of sequencing for both

pancreas genes and esophagus genes (linear model, p=1.3e-14 and p=6.1e-49

respectively However, both due to contamination being noted in some samples that are

sequenced a few days apart from a possible contaminating source and the SNP-based

evidence, we suspect the majority of the contamination occurred during library

preparation rather than the sequencing itself. Library preparation dates were not

documented (personal communication, GTEx Help Desk).

While the nucleic acid isolation date was only modestly associated with

contamination, physical contamination can easily occur at this stage. GTEx RNA

isolation was manually done in batches of 12 tissues, purposefully with a mix of donors

and tissues to minimize batch effects. Samples were individually cut and placed into

cryovials for homogenization, followed by further manipulations [34].

At the stage of library preparation or sequencing where our data indicates most

of the contamination occurred, there are multiple steps that could be implicated. The

library preparation was completed automatically in 96 well plates with a mix of tissues

and individuals to prevent batch effects [34]. Fluidic carryover could have occurred here.

At the sequencing level, a major concern is index contamination where index

oligonucleotides used for multiplexing can ligate to other sample transcripts, thus

contaminating the data after demultiplexing. Index based contamination is machine and

lane specific and can even occur at the creation of the indexes when multiple indexes

are purified on the same high-performance liquid chromatography column [35].

Additionally, if steps to clean libraries of free adapters/primers are not properly

executed, the remaining indexes can contaminate clusters in the flow cells [11]. Molecular

recombination of indexes during sequencing can also lead to faulty read assignment as

multiplex clusters can become contaminated by other samples that acquire the indices

of the native sample (index hopping). This has been demonstrated in scRNA-Seq, and

seems to occur more often on a HiSeq 4000 platform[28].

Specific to this study, GTEx samples were run on either HiSeq 2000 or 2500

machines in high output mode using cbot clustering. The flowcells were not patterned,

and therefore not prone to index-swapping (personal communication, GTEx Help Desk).

GTEx's use of custom i7 (later Illumina kit-based) dual indices also reduced the amount

of index hopping that can occur[34,35].

Contamination of scRNA-Seq, may or may not be occurring by yet an additional

mechanism. Ambient RNA, from disrupted ("broken") cells may be a major source of

contamination, as recently described[36]. If true, this would occur before the library

preparation and sequencing steps, would cause only a modest expression of the gene,

and would not be detectable by current doublet correcting mechanisms[13,37]. This

method of contamination might have implications for Human Cell Atlas studies, as we

demonstrated it clearly affects cross-tissue comparisons of common cell types in Tabula

Muris[33,38].

Using other bulk RNA-Seq data sets with similar sequencing methods and

scRNA-Seq data sets, with different preparation methods, we validated that it is

predominately contamination, not low-level transcription, which results in non-zero

expression values in inappropriate tissues/cells (Fig. 3d,e, Supplementary Fig. 3,

Supplementary Fig. 4, Supplementary Data 6). These extra data sets also uncover the

generalizability of this highly-expressed transcript contamination regardless of the labs

in which they take place or the methods employed.

So how big is the contamination problem? It depends on how the data is to be

used. Fortunately, in the GTEx data, the levels are overall low with only 2.85% of

samples having relatively high levels of *PRSS1* (TPM > 1,00). Thus, for many uses of

GTEx data, this level is irrelevant. However, for experiments that involve differential

expression profiling, these genes will repeatedly appear due to their variable levels of

contamination. We additionally note that the GTEx standard normalization pipeline

using PEER factors did not entirely eliminate this source of variation and numerous

eQTLs that were identified for the 18 genes described herein were located in incorrect

tissues (>40%). We caution that our results do not suggest that one should design a

study in which sample type (e.g. tissue type or disease state) is perfectly confounded

with library preparation or sequencing date. In fact, it was only because the GTEx study

sequenced samples from different individuals on the same date that we were able to

definitively show that contamination between samples occurred.

In other scenarios, this basal contamination may be more worrisome. Many

publications have reported rare, but variable gene expression in their samples claiming

their importance or disease-related behaviors[39]. Our findings call these reports into

question. The extent of cross-contamination, where one laboratories' samples are

prepped and sequenced at the same time as a different laboratories' unrelated samples

through a university core sequencing facility or sequencing company is unknown, but

likely frequent[40,41]. The xenomiR story, that rice miRNAs are found in human blood

through dietary means[42], was shown to result from library preparation

contamination[43,44]. For scRNA-Seq, this contamination falsely implies some gene

expression is ubiquitous across cells, which influences computational heterogeneity

analyses of cell types, beyond the known challenges of other biological and technical

artifacts[33,36,45]. Also, our work highlights the fact that work flows must be considered

carefully in very-low DNA mutation detection analysis in clinical cancer samples as

samples with higher tumor burdens may contaminate samples with lower tumor burdens

and falsely suggest treatment approaches[46,47]. In particular, GTEx data is available in

many outlets, including the UCSC Genome Browser. Variable, low-level expression of

PRSS1, *CELA3A* and others may falsely intrigue researchers, particularly within the

reported eQTLs.

We described low-level, variable expression contamination in the GTEx RNA-

Seq dataset. This variation was most noticeable for 18 highly-expressed, tissue-

enriched genes and strongly correlates with the library preparation and sequencing of

the samples. Similar contamination was observed in other bulk and scRNA-Seq data

sets suggesting a universality to this type of contamination. Evaluating low-level variable

gene expression in RNA sequencing data sets must be performed with precaution and

awareness of potential sample contamination.

Methods

Ethics Statement

All human data was publicly available or used with approval of the GTEx consortium.

Consent was obtained by those studies.

Retrieval of GTEx RNA-Seq dataset, FASTQ files, and sample data

The gene read counts of the RNA-Seq GTEx version 7 dataset (GTEx_Analysis_2016-

01-15_v7_RNASeQCv1.1.8_gene_reads.gct.gz) were downloaded from the GTEx

Portal (https://gtexportal.org/home/datasets), along with the de-identified sample

annotations (GTEx_v7_Annotations_SampleAttributesDS.txt). The FASTQ files of the

tissue samples and the variant call file (VCF) files of appropriate individuals were

downloaded from dbGaP (phs000424.v7.p2) with the required permissions.

Retrieval of Human Protein Atlas tissue enriched gene list

422 We obtained the HPA tissue enriched genes by downloading a CSV file from this filtered

423 site

424 (https://www.proteinatlas.org/search/tissue_specificity_rna:any;Tissue%20enriched+AN

425 D+sort_by:tissue+specific+score, visited on 6/21/18).

Second highest tissue expression of tissue enriched genes and expression in

single tissue studies

Using the GTEx portal (v7) we noted the TPM for the second highest expressing tissue

with greater than 60 samples. We then used GEO, SRA Run Selector, and Recount2

(https://jhubiostatistics.shinyapps.io/recount/) to find datasets where only one tissue was

sequenced, with the exception of the ovary sample which was obtained from the

Illumina BodyMap (16 tissue types, absent of pancreas)[48]. Due to our inability to find

salivary gland data we used prostate tissue, the third highest expressing tissue for

ZG16B. Heart (ERP009437), adipose tissue (SRP053101), lung (SRP032833), prostate

(SRP003611), stomach (ERP010889), and colon (SRP029880) TPMs were acquired

through Recount2 and were selected for normal samples. The testis data was

downloaded from GEO Omnibus (GSE103905). Vagina and ovary FASTQ data was

downloaded from SRA run selector (GSE68229 and GSE30611). FASTQ files were

mapped to the Genome Reference Consortium Human Build 38 (GRCh38) using the

software HISAT2 version 2.1.0 [49]. The output SAM files were turned into BAM files and

indexed using samtools version 1.9 [50,51]. Assembly was completed using StringTie

v1.3.4d and StringTie TPMs were used from the output [52,53].

Bulk sequencing processing

The acquired raw read counts were segmented into separate tissue subsets (48

tissues with ≥70 samples each) and their read counts were normalized using the

Variance Stabilizing Transformation feature in DESeq2 version 1.22.1 in R version 3.6.1

[18]. This method incorporates estimated size factors based on the median-ratio method,

and transformed by the dispersion-mean relationship. We then filtered the 56,202 genes

based on their mean expression (mean transformed count > 5) to reduce noise and

lessen the inflated effect of low expressing genes on correlations.

Identification of highly variable genes and clusters

All analyses were completed in R version 3.6.1. In each tissue, a threshold of a

>4 variance of variance stabilizing transformed normalized read counts was used as our

cut off for highly variable transcripts. These genes were then clustered using hierarchal

clustering on a distance generated by 1 - Kendall's rank-correlation coefficient. A tau

critical value was calculated based on the number of samples and genes expressed.

The correlation-based dendrogram was cut to produce gene clusters with average

within cluster correlation of at least the tau critical value.

Calculation of normalized expression scores

Normalized expression scores allow one to summarize the expression of a gene

cluster in a sample by the average normalized score of the genes in that cluster. These

normalized scores were calculated by subtracting the mean expression and dividing by

the median absolute deviation of the expression values for each gene across all

samples within a given tissue. The equation is as follows, where x is the VST

normalized expression of gene j in sample i, $t(i)$ is the tissue type for sample i, and J is

the number of genes in a given cluster:

$$S_i = \frac{1}{J} \sum_{j=1}^{J} \frac{x_{ij} - \bar{x}_j}{\underset{m:t(m)=t(i)}{Median} \left| x_{mj} - \bar{x}_j \right|}$$

A "pancreas" contamination score was defined as the average normalized score

using the genes *PRSS1, CELA3A, PNLIP,* and *CLPS*, across all tissues. An

"esophagus" normalized score was defined as the average normalized score using

genes *KRT4* and *KRT13* in a given tissue. All normalized scores were calculated

independently for each tissue, to account for tissue-specific between-sample variation in

the expression of these genes.

Linear mixed model analysis for pancreas gene contamination

Linear models were used to find associations between nucleic acid isolation or

sequencing on the same day as a potential contaminating tissue (source of gene

expression). These analyses were completed in R version 3.6.1 using the lmer function

in package lme4 (v1.1-21). In the linear mixed models we used all available tissues with

>40 samples not sequenced on the same day as a pancreas sample (15 tissues,

N=6,258). For the model, tissue was used as a covariate along with "pancreas

sequencing day" and "pancreas nucleic acid isolation day" were coded binarily as

True/False based on whether a given non-pancreas sample was sequenced or

underwent RNA isolation on the same day as a pancreas sample. Our model identified

an association between the pancreas contamination score with pancreas sequencing

day (T/F; 1/0), pancreas nucleic acid isolation day (T/F; 1/0), and tissue as covariates,

with the subject IDs included as a random effect. The equation for our model is below

where $t(i)$ gives the tissue type of sample i, $s(i)$ is defined as the sequencing day of

the sample, and $d(i)$ is the isolation date of sample i. Lastly T is the number of tissue

types, $u_{n(i)}$ is the random effect of subject ID for sample i, and ε_i is the error term for

said sample:

$$Y_i = \beta_0 + \beta_1 I\big(s(i) \in \{ s\big(m : t(m) = t_{pancreas}\big)\}\big)$$

$$+ \beta_2 I\big(d(i) \in \{ d\big(m : t(m) = t_{pancreas}\big)\}\big)$$

$$+ \sum_{k=3}^{T+1} \beta_k I(t(i) = k - 1) + u_{n(i)} + \varepsilon_i$$

Linear mixed model analysis for esophagus gene contamination

We repeated the above analysis with the esophagus mucosa, coding shared

sequencing day and shared nucleic acid isolation day similarly. In the linear mixed

model we used all tissues with a sample size >40 samples not sequenced on the same

day as the esophagus mucosa (8 tissues, N=3,917). The model used esophagus

nucleic acid isolation day (T/F; 1/0), esophagus sequencing day (T/F; 1/0), and tissue

as covariates along with subject IDs as a random effect on the esophagus

contamination score.

Base pair incongruency analysis

Base pair incongruency analysis required a contaminated tissue expression

FASTQ, a native tissue expression FASTQ, and the individual's VCF file. FASTQ files

were mapped to the Genome Reference Consortium Human Build 37 (hg19) using the

software HISAT2 version 2.1.0 [49]. The output SAM files were turned into BAM files and

indexed using samtools version 1.9 [50,51]. Preliminary analysis and development of

figures were generated using the Integrative Genome Viewer version 2.4.13 [54,55]).

Protein coding SNPs, rare variants, and personal variants (collectively referred to as

variants in this paper), were manually selected using IGV as a reference. Using the tool

bam-readcount version 0.8.0 in combination with a Python 3.6.2 script, a list of RNA-

Seq and genomic incongruencies were generated for the acquired sample BAM files.

PEER factor analysis

We obtained the GTEx RNA-Seq dataset from lung (N=427). The data underwent

trimmed mean of m-values (TMM) normalization and filtering out of lowly expressed

genes (< 0.1 TPM for 80% or more of the samples) before running PEER to identify

potential confounders [22]. Following GTEx's pipeline

(https://gtexportal.org/home/documentationPage#staticTextAnalysisMethods visited),

we then performed an inverse normal transformation (INT) on the expression values for

each gene in order to reduce the effect of outliers[23]. Normalized scores for each gene

are based on TMM-normalization, inverse-normal transformation, and scaling/centering

at zero. A linear regression was performed, with either raw inverse-normalized

expression values or expression normalized scores (after correcting for PEER factors)

as the outcomes and same-day pancreas (/esophagus) sequencing status as the

predictor. Beta estimates represent how many standard deviations greater the mean

expression of a gene is when samples are sequenced on the same day as a

contaminating tissue, even when accounting for variance explained by 60 PEER factors.

Cross-referencing eQTLs with contamination findings

We obtained and tallied eQTL reports for the 18 genes in Table 1 from the GTEx

eQTL browser (https://gtexportal.org visited on March 26, 2019). eQTLs were identified

by tissue association and conservatively placed in one of three categories: appropriate

expression, possible expression, and inappropriate expression. The appropriateness of

expression in any tissue was based on the evaluation of TPM levels in the tissue and

immunohistochemistry staining patterns as noted in the HPA[56].

Acquiring Human Protein Atlas, RNA Atlas RNA-Seq and METSIM Data

Using the R package recount version 1.8.2, we downloaded HPA RNA-Seq data,

accession ERP003613 [24]. The RNA Atlas was acquired by downloading their raw RNA-

seq counts from Gene Expression Omnibus (GSE120795)[25]. The HPA RNA-Seq was

performed across 27 tissues and the RNA Atlas was across 20 tissues. We filtered

samples down to the shared tissues of liver, lung, pancreas, prostate, skin, and

stomach as well as the potential sources of *PRL* contamination of pituitary, placenta and

uterus. Only sun exposed skin was used for GTEx analysis. TPM RNA-seq data from

the METSIM study was downloaded through the GEO Omnibus (GSE135134)[27].

Acquiring and Normalizing scRNA-Seq Data

We identified three publications with human scRNA-Seq data sets that all

contained endothelial and mesenchymal cells[29-31]. All three sets had median read

counts/cell >100,000. The processed read counts from the three studies were obtained

from the Gene Expression Omnibus (GEO) (GSE84133; GSE103322; GSE72056). For

each study we used the supplied cell type information to identify endothelial and

mesenchymal-type cells. Mesenchymal cells were labeled as "activated stellate"

(GSE84133), "fibroblast" (GSE103322), and "cancer-associated fibroblast" (GSE72056).

The GSE103322 and GSE72056 read data were generated using Smart-Seq2 on a

NextSeq 500 instrument[57]. Raw FASTQ data sets were normalized using RSEM to

calculate $E_{i,j} = log_2(TPM_{i,j}/10+1)$, where $TPM_{i,j}$ refers to transcript-per-million for gene i in

sample j [58]. The $TPM_{i,j}$ was then divided by 10, to be on the order of 100,000 transcripts

to account for single cell read depth and the effect of 0s in downstream analysis[29].

GSE84133 was generated using the inDrop method and sequenced on a HiSeq 2500

instrument[59]. Human islets were isolated using a modified Ricordi method which

includes collagenase, mechanical agitation, pumping and centrifugation[60] . To make

single cells, the islets were then centrifuged twice at 250 rpm and treated with TrypLE

Express before mechanical dispersion with a P100 pipette, followed by centrifugation at

500 rpm. The two cancer sample libraries (GSE103322, GSE72056) were generated

using Smart-Seq2 and sequenced on a NextSeq 500 instrument[57]. Read counts were

scaled using TPM/100 to equalize expression level counts to the other two studies. The

calculation of the percent contamination of endothelial cells by *INS* was the ratio of

average normalized *INS* read counts per endothelial cell divided by the normalized

average read count in beta cells and excluding the 9 cells with >1000 TPM/100.

565   We obtained the Tabula Muris read data from

566 https://figshare.com/articles/Robject_files_for_tissues_processed_by_Seurat/5821263

567 (visited on 8/8/19)[33]. We used the R package Seurat (v3.1.1) and selected endothelial

568 cells from nine organs/tissues (brain, fat, heart, kidney, limb muscle, liver, lung,

569 pancreas, trachea) based on the annotation of the seurat object in the metadata slot

570 cell_ontology_class labeled 'endothelial cell'[61]. tSNE plots were generated based on the

571 top 6 or top 10 PCs. The average number of PCs used by Tabula Muris per

572 tissue/organ to create tSNE maps was 12 and the median was 10. Images of protein

573 expression of contaminating pancreas genes were obtained from the HPA[56].

576 **Data availability**

577 No new sequencing data was created for this study. All data used in this study is

578 available through dbGap (https://www.ncbi.nlm.nih.gov/gap/), GEO

579 (https://www.ncbi.nlm.nih.gov/gds), recount2

580 (https://lcolladotor.github.io/project/recount2/) or Tabula Muris

581 (https://figshare.com/articles/Robject_files_for_tissues_processed_by_Seurat/5821263).

583 **Code availability**

584 Code for all analyses is deposited at GitHub

585 (https://github.com/mhalushka/gtex_contamination_code). Computational analyses

586 were done using public R packages except when specifically noted otherwise.

Acknowledgements

The authors thank the GTEx Help Desk, Kristen Ardlie, Sheila Dodge, and NDRI for

valuable assistance in understanding the technical data sets and islet isolation

methodology. M.K.H. was supported by grants 1R01HL137811, R01GM130564, and

P30CA006973 from the National Institutes of Health and 17GRNT33670405 from the

American Heart Association. T.O.N. was supported by grants R01GM130564 and

T32GM07814. D.E.A was supported by 1R01HL131573 and 1R01HL137811. M.N.M.

was supported by R01HL137811 and the University of Rochester CTSA award number

UL1TR002001. A.Z.R was supported by R01GM130564.

Contributions: M.K.H., M.N.M and A.Z.R conceived of the experiments and assisted

with the manuscripts. T.O.N. performed the experiments, analyzed the data and wrote

the manuscript. S.Y., R.X.V., V.P. and D.E.A. performed experiments and assisted on

the manuscript.

Conflicts of interest

602 The authors declare no conflicts of interest.

References

605   1   Consortium, G. T. The Genotype-Tissue Expression (GTEx) project. *Nat Genet* **45**, 580-585,
606   doi:10.1038/ng.2653 (2013).
607   2   Tomczak, K., Czerwinska, P. & Wiznerowicz, M. The Cancer Genome Atlas (TCGA): an
608   immeasurable source of knowledge. *Contemporary oncology* **19**, A68-77,
609   doi:10.5114/wo.2014.47136 (2015).
610   3   Zeisel, A. *et al.* Brain structure. Cell types in the mouse cortex and hippocampus revealed by
611   single-cell RNA-seq. *Science* **347**, 1138-1142, doi:10.1126/science.aaa1934 (2015).
612   4   Kumasaka, N., Knights, A. J. & Gaffney, D. J. Fine-mapping cellular QTLs with RASQUAL and
613   ATAC-seq. *Nat Genet* **48**, 206-213, doi:10.1038/ng.3467 (2016).
614   5   Gutman, D. A. *et al.* MR imaging predictors of molecular profile and survival: multi-institutional
615   study of the TCGA glioblastoma data set. *Radiology* **267**, 560-569, doi:10.1148/radiol.13120118
616   (2013).
617   6   Ewing, A. D. *et al.* Combining tumor genome simulation with crowdsourcing to benchmark
618   somatic single-nucleotide-variant detection. *Nature methods* **12**, 623-630,
619   doi:10.1038/nmeth.3407 (2015).
620   7   Okoniewski, M. J. & Miller, C. J. Hybridization interactions between probesets in short oligo
621   microarrays lead to spurious correlations. *BMC Bioinformatics* **7**, 276, doi:10.1186/1471-2105-7-
622   276 (2006).
623   8   van Dijk, E. L., Jaszczyszyn, Y. & Thermes, C. Library preparation methods for next-generation
624   sequencing: tone down the bias. *Experimental cell research* **322**, 12-20,
625   doi:10.1016/j.yexcr.2014.01.008 (2014).
626   9   Tuerk, A., Wiktorin, G. & Guler, S. Mixture models reveal multiple positional bias types in RNA-
627   Seq data and lead to accurate transcript concentration estimates. *PLoS computational biology*
628   **13**, e1005515, doi:10.1371/journal.pcbi.1005515 (2017).
629   10   Lusk, R. W. Diverse and widespread contamination evident in the unmapped depths of high
630   throughput sequencing data. *PLoS One* **9**, e110808, doi:10.1371/journal.pone.0110808 (2014).
631   11   Illumina. *Effects of Index Misassignment on Multiplexing and Downstream Analysis*,
632   <https://www.illumina.com/content/dam/illumina-
633   marketing/documents/products/whitepapers/index-hopping-white-paper-770-2017-004.pdf> (
634   12   Rosenberg, A. Z. *et al.* xMD-miRNA-seq to generate near in vivo miRNA expression estimates in
635   colon epithelial cells. *Scientific reports* **8**, 9783, doi:10.1038/s41598-018-28198-z (2018).
636   13   McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection in Single-Cell
637   RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**, 329-337 e324,
638   doi:10.1016/j.cels.2019.03.003 (2019).
639   14   Merchant, S., Wood, D. E. & Salzberg, S. L. Unexpected cross-species contamination in genome
640   sequencing projects. *PeerJ* **2**, e675, doi:10.7717/peerj.675 (2014).
641   15   Cibulskis, K. *et al.* ContEst: estimating cross-contamination of human samples in next-generation
642   sequencing data. *Bioinformatics* **27**, 2601-2602, doi:10.1093/bioinformatics/btr446 (2011).
643   16   Ma, X. *et al.* Analysis of error profiles in deep next-generation sequencing data. *Genome Biol* **20**,
644   50, doi:10.1186/s13059-019-1659-6 (2019).
645   17   McCall, M. N., Illei, P. B. & Halushka, M. K. Complex Sources of Variation in Tissue Expression
646   Data: Analysis of the GTEx Lung Transcriptome. *American journal of human genetics* **99**, 624-
647   635, doi:10.1016/j.ajhg.2016.07.007 (2016).
648   18   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
649   RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-8 (2014).
650   19   Muraro, M. J. *et al.* A Single-Cell Transcriptome Atlas of the Human Pancreas. *Cell Syst* **3**, 385-
651   394 e383, doi:10.1016/j.cels.2016.09.002 (2016).

20   Uhlen, M. *et al.* Proteomics. Tissue-based map of the human proteome. *Science* **347**, 1260419,
doi:10.1126/science.1260419 (2015).

21   Atlas, T. H. P. *The human tissue specific proteome*,
<https://www.proteinatlas.org/humanproteome/tissue/tissue+specific> (

22   Stegle, O., Parts, L., Piipari, M., Winn, J. & Durbin, R. Using probabilistic estimation of expression
residuals (PEER) to obtain increased power and interpretability of gene expression analyses.
Nature protocols **7**, 500-507, doi:10.1038/nprot.2011.457 (2012).

23   Consortium, G. T. *et al.* Genetic effects on gene expression across human tissues. *Nature* **550**,
204-213, doi:10.1038/nature24277 (2017).

24   Fagerberg, L. *et al.* Analysis of the human tissue-specific expression by genome-wide integration
of transcriptomics and antibody-based proteomics. *Molecular & cellular proteomics : MCP* **13**,
397-406, doi:10.1074/mcp.M113.035600 (2014).

25   Suntsova, M. *et al.* Atlas of RNA sequencing profiles for normal human tissues. *Sci Data* **6**, 36,
doi:10.1038/s41597-019-0043-4 (2019).

26   Chhibber, A. *et al.* Transcriptomic variation of pharmacogenes in multiple human tissues and
lymphoblastoid cell lines. *The pharmacogenomics journal* **17**, 137-145, doi:10.1038/tpj.2015.93
(2017).

27   Raulerson, C. K. *et al.* Adipose Tissue Gene Expression Associations Reveal Hundreds of
Candidate Genes for Cardiometabolic Traits. *American journal of human genetics* **105**, 773-787,
doi:10.1016/j.ajhg.2019.09.001 (2019).

28   Griffiths, J. A., Richard, A. C., Bach, K., Lun, A. T. L. & Marioni, J. C. Detection and removal of
barcode swapping in single-cell RNA-seq data. *Nature communications* **9**, 2667,
doi:10.1038/s41467-018-05083-x (2018).

29   Puram, S. V. *et al.* Single-Cell Transcriptomic Analysis of Primary and Metastatic Tumor
Ecosystems in Head and Neck Cancer. *Cell* **171**, 1611-1624 e1624, doi:10.1016/j.cell.2017.10.044
(2017).

30   Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by single-cell
RNA-seq. *Science* **352**, 189-196, doi:10.1126/science.aad0501 (2016).

31   Baron, M. *et al.* A Single-Cell Transcriptomic Map of the Human and Mouse Pancreas Reveals
Inter- and Intra-cell Population Structure. *Cell Syst* **3**, 346-360 e344,
doi:10.1016/j.cels.2016.08.011 (2016).

32   Chiou, J. *et al.* Single cell chromatin accessibility reveals pancreatic islet cell type- and state-
specific regulatory programs of diabetes risk. *BioRxiv*, doi:https://doi.org/10.1101/693671
(2019).

33   Tabula Muris, C. *et al.* Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris.
Nature **562**, 367-372, doi:10.1038/s41586-018-0590-4 (2018).

34   Consortium, G. T. Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis:
multitissue gene regulation in humans. *Science* **348**, 648-660, doi:10.1126/science.1262110
(2015).

35   Kircher, M., Sawyer, S. & Meyer, M. Double indexing overcomes inaccuracies in multiplex
sequencing on the Illumina platform. *Nucleic Acids Res* **40**, e3, doi:10.1093/nar/gkr771 (2012).

36   Young, M. D. & Behjati, S. SoupX removes ambient RNA contamination from droplet based
single cell RNA sequencing data. *BioRxiv*, doi:https://doi.org/10.1101/303727 (2018).

37   DePasquale, E. A. K. *et al. DoubletDecon: Cell-State Aware Removal of Single-Cell RNA-Seq*
Doublets (https://www.biorxiv.org/content/10.1101/364810v2, 2019).

38   Regev, A. *et al.* The Human Cell Atlas. *eLife* **6**, doi:10.7554/eLife.27041 (2017).

39   Witwer, K. W. & Halushka, M. K. Toward the promise of microRNAs - Enhancing reproducibility
and rigor in microRNA research. *RNA biology* **13**, 1103-1116,
doi:10.1080/15476286.2016.1236172 (2016).
40   Kryukov, K. & Imanishi, T. Human Contamination in Public Genome Assemblies. *PLoS One* **11**,
e0162424, doi:10.1371/journal.pone.0162424 (2016).
41   Longo, M. S., O'Neill, M. J. & O'Neill, R. J. Abundant human DNA contamination identified in non-
primate genome databases. *PLoS One* **6**, e16410, doi:10.1371/journal.pone.0016410 (2011).
42   Zhang, L. *et al.* Exogenous plant MIR168a specifically targets mammalian LDLRAP1: evidence of
cross-kingdom regulation by microRNA. *Cell Res* **22**, 107-126, doi:10.1038/cr.2011.158 (2012).
43   Tosar, J. P., Rovira, C., Naya, H. & Cayota, A. Mining of public sequencing databases supports a
non-dietary origin for putative foreign miRNAs: underestimated effects of contamination in NGS.
RNA **20**, 754-757, doi:10.1261/rna.044263.114 (2014).
44   Zhang, Y. *et al.* Analysis of plant-derived miRNAs in animal small RNA datasets. *BMC Genomics*
13, 381, doi:10.1186/1471-2164-13-381 (2012).
45   Li, H. *et al.* Reference component analysis of single-cell transcriptomes elucidates cellular
heterogeneity in human colorectal tumors. *Nat Genet* **49**, 708-718, doi:10.1038/ng.3818 (2017).
46   Tian, S. K. *et al.* Optimizing Workflows and Processing of Cytologic Samples for Comprehensive
Analysis by Next-Generation Sequencing: Memorial Sloan Kettering Cancer Center Experience.
Archives of pathology & laboratory medicine, doi:10.5858/arpa.2016-0108-RA (2016).
47   Van Allen, E. M. *et al.* Whole-exome sequencing and clinical interpretation of formalin-fixed,
paraffin-embedded tumor samples to guide precision cancer medicine. *Nature medicine* **20**,
682-688, doi:10.1038/nm.3559 (2014).
48   Collado-Torres, L., Nellore, A. & Jaffe, A. E. recount workflow: Accessing over 70,000 human
RNA-seq samples with Bioconductor. *F1000Research* **6**, 1558,
doi:10.12688/f1000research.12223.1 (2017).
49   Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory
requirements. *Nature methods* **12**, 357-360, doi:10.1038/nmeth.3317 (2015).
50   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079,
doi:10.1093/bioinformatics/btp352 (2009).
51   Li, H. A statistical framework for SNP calling, mutation discovery, association mapping and
population genetical parameter estimation from sequencing data. *Bioinformatics* **27**, 2987-2993,
doi:10.1093/bioinformatics/btr509 (2011).
52   Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from RNA-seq
reads. *Nature biotechnology* **33**, 290-295, doi:10.1038/nbt.3122 (2015).
53   Pertea, M., Kim, D., Pertea, G. M., Leek, J. T. & Salzberg, S. L. Transcript-level expression analysis
of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nature protocols* **11**, 1650-1667,
doi:10.1038/nprot.2016.095 (2016).
54   Thorvaldsdottir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics Viewer (IGV): high-
performance genomics data visualization and exploration. *Briefings in bioinformatics* **14**, 178-
192, doi:10.1093/bib/bbs017 (2013).
55   Robinson, J. T. *et al.* Integrative genomics viewer. *Nature biotechnology* **29**, 24-26,
doi:10.1038/nbt.1754 (2011).
56   Uhlen, M. *et al.* Towards a knowledge-based Human Protein Atlas. *Nature biotechnology* **28**,
1248-1250, doi:10.1038/nbt1210-1248 (2010).
57   Picelli, S. *et al.* Smart-seq2 for sensitive full-length transcriptome profiling in single cells. *Nature*
methods **10**, 1096-1098, doi:10.1038/nmeth.2639 (2013).

58   Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* **12**, 323, doi:10.1186/1471-2105-12-323
(2011).
59   Klein, A. M. *et al.* Droplet barcoding for single-cell transcriptomics applied to embryonic stem
cells. *Cell* **161**, 1187-1201, doi:10.1016/j.cell.2015.04.044 (2015).
60   Ricordi, C., Lacy, P. E., Finke, E. H., Olack, B. J. & Scharp, D. W. Automated method for isolation
of human pancreatic islets. *Diabetes* **37**, 413-420 (1988).
61   Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of single-cell
gene expression data. *Nature biotechnology* **33**, 495-502, doi:10.1038/nbt.3192 (2015).

Basal Contamination of Sequencing: Lessons from the GTEx dataset

Tim O. Nieuwenhuis et al.

Supplementary Information

Supplementary Figure 1. A correlation heatmap of the highly variable gene clusters in 343 transformed fibroblast samples. Red shows a positive correlation. Genes within the contamination cluster are given. The etiology of clusters A and B are unknown. Cluster C represents co-variable genes related to mitosis.

Normalization score of pancreas contamination

Pancreas Gene Cluster Identified in These Tissues

Pancreas Gene Cluster Not Identified in These Tissues

	Pancreas Cluster?	
	Yes	No
	TPM	TPM
CELA3A	9.06	5.43
CLPS	14.05	6.03
PRSS1	31.93	16.43
PNLIP	10.31	4.34

Supplementary Figure 2. Violin plots representing the pancreatic gene (*CELA3A,*
CLPS, *PRSS1*, *PNLIP*) normalization scores of samples sequenced the same day as a
pancreas sample and those that were not. The bottom right corner shows the median
TPM of pancreas genes in non-pancreas tissues. These were divided base on the
tissue having all four genes in their contamination cluster (N=26) or not (N=21).

Supplementary Figure 3. A scatterplot comparison of different contaminating genes between the GTEx adipose tissues (n = 797) and the adipose tissues harvested in the METSIM study (n = 434). We compared *CPA1* vs *PRSS1*, *KRT13* vs *KRT4*, *PGC* vs *LIPF*, and *MYBPC1* vs *CKM* as each of those gene pairs are highly expressed together in a contaminating tissue type. In GTEx the expression is consistently higher and has a stronger positive correlation compared to the METSIM study for three of those pairs Because skeletal muscle can be captured in adipose harvesting, *MYBPC1* and *CKM* also showed strong correlation in METSIM as a result of a direct contamination.

Supplementary Figure 4. Pancreatic acinar cell contamination drives endothelial cell diversity in Tabula Muris. **a** The major genes of the top 10 principal components (PCs) for nine groups of endothelial cells. * represents contaminating acinar cell expression. **b** tSNE plots based on 6 or 10 PCs demonstrate a large deviation of pancreas endothelial cells from a shared cluster to a unique cluster based on additional PCs. **c** In this black and white representation, a subset of pancreas cells are noted to move. **d** Representative images (HPA) of several highly-expressed acinar proteins that do not mark endothelial cells (black arrows) and *PECAM1*, a known endothelial cell marker.