N-Glycome regulation limits the transdifferentiation of endothelial cells into hematopoietic
stem cells

Authors:
Dionna M. Kasper[1, 2, 4], Yinyu Wu[1, 2, 4], Hanna K. Mandl[1, 2, 4], Kevin Salinas[1, 2, 4], Joey Ghersi[1, 2, 4],
Jared Hintzen[1, 2, 4], William Armero[1, 2, 4], Zhiheng He[1, 2, 4], Daniel W. Heindel[5,6], Eon Joo Park[3, 4],
William C. Sessa[3, 4], Lara K. Mahal[5,6], Karen K. Hirschi[1, 2, 4*], Stefania Nicoli[1,2,3,4*]

Affiliations:
[1]Yale Cardiovascular Research Center, Department of Internal Medicine, Section
of Cardiology, Yale University School of Medicine, New Haven, CT 06511, USA
[2]Department of Genetics, [3] Department of Pharmacology, Yale University School of Medicine,
New Haven, CT 06510, USA, [4]Vascular Biology & Therapeutics Program, Yale University
School of Medicine, New Haven, CT 06520, USA [5]Interdisciplinary Melanoma Cooperative
Group, Perlmutter Cancer Center New York University School of Medicine, New York, NY
10016, USA [6]Biomedical Chemistry Institute, Department of Chemistry, New York University,
New York, NY 10003, USA
*corresponding authors stefania.nicoli@yale.edu; karen.hirschi@yale.edu

Abstract:
Hematopoietic stem and progenitor cells (HSPCs) are required to establish and maintain the
adult blood system in vertebrates. During development, hemogenic endothelial cells undergo an
endothelial-to-hematopoietic transition (EHT) to generate HSPCs[1-4]. Growth factors and
epigenetic changes can promote EHT[1,3,5], but these mechanisms do not explain its tight
spatiotemporal regulation during development. Here, we show that microRNA (miR) miR-223-
mediated regulation of N-glycan biosynthesis intrinsically restrains EHT, representing the first
pathway that prevents excessive HSPC production. We find that miR-223 is uniquely expressed
in hemogenic endothelial cells undergoing EHT and in nascent HSPCs. Loss of miR-223
promotes the expansion of these cells in the zebrafish and mouse aorta-gonad-mesonephros
(AGM), where EHT occurs[6-8]. miR-223 targets *alg2* (α1,3/ α1,6 mannosyltransferase) and
st3gal2 (α2,3 sialyltransferase) for repression in the AGM endothelium. These two enzymes are

involved in the biosynthesis of N-glycans, a common co-translational modification[9,10] that
influences several pathophysiological processes[11,12], but has not yet been implicated in EHT.
Using an N-glycosensor, we demonstrate that vascular N-glycosylation increases during EHT,
and this process is disrupted upon loss of miR-223. Specifically, high-throughput glycome
analysis revealed terminal α1,3 mannose and α2,3 sialic acid modifications of membrane
proteins are altered upon loss of miR-223. Importantly, pharmacological manipulation targeting
these N-glycan types in wild-type embryos phenocopies the loss of miR-223 and enhances EHT
as well as HSPC production. Thus, the N-glycome plays a previously unappreciated role as an
intrinsic negative regulator of EHT, with specific mannose and sialic acid modifications serving
as key endothelial determinants of the hematopoietic fate.

Zebrafish endothelial-to-hematopoietic transition (EHT) has emerged as an instrumental
model to understand the formation and regulation of definitive hematopoietic stem and
progenitor cells (HSPCs) with long-term engraftment capability. As in their mammalian
counterparts, EHT occurs in endothelial cells that co-express vascular and hematopoietic genes
(hemogenic endothelial cells, hemECs). Through the progressive loss of endothelial gene
expression and concomitant increase in hematopoietic potential[1-4,13], hemECs give birth to
nascent HSPCs that delaminate from the vascular wall into the circulation to colonize secondary
hematopoietic organs[6-8]. This process is highly regulated, involving a precise interplay between
multiple developmentally-timed signaling pathways[1-5,13-16]. It is currently unclear whether such
precision is achieved exclusively by the regulation of positive hematopoietic regulators, or
whether yet-unknown inhibitory pathways play a role in EHT dynamics.

To discover underlying mechanisms of EHT inhibition, we investigated the function of
miR-223, a miRNA enriched in zebrafish endothelial cells; genetic deletion of miR-223 results in
HSPC expansion in the AGM[17]. We generated a transgenic reporter to examine miR-223
expression during EHT. Specifically, we used CRISPR/Cas9 genome editing to integrate the
Gal4 transcriptional activator[18] downstream of the endogenous miR-223 promoter (miR-
223:Gal4), and coupled it with a *Tg(5XUAS:eGFP)[nkuasgfp1a]* and an endothelial cell *Tg(kdrl:hras-*
mCherry)[ss896] marker. This approach fluorescently labels miR-223-expressing endothelial cells

with both GFP and mCherry (hereafter referred to as miR-223:GFP+ kdrl:mCH+) (Fig. 1a, b and

Supplementary Fig. 1a).

miR-223:GFP+ kdrl:mCH+ cells were dispersed in a salt and pepper pattern, with the

majority of cells positioned within the AGM (Fig. 1a-c). We found that the AGM population of

miR-223:GFP+ kdrl:mCH+ cells increased from the onset of EHT at 24 hours post fertilization

(hpf) to its peak at 32 hpf (Fig. 1c). In addition, they manifested a heterogeneous pattern of flat

to bulging morphologies, and underwent delamination from the AGM (Fig. 1a, Supplementary

Fig. 1b, c and Supplementary Movie 1). Importantly, miR-223:GFP+ kdrl:mCH+ cells co-

expressed the endothelial marker *kdrl* and hemEC markers *gata2b*[19] and *runx1*[20,21], as wells as

the nascent HSPC marker *cmyb*[22] (Fig. 1d), suggesting that miR-223 is expressed in hemECs

undergoing EHT.

To understand how miR-223 affects EHT, we analyzed the miR-223 mutant (miR-

223Δ/Δ) zebrafish AGM, and observed increased numbers of hemECs expressing *gata2b* at 24

hpf, as well as the downstream factor *runx1* at 27 hpf (Fig. 2a, b and Supplementary Fig. 2a,

b). Congruently, the kdrl:mCH+ cmyb:GFP+ nascent HSPC population[6] was enlarged for the

duration of EHT (Fig. 2a, c and Supplementary Fig. 2c), resulting in HSPC overexpansion at

subsequent hematopoietic sites[17]. Importantly, expression of the wild-type miR-223 gene in miR-

223:Gal4+ or in endothelial *fli1a+* cells rescued the HSPC overexpansion in miR-

223Δ/Δ zebrafish (Supplementary Fig. 2d-g), verifying that miR-223 functions in endothelial

cells of the AGM to limit HSPC production. Taken together, these findings support that miR-223

intrinsically restricts EHT in hemECs.

To further corroborate miR-223 function, we analyzed hemEC and HSPC populations

during mammalian EHT. We observed abundant miR-223 expression in sorted mouse hemECs

(Flk+cKit+CD45-SP+)[23] and endothelial cells (CD31+CD45-SP-) relative to non-endothelial

cells isolated from the embryonic day (e) 10.5 AGM, the primary site of definitive HSPC

production[2,7] (Fig. 2d, e). Mice with a global miR-223 knockout (KO)[24] displayed elevated

hemECs in the e10.5 AGM compared to wild-type (Fig. 2f). Subsequently, HSPCs (cKit+Sca+

Lin-) were expanded in miR-223 KO mice at secondary hematopoietic sites, namely the fetal

liver at e14.5 and the fetal bone marrow at e18.5 (Fig. 2d, f). These results establish miR-223 as

a conserved endothelial inhibitor of HSPC production.

Next, we sought to identify the miR-223-dependent mechanism limiting EHT in hemECs. miRNAs can fine tune gene expression by targeting mRNAs for decay, typically via complementary base-pairing to sites within 3' untranslated regions (3' UTRs), known as miRNA responsive elements (MREs)[25,26]. Therefore, we performed comparative mRNA profiling of zebrafish wild-type and miR-223Δ/Δ endothelial cells at the start of EHT[17], and validated the results by qRT-PCR (Fig. 3a). Amongst the top transcripts most upregulated upon loss of miR-223 and harboring a predicted miR-223 MRE were genes involved in generating N-glycosylated proteins (Fig. 3a, b and Supplementary Fig. 3a).

100   N-glycosylation involves the co-translational attachment and processing of oligosaccharides (glycans) onto asparagine (N) residues of secreted or membrane-bound proteins and regulates protein folding, stability, and activity[9,10] (Fig. 3b and Supplementary Fig. 3a). N-glycosylation begins in the endoplasmic reticulum (ER), where a preassembled dolichol lipid-linked sugar glycan core is transferred *en bloc* onto nascent polypeptide chains. The N-glycan is then progressively processed in the ER and Golgi apparatus as the protein traffics to the cell membrane. The putative miR-223 target *alg2* is an ER mannosyltransferase involved in adding α 1,3 / α 1,6-linked mannose moieties to the N-glycan core[9,27]. Mannose-binding lectin *lman2lb*, another candidate miR-223 target, promotes N-glycoprotein transport from the ER to the Golgi apparatus[28,29]. The putative miR-223 target *st3gal2*, is a sialyltransferase that adds α 2,3-linked sialic acid sugars to the N-glycan in the Golgi apparatus[30-32] (Fig. 3b and Supplementary Fig. 3a).

111   To determine whether miR-223 binds and regulates these three transcripts during EHT, we developed sensors[33] of miR-223 activity consisting of mCherry fused to a short *alg2*, *lman2lb*, or *st3gal2* 3'UTR fragment that contains the putative miR-223 MRE (Supplementary Fig. 3b). Expression of both the mCherry sensor and an eGFP that lacks a 3'UTR (for normalization) was driven from a bidirectional and endothelial-specific *fli1a* promoter. We injected these constructs into wild-type, miR-223Δ/+ and miR-223Δ/Δ zebrafish embryos and observed mCherry and eGFP expression in endothelial cells, as expected. Importantly, the mCherry:eGFP ratios from the *alg2* and *st3gal2* sensors were significantly higher in individual AGM endothelial cells in to in miR-223Δ/+ and miR-223Δ/Δ embryos compared to wild-type embryos during the peak of EHT at 32 hpf (Supplementary Fig. 3c-g). These data suggest that miR-223 directly binds and represses *alg2* and *st3gal2* transcripts in AGM endothelial cells. In contrast, the mCherry:eGFP

ratio from the *lman2lb* sensor was similar in individual AGM endothelial cells from wild-type

and mutant embryos, suggesting indirect regulation.

We generated a stable line of the *alg2* 3'UTR-miR-223 sensor for mutational analysis of

the MRE. CRISPR/Cas9-mediated mutation of the MRE led to an increased mCherry:eGFP ratio

in the AGM (Fig. 3c, d and Supplementary Fig. 3h), consistent with de-repression in miR-

223Δ/Δ . Further, mutation of the endogenous miR-223 MRE within the *alg2* 3'UTR

phenocopied the HSPC overexpansion observed in miR-223Δ/Δ zebrafish, both in the 32 hpf

AGM and the 54 hpf caudal hematopoietic tissue[34] (the hematopoietic site analogous to the

mammalian fetal liver) (Fig. 3e-g). Thus, these data suggest that miR-223 directly inhibits *alg2*

and/or *st3gal2* to limit HSPC production at the site of EHT, potentially via altered N-

glycosylation.

To investigate whether N-glycosylation changes during EHT, we generated an N-

glycosylation reporter. The reporter contains an N-glycosensor[35], consisting of firefly luciferase

fused to an ER translation sequence. ER translocation of firefly luciferase results in its N-

glycosylation and reduced bioluminescence[35] (Fig. 4a). The reporter also contains a renilla

luciferase (for normalization), and both genes are driven by the *fli1a* promoter (Fig. 4a).

Injection of the endothelial-specific N-glycosylation reporter into wild-type embryos revealed a

decrease in firefly luminescence from 24 hpf to 32 hpf, indicating that endothelial cell N-

glycosylation naturally increased from the onset to the peak of EHT (Fig. 4b). In addition, miR-

223Δ/Δ embryos showed increased firefly luminescence at 32 hpf compared to controls (Fig. 4c),

consistent with defective N-glycosylation of firefly luciferase upon dysregulation of miR-223.

Thus, N-glycosylation is regulated by miR-223 during EHT.

The increased luminescence of the N-glycosensor upon loss of miR-223 function could

reflect reduced N-glycan addition or altered N-glycan processing. To distinguish between these

possibilities, we first tested whether blocking N-glycan addition recapitulates HSPC

overexpansion in miR-223Δ/Δ embryos. We treated wild-type zebrafish during EHT with

tunicamycin (TM), an N-acetylglucosamine (GlcNAc)-1-phosphotransferase inhibitor that

prevents N-glycan core attachment to nascent polypeptides, leading to protein misfolding and ER

stress[36,37]. TM treatment increased firefly luminescence as expected[35], but did not result in an

overexpansion of *cmyb+* HSPCs at 32 hpf (Supplementary Fig. 4a, b). Likewise, *cmyb+*

expression did not increase upon treatment with thapsigargin, an N-glycosylation-independent

inducer of ER-stress[36] (Supplementary Fig. 4c). Thus, miR-223Δ/Δ hematopoietic defects do not

arise from an overall reduction in protein N-glycosylation or general ER stress.

To investigate whether N-glycan processing is disrupted by loss of miR-223, we used a

ratiometric lectin microarray strategy to analyze the glycomes of wild-type and mutant zebrafish

embryos[38]. We isolated membrane proteins from 32 hpf wild-type and miR-223Δ/Δ embryos,

and deciphered differences in specific sugar modifications based on their differential binding to

over 90 lectins or antibodies on the array[39]. Notably, glycoproteins from miR-223Δ/Δ embryos

had significantly increased levels of terminal α1,3-linked mannose modifications, as well as

decreased levels of α2,3-linked sialic acid and bisecting GlcNAc, compared to those from wild-

type (Fig. 4d, e). Thus, loss of miR-223 affects N-glycan processing.

Next, we determined whether compounds that alter N-glycan processing could promote

EHT and HSPC overexpansion, similar to loss of miR-223. Briefly, we treated wild-type

embryos undergoing EHT with swainsonine (SW) to enhance mannose-biased modification[37], or

a pan-sialyltransferase inhibitor (ST) to diminish terminal sialic acid incorporation[40] (Fig. 4 f-j

and Supplementary Fig. 4d). Remarkably, both SW and ST treatments of wild-type embryos

resulted in the overexpansion of kdrl:mCH+ cmyb:GFP+ cells and *cmyb+* nascent HSPCs in the

AGM at 32 hpf, recapitulating the miR-223Δ/Δ phenotype (Fig. 2c, Fig. 4g, h, j, and

Supplementary Fig. 4d). Taken together, we find that a distinct N-glycome established by miR-

223 limits the production of HSPCs from hemECs (Fig. 4k).

In summary, our data reveal an unanticipated role for the N-glycan biosynthesis pathway

in limiting HSPC production during EHT, thus illuminating how this process occurs. We found

that hemECs in the AGM possess an intrinsic mechanism to restrict EHT based on a "sugar

code", comprised of a specific level of cell surface glycans, α1,3 mannose and α2,3 sialic acid.

This regulation is mediated by miR-223-dependent repression of N-glycan biosynthesis genes,

which regulates the glycan repertoire (Fig. 4k).

Importantly, these findings lay the foundation for future mechanistic studies on how the

diverse array of hematopoietic regulators are limited in their activity by a common co-

translational modification, protein N-glycosylation. The discovery that EHT is regulated by a

specific glycan profile presents new exciting avenues for regulating *in vivo* and *ex-vivo* blood

stem cell production[41,42]. The wide variety of glycan metabolism modifiers and inhibitors could

be used to glyco-engineer cells, optimizing the production of glycoforms that facilitate the

reprogramming of somatic cells to HSPCs[43-45]. Thus, the application of our knowledge and

approaches could significantly improve HSPC production for therapeutic interventions of blood

disorders.

Methods

Animal Strains and Husbandry

Zebrafish were raised and maintained at 28.5˚C using standard methods (unless otherwise

indicated), and according to protocols approved by Yale University Institutional Animal Care

and Use Committee (# 2017-11473). AB was used as the wild type genetic background.

Zebrafish transgenic strains generated in this study are: *Tg(miR-223:KalTA4)^{ya362}*(wt

background)*; Tg(miR-223:KalTA4)^{ya363}* (miR-223^{ya303} background)*, Tg(fli1ep:mCherry-alg2

3'UTR sensor: eGFP)^{ya364}* (referred to as *alg2* 3'UTR-miR-223 MRE sensor). The following

lines have been described previously: miR-223^{ya303} and miR-223^{ya304} (referred to as miR-

223Δ/Δ)[17], *Tg(kdrl:gfp)^{la116}* (referred to as kdrl:GFP+)[46], *Tg(gata2b:KalTA4)^{sd32}* coupled with

Tg(UAS:lifeact-GFP)^{mu271} (referred to as gata2b:GFP+)[19]*; Tg(kdrl:hras-mCherry)^{s896}* (referred to

as kdrl:mCH)[6]*; Tg(kdrl:mCherry)^{is5} Tg(cmyb:GFP)^{zf169}* (referred to as kdrl:mcH+cmyb:GFP+)[6]*,

Tg(5xUAS:eGFP)^{nkuasgfp1a}* (referred to as UAS:GFP)[47]. For clarity, *Tg(miR-223:kalTA4)* is

referred to as miR-223:Gal4+ or when coupled with *Tg(5xUAS:eGFP)^{nkuasgfp1a}* is called miR-

223:GFP+. Zebrafish embryos and adults were genotyped as described[17]. See Table 1 for

genotyping primers.

All mouse experiments were approved by Yale University Institutional Animal Care and

Use Committee (#2017-11447). Mouse stains used in this study were wild-type (Jackson

#002014) and miR-223KO (Jackson #013198) in the C57BL/6 background. Mouse embryos

were generated from timed matings between females and males (>8 weeks). The morning of

detection of vaginal plug was defined as e0.5.

Generation of miR-223:Gal4 reporter zebrafish

To generate the miR-223:Gal4 reporter line, the improved Gal4 variant KalTA4[48] was inserted

into the endogenous miR-223^{+} (wild-type) or miR-223^{ya303} genomic locus on Chromosome 5

using CRISPR/Cas9 genomic editing according to published protocols with the following

modifications[18]. The KalTA4 integration site was designed approximately 50 bp upstream of the

miR-223 precursor stem loop genomic sequence (miRBase Accession # MI0001389). A ~300 bp

fragment surrounding this site was PCR amplified with flanking BamHI restriction sites and

cloned into the eGFP bait-E2A-KalTA4-pA plasmid[49] by T4 ligation according to the

manufacturer's protocols (NEB), to produce the miR-223-E2A-KalTA4 bait plasmid.

CRISPRScan[50,51] was then used to design a guide RNA (gRNA) that targets both the miR-223

genomic locus and miR-223-E2A-KalTA4 bait plasmid. The gRNA template for in vitro

transcription was PCR amplified and purified with Qiaquick PCR purification Kit (Qiagen). In

vitro transcription followed and was performed as described[52]. High gRNA targeting efficiency

at the miR-223 locus was confirmed by T7 endonuclease I (T7E1) in individual AB embryos

injected with 60 pg of miR-223 gRNA and 200 pg of Cas9 mRNA as before[52]. See Table 1 for

cloning and gRNA PCR primers.

To establish the stable miR-223:Gal4 reporter lines, *Tg(5xUAS:eGFP)[nkuasgfp1a]* transgenic

embryos were injected in the one-cell with 30 pg of miR-223-E2A-KalTA4 bait plasmid, 60 pg

of miR-223 gRNA, 200 pg of Cas9 mRNA. Injected embryos with eGFP fluorescent cells were

raised to adult hood and then outcrossed to establish the wildtype F1 founder *Tg(miR-*

223:KalTA4)[ya362]. F1 founders were then established in the *Tg(kdrl:hras-mCherry)[s896]*

background. The miR-223 mutant founder *Tg(miR-223:KalTA4)[ya363]* was independently

generated in a similar fashion except embryos from *Tg(5xUAS:eGFP)[nkuasgfp1a]* and miR-223[ya303]

outcrosses were injected, and the KalTA4 integration on the mutant allele was confirmed.

Quantitative RT-PCR

Embryos were disassociated into single cell suspensions and subjected to Fluorescence Activated

Cell Sorting (FACS) as described for transgenic zebrafish previously[53] or for mice below. 300-

500 ng of total RNA isolated from the indicated FAC-sorted populations was used in miRNA or

mRNA reverse transcription reactions and the resulting cDNA was used as template for SYBR

Green-based quantitative PCR as before[17]. qRT-PCR primers are listed in Table 1. U6 primers

(RNU6B, Hs_RNU6-2_1) were commercially provided from Qiagen.

The $2^{-\Delta\Delta CT}$ method was used to determine relative gene expression for quantitative RT-

PCR analyses. Mature miRNA expression was normalized to U6 snRNA levels and was relative

to the indicated control. mRNA levels were normalized to the beta actin housekeeping

gene, *actb1* and was relative to the indicated control. Statistical comparisons between replicate

pair ΔCT values for indicated groups were determined by a paired, two-tailed Student's t-test.

Immunofluorescence

Immunofluorescence was performed as follows to amplify GFP and mCherry expression in

zebrafish embryos imaged by confocal microscopy unless otherwise indicated. After overnight

4% formaldehyde fixation at 4˚C, 24, 27, 32, or 48 hpf embryos were washed with 1XPBS-

0.01%Tween-20 (PBSTw) 4 times for 5 minutes, and then permeabilized with 0.125% trypsin

(Millipore Sigma T4549) for 1, 3, 5, or 10 minutes, respectively. Embryos were washed in

blocking solution (0.8% Triton-X, 10% normal goat serum, 1% BSA, 0.01% sodium azide in

PBSTw) 3 times for 5 minutes, plus an additional 2-hour incubation with shaking. Antibody

concentrations used were 1:300 chicken anti-GFP (Abcam, ab13970) or rabbit anti-RFP

(Anitbodies Online, ABIN129578) primary antibodies and 1:400 Alexa Fluor 488 goat anti–

chicken IgG or Alexa Fluor 546 donkey anti-rabbit IgG (ThermoFisher) secondary antibodies.

Following each overnight antibody incubation at 4˚C, six washes for a total of 4 hours were

performed with blocking solution lacking goat serum at room temperature.

Immunofluorescent confocal micrographs were blinded and quantified for miR-

223:GFP+ kdrl:mCH+, kdrl:mCH+ gata2b:GFP, or kdrl:mCH+ cmyb:GFP cell numbers or *alg2*

3'UTR-miR223 MRE sensor expression (see below). Datasets that were normally distributed as

determined by the Shapiro-Wilk normality test were statistically evaluated by a Student's T-test;

otherwise, a non-parametric Mann-Whitney U test was applied.

Whole mount in situ hybridization

Whole mount in situ hybridization (WISH) was performed as previously described[17] using

hybridization probes for *gata2b*[19], *runx1*[54], and *cmyb*[55]. *gata2b* and *runx1* WISH staining was

quantified from equally-exposed, color-inverted bright field images with Image J as follows. The

Polygon Selection tool was used to outline the area of WISH staining along the entire length of

the yolk extension, corresponding to the AGM region. The mean WISH signal for the selected

area was obtained using the Measure tool and subtracted for mean background staining, which

was measured for the same area, but in a region of the trunk just above (dorsal) the AGM, which

lacks *gata2b* or *runx1* expression. Measurements from 2 biological replicates per genotype were

combined and an unpaired, two-tailed Student's t-test was applied to detect statistical differences

between conditions.

Images of *cmyb* WISH staining in the 32 hpf AGM and 54 hpf CHT for at least 12

embryos per replicate were blinded and then were categorized as having continuous (high) or

no/intermittent (low/medium) *cmyb* expression in the AGM region along the yolk extension.

Data was expressed as the percentage of embryos having high or low/medium staining,

respectively. Statistical differences between wild-type and miR-223Δ/Δ replicate pairs (collected

and stained on the same days) were determined by a paired, two-tailed Student's t-test.

Flow Cytometry in Mice

Mouse embryo dissection, cell isolation and flow cytometry of E10.5 AGM were performed as

previously described [56]. Briefly, developmental stage-matched miR-223KO and wild-type e10.5

AGM regions were freshly dissected and digested in 2mg/mL collagenase type II at 37 °C for

20 min in HBSS (Gibco) with 10% FBS (Gibco or Gemini) to obtain single-cell suspensions.

Cells were resuspended in DMEM (Gibco) with 10% FBS to a concentration of 2 x 10^6 cells/ml,

and incubated for 1 hour in 5 μg/mL Hoechst 33342 (Sigma) at 37 °C, followed by staining of

the following antibodies: CD45 (FITC, eBioscience 11-0451-85), c-Kit(APC, BD 553356),

Flk1(PE-Cy7, BD 561259), CD31 (PE, BD 553373). Single cell suspensions of e14.5 fetal liver

and e18.5 fetal bone marrow were immunostained with the following antibodies: c-Kit (PE,

eBioscience 12-1172-82), Sca-1 (FITC, eBioscience 11-5981-82), lineage antibody cocktail

(PerCP-Cy5.5, BD 561317). Cells were analyzed or sorted by flow cytometer LSRII or Aria (BD

Bioscience) respectively and flow cytometry data were analyzed with FlowJo software. Student's

t-test (unpaired, two tailed) was used to compare miR-223KO and wild-type relative cell

abundance, and data was charted as fold change relative to wild-type.

miR-223 expression constructs

miR-223 expression constructs (Supplementary Fig. 2) were constructed as previously described

with the following modifications[33]. To generate the pME-miR-223 middle entry cassette for

Gateway-compatible cloning, a 365 bp genomic sequence containing the miR-223 stem loop

precursor was PCR amplified with flanking KpnI and StuI sites (see Table 1), and the resulting

fragment was restriction digested and cloned into pME-miR using T4 ligation (NEB) according

to the manufacturer's protocol. Promoter entry cassettes p5E-*uasE1b* or p5E-*fli1a*[57] (Addgene

26026 and 31160) were utilized in LR multisite Gateway cloning reactions as before[33] to produce

UAS-mCherry-miR-223 and *fli1a*-mCherry-miR-223 constructs, respectively. Embryos were

injected in the one-cell with 25 pg each of either expression construct and Tol2 transposase

mRNA, and later selected for mCherry expression.

Endothelial miR-223 sensor assays & zebrafish generation

Endothelial N-glycogene 3'UTR-miR-223 MRE sensors were constructed as previously

described except the p3E entry vector was generated by subcloning a short (~88-117 bp) PCR

fragment of the *alg2*, *lman2lb*, or *st3gal2* 3'UTR that contains the miR-223 responsive element

using flanking SnaBI and SpeI restriction sites[33]. See Table 1 for N-glycogene 3'UTR cloning

primers. Mosaic expression of each sensor (Supplementary Fig. 3c, e-g) was achieved by

injecting single-cell embryos from miR-223Δ/+ in-crosses with 25 pg each of the sensor

construct and Tol2 transposase mRNA. For each expression construct, live injected embryos

were confocal imaged at ~32hpf using a constant exposure setting and then genotyped. Using

ImageJ on genotype-blinded maximum confocal projections, individual AGM endothelial cells

were first outlined using the Polygon Selection tool. Mean mCherry and eGFP intensities were

then quantified for the selected area using the Measure tool and were subtracted for mean image

background fluorescence in their respective channel. mCherry relative to eGFP intensities were

assessed for each genotype. Outlier values were removed using the ROUT method (Q= 1%) in

Graphpad Prism7. A Mann-Whitney U test was used for statistical comparisons.

To establish the stably-transmitting *Tg(fli1a:mCherry-alg2 3'UTR sensor: eGFP)*[ya364]

line (Fig. 3c), embryos injected with 25 pg eaph of Tol2 transposase and *alg2* 3'UTR-miR-223

MRE sensor construct were raised to adulthood and an F1 founder was the crossed into both

wild-type and miR-223Δ/Δ backgrounds. Sensor expression was quantified with Image J as

described above for 32 hpf fixed and immunostained embryos, except the Segmented Line tool

was used to measure mCherry and eGFP intensities along length of the dorsal aorta or posterior

cardinal vein vessel walls above the yolk extension. An unpaired, two-tailed Student's test was

used for statistical comparisons.

alg2 **3'UTR gRNA generation & injection**

CRISPRScan[50,51] was used to design two gRNAs to mutate the miR-223 responsive element

region in the *alg2* 3'UTR with CRISPR/Cas9 genome editing. gRNA preparation was performed

as above for miR-223:Gal4 zebrafish generation. gRNA templates were combined prior to in

vitro transcription. F0 embryos injected with 100 pg of the *alg2* 3'UTR gRNA mix and 200 pg of

Cas9 mRNA at the one-cell stage were used to confirm mutagenesis via T7 endonuclease I

(T7E1) assay and sequencing as described[52] or analyzed in phenotypic assays.

Endothelial N-glycosensor assays

The previously published firefly luciferase N-glycosensor[35] was modified to allow for its co-

expression with a renilla luciferase normalization control as follows. The DNA fragment

containing the EGFR ER translation sequence and Firefly luciferase gene with three N-

glycosylation sites was subcloned from the pcDNA3-ERlucT vector[35] into multiple cloning site

(MCS) A of the pIRESneo vector (Clontech) using NotI and BamHI sites. The resulting plasmid

was further modified by replacing the Neomycin resistant gene, located downstream of MCS A

and the internal ribosome entry site (IRES), with the Renilla luciferase gene from the pRL-CMV

vector (Promega) using SmaI and XbaI sites. The ERlucT-IRES-Rluc bicistronic expression

cassette was then PCR amplified using primers listed in Table 1 and then subcloned using the

pCR8/GW/TOPO TA Cloning Kit (Invitrogen), to produce the Gateway cloning-compatible

ERlucT-IRES-Rluc-pCR8 middle entry vector. To generate the endothelial N-glycosensor

expression construct, Gateway entry vectors p5E-*fli1a*[57] (Addgene 31160), ERlucT-IRES-Rluc-

pCR8, and p3E-793[33], were recombined into destination vector pDestTol2pA[58] using LR

Clonase II Multisite Gateway cloning (Invitrogen) according to the manufacturer's instructions.

For endothelial N-glycosensor assays, embryos were injected with 25 pg each of the

Tg(fli1a:ERlucT-IRES-Rluc:pDestTol2A) construct and Tol2 transposase into the one-cell and

grown to the indicated developmental stages. A Dual Luciferase Reporter Assay (Promega) was

then performed according to manufacturer's guidelines. Briefly, 25 dechorionated embryos per

condition were washed once with Dulbecco's PBS (Gibco) and then incubated shaking at 200

rpm in 25 µl of the provided 1X Passive Lysis Buffer for 30 minutes. Embryo extracts were

homogenized with a pellet pestle and spun at 13,000 rpm to remove cellular debris. Firefly and

Renilla bioluminescence readings were obtained from 20 µl of cleared lysate using the Synergy 2

plate reader and Gen 5 analysis software system (BioTek). Firefly bioluminescence was

normalized to renilla levels, and expressed as a fold change relative to indicated control.

Statistical differences of normalized firefly bioluminescence values between biological replicate

pairs were determined by a paired, two-tailed Student's t-test.

Lectin Microarray Glycomic Analysis

Lectin microarrays were prepared as previously described[39]. Slides were generated in the Mahal

laboratory using Nano-plotter v2.0 piezoelectric non-contact array printer (GeSiM) with a nano

A-J tip to print each lectin or antibody. Arrays were printed on Nexterion Slide H (Schott

Nexterion) under 50% relative humidity at a surface temperature of 12°C with lectins and

antibodies that were purchased from Vector Labs, R&D Systems, Santa Cruz, TCI, AbCam, E.Y.

Labs, or Sigma-Aldrich on the array. The recombinant lectins rGRFT, rCVN, and rSVN were

generous gifts from Dr. B. O'Keefe (NCI Frederick). SLBR-N, SLBR-H, and SLBR-B[59] were

gifts from Dr. B. Bensing (UCSF).

Membrane protein extracts applied to the array were isolated from 32 hpf wild-type and

miR-223Δ/Δ zebrafish embryos and fluorescently labeled as follows. 400 dechorionated embryos

per replicate sample were washed two times in deyolking buffer (55mM NaCl, 1.8mM KCl,

1.25mM NaHCO$_3$)[60]. The yolk was separated from embryos by gentle pipetting in 250-500 µl of

deyolking buffer, followed by 5 minutes of room temperature shaking at 850 rpm. Samples were

spun at 300 rcf for 30 seconds and the yolk-containing supernatant was discarded. To remove

any remaining yolk, the pelleted tissue was washed with Dulbecco's PBS (Gibco), and shook at

850 rpm for 2 minutes for a total of two times. After a 300g, 5 minute spin at 4°C, nearly all

supernatant was removed and tissue was incubated in 1 ml of cold hypotonic PBS solution

(10mM Na$_2$HPO$_4$, 1.8mM KH$_2$PO$_4$, 10mM KCl, 0.2mM MgCl$_2$) plus 1:100 protease inhibitor

cocktail set III (Millipore Sigma) for 10 minutes on ice to lyse cells. Samples were dounce-

homogenized 150 times with the B pestle on ice, and then spun at 500g for 5' at 4°C to remove

cell debris. Protein lysates were frozen with liquid nitrogen and stored at -80°C until used. To

obtain the membrane protein fraction, thawed lysates were spun at 100 kg for 1 hour at 4°C and

pellets were resuspended immediately in 100 µl of cold PBS (0.1M Na$_2$HPO$_4$, 0.15M NaCl, pH

7.2-7.4) with 1:100 protease inhibitors. Samples were fully homogenized by passing the sample

through a 25G needle 10-15 times on ice.

Protein concentrations for individual samples were determined using a BCA protein
assay following the manufacturer's protocol (Thermo Scientific Pierce). 50 µg of protein were
labeled with Alexa Fluor 555 and all reactions were quenched using 2M Tris (pH 6.8) with no
further dialysis. A mixed reference was prepared following the same protocol, but labeled with
Alexa Fluor 647. Samples for the array were prepared by mixing 5 µg of protein from each
individual sample with 5 µg of the mixed reference in 0.005% PBS-Tween (pH 7.4).

Prior to sample hybridization, the lectin microarray slides were blocked for an hour with
50 mM ethanolamine in 50 mM sodium borate buffer (pH 8.8) at room temperature. They were
then washed 3 times with 0.005% PBS-Tween (pH 7.4). Hybridization cassettes (Arrayit) were
utilized to isolate individual arrays on each slide (24 per slide) and prevent leakage from one
sample to the next. Samples were applied to individual arrays and one array per slide was
reserved for a reference vs reference. Samples were incubated on the array for 2 hours at 25 °C
with gentle agitation. After hybridization, the samples were carefully removed and the arrays
were washed 4 times with 0.005% PBS-Tween (pH 7.4) for 5 minutes followed by a final wash
with ddH2O. The arrays were scanned using a GenePix 4300A array scanner.

Background-subtracted median fluorescence intensities were extracted using GenePix Pro
v7.2. Lectins that did not meet signal to noise ratios <2 in >90% were removed from the data
prior to analysis. The data was then median-normalized for each dye used and the log2 of the
sample/reference ratio was determined for each technical replicate. The replicates were then
averaged for each lectin within each array. The log2 values were then used for statistical
analysis (Student's T-Test) and comparison of the sample types.

Pharmacological inhibitor treatments
Zebrafish embryos were exposed to chemical inhibitors of N-glycosylation at the following
concentrations, 0.75 µg/ml of tunicamycin (Millipore Sigma T7765), 5 µg/ml of swainsonine
(Millipore Sigma S9263), 200 µg/ml of 3Fax-Peracetyl Neu5Ac Sialyltransferase Inhibitor
(Millipore Sigma 566224); or to 0.4 µM of thapsigargin (Millipore Sigma 586006), an N-
glycosylation-independent inducer of ER stress. Equal concentrations of the DMSO vehicle were
applied as controls for each drug as indicated. All treatments were administered from 8-32 hpf in
standard embryo medium with 0.003% 1-phenyl-2-thiourea (PTU) to prevent pigmentation for
downstream analyses.

Image Acquisition

Zebrafish embryos imaged by confocal or bright-field microscopy were raised in 0.003% PTU
starting after the gastrulation stage to prevent pigmentation. Embryos imaged live by confocal
microscopy were anesthetized in 0.1% tricaine and mounted in 1% low melt agarose. The
majority of fluorescent images and timelapse movie were captured using an upright Leica
Microsystems SP5 confocal microscope using a 25X objective except for images in Fig. 3c of the
stable *alg2* 3'UTR-miR-223 MRE sensor line, which were captured with an upright Leica
Microsystems SP8 confocal microscope. Bright field images of whole mount in situ staining
were acquired with a Leica Microsystems M165FC stereomicroscope equipped with Leica
DFC295 camera.

Table 1: Primers used in this study

Primer Name	F/R	Primer Sequence (5'-3')	Assay	Ref
dre-miR-223 gRNA	F	taatacgactcactataggcacctggctcaacatgtcgtttt agagctagaa	gRNA PCR	this study
dre-miR-223 bait BamHI	F	ggccggatccccaaggaacccaatgaactg	miR-223 Gal4 bait cloning	this study
dre-miR-223 bait BamHI	R	ggccggatccctccctcttcctcctgcttt	miR-223 Gal4 bait cloning	this study
dre-miR-223	F	tgtcagtttgtcaaatacccc	qPCR	[17]
dre-*actb1*	F	gatctggcatcacaccttctac	qPCR	[17]
dre-*actb1*	R	tcttctctctgttggctttgg	qPCR	[17]
dre-*kdrl/flk1*	F	cgaacgtgaagtgacatacgg	qPCR	[16]
dre-*kdrl/flk1*	R	ccctctaccaaaccatgtgaaa	qPCR	[16]
dre-*gata2b*	F	accaccacactctggagac	qPCR	[19]
dre-*gata2b*	R	ctgttgcgtgtctgaatacc	qPCR	[19]
dre-*runx1*	F	cgtcttcacaaaccctcctcaa	qPCR	[16]
dre-*runx1*	R	gctttactgcttcatccggct	qPCR	[16]
dre-*cmyb*	F	tgctatccagagacactacagc	qPCR	[17]
dre-*cmyb*	R	agtgcctgttgtcccttcag	qPCR	[17]
dre-*alg2*	F	gcttacctgcgcatgatctac	qPCR	this study
dre-*alg2*	R	atacaggcgcttcagagcag	qPCR	this study
dre-*lman2lb*	F	agtggaacgcagtcctgaag	qPCR	this study
dre-*lman2lb*	R	tgatgctccttccttcatcc	qPCR	this study
dre-*st3gal2*	F	ctggtgggtgatgttgcag	qPCR	this study
dre-*st3gal2*	R	cacgaaccgtatggggaac	qPCR	this study
dre-miR-223	F	ggccaggcctccactcaaggtctcgcttt	expression	this study

StuI				construct cloning	
dre-miR-223 KpnI	R	ggccggtaccgagagtaacaggatcgcatgg		expression construct cloning	this study
dre-*alg2* 3'UTR	F	ccggtacgtagttaggacactgaaggtcatgtgctgtattg ggtttgaggttgttactgactgtgaatctcctcgtgctattga tattagactgaagacccgaactagtggcc		miR-223 sensor cloning	this study
dre-*alg2* 3'UTR	R	ggccactagttcgggtcttcagtctaatatcaatagcacga ggagattcacagtcagtaacaacctcaaacccaatacagc acatgaccttcagtgtcctaactacgtaccgg		miR-223 sensor cloning	this study
dre-*lman2lb* 3'UTR	F	ccggtacgtacagatatttgtgcgtttttctgtaatttttgtacc aattacactgaccttggcctttctccctgtgacagcatttaca gaggctaaagagttcaacgtcaactgtactagtggcc		miR-223 sensor cloning	this study
dre-*lman2lb* 3'UTR	R	ggccactagtacagttgacgttgaactctttagcctctgtaa atgctgtcacagggagaaaggccaaggtcagtgtaattg gtacaaaaattacagaaaaacgcacaaatatctgtacgtac cgg		miR-223 sensor cloning	this study
dre-*st3gal2* 3'UTR	F	ccggtacgtagatcattcctctgctctgaggaggagggat agattgatactcaataccagaaactgaaaaccactgacac acagtcactagtggcc		miR-223 sensor cloning	this study
dre-*st3gal2* 3'UTR	R	ggccactagtgactgtgtgtcagtggttttcagtttctggtat tgagtatcaatctatccctcctcctcagagcagaggaatga tctacgtaccgg		miR-223 sensor cloning	this study
ERlucT-IRES-Rluc	F	gccaccatgcgaccctcc		Nglyco sensor cloning	this study
ERlucT-IRES-Rluc	R	ttattgttcattttttgagaactcgctc		Nglyco sensor cloning	this study
dre-*alg2* 223MRE gRNA-1	F	taatacgactcactataggaaggtcatgtgctgtattgtttta gagctagaa		gRNA PCR	this study
dre-*alg2* 223MRE gRNA-2	F	taatacgactcactataggtctaatatcaatagcacggtttta gagctagaa		gRNA PCR	this study
gRNA universal primer	R	aaaagcaccgactcggtgccacttttttcaagttgataacgg actagccttattttaacttgctatttctagctctaaaac		gRNA PCR	[52]
dre-*alg2* 223MRE	F	cagcgtctcagcttttgtttag		genotyping	this study
dre-*alg2* 223MRE	R	accgttccatctggtcactc		genotyping	this study
dre-miR-223 geno	F	6FAM- ctgctgcaaacctgcattta		genotyping	this study
dre-miR-223 geno	R	gagagtaacaggatcgcatgg		genotyping	[17]

References

Clements, W. K. & Traver, D. Signalling pathways that control vertebrate haematopoietic stem cell specification. *Nat Rev Immunol* **13**, 336-348, doi:10.1038/nri3443 (2013).

Gritz, E. & Hirschi, K. K. Specification and function of hemogenic endothelium during embryogenesis. *Cell Mol Life Sci* **73**, 1547-1567, doi:10.1007/s00018-016-2134-0 (2016).

Medvinsky, A., Rybtsov, S. & Taoudi, S. Embryonic origin of the adult hematopoietic system: advances and questions. *Development* **138**, 1017-1031, doi:10.1242/dev.040998 (2011).

Jagannathan-Bogdan, M. & Zon, L. I. Hematopoiesis. *Development* **140**, 2463-2467, doi:10.1242/dev.083147 (2013).

Kasper, D. M. & Nicoli, S. Epigenetic and Epitranscriptomic Factors Make a Mark on Hematopoietic Stem Cell Development. *Curr Stem Cell Rep* **4**, 22-32 (2018).

Bertrand, J. Y. *et al.* Haematopoietic stem cells derive directly from aortic endothelium during development. *Nature* **464**, 108-111, doi:10.1038/nature08738 (2010).

Boisset, J. C. *et al.* In vivo imaging of haematopoietic cells emerging from the mouse aortic endothelium. *Nature* **464**, 116-120, doi:10.1038/nature08764 (2010).

Kissa, K. & Herbomel, P. Blood stem cells emerge from aortic endothelium by a novel type of cell transition. *Nature* **464**, 112-115, doi:10.1038/nature08761 (2010).

Stanley, P., Taniguchi, N. & Aebi, M. in *Essentials of Glycobiology* (eds rd *et al.*) 99-111 (2015).

Moremen, K. W., Tiemeyer, M. & Nairn, A. V. Vertebrate protein glycosylation: diversity, synthesis and function. *Nat Rev Mol Cell Biol* **13**, 448-462, doi:10.1038/nrm3383 (2012).

Pinho, S. S. & Reis, C. A. Glycosylation in cancer: mechanisms and clinical implications. *Nat Rev Cancer* **15**, 540-555, doi:10.1038/nrc3982 (2015).

Lanctot, P. M., Gage, F. H. & Varki, A. P. The glycans of stem cells. *Curr Opin Chem Biol* **11**, 373-380, doi:10.1016/j.cbpa.2007.05.032 (2007).

McKinney-Freeman, S. *et al.* The transcriptional landscape of hematopoietic stem cell ontogeny. *Cell Stem Cell* **11**, 701-714, doi:10.1016/j.stem.2012.07.018 (2012).

Gerri, C., Marass, M., Rossi, A. & Stainier, D. Y. R. Hif-1alpha and Hif-2alpha regulate hemogenic endothelium and hematopoietic stem cell formation in zebrafish. *Blood* **131**, 963-973, doi:10.1182/blood-2017-07-797795 (2018).

15   Gore, A. V. *et al.* Epigenetic regulation of hematopoiesis by DNA methylation. *Elife* **5**,
e11813, doi:10.7554/eLife.11813 (2016).

16   North, T. E. *et al.* Hematopoietic stem cell development is dependent on blood flow. *Cell*
137, 736-748, doi:10.1016/j.cell.2009.04.023 (2009).

17   Kasper, D. M. *et al.* MicroRNAs Establish Uniform Traits during the Architecture of
Vertebrate Embryos. *Dev Cell* **40**, 552-565 e555, doi:10.1016/j.devcel.2017.02.021
(2017).

18   Auer, T. O., Duroure, K., Concordet, J. P. & Del Bene, F. CRISPR/Cas9-mediated
conversion of eGFP- into Gal4-transgenic lines in zebrafish. *Nat Protoc* **9**, 2823-2840,
doi:10.1038/nprot.2014.187 (2014).

19   Butko, E. *et al.* Gata2b is a restricted early regulator of hemogenic endothelium in the
zebrafish embryo. *Development* **142**, 1050-1061, doi:10.1242/dev.119180 (2015).

20   Chen, M. J., Yokomizo, T., Zeigler, B. M., Dzierzak, E. & Speck, N. A. Runx1 is
required for the endothelial to haematopoietic cell transition but not thereafter. *Nature*
457, 887-891, doi:10.1038/nature07619 (2009).

21   Wilkinson, R. N. *et al.* Hedgehog and Bmp polarize hematopoietic stem cell emergence
in the zebrafish dorsal aorta. *Dev Cell* **16**, 909-916, doi:10.1016/j.devcel.2009.04.014
(2009).

22   Zhang, Y., Jin, H., Li, L., Qin, F. X. & Wen, Z. cMyb regulates hematopoietic
stem/progenitor cell mobilization during zebrafish hematopoiesis. *Blood* **118**, 4093-4101,
doi:10.1182/blood-2011-03-342501 (2011).

23   Marcelo, K. L. *et al.* Hemogenic endothelial cell specification requires c-Kit, Notch
signaling, and p27-mediated cell-cycle control. *Dev Cell* **27**, 504-515,
doi:10.1016/j.devcel.2013.11.004 (2013).

24   Johnnidis, J. B. *et al.* Regulation of progenitor cell proliferation and granulocyte function
by microRNA-223. *Nature* **451**, 1125-1129, doi:10.1038/nature06607 (2008).

25   Baek, D. *et al.* The impact of microRNAs on protein output. *Nature* **455**, 64-71,
doi:10.1038/nature07242 (2008).

26   Bartel, D. P. Metazoan MicroRNAs. *Cell* **173**, 20-51, doi:10.1016/j.cell.2018.03.006
(2018).

27   Huffaker, T. C. & Robbins, P. W. Yeast mutants deficient in protein glycosylation. *Proc*
Natl Acad Sci U S A **80**, 7466-7470 (1983).

28   Kamiya, Y. *et al.* Molecular basis of sugar recognition by the human L-type lectins
ERGIC-53, VIPL, and VIP36. *J Biol Chem* **283**, 1857-1861,
doi:10.1074/jbc.M709384200 (2008).

29 Gomez-Navarro, N. & Miller, E. Protein sorting at the ER-Golgi interface. *J Cell Biol*
215, 769-778, doi:10.1083/jcb.201610031 (2016).

30 Hemmoranta, H. *et al.* N-glycan structures and associated gene expression reflect the
characteristic N-glycosylation pattern of human hematopoietic stem and progenitor cells.
Exp Hematol **35**, 1279-1292, doi:10.1016/j.exphem.2007.05.006 (2007).

31 Sturgill, E. R. *et al.* Biosynthesis of the major brain gangliosides GD1a and GT1b.
Glycobiology **22**, 1289-1301, doi:10.1093/glycob/cws103 (2012).

32 Giordanengo, V. *et al.* Cloning and expression of cDNA for a human Gal(beta1-
3)GalNAc alpha2,3-sialyltransferase from the CEM T-cell line. *Eur J Biochem* **247**, 558-
566 (1997).

33 Nicoli, S. *et al.* MicroRNA-mediated integration of haemodynamics and Vegf signalling
during angiogenesis. *Nature* **464**, 1196-1200, doi:10.1038/nature08889 (2010).

34 Murayama, E. *et al.* Tracing hematopoietic precursor migration to successive
hematopoietic organs during zebrafish development. *Immunity* **25**, 963-975,
doi:10.1016/j.immuni.2006.10.015 (2006).

35 Contessa, J. N. *et al.* Molecular imaging of N-linked glycosylation suggests glycan
biosynthesis is a novel target for cancer therapy. *Clin Cancer Res* **16**, 3205-3214,
doi:10.1158/1078-0432.CCR-09-3331 (2010).

36 DeRossi, C. *et al.* trappc11 is required for protein glycosylation in zebrafish and humans.
Mol Biol Cell **27**, 1220-1234, doi:10.1091/mbc.E15-08-0557 (2016).

37 Esko, J. D., Bertozzi, C. & Schnaar, R. L. in *Essentials of Glycobiology* (eds rd *et al.*)
701-712 (2015).

38 Pilobello, K. T., Slawek, D. E. & Mahal, L. K. A ratiometric lectin microarray approach
to analysis of the dynamic mammalian glycome. *Proc Natl Acad Sci U S A* **104**, 11534-
11539, doi:10.1073/pnas.0704954104 (2007).

39 Pilobello, K. T., Agrawal, P., Rouse, R. & Mahal, L. K. Advances in lectin microarray
technology: optimized protocols for piezoelectric print conditions. *Curr Protoc Chem*
Biol **5**, 1-23, doi:10.1002/9780470559277.ch120035 (2013).

40 Wang, L., Liu, Y., Wu, L. & Sun, X. L. Sialyltransferase inhibition and recent advances.
Biochim Biophys Acta **1864**, 143-153, doi:10.1016/j.bbapap.2015.07.007 (2016).

41 Perlin, J. R., Robertson, A. L. & Zon, L. I. Efforts to enhance blood stem cell
engraftment: Recent insights from zebrafish hematopoiesis. *J Exp Med* **214**, 2817-2827,
doi:10.1084/jem.20171069 (2017).

42   Rowe, R. G., Mandelbaum, J., Zon, L. I. & Daley, G. Q. Engineering Hematopoietic
Stem Cells: Lessons from Development. *Cell Stem Cell* **18**, 707-720,
doi:10.1016/j.stem.2016.05.016 (2016).

43   Sminia, T. J., Zuilhof, H. & Wennekes, T. Getting a grip on glycans: A current overview
of the metabolic oligosaccharide engineering toolbox. *Carbohydr Res* **435**, 121-141,
doi:10.1016/j.carres.2016.09.007 (2016).

44   Zhang, P. *et al.* Challenges of glycosylation analysis and control: an integrated approach
to producing optimal and consistent therapeutic drugs. *Drug Discov Today* **21**, 740-765,
doi:10.1016/j.drudis.2016.01.006 (2016).

45   Boheler, K. R. & Gundry, R. L. Concise Review: Cell Surface N-Linked Glycoproteins
as Potential Stem Cell Markers and Drug Targets. *Stem Cells Transl Med* **6**, 131-138,
doi:10.5966/sctm.2016-0109 (2017).

46   Choi, J. *et al.* FoxH1 negatively modulates flk1 gene expression and vascular formation
in zebrafish. *Dev Biol* **304**, 735-744, doi:10.1016/j.ydbio.2007.01.023 (2007).

47   Asakawa, K. *et al.* Genetic dissection of neural circuits by Tol2 transposon-mediated
Gal4 gene and enhancer trapping in zebrafish. *Proc Natl Acad Sci U S A* **105**, 1255-1260,
doi:10.1073/pnas.0704963105 (2008).

48   Distel, M., Wullimann, M. F. & Koster, R. W. Optimized Gal4 genetics for permanent
gene expression mapping in zebrafish. *Proc Natl Acad Sci U S A* **106**, 13365-13370,
doi:10.1073/pnas.0903060106 (2009).

49   Auer, T. O., Duroure, K., De Cian, A., Concordet, J. P. & Del Bene, F. Highly efficient
CRISPR/Cas9-mediated knock-in in zebrafish by homology-independent DNA repair.
Genome Res **24**, 142-153, doi:10.1101/gr.161638.113 (2014).

50   Moreno-Mateos, M. A. *et al.* CRISPRscan: designing highly efficient sgRNAs for
CRISPR-Cas9 targeting in vivo. *Nat Methods* **12**, 982-988, doi:10.1038/nmeth.3543
(2015).

51   Vejnar, C. E., Moreno-Mateos, M. A., Cifuentes, D., Bazzini, A. A. & Giraldez, A. J.
Optimized CRISPR-Cas9 System for Genome Editing in Zebrafish. *Cold Spring Harb*
Protoc **2016**, doi:10.1101/pdb.prot086850 (2016).

52   Narayanan, A. *et al.* In vivo mutagenesis of miRNA gene families using a scalable
multiplexed CRISPR/Cas9 nuclease system. *Sci Rep* **6**, 32386, doi:10.1038/srep32386
(2016).

53   Ristori, E. & Nicoli, S. miRNAs expression profile in zebrafish developing vessels.
Methods Mol Biol **1214**, 129-150, doi:10.1007/978-1-4939-1462-3_7 (2015).

54   Burns, C. E., Traver, D., Mayhall, E., Shepard, J. L. & Zon, L. I. Hematopoietic stem cell
581   fate is established by the Notch-Runx pathway. *Genes Dev* **19**, 2331-2342,
582   doi:10.1101/gad.1337005 (2005).

583   55   Thompson, M. A. *et al.* The cloche and spadetail genes differentially affect
584   hematopoiesis and vasculogenesis. *Dev Biol* **197**, 248-269, doi:10.1006/dbio.1998.8887
585   (1998).

586   56   Fang, J. S., Gritz, E. C., Marcelo, K. L. & Hirschi, K. K. Isolation of Murine Embryonic
587   Hemogenic Endothelial Cells. *J Vis Exp*, doi:10.3791/54150 (2016).

588   57   Villefranc, J. A., Amigo, J. & Lawson, N. D. Gateway compatible vectors for analysis of
589   gene function in the zebrafish. *Dev Dyn* **236**, 3077-3087, doi:10.1002/dvdy.21354 (2007).

590   58   Kwan, K. M. *et al.* The Tol2kit: a multisite gateway-based construction kit for Tol2
591   transposon transgenesis constructs. *Dev Dyn* **236**, 3088-3099, doi:10.1002/dvdy.21343
592   (2007).

593   59   Bensing, B. A., Li, Q., Park, D., Lebrilla, C. B. & Sullam, P. M. Streptococcal Siglec-like
594   adhesins recognize different subsets of human plasma glycoproteins: implications for
595   infective endocarditis. *Glycobiology* **28**, 601-611, doi:10.1093/glycob/cwy052 (2018).

596   60   Link, V., Shevchenko, A. & Heisenberg, C. P. Proteomics of early zebrafish embryos.
597   *BMC Dev Biol* **6**, 1, doi:10.1186/1471-213X-6-1 (2006).
598
599

Author Contributions

D.M.K. and S.N. designed and conducted experiments, analyzed data, and wrote the manuscript. H.K.M, K.S., J.G., J.H., and W.A. carried out zebrafish experiments and analyzed data. K.K.H., Y.W. and Z.H. designed and performed mouse phenotypic experiments and analyses. L.K.M. and D.W.H. conducted lectin microarray analysis. W.C.S. and EJ. P. created and shared the N-glycosensor. All authors edited the paper.

Acknowledgments

The authors would like to thank Meredith Cavanaugh for fish husbandry and care. We would also like to acknowledge Angela Anderson, Valentina Greco, Daniel DiMaio, Liana C. Boraas, and Siyuan Chen for critical reading of the manuscript. This work was supported by grants from the NIH (R01HL130246 and R56DK118728 to S.N. and F32HL132475, U54DK106857, and 1K99HL141687 to D.K.

Competing interests

The authors declare no competing interests.

Corresponding authors
Correspondence to Stefania Nicoli and Karen K. Hirschi.

620   **Figure 1**

c

d

621
Figure 1. miR-223 is expressed in hemogenic endothelial cells undergoing EHT
a, *top,* Schematic of miR-223:Gal4 reporter integrated into the endogenous miR-223 locus. Gal4
activates UAS:GFP to fluorescently mark miR-223-expressing cells (miR-223:GFP+). *bottom,*
Lateral maximum confocal projections of the zebrafish AGM. White and yellow arrows point to
flat and budding miR-223:GFP+; kdrl:mCH+ endothelial cells in the VDA, respectively. **b,**
Overview of zebrafish EHT. In the AGM region, flattened endothelial cells (red) gain hemogenic
potential (orange), round up, and bud as nascent HSPCs (yellow) from the VDA wall into the
PCV circulation from 24-48 hpf. Cells outlined in green or red represent miR-223:GFP+
kdrl:mCH+ or miR-223:GFP- kdrl:mCH+ endothelial cells, respectively. **c,** Mean ± SEM
number of miR-223:GFP+; kdrl:mCH+ endothelial cells in the VDA of the AGM compared to
other regions in the DA and in the PCV (n = 36-43 embryos). Significance determined by
unpaired, two-tailed Mann-Whitney U test. **d,** Mean ± SEM fold change expression in miR-
223:GFP+ kdrl:mCH+ endothelial cells relative to miR-223:GFP- kdrl:mCH- non-endothelial
cells at 27 hpf by qRT-PCR (n = 3 technical replicates). Significance determined by a paired,
two-tailed, Student's t-test. For all plots, significance calculations are represented as [**]p ≤
0.01, [***]p ≤ 0.01, [****]p ≤ 0.0001 Abbreviations: AGM, aorta-gonad mesonephros; DA, dorsal
aorta; EC, endothelial cell; EHT, endothelial-to-hematopoietic transition; hemEC, hemogenic
endothelial cell, HSPC, hematopoietic stem/progenitor cell; PCV, posterior cardinal vein; VDA,
ventral dorsal aorta
641
642

643   **Figure 2.**

644
Figure 2. miR-223 is a conserved endothelial inhibitor of EHT and HSPC production
a, Schematic illustrates the EHT defects observed in the zebrafish miR-223Δ/Δ VDA of the
AGM compared to wild-type. Cell types involved in EHT are color-coded according to their
defining markers. **b,** Mean ± SEM number of kdrl:mCH+ gata2b:GFP+ cells in the VDA of the
zebrafish AGM during EHT (n = 4 embryos). **c,** Mean ± SEM number of kdrl:mCH+
cmyb:GFP+ cells in the VDA of the zebrafish AGM during EHT (n = 14-36 embryos). 36 hpf
data was previously published[17] and re-plotted for reference. **d**, Schematic depicts mouse
embryonic hematopoietic sites. HSPCs are produced in the AGM and mature as they transit to
the fetal liver, and then fetal bone marrow. **e,** Mean ± SEM fold change of mature miR-223
expression in wild-type FAC-sorted mouse endothelial cells (CD31+CD45-SP-) and hemECs
(Flk1+cKit+CD45- SP+) relative to non-endothelial cells (CD31-CD45-) in the e10.5 AGM (n =
3 technical replicates). **f,** Mean ± SEM percentage of hemECs (Flk+cKit+CD45- SP+) (n = 6
replicates) or HSPCs (cKit+Sca+ Lin-) at indicated hematopoietic sites at e10.5 (n = 5-8
replicates), e14.5 (n = 16 replicates), and e18.5 (n = 7 replicates) in the miR-223KO relative to
wild-type. For all plots, significance calculations are represented as *p ≤ 0.05, **p ≤ 0.01, ***p ≤
0.01, ****p ≤ 0.0001, two-tailed Student's t-test. Abbreviations: AGM, aorta-gonad mesonephros;
EC, endothelial cell; EHT, endothelial-to-hematopoietic transition; FBM, fetal bone marrow,
FBM; FL, fetal liver; hemEC, hemogenic endothelial cells, HSPC, hematopoietic
stem/progenitor cell; PCV, posterior cardinal vein; VDA, ventral dorsal aorta
664
665

Figure 3.

Figure 3. miR-223 targets N-glycogenes to limit HSPC production

a, Mean ± SEM fold change expression of N-glycogenes putatively targeted by miR-223 in miR-223Δ/Δ relative to wild-type kdrl:GFP+ endothelial cells at 27 hpf by qRT-PCR (n = 6-7 biological replicates). **b,** Schematic shows the biosynthesis of a hypothetical N-glycan (also see text and Supplementary Fig. 3a). Solid and dotted arrows indicate single and multiple steps occurring during the process, respectively. Only N-glycogenes putatively targeted by miR-223 are indicated and are color-coded based on their function in the ER (green) or Golgi (orange) during N-glycosylation. **c,** Lateral confocal Z-projections of the stable *Tg(fli1a:mCherry-alg2 3'UTR sensor: eGFP)* transgenic line in the zebrafish AGM at 32 hpf. Arrows indicate mCherry de-repression in endothelial cells along the VDA wall of embryos deficient for miR-223 or injected with Cas9 mRNA and a gRNA that disrupts miR-223 binding to the *alg2* 3'UTR-miR-223 MRE sensor. **d,** Mean ± SEM mCherry intensity normalized to eGFP along the length of the VDA wall in the AGM of 32 hpf embryos stably expressing *alg2* 3'UTR-miR-223 MRE sensor (n = 8-11 embryos) **e,** Representative expression of HSPC marker *cmyb* in the AGM at 32 hpf or CHT at 54 hpf (arrows) of zebrafish embryos injected with a gRNA that disrupts miR-223 binding to the endogenous *alg2* 3'UTR by whole mount in situ hybridization. **f, g** Mean ± SEM percentage of embryos with high or low/medium *cmyb* expression in the 32 hpf DA (**f**) or 54 hpf CHT (**g**) (n = 3 biological replicates). For all plots, significance calculations are represented as [*]p ≤ 0.05, [**]p ≤ 0.01, [***]p ≤ 0.01, [****]p ≤ 0.0001, two-tailed Student's t-test. Abbreviations: AGM, aorta-gonad mesonephros; DA, dorsal aorta; Dol, dolichol lipid; CHT, caudal hematopoietic tissue, GlcNAc, N-acetylglucosamine; gRNA, guide RNA; HSPC, hematopoietic stem/progenitor cell; MRE, miRNA responsive element; PCV, posterior cardinal vein; ER, endoplasmic reticulum; VDA, ventral dorsal aorta

Figure 4. miR-223 regulates the endothelial N-glycome during EHT.

a, Design of the N-glycosensor construct. The endothelial *fli1a* promoter drives co-expression of firefly luciferase, tagged with an ER translation sequence, and an un-localized renilla luciferase normalization control. Firefly luciferase localization to the ER permits glycosylation of its three naturally occurring N-glycosylation consensus sites, consisting of asparagine-X-serine/threonine (N-X-S/T), where X is any amino acid except proline. Firefly bioluminescence is affected by the extent of ectopic glycan addition as indicated. **b,** Mean ± SEM firefly normalized to renilla bioluminescence in wild-type endothelial cells at 24 hpf relative to 32 hpf. **c,** Mean ± SEM firefly normalized to renilla bioluminescence in miR-223Δ/Δ relative to wild-type endothelial cells at 32 hpf. **d,** Volcano plot depicts $\log_2$ fold change of glycans in 32 hpf miR-223Δ/Δ (n = 8 biological replicates) relative to wild-type (n = 5 biological replicates) by ratiometric lectin microarray. Dots represent discrete glycan moieties recognized by individual lectins. Glycan types that were significantly changed and recognized by multiple lectins are indicated, including α1,3 mannose (green), biantennary N-glycan stuctures with bisecting GlcNAc (blue), and α2,3 sialic acid (purple). Dotted treadline indicates a significant $-\log_{10}(0.05)$ p-value. **e,** Structures for N-glycans highlighted in **d**. Shaded boxes indicate binding specificities of associated lectins and are color coded accordingly. **f, i,** N-glycan structures depict the inhibitory activities of SW and ST. SW is an α-Mannosidase II inhibitor that removes the α1,3 and α1,6 mannoses indicated in the diagram (**f**). ST is a pan-sialyltransferase inhibitor that prevents terminal sialic acid modification (**i**). **g,** Representative images depict the zebrafish AGM at 32 hpf upon treatment with 5 µg/ml SW compared to 0.2% DMSO control from 8-32 hpf. White arrows in confocal projections (*left*) point to representative kdrl:mCh+ cmyb:gfp+ cells undergoing EHT in the VDA. Black arrows in bright field images (*right*) show *cmyb* expression in the AGM by whole mount in situ hybridization. **h, j,** Mean ± SEM number of kdrl:mCh+ cmyb:gfp+ cells in the VDA of embryos treated with SW (**h,** *left*) or ST (**j,** *left*) compared to the respective DMSO control at 32 hpf (n = 10 embryos). Mean ± SEM percentage of embryos with high or low/medium *cmyb* expression upon treatment with SW (**h,** *right*) (n = 4 biological replicates) or ST (**j,** *right*) compared to the respective DMSO control (n = 5 biological replicates). **k,** Model illustrates how miR-223-dependent regulation of the endothelial N-glycome restricts EHT and HSPC production. α1,3 mannose and α2,3 sialic acid glycans on membrane proteins are altered upon loss of miR-223 activity as shown. This altered "sugar code" promotes hemEC transition to HSPCs. miR-223+ expressing and non-expressing cells are outlined in green and red, respectively. Only relevant N-glycan moieties are depicted. For all plots, significance calculations are represented as $^{*}p \leq 0.05$, $^{**}p \leq 0.01$, two-tailed Student's t-test. Abbreviations: AGM, aorta-gonad mesonephros; DA, dorsal aorta; EHT, endothelial-to-hematopoietic transition; ER, endoplasmic reticulum; GlcNAc, N-acetylglucosamine; HSPC, hematopoietic stem/progenitor cell; IRES, internal ribosome entry site; PCV, posterior cardinal vein; ST, sialyltransferase inhibitor; SW, swainosine; VDA, ventral dorsal aorta

742   **Supplementary Figure 1.**

a

b

c

743
Supplementary Figure 1. miR-223-expressing cells delaminate from AGM
a, Mean ± SEM fold change of mature miR-223 expression in *Tg(miR-223:Gal4)$^{ya362/+}$* relative
to non-transgenic wild-type embryos at 32 hpf by qRT-PCR (n = 4 biological replicates). **b,**
Lateral maximum confocal Z-projection of time-lapse movie frames during EHT. Arrows point
to two miR-223:GFP+; kdrl:mCH+ cells budding from the VDA in the AGM. Relative time is
indicated in hr:min:sec (see also Supplementary Movie 1). **c,** Mean ± SEM percentage of miR-
223:GFP+ kdrl:mCH+ endothelial cells/ embryo that are within the VDA wall or appear to have
"budded" from the VDA into the sub-aortic space between the DA and PCV at 32 hpf (n = 36
embryos). For all plots, significance calculations are represented as n.s. (not significant, p > 0.05),
****p ≤ 0.0001, two-tailed Student's t-test. Abbreviations: AGM, aorta-gonad mesonephros; DA,
dorsal aorta; hemEC, hemogenic endothelial cells, PCV, posterior cardinal vein; VDA, ventral
dorsal aorta
756
757

Supplementary Figure 2.

761   *Supplementary Figure 2 Legend (next page)*

Supplementary Figure 2. miR-223 restricts EHT and HSPC production.
a, Representative expression of *gata2b* (*top*) and *runx1* (*bottom*) in the wild-type or miR-223Δ/Δ AGM by WISH. **b,** Quantification of *gata2b* (*left*) and *runx1* (*right*) WISH staining in **a.** Bar plots show normalized mean WISH signal ± SEM (n values are indicated in **a**). **c,** Representative confocal projections for Fig. 2c quantification. Arrows point to representative examples of kdrl:mCH+ cmyb:GFP+ cells undergoing EHT from the VDA in the zebrafish AGM. **d,** Schematic of *Tg(UAS:mCherry-miR-223)* construct to re-express miR-223 using miR-223:Gal4+ line via Tol2-mediated transgenesis. A UAS promoter drives expression of the pre-miR-223 precursor within an intron of mCherry. Splicing of this transcript liberates the pre-miR-223 stem loop, allowing it to be further processed into its mature and active form. **e,** Images show representative *cmyb* expression in the AGM (arrows) of miR-223:Gal4; miR-223Δ/Δ embryos injected with *Tg(UAS:mCherry-miR-223)* in **d** and uninjected controls at 32 hpf by WISH. Re-introduction of miR-223 into miR-223:Gal4+ cells rescues miR-223Δ/Δ *cmyb+* HSPC overexpansion. **f,** Construct as in **d**, except the endothelial *fli1a* enhancer/promoter drives miR-223 expression. **g,** Images show representative *cmyb* expression in the AGM (arrows) of miR-223Δ/Δ embryos injected with *Tg(fli1a ep:mCherry-miR-223)* in **f** and uninjected controls at 32 hpf by WISH. Re-introduction of miR-223 into endothelial cells rescues miR-223Δ/Δ *cmyb+* HSPC overexpansion. For all plots, significance calculations are represented as **p ≤ 0.01, ****p ≤ 0.0001, unpaired two-tailed Student's t-test. Abbreviations: AGM, aorta-gonad mesonephros; DA, dorsal aorta; EHT, endothelial-to-hematopoietic transition; HSPC, hematopoietic stem/progenitor cell; inj, injected; unj, uninjected; SA, splice acceptor site; SD, splice donor site; VDA, ventral dorsal aorta; WISH, whole mount in situ hybridization.

Supplementary Figure 3. miR-223 targets N-glycosylation genes in the AGM during EHT.
a, Overview of N-glycan biosynthesis. Putative miR-223 target genes are indicated and color-coded based on their function in the ER (green) or Golgi (orange) during N-glycosylation. Black arrowheads point to Alg2 and St3Gal2 glycosyltransferase activities on a hypothetical N-glycan core after its addition to a polypeptide chain, and to the ER-Golgi transport function of Lman2lb. (See also text and Fig. 3b). **b,** Endothelial N-glycogene-3'UTR-miR-223 MRE sensor construct to test direct miR-223 targeting of N-glycogenes. The *fli1a* enhancer/promoter drives co-expression of mCherry controlled by a ~100-125 bp N-glycogene 3'UTR fragment containing the miR-223 MRE, and an unregulated eGFP normalization control. **c,** Schematic illustrates de-repression of endogenous *alg2* or N-glycogene sensor mRNAs upon loss of miR-223 activity. **d,** Representative confocal micrographs showing Tol2-mediated mosaic integration of the *alg2* 3'UTR-miR-223 MRE sensor at 32 hpf in the zebrafish AGM. White arrowheads point to mCherry repression in endothelial cells along the VDA wall. Upon miR-223 depletion, mCherry is de-repressed in a subset of these cells (yellow arrows), consistent with the salt and pepper miR-223 expression pattern (See Fig. 1a). **e-g,** Box-plots show mCherry intensity normalized to eGFP for AGM endothelial cells in 32 hpf embryos mosaically-expressing the *alg2* (**e**), *lman2lb* (**f**), or *st3gal2* (**g**) 3'UTR-miR-223 MRE sensors. Embryos were derived from miR-223Δ/+ in-crosses. n values are indicated and represent the number of endothelial cells analyzed. Significance was determined by two-tailed Mann-Whitney U test. **h,** Mean ± SEM mCherry intensity normalized to eGFP along the length of the axial vessel walls in the AGM of 32 hpf embryos stably-expressing the *alg2* 3'UTR-miR-223 MRE sensor. Axial vessel walls include the dorsal (D) and ventral (V) walls of the DA (red) or PCV (blue), referred to as DDA, VDA, DCV, and VCV, respectively, as depicted in the graph schematic (n = 8-11 embryos, two-tailed Student's t-test.) For all plots, significance calculations are represented as n.s. (not significant, $p > 0.05$), $^*p \leq 0.05$, $^{**}p \leq 0.01$, $^{***}p \leq 0.001$, $^{****}p \leq 0.0001$. Abbreviations: AGM, aorta-gonad mesonephros; DA, dorsal aorta; ER, endopasmic reticulum; GlcNAc, N-acetylglucosamine; gRNA, guide RNA; MRE, miRNA responsive element; PCV, posterior cardinal vein

819 **Supplementary Figure 4.**

820
Supplementary Figure 4. Hypoglycosylation and ER-stress do not lead to HSPC
overexpansion.
a, Mean ± SEM firefly normalized to renilla bioluminescence in endothelial cells of wild-type
embryos injected with the N-glycosensor (see Fig. 3a) and treated with TM relative to DMSO
control (n = 6 biological replicates) at 32 hpf (n = 6 biological replicates). Significance is
represented as *p ≤ 0.05, paired, two-tailed Student's t-test. **b, c** Images show representative *cmyb*
expression in the AGM (arrows) of 32 hpf wild-type embryos treated with TM (**b**), TG (**c**), and
the respective DMSO control by WISH. **d,** Representative images depict the zebrafish AGM at
32 hpf upon treatment with 200 µg/ml ST compared to 2% DMSO control from 8-32 hpf. White
arrows in confocal projections (*left*) point to representative kdrl:mCh+ cmyb:gfp+ cells
undergoing EHT in the VDA. Black arrows in bright field images (*right*) show representative
cmyb expression in the AGM by WISH. Abbreviations: AGM, aorta-gonad mesonephros; DA,
dorsal aorta; PCV, posterior cardinal vein; TG, thapsigargin; TM, tunicamycin; VDA, ventral
dorsal aorta; WISH, whole mount in situ hybridization
835

836 **Supplementary Movie 1. miR-223-expressing cells delaminate from AGM**
837 Lateral maximum confocal Z-projection of a time-lapse movie during EHT. Two miR-
838 223:GFP+; kdrl:mCH+ cells are shown budding from the ventral dorsal aorta in the AGM.
839 Relative time is indicated in hr:min:sec (see also Supplementary Fig. 1b).