Exploring the link between adenine DNA methylation and 3D genome organization in the parasite *Trichomonas vaginalis*

Ayelen Lizarraga [1]; Zach Klapholz O'Brown[2,3]; Lara Roach[2,3]; Eric Lieberman Greer [2,3]; Pablo H. Strobl-Mazzulla [4]; Natalia de Miguel [1]*

1 Laboratorio de Parásitos Anaerobios, Instituto Tecnológico Chascomús (INTECH), CONICET-UNSAM, Chascomús B7130IWA, Argentina

2 Division of Newborn Medicine, Children's Hospital Boston, 300 Longwood Avenue, Boston, MA 02115, USA

3 Department of Pediatrics, Harvard Medical School, Boston, MA 02115, USA

4 Laboratorio de Biología del Desarrollo, Instituto Tecnológico Chascomús (INTECH), CONICET-UNSAM, Chascomús B7130IWA, Argentina

*Corresponding author.

Email: ndemiguel@intech.gov.ar (NdM)

ABSTRACT

Trichomonas vaginalis is a common sexually transmitted parasite that colonizes

the human urogenital tract causing infections that range from asymptomatic to highly

inflammatory. Recent works have highlighted the importance of histone modifications in

the regulation of transcription and parasite pathogenesis. However, the nature of DNA

methylation in the parasite remained unexplored. Using a combination of immunological

techniques and UHPLC, we analyzed the abundance of DNA methylation in strains with

differential pathogenicity demonstrating that N6-methyladenine (6mA), and not 5-

methylcytosine (5mC), is the main DNA methylation mark in *T. vaginalis*. Genome-wide

distribution of 6mA reveals that this mark is enriched at intergenic regions, with a

preference for certain superfamilies of DNA transposable elements. We show that 6mA

in *T. vaginalis* is associated with silencing when present on genes. Interestingly,

bioinformatics analysis revealed the presence of transcriptionally active or repressive

intervals flanked by 6mA-enriched regions and results from chromatin conformation

capture (3C) experiments suggest these 6mA flanked regions are in close spatial

proximity. These associations were disrupted when parasites were treated with the

demethylation activator ascorbic acid. This finding revealed the newly role for 6mA in

modulating 3D chromatin structure and gene expression in this deep-branching

eukaryote.

AUTHOR SUMMARY

Trichomonas vaginalis is a common sexually transmitted parasite that colonizes the human urogenital tract causing infections that range from asymptomatic to highly inflammatory. Chronic infections have been associated with high risk pregnancies, increased risk of acquiring HIV and higher susceptibility to developing cervical or prostatic cancer. Despite their importance in other organisms, the epigenetic mechanisms involved in gene regulation in the parasite remain poorly understood. Recent works have highlighted the importance of histone modifications in the regulation of transcription and parasite pathogenesis. However, nothing is known about the methylation status of *T. vaginalis* genomic DNA to this day. In this work we demonstrate that N6-methyladenine (6mA), and not 5-methylcytosine (5mC), is the main methylation mark in *T. vaginalis* DNA. We explore the genome-wide distribution of 6mA and show this mark has a preference for intergenic regions and is associated with silencing when present on genes. We find the presence of transcriptionally active or repressive intervals flanked by 6mA-enriched regions and our results suggest these 6mA flanked regions are in close spatial proximity. Our findings describe for the first time the presence of DNA methylation in *T. vaginalis* and reveal a new role for 6mA in modulating 3D chromatin architecture.

INTRODUCTION

Trichomonas vaginalis is a flagellated protozoan parasite responsible for trichomoniasis, the most frequent non-viral sexually transmitted infection. Although generally asymptomatic [1], *T. vaginalis* infection can cause numerous pathologies like vaginitis, urethritis, prostatitis and various complications when infection is acquired during pregnancy [2, 3]. Chronic infection has been associated with an increased risk of acquiring and transmitting HIV [4] and higher susceptibility to developing cervical or prostatic cancer [5, 6].

With ~160Mb distributed over 6 chromosomes, the genome of *T. vaginalis* is the largest protozoan parasite genome sequenced to date [7]. Unfortunately, the highly repetitive nature of the genome (up to 65% comprised of repetitive elements) prevented the complete assembly of the scaffolds and subsequent studies of *T. vaginalis* chromosome architecture [8]. Analysis of the age distribution of gene families coupled with the large genome size and low repeat polymorphism indicates the genome underwent one or more large-scale genome duplication events [8]. Intriguingly, only half of the ~46,000 protein-coding genes appear to be expressed [9, 10] and individual members within gene families show differential regulation in response to different stimuli [9-11]. These results indicate that extrinsic and intrinsic factors accurately control the fine tuning of gene expression in this extracellular parasite.

Studies on *T. vaginalis* transcriptional regulation mostly focused on identifying cis-regulatory elements and their cognate transcription factors. Few core promoter elements or transcription factors have been identified to date [12]. The metazoan-like Initiator (Inr) element [13] present in ~75% of all protein coding genes [8] was found to be the main core promoter element responsible for directing basal transcription in *T.*

vaginalis. A novel initiator binding protein, IBP39, was identified and demonstrated to specifically recognize *T. vaginalis* Inr and directly interact with RNAPII [13, 14]. Additionally, several conserved motifs that resemble metazoan Myb recognition elements [15] as well as proteins containing a Myb DNA binding domain, like novel transcription factor M3BP [15], have also been found in *T. vaginalis* [12]. This Myb recognition elements can act as cis-regulatory elements directing the transcription start site selection [15] and transcription of specific genes [16]. However, little is known about other mechanisms of gene regulation in this parasite [12]. Epigenetic mechanism regulates gene expression by modifying chromatin accessibility to transcription factors and other components of the transcriptional machinery [17]. Specifically, histone modifications have been shown to be of importance in the regulation of gene expression and antigenic variation of other unicellular parasites [18]. Recent studies have demonstrated that these modifications are present in *T. vaginalis* and play an essential role in transcriptional regulation and parasite pathogenesis [19, 20].

DNA methylation is one of the major epigenetic mechanism involved in gene regulation. The most common DNA modification in eukaryotes is 5-methylcytosine (5mC), an epigenetic mark usually associated with gene silencing [21] that participates in a wide variety of processes such as imprinting, X chromosome inactivation, and silencing of transposable elements [22]. Though common among higher eukaryotes, 5mC is not universally present in unicellular organisms [18] and thus few protozoan parasites with detectable levels of 5mC in their genome have been identified and its function fully characterized [18, 23-25]. By contrast, N6-methyladenine (6mA) is the most prevalent DNA modification in prokaryotes [26] and early reports debated the existence and abundance of this mark in eukaryotes [27]. However, in recent years numerous reports have greatly expanded the list of organisms with 6mA to include flies [28], nematodes

[29], green algae [30] and even vertebrates [31-33]. The abundance of this mark varies

dramatically between species, with levels of 6mA that range from 6-7 parts per million

(p.p.m.) of total adenines in mouse embryonic stem cells [31] up to 2.7% of total adenines

in early-diverging fungi [34]. Likewise, this variation between organisms was also

evidenced in their genomic distribution, suggesting that 6mA might have specie-specific

functions[35]. Intriguingly, in some cases the role of 6mA has been associated with active

transcription in *Arabidopsis* [36], *Drosophila* [28], *Chlamydomonas* [30] and early-

diverging fungi [34]. Only two reports describe 6mA as a repressive mark similar to 5mC

[31, 37].

Despite its importance in other organisms, the methylation status of *T. vaginalis*

DNA remained unexplored. In this work, we use a combination of immunological

techniques and UHPLC coupled with mass-spectrometry to determine for the first time

the presence of 6mA and 5mC in the gDNA of *T. vaginalis*. We interrogate the

distribution of 6mA across the genome using an adapted methylated DNA

immunoprecipitation assay followed by high-throughput sequencing (MeDIP-seq).

Interestingly, we found that 6mA is enriched at intergenic regions, with a preference for

repetitive elements. Comparison with published RNA-seq data of the same strain [38]

suggests 6mA is associated with silenced transcription when found on gene bodies.

Finally, results from chromatin conformation capture experiments suggest that 6mA is

associated with chromatin loop formation, raising the intriguing possibility that 6mA

could have a role in modulating 3D genome architecture. This is the first time DNA

methylation is described in *T. vaginalis* and highlights the importance of epigenetic gene

regulation in this deep-branching eukaryote.

RESULTS

Identification of 6mA and 5mC in _T. vaginalis_ genomic DNA

Given the importance of 5mC and 6mA in other organisms, we set out to

investigate the presence of these marks in _T. vaginalis_. We first evaluated the presence of

5mC in _T. vaginalis_ DNA by a dot blot assay using genomic DNA extracted from two _T._

vaginalis strains, B7268 and G3, with high and low adherence capacity, respectively (Fig.

1A). Unexpectedly, we failed to detect 5mC in either strain, contrary to our chicken

gDNA control (Fig. 1A). In agreement, we were unable to detect this mark by

immunofluorescence in the nucleus of the parasite (Fig. 1B). These data suggest the

absence, or very low abundance, of 5mC mark in the DNA of _T. vaginalis_. On the other

hand, we revealed by immunofluorescence the presence of m6A mark both in the nuclei

and cytoplasm (Fig. 1C). Importantly, parasites treatment with RNAse revealed 6mA

exclusively detected in the nucleus demonstrating that this epigenetic mark is present in

the RNA and DNA of _T. vaginalis_. Similarly, using restriction enzymes MboI and DpnI

which digest unmethylated and methylated GATC sequence respectively [39], we found

that _T. vaginalis_ genome possesses both methylated and un-methylated adenines (Fig.

1D). In order to determine whether parasite adherence could be correlated with 6mA

abundance, we performed dot blot assays (Fig. 1E) and UHPLC followed by mass

spectrometry (Fig. 1F) using gDNA from an adherent (B7268) and a less-adherent (G3)

strain. However, we found no difference on the abundance of 6mA levels among them.

Importantly, we demonstrated that 6mA is present in 2.5% of all adenines, but extremely

low levels of 5mC have been detected (0.004-0.008% of all cytosines) by UHPLC

followed by mass spectrometry (Fig. 1F). It is important to note that, to exclude the

possibility of signal contamination from _T. vaginalis_ symbiont _M. hominis_[40]_,_ parasite

cultures were subjected to antibiotic treatment and checked for bacteria contamination

prior to the experiments (Sup. Fig. 1), confirming 5mC and 6mA detection came

exclusively from *T. vaginalis* genomic DNA. Taken together, our results demonstrated

that 6mA, but not 5mC, is the predominant DNA methylation mark in *T. vaginalis*.

To be considered an epigenetic mark, 6mA needs to be maintained after DNA

replication. Our observations using hydroxyurea synchronized cultures revealed the

expected decrease in 6mA levels in S phase due to DNA replication, which remained

during the G2/M phase of the cell cycle, and restored original abundance after cell

division (Sup. Fig. 2). These observations support the presence of active

methyltransferase(s) responsible for maintaining 6mA levels after cell division.

Genome-wide mapping of 6mA in *T. vaginalis*

To determine the function of DNA methylation in *T. vaginalis* it is essential to

identify its genomic distribution. To this end, we performed an adapted methylated DNA

immunoprecipitation technique using specific antibodies followed by high-throughput

sequencing (MeDIP-seq) [41] on the adherent strain B7268 (Fig. 2A). Unfortunately, due

to the extremely low levels of 5mC in *T. vaginalis* we were unable to get good quality

reads from two independent MeDIP-seq experiments (data not shown). Therefore, we

decided to focus exclusively on 6mA for further analysis. To this end, two independent

high-throughput sequencing experiments, each containing pooled DNA from three

independent immunoprecipitation assays, were performed. In order to evaluate specificity

and enrichments of 6mA in the immunoprecipitated fraction when using a specific anti-

6mA antibody, a dot blot assay was performed using an IgG control antibody (Fig. 2B).

After quality filtering (q-value < 0,01, FE > 2) we obtained a total of 10388 6mA

peaks found in both replicates, corresponding to 92,1% and 75,6% of all peaks present in

each independent MeDIP-seq experiment (Fig. 2C, Sup. Table 1). The most prevalent

motif, found in 4692 peaks, was CAATACCC (p-value = 1e-1642, Sup. Fig. 3A).

Remarkably, peak visualization in a genome browser suggested a clear enrichment of

peaks in intergenic regions (Fig. 2D). Indeed, peak annotation revealed that 94% of all

MeDIP-seq peaks (9746 6mA peaks) were intergenic (Fig. 3A). Importantly, a shuffled

distribution of peaks across *T. vaginalis* genome would result in an even distribution

between genes and intergenic regions (Fig. 3A), confirming 6mA is truly enriched in

intergenic regions.

Among the 642 6mA peaks located in genes, most of them were distributed

between the coding region (48%) and the transcription termination sites (TTS) (43%),

with only 9% found in the transcription start sites (TSS) (Fig. 3B). Comparing those

numbers to the percentages that would occur if the peaks had a shuffled distribution (22%

in the TTS, 60% in the coding region and 12% in the TSS) our results revealed an

enrichment of 6mA in the TTS when located in genes (Fig. 3B). In order to understand

the functional role of 6mA, we retrieved the IDs of the methylated genes (Sup. Table 2)

and found 70% belong to different families of transposable elements, with the majority

(61%) belonging to families previously described in *T. vaginalis* according to Repbase

(Fig. 3C). The rest of the identified genes could be divided into genes with conserved

domains that allow the assignment of a predicted role (8%) and hypothetical proteins with

unknown functions (22%) found exclusively in *T. vaginalis* (Fig. 3C). Intriguingly, the

observed enrichment of 6mA peaks in the TTS was found exclusively in transposable

element (TE) genes, suggesting a particular role for methylation at the 3' of TE genes.

Gene ontology analysis of the methylated genes (excluding genes from TEs) revealed an

enrichment of genes involved in protein modification and phosphorylation (p-value <

0.05) (Sup. Table 3). However, it is important to take into account that most of non-TE

genes have no identifiable domain (hypothetical proteins) and therefore were not

considered into the GO enrichment analysis.

Considering that 6mA has been previously described as an epigenetic mark of

transposable elements in other organisms [28, 31] and 70% of *T. vaginalis* methylated

genes belong to TEs, we analyzed if intergenic peaks were also located within TEs.

Analysis using RepeatMasker revealed that out of 94% of intergenic peaks, 57.4% were

found in *T. vaginalis* repetitive elements (RE) previously described in Repbase. Most of

the peaks were found in Mutator, Kolobok, and Polinton superfamilies of *T. vaginalis*

DNA transposons (Fig. 3D). However, if we consider the numbers of TEs within the

genome, repeat element DNA7 and TE superfamilies P and Kolobok were the most

enriched in the MedIP-seq (Sup. Table 4), indicating a preference of intergenic 6mA for

certain types of REs and raising the question of whether 6mA could act as a regulator of

specific TEs.

6mA methylation and gene expression

6mA has been found to be an epigenetic mark of active genes in *Arabidopsis*[36],

Drosophila [28] and *Chlamydomonas* [30] or repressed genes in mouse embryonic stem

cells [31] and rice [37]. To examine whether this mark has a role in transcription in *T.*

vaginalis, we made use of publicly available RNA-seq data from B7268 strain [38]. To

test if this data was applicable for our analysis, we divided the genes into low (RPKM =<

1), moderate (1 < RPKM =< 20), and high (RPKM > 20) expression, and selected some

well characterized genes from each group to measure their relative expression levels via

RT-qPCR. Our results showed the relative expression of each gene was in accordance to

what had been described in bibliography [20, 42] and more importantly, we could order

the genes based on their relative expression obtained by RT-qPCR similarly as they have

been ordered based on their RPKM (Sup. Table 5), validating the use of the RNA-seq in

our analysis.

We next looked into the expression levels of the methylated genes using RNA-

seq data available and maintaining the same criteria for low, moderate and high

expression. Considering that the majority of those genes belong to *T. vaginalis*

transposable elements (Fig. 3C), it was not surprising that gene expression analysis

revealed over 80% of all methylated genes are poorly expressed, with 14% possessing

moderate expression and only a small percentage (4%) high expression (Fig. 4A).

However, when we considered the expression levels of the methylated genes without

taking into account TE genes, poorly expressed genes were still the most abundant group

(over 60%, data not shown), suggesting 6mA could be a sign of repressed expression

when found on genes.

As described previously, most of the MeDIP-seq peaks have an intergenic

localization (Fig. 3A). In order to evaluate if 6mA could be affecting the expression of

adjacent genes as has been described for 6mA in mouse embryonic stem cells [31], we

analyzed the expression level of the genes closest to intergenic methylation (Fig. 4B). We

obtained a total of 6000 genes found near 6714 peaks. However, due to the fragmented

nature of *T. vaginalis* reference genome not all the intergenic peaks could be considered,

and it is likely that the real number of genes near intergenic 6mA could be higher.

Surprisingly, in contrast with our observations regarding the expression levels of

methylated genes (Fig. 4A), we found a higher percentage of expressed genes (13,5%

with moderate and 16,4% with high expression) near intergenic methylation compared to

poorly expressed genes (Fig. 4B). In fact, genes with intergenic 6mA peaks nearby were

expressed at significantly higher levels than genes in unmethylated regions (Fig. 4C).

Finally, we examined the association between the 6mA peaks distance and expression

level of associated genes. We have observed that intergenic 6mA is primarily located

between -1500pb and +1000bp of the nearest genes regardless of their expression level

(Fig. 4D). Intriguingly, not expressed genes are found significantly closer to their nearest

upstream or downstream intergenic peak than expressed genes (Sup. Fig. 3B-C). These

observations suggest that the effect of 6mA on gene expression in *T. vaginalis* depends

on its relative position to the corresponding gene.

6mA delimits chromatin loops

During the analysis of the expression levels of genes near intergenic methylation

regions, we noticed various instances of intergenic 6mA peaks flanking sections of the

genome where all the genes contained within had similar expression levels. A closer look

revealed the presence of 1046 such intervals with more than one gene: 452 containing not

expressed genes (repressive intervals) and 594 containing expressed genes (active

intervals) (Fig. 5A). These intervals varied in the number of genes they contained (from

2 to up to 36 genes) (Fig. 5B) as well as in length, with some spanning over 20kb (Sup.

Table 6). As can be observed in Figure 5B, intervals containing a greater number of genes

tends to be silenced. In an opposite manner, most of the intervals containing fewer genes

are actively transcribed.

In other organisms it is well known that the three-dimensional genome

organization within the nucleus is an important element in the regulation of gene

expression [43]. In particular, chromatin looping is a type of intrachromosomal

interaction that has been shown to influence gene expression [44]. A recent work in

mammalian cells demonstrated that 5mC was capable of regulating the expression of two

genes 10 kbp apart via chromatin loop formation [45]. Based on these antecedents and

our bioinformatics observations, we performed a chromatin conformation capture (3C)

assay (Fig. 6A) to determine if the observed intervals, delimited by intergenic 6mA, could

be forming this type of chromatin architecture. In order to select intervals to analyze by

3C assay, we filtered out all the intervals with fewer than 5 genes and those mostly

comprised of TE genes, obtaining a list of 80 active and 5 repressive intervals.

Surprisingly, we found several intervals containing BspA family members: 2 of the

repressive intervals contained 4 BspA genes each, while 6 active intervals contained one

BspA gene per interval. In bacteria, the BspA family of cell surface proteins has been

shown to be involved in the colonization of the oral mucosa and triggering of host immune

response and works on *T. vaginalis* BspAs suggest these proteins might have a similar

function in the parasite [46]. Therefore, we decided to perform 3C on those active and

repressive intervals that contained BspA genes (Sup. Table 8). Unfortunately, the

repetitive nature of *T. vaginalis* genome presented a challenge for the design of specific

3C PCR primers, limiting the amount of intervals to be considered for the experiment to

repressive intervals p5512-5513 and p2296-2297, and active intervals p4130-4131, p560-

561 and p271-272. Amongst this set of regions, we were unable to amplify either p560-

561 or p271-272. On the other hand, PCR amplification of 3C product using the primer

pairs for p5512-5513 and p4130-4131 showed bands of the expected size for each interval

that were absent when non-crosslinked control or undigested gDNA were used as

template (1-2, Fig. 6B). The intensity of these bands was higher than the ones obtained

using an alternative set of primers meant to interrogate the interaction between one

extreme of p5512-5513 and p4130-4131 and a flanking region located at a greater

distance (1-3, Fig. 6B). Furthermore, sequencing of each 3C band verified that the

fragments belonged to the expected cross-ligation products (Sup. Fig. 4). Taken together,

our results suggest the presence of chromatin looping at these regions. Additionally, we

manage to amplify the interval p2296-2297 (Sup. Fig. 4C), obtaining a band of the

expected size. Although we were not able to achieve reproducible results with this primer

pair, band sequencing confirmed that the identity of the fragment was the expected cross-

ligation product (Sup. Fig. 4D), suggesting the formation of a chromatin loop.

To unambiguously demonstrate that 6mA is responsible of regulate the formation

of these structures, we performed a 3C assay of parasites treated with ascorbic acid (AA),

which has been previously shown to favor DNA demethylation in mammalian cells [47].

Dot blot assay of parasites treated with AA confirmed a diminution of 6mA mark of

parasites treated with 100 mM AA compared to non-treated parasites (Fig. 6C).

Excitingly, parasite treated with AA were unable to form both p5512-5513 and p4130-

4131 loops using 3C analyzes (Fig. 6D). Consistent with this finding, our observations

suggests 6mA could indeed be a mark associated with chromatin looping and regulate the

transcriptional activity of gene-associated blocks in *T. vaginalis*.

DISCUSSION

Epigenetics studies and it relationship with gene expression in *T. vaginalis* are still

in their infancy. To date, only the contribution of histone modifications to the regulation

of gene expression in the parasite has been described [19, 20]. In the present study, we

describe for the first time the major presence of 6mA modification in the genome of this

unicellular parasite. Here, we present a whole genome analysis of 6mA distribution that

provides insight into the possible contributions of this modification to 3D genome

architecture and gene expression.

In recent years, the role of 6mA has been expanded from a primarily prokaryotic

DNA modification to an important epigenetic mark present in several multicellular and

unicellular eukaryotic organisms [35]. In this study, we use antibody-dependent and

independent methods to demonstrate that 6mA is an abundant DNA modification in *T. vaginalis* genome. 6mA levels in the parasite are comparable to those found in early-diverging fungi [34], making *T. vaginalis* one of the eukaryotic organisms with the highest levels of 6mA reported to date. It is interesting to note that just like early-diverging fungi [34], 5mC in *T. vaginalis* was nearly undetectable. Unfortunately, detection methods that rely on the use of antibodies can give false positives when 5mC levels are low[48], which could explain the low-quality reads obtained from the 5mC MeDIP-seq experiment. These results support our conclusion that 6mA, and not 5mC, is the main DNA modification in *T. vaginalis* genomic DNA.

To be considered an epigenetic mark, 6mA needs to be maintained after DNA replication. In *Tetrahymena*, maintenance methylation occurs quickly after DNA replication [49]. Similarly, 6mA in *Chlamydomonas* is installed shortly after DNA replication and is stably maintained during cell proliferation [30]. Although the precise moment of 6mA deposition in the newly synthetized DNA strand in the parasite remains elusive, our observations indicate maintenance of 6mA methylation occurs after cell division and supports the presence of active 6mA methyltransferase(s) in the parasite. It should be noted that although phylogenetic analysis revealed the presence of a group of parasite specific 6mA DNA methyltransferases encoded in the genome database (TrichDB) [50], after several attempts of cloning and transfection to perform gain of function studies, we have not been able to validate these candidates by functional analyses (data not shown). On the other hand, while DNA 6mA demethylases have been identified in several organisms [28, 29, 31, 37], no classical TET or AlkB homologs were identified in *T. vaginalis* genome [50]. This could be indicative of the presence of non-conserved enzyme, a lack of active 6mA demethylation or the use of an alternative pathway, such as the base excision-repair DNA demethylation pathway found in plants [51]. It will be

of interest to take a closer look at the DNA glycosylases found in *T. vaginalis* genome

database and evaluate if they could be involved in active 6mA DNA demethylation.

Identifying the proteins responsible for the regulation of 6mA levels in the parasite will

further our understanding of the mechanisms and functions of this mark.

Genome-wide distribution of 6mA varies according to the species being

considered [28-31]. An evolutionary conservation for 6mA in unicellular organisms has

been proposed due to the similarities in 6mA distribution pattern and function between

green algae and *Tetrahymena* [49]. However, using a MeDIP-seq approach we found the

distribution pattern in *T. vaginalis* resembles more that of 6mA in *Drosophila* and mouse

than other unicellular eukaryotes. Methylation peaks in *T. vaginalis* are enriched in

intergenic regions and seem to have a preference for certain types of DNA TE

superfamilies, reminiscent of the preference for young LINE-1 transposons described in

mouse [31]. In Drosophila, some 6mA peaks are enriched in the gene bodies of

transposons [28]. Similarly, we found that the majority of the methylated genes in the

parasite belong to transposable elements, although 6mA peaks are preferentially located

in the TTS of these TE genes (data not shown). This predilection for certain types of TEs

and the specific localization of peaks in transposon genes raises the question of whether

6mA could act as a regulator of specific TEs in *T. vaginalis*.

Like its genomic distribution, 6mA association with transcription is dependent of

the organism in question. It has been demonstrated that 6mA acts as a mark of active

transcription in *Arabidopsis* [36], *Drosophila* [28], green algae [30] and fungi [34], while

it is a mark of transcriptional repression in mouse [31] and rice [37]. In this study, 6mA

seems to be associated with silencing when found on genes. Importantly, we cannot rule

out the possibility of 6mA acting in conjunction with other components of the epigenetic

code to regulate gene expression in a context-dependent manner. In future studies, it will

be interesting to determine whether any 6mA binding proteins exist in *T. vaginalis* which

could help regulate the response of chromatin to this modification. An integrative analysis

is necessary to fully understand how epigenetics shapes the transcriptional landscape of

T. vaginalis.

Chromatin looping is a common intrachromosomal protein-mediated interaction

capable of influencing gene expression. The function of these interactions is determined

by the nature of the sequences brought together and the proteins that interact with them

[52]. For example, promoter-enhancer loops are strongly associated with gene activation

[53] and it has been proposed that the presence of chromatin loops may contribute to the

co-regulation of specific block of genes by increasing the local concentration of the

transcription machinery, resulting in the formation of structures similar to transcription

factories [54]. Alternatively, chromatin looping has also been implicated in

transcriptional repression in *Drosophila* [55] and mammals [56]. In agreement with this

notion, our bioinformatics analysis revealed the presence of active and repressive

intervals of varying length flanked by 6mA. Interestingly, we found some of them

containing one or more members of the BspA-like gene family of surface proteins which

might be involved in parasite pathogenesis [57]. In *P. falciparum*, chromosomes

containing virulence genes of the var family form loops that permit the perinuclear

colocalization of the var genes [58], and a recent work demonstrated that this clustering

of virulent genes is a common feature among other *Plasmodium* species [59]. A chromatin

conformation capture assay confirmed that the methylated extremes in both

transcriptionally active and repressive block of genes, containing BspA genes, are in close

spatial proximity. Since clustered organization of virulence genes allows coordination of

gene expression in *Plasmodium* [59], more studies are needed to evaluate if the same

could be said for *T. vaginalis*.

The association of 6mA with chromatin looping is noteworthy because it points

to a novel function for this epigenetic mark in eukaryotes with scarce 5mC levels. Given

that our finding demonstrated that chromatin structure may regulate, positively or

negatively, specific block of genes; these finding open new avenues for understanding

the role of m6A writer, erasers and reader that may play an important role on the

transcriptional regulation of *T. vaginalis*.

MATERIALS AND METHODS

Parasites, cell cultures and media

Trichomonas vaginalis strains B7268[60] and G3 (ATCC PRA-98, Kent, UK) were

cultured at 37°C in TYM medium supplemented with 10% horse serum, penicillin and

streptomycin (Invitrogen) [61]. To obtain *M. hominis* free strains, parasite cultures were

treated for two weeks with the recommended working concentration of BM-Cyclin

(Roche). *M. hominis* clearance was confirmed by PCR as described previously [62].

6mA and 5mC immunolocalization experiments

Parasites in the absence of host cells were incubated at 37°C on glass coverslips as

previously described [63] for 4 hours, then fixed and followed by permeabilization in cold

methanol for 10 min. Cells were washed with PBS and incubated at 37°C with a 1:100

dilution of RNase A (QIAGEN) in PBS for 3 hours. Non-RNase A treated control

parasites were incubated for 3 hours in PBS. Parasites were washed once again, blocked

with 5% FBS in PBS for 30 min and incubated overnight with a 1:100 dilution of anti-

6mA or anti-5mC (Abcam, USA) primary antibody in PBS containing 2% FBS. Parasites

were washed and then incubated with a 1:5000 dilution of Alexa Fluor conjugated

secondary antibody (Molecular Probes). Nuclei was stained with DAPI (2 mg/ml) and

coverslips were mounted onto microscope slips using Fluoromount Aqueous Mounting

Medium (Sigma-Aldrich). All observations were performed on a Nikon E600

epifluorescence microscope. Adobe Photoshop (Adobe Systems) was used for image

processing.

Genomic DNA extraction

435   *T. vaginalis* cultures were lysed using a solution of 8 M urea, 2% sarkosyl, 0.15 M NaCl,

436   0.001 M EDTA, and 0.1 M Tris HCL pH7.5 [64]. RNase A was added to a final

437   concentration of 0.1 mg/ml and lysates were incubated at 37°C for 2-3 hours. Total DNA

438   was extracted using phenol/chloroform/isoamyl alcohol (Sigma-Aldrich), precipitated

439   with isopropanol, and resuspended in nuclease-free water.

Dot Blot Assays

RNA free genomic DNA samples were diluted to the desired concentration and 1 µl of

each sample was spotted on Amersham Hybond-N+ membranes (GE Healthcare). For

anti-5mC blots, DNA was denatured for 10 minutes at 95°C. Membranes were allowed

to air dry and then DNA was crosslinked for 3 minutes twice in a Hoefer UVC 500

Crosslinker (Amersham Biosciences). Membranes were stained with 0.02% methylene

blue in 0.3 M AcNA pH 5.2 solution for DNA loading control and then washed with

distilled water. The membranes were blocked for 1 hour in 5% milk TBS-T and incubated

overnight at 4°C with 1:1000 anti-6mA primary antibody dilution or 1:500 anti-5mC

primary antibody dilution in 5% milk TBS-T. Membranes were washed three times for

10 min with TBS-T and incubated with a 1:10000 dilution of AP-conjugated secondary

antibody in 5% milk TBS-T for 1 hour at room temperature. After three washes of 10 min

with TBS-T, blots were developed using NBT and BCIP (Roche).

Restriction enzyme digestion assay

Restriction enzyme digestion was performed by treating 1 µg of genomic DNA with 1 µl

of 10 U/µl MboI or DpnI restriction enzyme (Thermo Scientific) at 25°C overnight.

Sample with no restriction enzyme was used as control. Enzyme was heat inactivated and

the product mixture was separated by agarose gel electrophoresis. Gels were visualized

with a UV transilluminator.

Quantification of 6mA by UHPLC-QQQ-MS/MS Analysis

6mA abundance was quantified as previously described [29]. 1.5 µg of genomic DNA in

30 ml ddH2O was digested to free nucleosides using 5 U of DNA Degradase Plus (Zymo

Research) in 25 µl reactions incubated for 2 hours at 37°C. The diluted solution was filtered

through 0.22 mm filter and 10 ml solution was injected into LC-MS/MS. The nucleosides

were separated by reverse phase ultra-high performance liquid chromatography (UHPLC)

on a C18 column (Agilent), with online mass spectrometry detection using Agilent 6460

QQQ triple-quadrupole tandem mass spectrometer (MS/MS) set to multiple reaction

monitoring (MRM) in positive electrospray ionization mode. Nucleosides were quantified

using the nucleoside precursor ion to base ion mass transitions of 266.1-150.0 for 6mA

and 252.1-136.0 for A. Quantification of the ratio 6mA/A was performed using the

calibration curves obtained from nucleoside standards running at the same time.

Measurement of 6mA and dA levels in the DNA degradase enzyme mix alone served as

an additional control for background levels of nucleic acid contributed by the digestion

enzymes. The low levels of 6mA and dA in the DNA degradase enzyme mix were

subtracted from the final measurements. Because *T. vaginalis* have relatively high levels

of 6mA, the low levels of methylated and unmethylated DNA in the reaction mix were a

negligible fraction of the total digested DNA samples.

Cell cycle synchronization

478   Parasites (5 x 10^5 cells/ml) were incubated overnight at 37°C in the presence of 100 μM

479   hydroxyurea (HU). The following day cells were resuspended in fresh medium without

480   HU, divided into 8 ml aliquots and left to grow at 37°C. An aliquot was taken every two

481   hours and was used for flow cytometry analysis to determine cell cycle stage. The sample

482   left was used for genomic DNA extraction to determine abundance of 6mA.

Flow cytometry

Parasites were treated as described previously [65]. Parasites were fixed in 5 ml of ice-

cold 100% EtOH, and incubated at 4°C overnight. Each sample was washed once in 1 ml

PBS containing 2% v/v bovine serum (BS) and resuspended in 1 ml PBS containing 180

μg/ml RNase A and 2% v/v BS. After 30 min incubation at 37°C samples were stained

with 25 μg/ml propidium iodide (PI) solution and incubated an additional 30 min at 37°C.

Samples were analyzed using a fluorescence-activated cell sorter (BD FACSCalibur; BD

Biosciences, San José, CA, USA) with appropriate filter sets. Data was analyzed using

FlowJo 7.6 software.

MeDIP-seq

20 μg of purified genomic DNA from *T. vaginalis* strain B7268 [60] was diluted in 400

μl 1X IP buffer (10 mM Na-Phosphate pH 7, 140 mM NaCl, 0.05% Triton X-100) and

fragmented to 100-500 bp using a water-bath sonicator (Amp: 50%, Cycle: 30 sec ON

and 30 sec OFF, Time: 20 min). Sample was divided into aliquots containing 6-7 μg of

sonicated DNA and heat-denatured. 1 μg was saved and stored at -20°C to use as input

control, the rest was immunoprecipitated overnight at 4°C using 5 μg of anti-6mA

antibody (Abcam) bound to protein A magnetic beads (Invitrogen). After three washes

with 1X IP buffer, beads containing bound methylated DNA were resuspended in 250 μl

digestion buffer (50 mM Tris pH 8, 10 mM EDTA, 0.5% SDS) with 100 μg Proteinase K

(Roche) and incubated at 65°C for 4 h. Eluted DNA and input control samples were

purified using phenol/choloroform/isoamyl alchohol extraction followed by EtOH

precipitation. MeDIP samples were sent to Applied Biological Materials Inc. for library

preparation and paired-end sequencing. Two independent high-throughput sequencing

experiments, each containing pooled DNA from three independent immunoprecipitation,

were performed.

Bioinformatics analysis of MeDIP-seq data

After the adaptors were trimmed, alignment of high-quality fastq reads to the reference

genome (GCF 000002825.2 ASM282v1) was performed using Bowtie2 [66] with –qc-

filter parameter and all other parameters at default settings. The BAM files were used as

input for MACS2[67], which was run with the –g parameter set at 1.76e8 and all other

parameters at default settings. Input reads were used as control sample for peak calling.

Peaks were then filtered by q-value (< 0.01) and fold enrichment (> 2). Only peaks with

more than 75% overlap between replicates were considered for further analysis.

HOMER2 [68] was used for peak annotation. For repetitive element annotation,

RepeatMasker v4.0. [69] was used with a custom library of *T. vaginalis* specific RE taken

from Repbase [70]. GO enrichment analysis was performed using the Results Analysis

feature at TrichDB with default settings.

Chromatin conformation capture assay

Crosslinking of DNA was achieved by incubation of 50 ml (~1 x 10^6 cells/ml) of *T.*

vaginalis strain B7268 [60] culture with 1% formaldehyde for 15 min at room

temperature. Crosslinking was stopped by the addition of 125 mM glycine, followed by

5 min incubation at room temperature and then on ice for 15 min. Cells were harvested,

washed in 10 ml PBS/PI and resuspended in 1.4 ml of cold lysis buffer (10 mM Tris-HCl

pH 8, 10 mM NaCl, 0.2% NP-40, 1X PI) followed by homogenization with a dounce

homogenizer pestle A. After centrifugation, the pellet was washed once with 1.25X

Buffer H (Promega) and resuspended in 500 ul of 1.25X Buffer H (Promega). A total of

7.5 μl of SDS 20% was added and incubated for 40 min at 65°C and then for 20 min at

37°C. SDS was quenched with 50 μl of 20% Triton x-100 at 37°C for 1 h. 400U of EcoRI

(Promega) was added and incubated overnight at 37°C followed by enzyme heat

inactivation with 1% SDS at 65 °C for 20 min. Approximately 30 μl of digested sample

was set aside to test restriction enzyme efficiency by electrophoresis and PCR. Digestion

product was cooled and transferred to a pre-chilled 15 ml tube containing 4 ml 1.1X

Ligase Buffer (Promega) and 1% Triton X-100. After 1 h incubation at 37°C the sample

was cooled down and mixed with 30 Weiss Units of T4 DNA Ligase (Promega) and

incubated for 4 h in a water bath at 16°C. Crosslinking was reversed overnight at 65 °C

in the presence of 300 μg of Proteinase K (Roche). DNA was purified using

phenol/choloroform/isoamyl alchohol extraction followed by EtOH precipitation. The

sequences of primers used in 3C assays are shown in Sup. Table 8. 3C PCR fragments

were purified and sequenced to verify the identity of the cross-ligation product.

ACCESSION NUMBERS

All sequencing data that support the findings of this study have been deposited in the

NCBI Sequence Read Archive (SRA) and are accessible through the SRA accession

number PRJNA526331. All other relevant data are available from the corresponding

author on request

549   ACKNOWLEDGEMENTS

550   We thank Dr. Pablo Manavella and our colleagues in the lab for helpful discussions.

552   FUNDING

This research was supported with a grant from the ANPCyT grant BID PICT 2015-2118

(NdM). Work in the Greer lab was supported by NIH grants DP2AG055947 and

R21HG010066 (ELG) . NdM and PHS are researchers from the National Council of

Research (CONICET) and UNSAM. AL is a PhD fellow from CONICET. The funders

had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

560   CONFLICT OF INTEREST

561   The authors declare no competing interests.

563   REFERENCES

1.   Kissinger P. Trichomonas vaginalis: a review of epidemiologic, clinical and

treatment issues. BMC Infect Dis. 2015;15:307. Epub 2015/08/06. doi: 10.1186/s12879-

015-1055-0. PubMed PMID: 26242185; PubMed Central PMCID: PMCPMC4525749.

2.   Silver BJ, Guy RJ, Wand H, Ward J, Rumbold AR, Fairley CK, et al. Incidence

of curable sexually transmissible infections among adolescents and young adults in

remote Australian Aboriginal communities: analysis of longitudinal clinical service data.

Sex Transm Infect. 2015;91(2):135-41. Epub 2014/11/06. doi: 10.1136/sextrans-2014-

051617. PubMed PMID: 25371420.

3.   Swygard H, Sena AC, Hobbs MM, Cohen MS. Trichomoniasis: clinical

manifestations, diagnosis and management. Sex Transm Infect. 2004;80(2):91-5. Epub

2004/04/01. PubMed PMID: 15054166; PubMed Central PMCID: PMCPMC1744792.

4.   Graham SM, Holte SE, Peshu NM, Richardson BA, Panteleeff DD, Jaoko WG, et

al. Initiation of antiretroviral therapy leads to a rapid decline in cervical and vaginal HIV-

1   shedding.   AIDS.   2007;21(4):501-7.   Epub   2007/02/16.   doi:

10.1097/QAD.0b013e32801424bd. PubMed PMID: 17301569.

5.   Gander S, Scholten V, Osswald I, Sutton M, van Wylick R. Cervical dysplasia

and associated risk factors in a juvenile detainee population. J Pediatr Adolesc Gynecol.

2009;22(6):351-5. Epub 2009/07/14. doi: 10.1016/j.jpag.2009.01.070. PubMed PMID:

19592281.

6.   Twu O, Dessi D, Vu A, Mercer F, Stevens GC, de Miguel N, et al. Trichomonas

vaginalis homolog of macrophage migration inhibitory factor induces prostate cell

growth, invasiveness, and inflammatory responses. Proc Natl Acad Sci U S A.

2014;111(22):8179-84. Epub 2014/05/21. doi: 10.1073/pnas.1321884111. PubMed

PMID: 24843155; PubMed Central PMCID: PMCPMC4050605.

7.   Lv Z, Wu Z, Zhang L, Ji P, Cai Y, Luo S, et al. Genome mining offers a new

starting point for parasitology research. Parasitol Res. 2015;114(2):399-409. Epub

2015/01/08. doi: 10.1007/s00436-014-4299-5. PubMed PMID: 25563615.

8.   Carlton JM, Hirt RP, Silva JC, Delcher AL, Schatz M, Zhao Q, et al. Draft genome

sequence of the sexually transmitted pathogen Trichomonas vaginalis. Science.

2007;315(5809):207-12. Epub 2007/01/16. doi: 10.1126/science.1132894. PubMed

PMID: 17218520; PubMed Central PMCID: PMCPMC2080659.

9.   Gould SB, Woehle C, Kusdian G, Landan G, Tachezy J, Zimorski V, et al. Deep

sequencing of Trichomonas vaginalis during the early infection of vaginal epithelial cells

and amoeboid transition. Int J Parasitol. 2013;43(9):707-19. Epub 2013/05/22. doi:

10.1016/j.ijpara.2013.04.002. PubMed PMID: 23688716.

10.   Huang KY, Chen YY, Fang YK, Cheng WH, Cheng CC, Chen YC, et al. Adaptive

responses to glucose restriction enhance cell survival, antioxidant capability, and

autophagy of the protozoan parasite Trichomonas vaginalis. Biochim Biophys Acta.

2014;1840(1):53-64. Epub 2013/08/21. doi: 10.1016/j.bbagen.2013.08.008. PubMed

PMID: 23958562.

11.   Horvathova L, Safarikova L, Basler M, Hrdy I, Campo NB, Shin JW, et al.

Transcriptomic identification of iron-regulated and iron-independent gene copies within

the heavily duplicated Trichomonas vaginalis genome. Genome Biol Evol.

2012;4(10):1017-29. Epub 2012/09/15. doi: 10.1093/gbe/evs078. PubMed PMID:

22975721; PubMed Central PMCID: PMCPMC3490414.

12.   Smith A, Johnson P. Gene expression in the unicellular eukaryote Trichomonas

vaginalis. Res Microbiol. 2011;162(6):646-54. Epub 2011/04/23. doi:

10.1016/j.resmic.2011.04.007. PubMed PMID: 21511031.

13.   Liston DR, Lau AO, Ortiz D, Smale ST, Johnson PJ. Initiator recognition in a

primitive eukaryote: IBP39, an initiator-binding protein from Trichomonas vaginalis.

Mol Cell Biol. 2001;21(22):7872-82. Epub 2001/10/18. doi: 10.1128/MCB.21.22.7872-

7882.2001. PubMed PMID: 11604521; PubMed Central PMCID: PMCPMC99959.

14.   Schumacher MA, Lau AO, Johnson PJ. Structural basis of core promoter

recognition in a primitive eukaryote. Cell. 2003;115(4):413-24. Epub 2003/11/19.

PubMed PMID: 14622596.

15.   Smith AJ, Chudnovsky L, Simoes-Barbosa A, Delgadillo-Correa MG, Jonsson

ZO, Wohlschlegel JA, et al. Novel core promoter elements and a cognate transcription

factor in the divergent unicellular eukaryote Trichomonas vaginalis. Mol Cell Biol.

2011;31(7):1444-58. Epub 2011/01/20. doi: 10.1128/MCB.00745-10. PubMed PMID:

21245378; PubMed Central PMCID: PMCPMC3135286.

16.   Ong SJ, Huang SC, Liu HW, Tai JH. Involvement of multiple DNA elements in

iron-inducible transcription of the ap65-1 gene in the protozoan parasite Trichomonas

vaginalis. Mol Microbiol. 2004;52(6):1721-30. Epub 2004/06/10. doi: 10.1111/j.1365-

2958.2004.04088.x. PubMed PMID: 15186420.

17.   Golbabapour S, Abdulla MA, Hajrezaei M. A concise review on epigenetic

regulation: insight into molecular mechanisms. Int J Mol Sci. 2011;12(12):8661-94. Epub

2012/01/25. doi: 10.3390/ijms12128661. PubMed PMID: 22272098; PubMed Central

PMCID: PMCPMC3257095.

18.   Croken MM, Nardelli SC, Kim K. Chromatin modifications, epigenetics, and how

protozoan parasites regulate their lives. Trends Parasitol. 2012;28(5):202-13. Epub

2012/04/07. doi: 10.1016/j.pt.2012.02.009. PubMed PMID: 22480826; PubMed Central

PMCID: PMCPMC3340475.

19. Song MJ, Kim M, Choi Y, Yi MH, Kim J, Park SJ, et al. Epigenome mapping highlights chromatin-mediated gene regulation in the protozoan parasite Trichomonas vaginalis. Sci Rep. 2017;7:45365. Epub 2017/03/28. doi: 10.1038/srep45365. PubMed PMID: 28345651; PubMed Central PMCID: PMCPMC5366954.

20. Pachano T, Nievas YR, Lizarraga A, Johnson PJ, Strobl-Mazzulla PH, de Miguel N. Epigenetics regulates transcription and pathogenesis in the parasite Trichomonas vaginalis. Cell Microbiol. 2017;19(6). Epub 2017/01/06. doi: 10.1111/cmi.12716. PubMed PMID: 28054438.

21. Chen K, Zhao BS, He C. Nucleic Acid Modifications in Regulation of Gene Expression. Cell Chem Biol. 2016;23(1):74-85. Epub 2016/03/05. doi: 10.1016/j.chembiol.2015.11.007. PubMed PMID: 26933737; PubMed Central PMCID: PMCPMC4779186.

22. Jones PA, Takai D. The role of DNA methylation in mammalian epigenetics. Science. 2001;293(5532):1068-70. Epub 2001/08/11. doi: 10.1126/science.1063852. PubMed PMID: 11498573.

23. Wei H, Jiang S, Chen L, He C, Wu S, Peng H. Characterization of Cytosine Methylation and the DNA Methyltransferases of Toxoplasma gondii. Int J Biol Sci. 2017;13(4):458-70. Epub 2017/05/23. doi: 10.7150/ijbs.18644. PubMed PMID: 28529454; PubMed Central PMCID: PMCPMC5436566.

24. Ponts N, Fu L, Harris EY, Zhang J, Chung DW, Cervantes MC, et al. Genome-wide mapping of DNA methylation in the human malaria parasite Plasmodium falciparum. Cell Host Microbe. 2013;14(6):696-706. Epub 2013/12/18. doi: 10.1016/j.chom.2013.11.007. PubMed PMID: 24331467; PubMed Central PMCID: PMCPMC3931529.

25.   Fisher O, Siman-Tov R, Ankri S. Characterization of cytosine methylated regions

and 5-cytosine DNA methyltransferase (Ehmeth) in the protozoan parasite Entamoeba

histolytica.   Nucleic   Acids   Res.   2004;32(1):287-97.   Epub   2004/01/13.   doi:

10.1093/nar/gkh161.   PubMed   PMID: 14715927;   PubMed   Central   PMCID:

PMCPMC373271.

26.   Luo GZ, Blanco MA, Greer EL, He C, Shi Y. DNA N(6)-methyladenine: a new

epigenetic mark in eukaryotes? Nat Rev Mol Cell Biol. 2015;16(12):705-10. Epub

2015/10/29. doi: 10.1038/nrm4076. PubMed   PMID: 26507168;   PubMed   Central

PMCID: PMCPMC4763336.

27.   Vanyushin BF, Tkacheva SG, Belozersky AN. Rare bases in animal DNA. Nature.

1970;225(5236):948-9. Epub 1970/03/07. PubMed PMID: 4391887.

28.   Zhang G, Huang H, Liu D, Cheng Y, Liu X, Zhang W, et al. N6-methyladenine

DNA modification in Drosophila. Cell. 2015;161(4):893-906. Epub 2015/05/06. doi:

10.1016/j.cell.2015.04.018. PubMed PMID: 25936838.

29.   Greer EL, Blanco MA, Gu L, Sendinc E, Liu J, Aristizabal-Corrales D, et al. DNA

Methylation on N6-Adenine in C. elegans. Cell. 2015;161(4):868-78. Epub 2015/05/06.

doi: 10.1016/j.cell.2015.04.005. PubMed PMID: 25936839; PubMed Central PMCID:

PMCPMC4427530.

30.   Fu Y, Luo GZ, Chen K, Deng X, Yu M, Han D, et al. N6-methyldeoxyadenosine

marks active transcription start sites in Chlamydomonas. Cell. 2015;161(4):879-92. Epub

2015/05/06.   doi: 10.1016/j.cell.2015.04.010.   PubMed   PMID: 25936837;   PubMed

Central PMCID: PMCPMC4427561.

31.   Wu TP, Wang T, Seetin MG, Lai Y, Zhu S, Lin K, et al. DNA methylation on

N(6)-adenine in mammalian embryonic stem cells. Nature. 2016;532(7599):329-33.

Epub 2016/03/31. doi: 10.1038/nature17640. PubMed PMID: 27027282; PubMed

Central PMCID: PMCPMC4977844.

32.   Liu J, Zhu Y, Luo GZ, Wang X, Yue Y, Wang X, et al. Abundant DNA 6mA

methylation during early embryogenesis of zebrafish and pig. Nat Commun.

2016;7:13052. Epub 2016/10/08. doi: 10.1038/ncomms13052. PubMed PMID:

27713410; PubMed Central PMCID: PMCPMC5059759.

33.   Koziol MJ, Bradshaw CR, Allen GE, Costa ASH, Frezza C, Gurdon JB.

Identification of methylated deoxyadenosines in vertebrates reveals diversity in DNA

modifications. Nat Struct Mol Biol. 2016;23(1):24-30. Epub 2015/12/23. doi:

10.1038/nsmb.3145. PubMed PMID: 26689968; PubMed Central PMCID:

PMCPMC4941928.

34.   Mondo SJ, Dannebaum RO, Kuo RC, Louie KB, Bewick AJ, LaButti K, et al.

Widespread adenine N6-methylation of active genes in fungi. Nat Genet. 2017;49(6):964-

8. Epub 2017/05/10. doi: 10.1038/ng.3859. PubMed PMID: 28481340.

35.   Meyer KD, Jaffrey SR. Expanding the diversity of DNA base modifications with

N(6)-methyldeoxyadenosine. Genome Biol. 2016;17:5. Epub 2016/01/15. doi:

10.1186/s13059-016-0874-7. PubMed PMID: 26762117; PubMed Central PMCID:

PMCPMC4712593.

36.   Liang Z, Shen L, Cui X, Bao S, Geng Y, Yu G, et al. DNA N(6)-Adenine

Methylation in Arabidopsis thaliana. Dev Cell. 2018;45(3):406-16 e3. Epub 2018/04/17.

doi: 10.1016/j.devcel.2018.03.012. PubMed PMID: 29656930.

37.   Zhou C, Wang C, Liu H, Zhou Q, Liu Q, Guo Y, et al. Identification and analysis

of adenine N(6)-methylation sites in the rice genome. Nat Plants. 2018;4(8):554-63. Epub

2018/08/01. doi: 10.1038/s41477-018-0214-x. PubMed PMID: 30061746.

38.   Bradic M, Warring SD, Tooley GE, Scheid P, Secor WE, Land KM, et al. Genetic

Indicators of Drug Resistance in the Highly Repetitive Genome of Trichomonas

vaginalis. Genome Biol Evol. 2017;9(6):1658-72. Epub 2017/06/22. doi:

10.1093/gbe/evx110. PubMed PMID: 28633446; PubMed Central PMCID:

PMCPMC5522705.

39.   Piknova M, Pristas P, Javorsky P, Kasperowic A, Michalowski T. GATC-specific

restriction and modification systems in treponemes. Lett Appl Microbiol.

2004;38(4):311-4. Epub 2004/06/25. PubMed PMID: 15214731.

40.   Fichorova R, Fraga J, Rappelli P, Fiori PL. Trichomonas vaginalis infection in

symbiosis with Trichomonasvirus and Mycoplasma. Res Microbiol. 2017;168(9-10):882-

91. Epub 2017/04/04. doi: 10.1016/j.resmic.2017.03.005. PubMed PMID: 28366838.

41.   Weber M, Davies JJ, Wittig D, Oakeley EJ, Haase M, Lam WL, et al.

Chromosome-wide and promoter-specific analyses identify sites of differential DNA

methylation in normal and transformed human cells. Nat Genet. 2005;37(8):853-62. Epub

2005/07/12. doi: 10.1038/ng1598. PubMed PMID: 16007088.

42.   Leitsch D, Janssen BD, Kolarich D, Johnson PJ, Duchene M. Trichomonas

vaginalis flavin reductase 1 and its role in metronidazole resistance. Mol Microbiol.

2014;91(1):198-208. Epub 2013/11/22. doi: 10.1111/mmi.12455. PubMed PMID:

24256032; PubMed Central PMCID: PMCPMC4437529.

43.   Denker A, de Laat W. The second decade of 3C technologies: detailed insights

into nuclear organization. Genes Dev. 2016;30(12):1357-82. Epub 2016/06/25. doi:

10.1101/gad.281964.116. PubMed PMID: 27340173; PubMed Central PMCID:

PMCPMC4926860.

44.   Rao SS, Huntley MH, Durand NC, Stamenova EK, Bochkov ID, Robinson JT, et

al. A 3D map of the human genome at kilobase resolution reveals principles of chromatin

looping. Cell. 2014;159(7):1665-80. Epub 2014/12/17. doi: 10.1016/j.cell.2014.11.021.

PubMed PMID: 25497547; PubMed Central PMCID: PMCPMC5635824.

45.   Oyinlade O, Wei S, Kammers K, Liu S, Wang S, Ma D, et al. Analysis of KLF4

regulated genes in cancer cells reveals a role of DNA methylation in promoter- enhancer

interactions.   Epigenetics.   2018;13(7):751-68.   Epub   2018/07/31.   doi:

10.1080/15592294.2018.1504592. PubMed PMID: 30058478; PubMed Central PMCID:

PMCPMC6224223.

46.   Noel CJ, Diaz N, Sicheritz-Ponten T, Safarikova L, Tachezy J, Tang P, et al.

Trichomonas vaginalis vast BspA-like gene family: evidence for functional diversity

from structural organisation and transcriptomics. BMC Genomics. 2010;11:99. Epub

2010/02/11. doi: 10.1186/1471-2164-11-99. PubMed PMID: 20144183; PubMed Central

PMCID: PMCPMC2843621.

47.   Yu XX, Liu YH, Liu XM, Wang PC, Liu S, Miao JK, et al. Ascorbic acid induces

global epigenetic reprogramming to promote meiotic maturation and developmental

competence of porcine oocytes. Sci Rep. 2018;8(1):6132. Epub 2018/04/19. doi:

10.1038/s41598-018-24395-y. PubMed PMID: 29666467; PubMed Central PMCID:

PMCPMC5904140.

48.   Singh A, Vancura A, Woycicki RK, Hogan DJ, Hendrick AG, Nowacki M.

Determination of the presence of 5-methylcytosine in Paramecium tetraurelia. PLoS One.

2018;13(10):e0206667. Epub 2018/11/01. doi: 10.1371/journal.pone.0206667. PubMed

PMID: 30379964; PubMed Central PMCID: PMCPMC6209305 by Storm Therapeutics

Limited. There are no patents, products in development or marketed products to declare.

This does not alter our adherence to all the PLOS ONE policies on sharing data and

materials.

49.　　Wang Y, Chen X, Sheng Y, Liu Y, Gao S. N6-adenine DNA methylation is

associated with the linker DNA of H2A.Z-containing well-positioned nucleosomes in Pol

II-transcribed genes in Tetrahymena. Nucleic Acids Res. 2017;45(20):11594-606. Epub

2017/10/17. doi: 10.1093/nar/gkx883. PubMed PMID: 29036602; PubMed Central

PMCID: PMCPMC5714169.

50.　　Iyer LM, Zhang D, Aravind L. Adenine methylation in eukaryotes: Apprehending

the complex evolutionary history and functional potential of an epigenetic modification.

Bioessays. 2016;38(1):27-40. Epub 2015/12/15. doi: 10.1002/bies.201500104. PubMed

PMID: 26660621; PubMed Central PMCID: PMCPMC4738411.

51.　　Zhu JK. Active DNA demethylation mediated by DNA glycosylases. Annu Rev

Genet. 2009;43:143-66. Epub 2009/08/08. doi: 10.1146/annurev-genet-102108-134205.

PubMed PMID: 19659441; PubMed Central PMCID: PMCPMC3137514.

52.　　Ong CT, Corces VG. CTCF: an architectural protein bridging genome topology

and function. Nat Rev Genet. 2014;15(4):234-46. Epub 2014/03/13. doi:

10.1038/nrg3663. PubMed PMID: 24614316; PubMed Central PMCID:

PMCPMC4610363.

53.　　Rao SSP, Huang SC, Glenn St Hilaire B, Engreitz JM, Perez EM, Kieffer-Kwon

KR, et al. Cohesin Loss Eliminates All Loop Domains. Cell. 2017;171(2):305-20 e24.

Epub 2017/10/07. doi: 10.1016/j.cell.2017.09.026. PubMed PMID: 28985562; PubMed

Central PMCID: PMCPMC5846482.

54.　　Rowley MJ, Corces VG. Organizational principles of 3D genome architecture.

Nat Rev Genet. 2018;19(12):789-800. Epub 2018/10/28. doi: 10.1038/s41576-018-0060-

8. PubMed PMID: 30367165; PubMed Central PMCID: PMCPMC6312108.

55.　　Eagen KP, Aiden EL, Kornberg RD. Polycomb-mediated chromatin loops

revealed by a subkilobase-resolution chromatin interaction map. Proc Natl Acad Sci U S

A. 2017;114(33):8764-9. Epub 2017/08/03. doi: 10.1073/pnas.1701291114. PubMed

PMID: 28765367; PubMed Central PMCID: PMCPMC5565414.

56.   Guo Y, Perez AA, Hazelett DJ, Coetzee GA, Rhie SK, Farnham PJ. CRISPR-

mediated deletion of prostate cancer risk-associated CTCF loop anchors identifies

repressive chromatin loops. Genome Biol. 2018;19(1):160. Epub 2018/10/10. doi:

10.1186/s13059-018-1531-0. PubMed PMID: 30296942; PubMed Central PMCID:

PMCPMC6176514.

57.   Hirt RP. Trichomonas vaginalis virulence factors: an integrative overview. Sex

Transm Infect. 2013;89(6):439-43. Epub 2013/05/23. doi: 10.1136/sextrans-2013-

051105. PubMed PMID: 23694938; PubMed Central PMCID: PMCPMC3749517.

58.   Ay F, Bunnik EM, Varoquaux N, Bol SM, Prudhomme J, Vert JP, et al. Three-

dimensional modeling of the P. falciparum genome during the erythrocytic cycle reveals

a strong connection between genome architecture and gene expression. Genome Res.

2014;24(6):974-88. Epub 2014/03/29. doi: 10.1101/gr.169417.113. PubMed PMID:

24671853; PubMed Central PMCID: PMCPMC4032861.

59.   Bunnik EM, Cook KB, Varoquaux N, Batugedara G, Prudhomme J, Cort A, et al.

Changes in genome organization of parasite-specific gene families during the

Plasmodium transmission stages. Nat Commun. 2018;9(1):1910. Epub 2018/05/17. doi:

10.1038/s41467-018-04295-5. PubMed PMID: 29765020; PubMed Central PMCID:

PMCPMC5954139.

60.   Upcroft JA, Upcroft P. Drug susceptibility testing of anaerobic protozoa.

Antimicrob Agents Chemother. 2001;45(6):1810-4. Epub 2001/05/17. doi:

10.1128/AAC.45.6.1810-1814.2001. PubMed PMID: 11353630; PubMed Central

PMCID: PMCPMC90550.

61.   Clark CG, Diamond LS. Methods for cultivation of luminal parasitic protists of

clinical importance. Clin Microbiol Rev. 2002;15(3):329-41. Epub 2002/07/05. PubMed

PMID: 12097242; PubMed Central PMCID: PMCPMC118080.

62.   Blanchard A, Yanez A, Dybvig K, Watson HL, Griffiths G, Cassell GH.

Evaluation of intraspecies genetic variation within the 16S rRNA gene of Mycoplasma

hominis and detection by polymerase chain reaction. J Clin Microbiol. 1993;31(5):1358-

61. Epub 1993/05/01. PubMed PMID: 7684753; PubMed Central PMCID:

PMCPMC262938.

63.   de Miguel N, Lustig G, Twu O, Chattopadhyay A, Wohlschlegel JA, Johnson PJ.

Proteome analysis of the surface of Trichomonas vaginalis reveals novel proteins and

strain-dependent differential expression. Mol Cell Proteomics. 2010;9(7):1554-66. Epub

2010/05/15. doi: 10.1074/mcp.M000022-MCP201. PubMed PMID: 20467041; PubMed

Central PMCID: PMCPMC2938091.

64.   Mercer F, Diala FG, Chen YP, Molgora BM, Ng SH, Johnson PJ. Leukocyte Lysis

and Cytokine Induction by the Human Sexually Transmitted Parasite Trichomonas

vaginalis. PLoS Negl Trop Dis. 2016;10(8):e0004913. Epub 2016/08/17. doi:

10.1371/journal.pntd.0004913. PubMed PMID: 27529696; PubMed Central PMCID:

PMCPMC4986988.

65.   Iriarte LS, Midlej V, Frontera LS, Moros Duarte D, Barbeito CG, de Souza W, et

al. TfVPS32 Regulates Cell Division in the Parasite Tritrichomonas foetus. J Eukaryot

Microbiol. 2018;65(1):28-37. Epub 2017/05/10. doi: 10.1111/jeu.12424. PubMed PMID:

28477402.

66.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat

Methods. 2012;9(4):357-9. Epub 2012/03/06. doi: 10.1038/nmeth.1923. PubMed PMID:

22388286; PubMed Central PMCID: PMCPMC3322381.

67.   Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-

based analysis of ChIP-Seq (MACS). Genome Biol. 2008;9(9):R137. Epub 2008/09/19.

doi: 10.1186/gb-2008-9-9-r137. PubMed PMID: 18798982; PubMed Central PMCID:

PMCPMC2592715.

68.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple

combinations of lineage-determining transcription factors prime cis-regulatory elements

required for macrophage and B cell identities. Mol Cell. 2010;38(4):576-89. Epub

2010/06/02. doi: 10.1016/j.molcel.2010.05.004. PubMed PMID: 20513432; PubMed

Central PMCID: PMCPMC2898526.

69.   Smit A, Hubley, R & Green, P. RepeatMasker Open-4.0. 2013-2015

70.   Bao W, Kojima KK, Kohany O. Repbase Update, a database of repetitive

elements in eukaryotic genomes. Mob DNA. 2015;6:11. Epub 2015/06/06. doi:

10.1186/s13100-015-0041-9. PubMed PMID: 26045719; PubMed Central PMCID:

PMCPMC4455052.

FIGURE LEGENDS

Figure 1: DNA methylation in *T. vaginalis* genome. A) Dot blot assay against 5mC

shows lack of signal in the genomic DNA of highly-adherent strain B7268 and less-

adherent strain G3. Left: gDNA dots stained with methylene blue for loading control.

Right: 5mC signal. Chicken gDNA was used as positive control. B) Immunofluorescence

assay using specific antibodies fails to detect 5mC in *T. vaginalis* nucleus. Nucleus was

stained with DAPI. C) Immunofluorescence assay using anti-6mA specific antibody

evidenced this modification in the cytoplasm (RNA) and nucleus (gDNA), or exclusively

in the nucleus of RNase-treated parasites. Nucleus was stained with DAPI. D)

Methylation sensitive RE assay shows the presence of methylated and un-methylated

adenines in *T. vaginalis* gDNA. Lane 1: MW. Lane 2: undigested gDNA. Lane 3: MboI

treated gDNA (cuts GATC sites). Lane 4: DpnI treated gDNA (cuts G6mATC sites). E)

Anti-6mA dot blot assay of RNA-free gDNA shows similar abundance of 6mA in the

genomic DNA of strain B7268 and strain G3. Left: gDNA dots stained with methylene

blue for loading control. Right: 6mA signal. F) Quantification of methylated bases in the

gDNA of adherent strain B7268 and less-adherent strain G3 using UHPLC followed by

mass spectrometry shows high levels of 6mA and low levels of 5mC, but not differences

among stains for both modifications. Each column represents the mean and SD of three

independent experiments per group. Graphic is shown in logarithmic scale.

Figure 2: 6mA MeDIP-seq of adherent strain B7268. A) Schematic representation of

the adapted MeDIP-seq protocol. B) Dot blot assay of the different MeDIP fractions

shows enrichment of 6mA-containing fragments in the 6mA-IP fraction compared to the

input control. Lack of signal in the IgG-IP fraction shows the specificity of 6mA antibody.

C) Workflow showing final peak count obtained from two independent 6mA MeDIP-seq

experiments. Peaks were filtered by q-value (<0.01) and fold enrichment (>2). Only peaks

with at least 75% overlap between experiments were considered for further study. D)

Genome browser visualization of 6mA peaks. Representative image of three different

contigs (DS113177, DS113178 and DS113181) shows 6mA possesses a primarily

intergenic localization. To row: 6mA peaks. Bottom row: *T. vaginalis* genome

Figure 3: Genome distribution of 6mA peaks. A) 6mA is enriched at intergenic regions.

Annotated peaks were grouped according to their genomic localization: peaks that fall

within the transcription start sites (TSS), coding region and transcription termination sites

(TTS) were considered gene peaks and grouped together, the rest of the peaks were

considered intergenic. MeDIP: Percentages of intergenic and gene peaks according

MeDIP-seq analysis. Shuffled: MeDIP-seq peaks were randomly distributed across the

genome and the percentage of gene and intergenic peaks was calculated. B) Enrichment

of 6mA at the TTS region of methylated genes. MeDIP: Percentages of peaks within the

TSS, CDS and TTS regions of methylated genes according MeDIP-seq analysis.

Shuffled: Peaks randomly distributed across the genome. C) Identity of *T. vaginalis*

methylated genes. A total of 642 MeDIP peaks fall within 512 genes: 61% belong to

transposable elements described in the parasite (*T. vginalis* TEs), 9% of genes possess

sequence similarity to repetitive elements found in other organisms (other TEs), 22% are

T. vaginalis specific hypothetical proteins with unknown functions (hypothetical

proteins), 8% are genes with sequence similarity to genes from other organisms (genes

with known domains). D) Identity of *T. vaginalis* methylated repetitive elements. Only

intergenic peaks with more than 50% overlap with a *T. vaginalis* RE were considered,

obtaining a total of 5596 peaks. Most represented DNA transposon superfamilies included

Mutator (MuDR) with 24% of all peaks, Polinton (27%) and Kolobok (16%). Less

represented TE superfamilies included hAT (8%), Mariner (5%) and P (8%). 10% of the

peaks were found in *T. vaginalis* DNA repeat DNA7.

Figure 4: Expression levels of genes closest to 6mA DNA methylation. A) Intragenic

6mA methylation is associated with low expression. 82% of the 512 identified methylated

genes are poorly expressed (RPKM =< 1), 14% moderately expressed (1 < RPKM =<

20), and 4% highly expressed (RPKM > 20). B) Percentage of genes with different

expression levels near intergenic 6mA peaks. Of the 6000 genes closest to intergenic

methylation, 3023 have low expression, 2106 have moderate expression and 871 genes

have high expression, corresponding to 7,9%, 13,5% and 16,4% of all *T. vaginalis* poorly,

moderately and highly expressed genes respectively. C) Bar plot comparing the median

and interquartile range of RPKM expression of genes near intergenic 6mA and genes in

non-methylated regions shows genes near intergenic methylation possess a significantly

higher expression than genes in non-methylated regions. The p-value was calculated by

a two-sided Mann-Whitney test (P<0.0001). D) Distribution of methylation frequency

show intergenic 6mA is primarily located between -1500pb and +1000bp of the nearest

not expressed (RPKM =< 1) or expressed (RPKM > 1) genes. Methylation frequency of

intergenic peaks was calculated and plotted against the distance to the nearest gene, with

0 corresponding to peaks that fall within genes.

Figure 5: Chromatin conformation capture assay. A) Representation of intervals

delimited by 6mA peaks comprised of genes with the similar level of expression. 1046

intervals were found, 452 containing not expressed genes (red) and 592 containing

expressed genes (green). B) Quantification of intervals flanked by 6mA containing

different numbers of expressed (grey) and non-expressed (white) genes.

Figure 6: 3C assay of parasites treated with ascorbic acid. A) Schematic

representation of the chromatin conformation capture assay (3C). B) 3C assay shows

evidence of loop formation in *T. vaginalis* genome. Lanes 1-4: serial dilutions of

uncrosslinked control template. Lanes 5: undigested gDNA. Lanes 6-9: serial dilutions of

3C template. Loop p5512-5513: amplification of a region containing a block of

transcriptionally-repressed genes. Loop p4130-4131: amplification of a region containing

a block of transcriptionally-active genes. Non-digested control: amplification of a region

without a restriction enzyme site in between the utilized primers. C) Quantification of

6mA levels of parasites treated with different concentrations of ascorbic acid (AA) shows

diminution of 6mA abundance in parasites treated with 100 mM AA compared to

untreated parasites. D) Parasite AA exposition prevents the formation of both p5512-5513

and p4130-4131 loops. Control: 3C assay of non-treated parasites. AA: 3C assay

performed with parasites treated with 100mM of AA. Same amount of template was used

for PCR amplification using primers for repressive loop p5512-5513 and active loop

p4130-4131. Non-digested control: amplification of a region without a restriction enzyme

site in between the utilized primers.

Supplementary Figure. 1: *T. vaginalis* cultures free of *M. hominis* contamination.

PCR using *M. hominis* specific primers. Lane 1: gDNA of parasites from strain G3

subjected to BM-cyclin treatment. Lane 2: gDNA of parasites from strain B7268

subjected to BM-cyclin treatment. Lane 3: negative control. Lane 4: gDNA of non-treated

parasites as positive control of *M. hominis* contamination.

Supplementary Figure 2: 6mA methylation levels during *T. vaginalis* cell cycle

progression. Samples from a hydroxyurea synchronized culture were taken every two

hours post-treatment. Part of the sample was used for cell cycle stage determination by

flow cytometry (A), the rest was used for gDNA extraction to measure 6mA levels using

a dot blot assay (B). Representative image of two independent experiments is shown. C)

Quantification of dot blot signal.

Supplementary Figure 3: MeDIP-seq analysis. A) Most prevalent methylation motif

identified by HOMER2, found in 4692 peaks (p-value = 1e-1642). B) Boxplot comparing

the median and interquartile range of gene distance to intergenic peaks found at the 5' of

expressed and not expressed genes. The p-value was calculated by a two-sided Mann-

Whitney test (P<0.0001) C). Boxplot comparing the median and interquartile range of

gene distance to intergenic peaks found at the 3' of expressed and not expressed genes.

The p-value was calculated by a two-tailed unpaired Student's t test (P < 0.05). D)

MeDIP-qPCR of four genes selected from MeDIP-seq demonstrated enrichment in 6mA

immunoprecipated sample (6mA-IP) compared to sample immunoprecipitated with IgG

control antibody (Ig-IP).

Supplementary Figure 4: Sanger sequencing of 3C PCR fragments. Amplified 3C

fragments of the appropriate length were purified and sequenced to corroborate they

belonged to the expected cross-ligation product. A) Sequencing of active loop p4130-

4131. B) Sequencing of repressive loop p5512-5513. EcoRI site is highlighted. C) PCR

amplification of active loop p2296-2297 shows band of the expected size. Lane 1:

undigested gDNA. Lane 2: 3C template. Lane 3: MW. D) Sanger sequencing confirms

the amplified fragment belongs to the expected cross-ligation product. EcoRI site is

highlighted.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A) Crosslinking → RE digestion → Intramolecular ligation → Reverse crosslink → PCR detection

B) Loop p4130-4131 (7 genes – expressed)
1-2
1-3

Loop p5512-5513 (11 genes – not expressed)
1-2
1-3

Non-digested control

Control   3C

C) 6mA levels: 0 mM, 10 mM, 50 mM, 100 mM

D) B7268 3C PCR
Loop p4130-4131
Loop p5512-5513
Non-digested control
Control   AA

Figure 6