Title: Reliability, sensitivity and predictive value of fMRI during multiple object

tracking as a marker of cognitive training gain in combination with tDCS in stroke

survivors

Authors: Knut K. Kolskår[a,b,c,*], Geneviève Richard[a,b,c], Dag Alnæs[a], Erlend S.Dørum[a,b,c],

Anne-Marthe Sanders[a,b,c], Kristine M. Ulrichsen[a,b,c,], Jennifer Monereo Sánchez[a], Hege Ihle-

Hansen[d], Jan E. Nordvik[e], Lars T. Westlye[a,c,f*]

[a] NORMENT, Division of Mental Health and Addiction, Oslo University Hospital & Institute

of Clinical Medicine, University of Oslo, Norway

[b] Sunnaas Rehabilitation Hospital HT, Nesodden, Norway

[c] Department of Psychology, University of Oslo, Norway

[d] Department of Geriatric Medicine, Oslo University Hospital, Oslo, Norway

[e] Catosenteret Rehabilitation Center, Son, Norway

[f] KG Jebsen Centre for Neurodevelopmental Disorders, University of Oslo, Norway

* Corresponding authors: Knut K. Kolskår (kolskaar@gmail.com, +47 995 97 425) & Lars T.

Westlye (l.t.westlye@psykologi.uio.no), Department of Psychology, University of Oslo, Pb.

1094, Blindern, 0317 OSLO, Norway, phone: +47 22845000, Fax: +47 22845001

Acknowledgements: This work was supported by the South-Eastern Norway Regional Health

Authority (2014097, 2015044, 2015073), the Norwegian ExtraFoundation for Health and

Rehabilitation (2015/FO5146), the Research Council of Norway (249795, 262372) and the

European Research Council under the European Union's Horizon 2020 research and

Innovation program (ERC StG, Grant 802998).

Data availability: The data that support the findings of this study are openly available in

https://osf.io/ndpuz/, at DOI 10.17605/OSF.IO/NDPUZ

Conflict of Interest: We declare no competing financial interests. We obtained appropriate

ethical approval from the local ethics committee, and all procedures were in line with the

declaration of Helsinki. Relevant data have been made available at osf.io:

https://osf.io/ndpuz/.

Keywords: stroke, rehabilitation, fMRI, cognitive training, tDCS

Abstract

Computerized cognitive training (CCT) combined with transcranial direct current stimulation

(tDCS) has showed some promise in alleviating cognitive impairments in patients with brain

disorders, but the robustness and possible mechanisms are unclear. In this prospective double-

blind randomized clinical trial, we investigated the feasibility and effectiveness of combining

CCT and tDCS, and tested the predictive value of and training-related changes in fMRI-based

brain activation during attentive performance (multiple object tracking) obtained before,

during and after the three-weeks intervention in chronic stroke patients (> 6 months since

hospital admission). Patients were randomized to one of two groups, receiving CCT and

either (1) tDCS targeting left dorsolateral prefrontal cortex (1 mA), or (2) sham tDCS, with

40s active stimulation (1 mA) before fade out of the current. 77 patients were enrolled in the

study, 54 completed the cognitive training, and 48 completed all training and MRI sessions.

We found significant improvement in performance across all trained tasks, but no additional

gain of tDCS. fMRI-based brain activation showed high reliability, and higher cognitive

performance was associated with increased tracking-related activation in the dorsal attention

network (DAN) and default mode network (DMN) as well as anterior cingulate after

compared to before the intervention. We found no significant associations between cognitive

gain and brain activation measured before training or in the difference in activation after

intervention. Combined, these results show significant training effects on trained cognitive

tasks in stroke survivors, with no clear evidence of additional gain of concurrent tDCS.

Introduction

While stroke mortality has decreased during the last decades in western countries, it remains a leading cause of disabilities (Feigin et al., 2017), and many stroke survivors experience persistent cognitive sequelae. Indeed, prevalence of neurocognitive disorders in post stroke survivors have been reported to as high as 54% (Barbay et al., 2018), and may increase risk of dementia (Pendlebury, 2012). Reduced working memory capacity is among the most common impairments (Teasell, 1992; Zinn et al., 2007), with high prevalence at the chronic stage (Mahon et al., 2017; Schaapsmeerders et al., 2013), and potentially large impact on daily functioning (Synhaeve et al., 2014). Stroke lesions influence brain activation as measured using brain imaging (Altamura et al., 2009), and working memory deficits have been linked to aberrant brain activation during task engagement (Ziemus et al., 2007).

Computerized cognitive training (CCT) has shown some promise in augmenting cognitive rehabilitation. A recent meta-analysis revealed small but significant improvements on cognitive functioning in response to CCT in healthy participants (Au et al., 2015). Similar findings have been reported in patients with Parkinson's disease (Leung et al., 2015), mild cognitive impairment (Maffei et al., 2017), brain injury (Weicker et al., 2016), and stroke (De Luca et al., 2018b; Peers et al., 2018), as well as schizophrenia (Prikken et al., 2018) and major depression (Motter et al., 2016). However, the underlying mechanisms remain elusive.

Furthermore, non-invasive electrical stimulation of the brain has been suggested to augment training benefits (Kang et al., 2015; Marquez et al., 2015). Indeed, evidence suggests amplified motor recovery post-stroke, when physical therapy is paired with transcranial direct current stimulation (tDCS) (Kang et al., 2015), and the beneficial effects have been suggested to generalize to a cognitive rehabilitation context (Andrews et al., 2011; Fregni et al., 2005; Lawrence et al., 2018). Despite some positive findings, the overall effectiveness of tDCS has been questioned (Medina and Cason, 2017). Also, the potential mechanisms, if any, are

unclear, yet enhancing neuronal excitability by lowering the membrane-potential has been

suggested as one of the candidate means (Giordano et al., 2017).

Functional MRI has become an increasingly used non-invasive brain imaging method

linking behavior to brain activation patterns. Studies investigating alterations in brain

activation in response to CCT have reported mixed results. In healthy controls, training gains

have been associated with decreased activity in medial and dorsal frontal areas, including

anterior cingulate (Heinzel et al., 2016; Miró-Padilla et al., 2018; Motes et al., 2018), with

increased efficiency reflected in lowered activity as a suggested cognitive mechanism

(Constantinidis and Klingberg, 2016). For patient cohorts the current evidence is sparser, but

increased activity in frontal and parietal areas in response to cognitive training has been

reported in patients with schizophrenia (Haut et al., 2010; Ramsay and MacDonald, 2015;

Subramaniam et al., 2014). In stroke patients, CCT has been associated with increased

functional connectivity (FC) between hippocampus, frontal and parietal areas, which was

associated with improved cognitive performance (Lin et al., 2014; Yang et al., 2014).

Despite some promising findings, the effects of CCT (Buitenweg et al., 2018; De Luca

et al., 2018a; Melby-Lervåg and Hulme, 2013; Melby-Lervåg et al., 2016) and tDCS

(Dedoncker et al., 2016; Hill et al., 2016) are still unclear. More knowledge about the origins

of individual differences in training gains is key to evaluate and improve selection to fully

realize the clinical potential of these interventions. Investigations of the neural underpinnings

of cognitive improvement in response to CCT and tDCS may help clarify the benefits and

limitations for future clinical application. It is therefore key to better understand mechanisms

besides behavioral transfer-effects before evaluating the full effect of these programs.

To this end, we investigated differences in training-related gains of a commonly used

working memory computerized training program in combination with tDCS, and tested the

predictive value and sensitivity of fMRI-based brain activation as a marker for training-

related gain in task performance. Specifically, we estimated activation patterns during

multiple object tracking (MOT) at baseline, before initiating training (on average 4 weeks

after baseline measure), and after a three-week intervention. The multiple object tracking

(MOT) task offers a promising tool for assessing and manipulate converging activation

patterns across attention and working memory tasks in both healthy controls and stroke

patients (Dørum et al., 2016; Pylyshyn and Strom, 1988; Walle et al., 2019). MOT robustly

recruit brain areas (Alnæs et al., 2014) comparable with activations associated with Cogmed

training (Olesen et al., 2004), including canonical task-positive and task-negative brain

networks. To assess specific associations with tDCS and following a double-blind design, in

addition to CCT each participant was pseudo-randomized into two groups: 50% (n=27)

receiving active tDCS directed towards the left dorsolateral prefrontal cortex and 50% (n=27)

receiving sham stimulation.

We tested the following hypotheses: (1) Variability in cognitive training gain is

reflected in differential slopes of increased performance across participants, (2) level of

improvement is affected by tDCS stimulation protocol, (3) fMRI-based brain activation

patterns prior to training are predictive of training gain, and (4) individual differences in task

improvement are reflected in differential patterns of brain activation after the three-weeks

intervention, in particular involving frontal and parietal brain regions.

Methods

Sample and exclusion criteria

Figure 1 depicts the study recruitment pipeline. Participants were recruited from two major

hospitals located in Oslo, Norway (Stroke unit at Oslo University hospital, and Geriatric

department at Diakonhjemmet Hospital, Oslo, Norway). Hospital staff identified suitable

participants. Inclusion criterion was identifiable stroke with ischemic or hemorrhagic

etiology. Patients with transient ischemic attacks (TIA) were not included. Exclusion criteria

included MRI contraindications, other neurological diseases diagnosed prior to the stroke,

severe psychiatric diagnosis including bipolar disorder, schizophrenia or drug abuse.

Approximately 900 invitations were sent out, and 250 responded; among those, 77 patients

were eligible for participation and were initially included in the study (Ulrichsen et al., 2019).

19 participants withdrew from the study prior to (n=14) or during (n=5) the cognitive training

due to the labor-intensity of the intervention. Four participants were excluded during the

course of the intervention due to newly occurring medical issues violating the inclusion

criteria. Two participants were included without radiological findings when admitted to the

stroke unit, but displayed neurological symptoms. Initial MRI diffusion weighted imaging

revealed possible acute infarctions and the patients were treated with thrombolysis.

Subsequent T2-weighted imaging revealed no certain radiological findings, but symptoms

persisted upon discharge and they were diagnosed by a neurologist with ICD-10 code I63.9

Cerebral infarction, and therefore included. In total, 54 participants completed the cognitive

training program. For timepoint 1, one participant had corrupted behavioral data and two

participants were not able to complete the task during the fMRI assessment, and were

excluded from all MRI analysis. For MRI timepoint 2 and 3, one participant had poor

alignment to the MNI-template, and two participants were not able to complete the task,

yielding a total of 51 participants at baseline and 48 with full MRI at all timepoints. In this

final sample, 26 participants received active stimulation, and 22 received sham stimulation.

The study was approved by the Regional Committee for Medical and Health Research

Ethics South-East Norway (2014/694; 2015/1282), and all participants provided written

informed consent.

178
179 **Figure 1**. Flowchart displaying inclusion pipeline
180

Patient characteristics and lesion demarcation

Sample characteristics for the participants completing the intervention (n=54) are described in

Table 1, including National Institutes of Health Stroke Scale (NIHSS) (Goldstein et al., 1989),

Trial of Org 10172 in Acute Stroke Treatment (TOAST) (Adams et al., 1993), Mini-Mental

State Exam (MMSE) (Strobel and Engedal, 2008) and IQ derived from Wechsler Abbreviated

Scale of Intelligence (WASI) (Ørbeck and Sundet, 2007). Briefly, the sample mean age was

69 years (sd=7.3), comprised 74% males, and average IQ of 110 (sd=16). Two sample t-tests

revealed no significant differences between the active and sham tDCS-groups.

For each participant, lesion demarcation was performed by a trained assistant, utilizing

the semi-automated toolbox clusterize, implemented for SPM8 (Clas et al., 2012; de Haan et

al., 2015), and guided by radiological descriptions. Figure 2 displays a probabilistic lesion

map across participants. Supplementary Figure 1 provides an overview of individual lesions.

196
197 **Figure 2.** Heatmap displaying lesion overlap across included participants. The color scale
198 reflects the number of overlapping lesions in each voxel (ranging from 0 to 7).
199

	Mean	sd	Min	Max	tDCS grouping t/p
Participant descriptives (current)					
Age at inclusion	69.13	7.374	47	81	**0.11 / 0.907**
Males (%)	74.07	-			**23 / 17 males**
Time between stroke and inclusion (months)	25.74	9.17	6	45	**0.36 / 0.720**
Self-reported education in years	14.39	3.71	9	30	**-0.59 / 0.559**
MMSE at inclusion	27.98	1.855	22	30	**0.81 / 0.419**
IQ	110.45	16.906	66	136	**-0.41 / 0.687**
Days between inclusion and training	33.78	12.986	19	74	**0.82 / 0.419**
Days between MRI assessment pre/post training	31.72	6.257	19	49	**1.99 / 0.052**
Clinical variables during hospitalization					
NIHSS at discharge	1.33	1.53	0	7	
TOAST classification	Large vessel disease n=20 Small vessel disease n=18 Cardioembolic disease n=6 Other n=10				

200 Table 1. Sample characteristics. Two-sample t-tests did not reveal any significant differences
201 between the active and sham group regarding participant characteristics
202

Study protocol

The protocol have previously been described in detail (Richard et al., 2020). Briefly, after

inclusion the participants completed MRI assessment at three timepoints: timepoint 1, at

timepoint 2 on average 3-4 weeks after timepoint 1 without additional contact to enable a

double baseline for comparison, and at timepoint 3, after the participant completed the

training protocol.

Cognitive training

The patients performed a computerized working memory training program (Cogmed QM,

Cogmed Systems AB, Stockholm, Sweden), comprising 12 auditory-verbal and visual-spatial

working memory tasks. The implementation of the protocol has previously been reported in

detail (Richard et al., 2020). Briefly, the training comprised five weekly training sessions over

3-4 weeks, with 17 training sessions for each participant in total. Each participant completed

two weekly tDCS stimulation sessions at the hospital, with a total of 6 tDCS sessions with a

minimum of 48 hours between each stimulation. The remaining Cogmed sessions was

completed at home. Level of Cogmed task difficulty was automatically adjusted according to

each participant's performance level. The initial two first sessions for each task was discarded

from statistical analysis to ensure each participant was at a desired level of difficulty. The

same task setup was used for all participants, and tasks completed three times or less over the

course of training were excluded. Eight Cogmed subtests were included in further analysis,

including Grid, Cube, Digits, Hidden objects, Sort, Rotation, Twist, and 3D-cube.

tDCS

Participants were randomly assigned to either active or sham condition, using codes provided

by the manufacturer and implemented by an in-house Matlab script, pseudo-randomizing

participants while maintaining balance in groups of 20. tDCS stimulation was applied at six

occasions for each participant, with a minimum of 48 hours between stimulations, aiming at

an average of two stimulations per week. Stimulation current was 1000 μA, stimulation time

was 20 minutes for the active group with a ramp-up time of 120 seconds and fade-out time of

30s. The current intensity was limited to 1mA to limit the risk of adverse events due to the

electrical stimulation. The sham stimulation consisted of ramp-up followed by 40 seconds of

active stimulation and then fade-out, following factory settings. Stimulation was delivered via

a battery-driven direct current stimulator (neuroConn DC stimulator plus, Germany), through

5 x 7 cm rubber pads, yielding a 28.57 μA/cm2 current density at skin surface. Impedance

threshold at start-up of stimulation was < 20 kΩ, and was regulated by the device with an

absolute threshold at 40 kΩ for automatically terminating stimulation. The pads were covered

with high-conductive gel (Abralyt HiCl, Falk Minow Services Herrsching, Germany) to

enhance conductance.

Electrode location was determined by following the 10-20 system, where the anodal

electrode was located at F3 corresponding to left DLPFC and cathodal at O2 corresponding to

right occipital/cerebellum, and fixated with rubber bands.

fMRI task

Participants performed a blocked version of the MOT (Pylyshyn and Storm, 1988), as

previously described in detail (Alnæs et al., 2015b). Briefly, the participants were presented

with ten identical circles on a grey background. All objects were initially blue, before zero

(passive viewing), one (load 1) or two (load 2) of the objects turned red for 2.5 seconds,

designating them as targets. After targets had returned to blue, the objects started moving

randomly around the screen for 12 seconds. After the 12 second tracking period, the objects

became stationary, one of the objects turned green, and the participants were instructed to

respond "yes" or "no" to whether or not the green probe had been designated as a target in

that trial. Participants were instructed to keep fixation on a fixation point at the center of the

screen during both attentive tracking and passive viewing. The task consisted of 24 semi-

randomized trials in total, ensuring the same condition was not repeated consecutively.

MRI acquisition

MRI data was acquired with a 3T GE 750 Discovery MRI scanner (32-channel head coil)

located at Oslo University Hospital. T1-weighted images was acquired using a 3D IR-

prepared FSPGR (BRAVO) with the following parameters: scan time: 4:43 minutes and

included 188 sagittal slices, repetition time (TR): 8.16 ms, echo time: 3.18 ms, flip angle: 12º,

voxel size: $1 \times 1 \times 1$ mm, field of view: 256 x 256 mm,

Functional MRI sequences was acquired with a BOLD-sensitive gradient echo planar

sequence, with TR of 2.25 s, TE 30 ms, field of view 256 x 256 mm, slice thickness 3 mm

with 0.5 mm gap acquired ascending interleaved, 90º flip angle, with R/L orientation. The

initial first five volumes were removed before analysis.

fMRI data processing

fMRI data was processed using FMRI Expert Analysis Tool (FEAT) Version 6.00, from

FMRIB's Software Library (FSL; Jenkinson et al., 2012; Smith et al., 2004), and included the

following steps: correction for motion using MCFLIRT (Jenkinson et al., 2002), linear trend

removal and high-pass filtering (0.01 Hz), removal of non-brain tissue using BET (Smith,

2002), spatial smoothing with a Gaussian kernel of full width at half maximum (FWHM) of 6

mm (SUSAN ; Smith and Brady, 1997), and linear registration using FLIRT (Jenkinson and

Smith, 2001) to Montreal Neurological Institute (MNI) 152 standard space using the T1-

weighted scan as an intermediate. In the same manner, normalization parameters were

estimated for each participants T2-weighted image, and applied to the demarked lesion.

Statistical analysis of behavioral data

Cogmed main effects of time: To investigate the trajectory of Cogmed performance across

participants and time points, linear mixed effects (lme) models were estimated for each test,

using the lme function from the nlme-package (Pinheiro et al., 2017) in R (Team, 2016). Task

performance was entered as dependent variable, with training session as independent variable,

and age, sex, tDCS-group, educational level in years (self-reported), and interaction between

training session and tDCS, as fixed factors, and participant as random factor. To assess

normality in the data according to criteria for lme, we plotted distribution of random effects

intercepts, see supplementary Figure 7. Visual inspection did not indicate severe violations of

the criteria.

291   *Cogmed individual trajectories:* To quantify individual trajectories in Cogmed

performance across the training-period, we estimated for each test and each participant a

linear model with performance as dependent variable and session number as independent

variable, yielding a beta-estimate (slope) for each test for each participant reflecting changes

in performance over time. As a measure of overall performance for each participant, we used

the average task performance across sessions for each test for each participant. To investigate

task homogeneity in beta-estimates and average performance we performed multivariate

outlier detection with the aq.plot function in the mvoutliers package in R (P. Filzmoser and

Gschwandtner, 2018), with an alpha of 0.001 and 90% of the sample to estimate the minimum

covariance determinant. Two subtests ("hidden objects" and "Digits") were found to have a

high number of outliers compared to the other subtests, and were excluded from further

analysis (See Supplementary Figure 2 for details).

303   To derive a common score for average Cogmed performance and gain, we performed

a principal component analysis (PCA) on beta-estimates and average performance,

respectively. All scores were zero-centered and standardized prior to running the PCA. We

used the first factor from the beta PCA as a *Cogmed change score*, and the first factor from

the average performance PCA as a *Cogmed average performance score (*see Supplementary

Figure 3 for scree-plot derived from the PCA performed on both with and without the

excluded tasks*).*

To explore potential association between initial cognitive capacity and Cogmed

change score, we correlated performance on the cognitive performance at baseline assessed

with the Cabpad battery (https://www.cognisoft.info) at TP1, described in detail previously

(Richard et al., 2019) with the Cogmed change score. To test for associations between initial

cognitive performance and response to tDCS we estimated linear models using the same

baseline cognitive performance from Cabpad as the dependent variable, and Cogmed change

score and tDCS group as independent variables. Models were estimated using the lm function

in the stats package in R (Team, 2013).

To assess the association between Cogmed scores and lesion severity, we estimated

linear models for both Cogmed change and average performance scores, using lesion volume,

number of lesions and their interaction as explanatory variables. To assess impact of duration

from the stroke to inclusion on gains from the training, we correlated time since injury

measured in months with the Cogmed change score as well as the Cogmed average

performance score.

fMRI behavioral analysis: To investigate performance during the MOT task across

timepoints, linear mixed effects (lme) models were estimated for both load 1 and load 2, using

the lme function from the nlme-package (Pinheiro et al., 2017) in R. Task performance was

entered as dependent variable, with training session as independent variable, and participant

as random factor. To assess for pairwise differences across the timepoints, post hoc analysis

was performed using the ghlt function from the multcomp package (Hothorn et al., 2008) in

R, and corrected for multiple comparisons using Bonferroni correction.

fMRI data analysis

First-level GLMs for each participant and scanning session were estimated using FEAT

(Woolrich et al., 2001), including regressors for passive viewing, load 1, load 2, responses, as

well as the standard and extended motion parameters estimated during preprocessing. The

following contrasts were estimated: passive viewing, load 1, load 2, tracking ([load 1 + load

2] – passive viewing) and load (load 2 – load 1).

To estimate main effects of experimental conditions across individuals, contrasts of

parameter estimates (COPEs) from each participant's first scan session (baseline) were

submitted to randomise (Winkler et al., 2014) performing a one sample t-test for each

condition separately. Multiple comparisons were corrected for using permutation testing and

threshold free cluster enhancement (TFCE (Smith and Nichols, 2009)).

To estimate associations with task performance with brain activation during task

engagement, individual COPE maps for load 1 and load 2 were stacked separately, and

submitted to randomise, including sex, age and task performance as covariates.

To test for differences in brain activation between scan sessions, we first performed

fixed effects analysis for each participant separately, including all COPEs from first level

GLM, specifying the contrasts session 3 minus session 2, session 3 minus session 1, and

session 2 minus session 1. Next, all individual COPE maps were submitted to randomise to

test for group-level main effects while correcting for multiple comparisons using permutation

testing and threshold free cluster enhancement (TFCE).

To quantify within-subject reliability in brain activation across time we estimated

intraclass correlation coefficient (ICC) for each voxel between assessment two and three

utilizing the variance from one-way random ANOVA, for the contrasts passive viewing, load

1 and load 2, as implemented in the ICCest function from the ICC R package (Wolak et al.,

2012).

To test for associations between Cogmed change and average performance scores and

brain activation, individual level COPE maps for timepoint 1 and the difference between

timepoint 3 and 2 including the conditions passive viewing, load1, load 2, tracking as well as

load were stacked separately and submitted to randomise. The model included sex, age, and

either Cogmed change score or Cogmed average performance score (obtained from the PCA).

In the same manner to explore differences associated with tDCS stimulation protocol, the

same COPE´s from the difference between timepoint 3 and 2 including all conditions were

submitted to randomise, with the model including sex, age and tDCS-condition. All models

were corrected for multiple comparisons across space using 5000 permutations and

thresholded using TFCE.

Results

Cogmed performance

Figure 3 shows Cogmed performance over time for each group and Table 2 displays

corresponding summary statistics. Lme revealed a significant group-level improvement on all

Cogmed tests included in the analysis. In addition, we found a significant association with age

on mean performance across all tests, indicating lower average performance with higher age.

We found no significant associations between Cogmed performance and sex, tDCS group

(sham/experimental), or years of education. We found a significant interaction between time

and tDCS on the twist task, suggesting augmented training gain in the participants in the

active tDCS stimulation group. However, after removal of one outlier defined by visual

inspection, the result did not reach significance (see Supplementary Table 1 and

Supplementary Figure 4 for corresponding statistics and visualization).

382
383 **Figure 3.** Individual level task performance during the course of the intervention period for
384 each group (tDCS sham/active). The fit lines are based on a linear fit within each group.
385
386

	Grid	Sort	Digits	Cube	Hidden objects	Twist	3D-Cube	Rotation
Session	**8.66** *(<0.001)*	**6.40** *(<0.001)*	**12.79** *(<0.001)*	**7.44** *(<0.001)*	**11.03** *(<0.001)*	**6.45** *(<0.001)*	**3.16** *(0.002)*	**2.21** *(0.028)*
Age	**-3.05** *(0.004)*	**-2.11** *(0.041)*	**-6.07** *(<0.001)*	**-2.54** *(0.015)*	**-4.86** *(<0.001)*	**-3.62** *(0.001)*	**-3.90** *(<0.001)*	**-3.41** *(0.001)*
Sex	-0.89 (0.377)	-0.33 (0.741)	-2.29 (0.027)	-1.22 (0.23)	-2.25 (0.029)	-1.46 (0.15)	-1.42 (0.162)	-1.34 (0.187)
tDCS	-0.65 (0.521)	-1.03 (0.309)	-0.06 (0.951)	-0.53 (0.599)	0.12 (0.908)	1.00 (0.324)	-0.82 (0.414)	-0.81 (0.422)
Education	0.06 (0.951)	0.96 (0.34)	0.90 (0.374)	0.14 (0.886)	1.57 (0.124)	0.01 (0.991)	-0.15 (0.884)	0.33 (0.744)
Session x tDCS	0.2 (0.842)	-0.39 (0.7)	-1.53 (0.128)	0.46 (0.644)	-1.79 (0.074)	**-2.83** **(0.005)**	1.72 (0.087)	1.23 (0.22)

388 **Table 2**. Summary statistics (t/(p))from linear mixed effects models testing for associations
389 between task performance and session, age, sex, tDCS, education and the interaction between
390 session and tDCS. t-vaules along with raw p-vaules are displayed, significant findings are
391 highlighted (FDR-corrected, alpha = 0.05).
392
393

We found no significant correlation between Cogmed average performance and Cogmed

change score (r=.17, t=1.29, p=.2). We found a significant association between lesion volume

and Cogmed change score (t -2.5, p=.017), indicating a negative correlation between lesion

volume and training response, which did not retain significance after removal of one outlier

(t=-.58, p=.563). See Supplementary Table 2 for corresponding statistics. Furthermore,

results revealed no significant correlations between time since injury and Cogmed average

performance (r= -0.19, t = -1.38, p = 0.173) nor Cogmed change score (-0.19, t = -1.35, p =

0.181).

Linear models revealed no significant associations between cognitive capacity as

measured using Cabpad at baseline and gain in Cogmed performance. Furthermore, there

were no significant interactions between Cabpad score at baseline and tDCS on Cogmed

change score. See supplementary figure 6 and supplementary table 4 for corresponding

statistics.

Behavioral fMRI results

LME analysis revealed main effect of time on task performance during both load 1 and load

2. Post hoc analysis revealed significant differences between timepoint 3 and 1, as well as

between timepoint 2 and 1, but not between timepoint 3 and 2. See Table 3 and Figure 4 for

details.

	Main effect		Tp2 - Tp1		Tp3 – Tp1		Tp3 – Tp 2	
Model	**F**	**p**	**t**	**p**	**t**	**p**	**t**	**p**
Load 1	**9.22**	**2.20e-04**	**3.52**	**1.28e-03**	**3.89**	**3.03e-04**	0.36	1.00
Load 2	**8.24**	**5.04e-04**	2.31	0.06	**4.05**	**1.56e-04**	1.73	0.25

415 **Table 3.** Summary statistics from linear mixed effects models testing for associations
416 between task performance and session for the MOT task. Bonferroni-corrected p-values are
417 displayed, significant results are highlighted.
418

419
420 **Figure 4.** Task performance during the MOT task at the three different timepoints. Significant
421 differences in group means is marked *, Table 3 displays corresponding statistics.
422

Association between task performance and fMRI activation.

Linear models revealed no significant association between in-scanner task performance and

fMRI activation during neither load 1 or load 2 during the first assessment (TP 1). In-scanner

behavioral performance was therefore not included in further analysis.

fMRI main effects of task

Figure 5 shows main effects of task condition at baseline, and Table 4 summarizes anatomical

locations. Passive viewing was associated with increased activity in the dorsal attention

network (DAN) and the inferior and superior occipital cortex, and decreased activation in pre

and post-central gyrus, supplementary motor area (SMA), anterior cingulate, as well as

default mode network (DMN), including posterior cingulate gyrus (PCC), medial prefrontal

gyrus (mPFC), and superior and medial temporal gyrus (S/MTG).

Load 1 was associated with increased activity in DAN, extending into the inferior and

superior occipital cortex, cerebellum, and precentral gyrus, in the cingulo-opercular (CO)

network, including thalamus, frontal operculum and insula, as well as in the ventral attention

(VAN) network including temporoparietal junction and supramarginal gyrus, middle frontal

gyrus, and frontal operculum. Decreased activity was seen in DMN nodes including PCC,

mPFC and S/MTG. Load 2 showed a similar pattern to Load 1, with additional increased

activation in right anterior middle frontal gyrus. Tracking over passive viewing revealed same

activation pattern as load 1 and load 2, with additional bilateral activation of anterior middle

frontal gyrus. Load-related (load 2 –load1) increases in activation were seen in superior and

inferior parietal lobe, as well as right anterior middle frontal gyrus, with no significant load-

dependent decreases in activation.

447
448 **Figure 5.** Main effects from task-fMRI for (A) passive viewing > baseline, (B) load 1 >
449 baseline, (C) load 2 > baseline, (D) tracking > passive viewing, and (E) increased cognitive
450 load (load 2 > load 1). Panel A display corresponding coordinates in MNI, and left / right
451 orientation. P-values (1-p) have been multiplied with the sign of the effect in relation to the
452 main contrast.
453

Condition	Direction	size	X	Y	Z	Area
PV	pos	4029	33	19	30	Lateral occipital cortex
	pos	3861	55	18	32	Lateral occipital cortex
	pos	1784	29	41	61	Superior parietal lobe
	neg	64337	33	35	21	Cerebellum
Load1	pos	57023	59	29	8	Cerebellum
	neg	12615	33	39	23	Cerebellum
	neg	1494	19	62	24	Middle temporal gyrus
	neg	256	65	53	42	Central opercular cortex
	neg	153	20	78	48	Middle frontal gyrus
	neg	19	22	71	50	Inferior frontal gyrus
Load2	pos	66242	59	29	8	Cerebellum
	pos	411	65	71	37	Frontal operculum
	pos	8	65	80	55	Middle frontal gyrus
	neg	10269	31	40	26	Occipital fusiform cortex
	neg	694	18	63	26	Superior temporal gyrus
Tracking	pos	101656	59	29	8	Cerebellum
Load	pos	4590	24	42	55	Inferior parietal lobe
	pos	1709	67	42	56	Inferior parietal lobe
	pos	245	24	79	54	Middle frontal gyrus

455   **Table 4.** Clusters associated with different MOT condition at baseline (all p<.05, corrected)
456
457

458   *fMRI – reliability*

459   Figure 6 shows voxel-wise ICC for COPE values for passive viewing, load 1 and load 2,

460   between timepoint 2 and 3. The results revealed ICC>0.4 across major portions of the brain,

461   including occipital, temporal, parietal lobe for all conditions, as well as and a large part of the

462   frontal lobes for load 1.

464
465 **Figure 6.** Voxel-wise ICC for condition passive viewing (A), load 1 (B), and load 2 (C).
466 Thresholded at ICC > 0.4.
467

468 *Pairwise comparison of fMRI estimates across timepoints*

469 Comparison of fMRI signal between the timepoints revealed no significant differences

470 between timepoint 3 and 1, timepoint 2 and 1 as well timepoint 3 and 2, for neither passive

471 viewing, load 1, load 2, tracking nor load, indicating fluctuating patterns, but not in a time-

472 dependent manner.

Cogmed –fMRI associations

We found no significant associations between the Cogmed change score and brain activation

at timepoint 1, or between Cogmed change score and the difference in brain activation

between timepoint 2 and 3.

Furthermore, results revealed no significant associations between the Cogmed average

score and brain activation at timepoint 1. However, between timepoint 2 and 3 we found a

significant association between Cogmed average performance score and the difference in

activation during attentive tracking, indicating larger activation increases in the anterior and

posterior parts of the cingulate cortex and left precuneus, superior parietal lobe, lateral

occipital cortex, as well as cerebellum with greater average Cogmed performance. See Table

5 and Figure 7 for corresponding statistics. Using a Cogmed average performance score based

on all 8 subtests (from the PCA, now including two sub-tests previously excluded), revealed

slightly weaker but highly similar effects in the (Supplementary Figure 5 and Table 3),

suggesting that excluding the two subtests did not bias the results.

	Direction	Size	x	y	z	Location
Frontal	pos	120	30.1	64.4	61.4	Middle frontal gyrus
	pos	46	59.5	63.9	60.5	MFG L
Limbic	pos	1269	41.7	74.2	53.3	ACC
	pos	261	43.8	52.3	53.8	PCC
Parietal	pos	83	23	36.8	61.9	SPL
	pos	207	55	30.2	48.7	Precuneus cortex
Occipital	pos	117	21.5	25.9	39.3	Lateral occipital cortex
	pos	100	61.6	22.7	44.4	Lateral occipital cortex
	pos	33	35.8	15.3	36.3	Occipital pole
Cerebellar	pos	54	60	26	13	Cerebellum
	pos	16	41.1	37.2	23.5	Cerebellum

493 **Table 5.** Cogmed average score clusters associated with difference in activation between
494 timepoint 2 and 3. (all p<.05, corrected).

496
497 **Figure 7.** Association between Cogmed average performance score, and difference in task-
498 related activation (L1 & L2 > PV) between timepoint 3 and 2.
499

tDCS –fMRI associations

Results revealed no difference in fMRI-activation between the active and sham tDCS

stimulation groups.

Discussion

In this study, we tested the predictive value of fMRI-based brain activation on performance

gains following a commercially available working memory training package. Additionally,

we investigated whether active tDCS combined with Cogmed induced neural alterations

measured with task-based fMRI.

The results revealed significant increase in performance on the trained tasks, with no

additional benefits from tDCS on training gain. Our analysis revealed robust activation in

response to the MOT task in line with previous findings. It is important to note that the

current study investigated possible markers for individual training outcome, and a

comprehensive evaluation of the potential behavioral and clinical transfer effects is beyond

the scope of the current study. Importantly, investigation of reliability across timepoints

supported the feasibility of utilizing task-fMRI for delineating neural correlates of behavioral

training in a clinical group. Following the promising reliability in brain activation across the

different fMRI assessments, we found no significant association between cognitive

performance gain during the cognitive training and brain activation prior to training, or with

brain activation change between pre- and post-training. However, higher average performance

across the training period was associated with higher load-dependent increase in activation in

the precuneus, and furthermore associated with increased activation in the DAN and DMN as

well as the ACC during active task performance after compared to before intervention.

523   Tailoring behavioral interventions for alleviating cognitive difficulties following

stroke calls for investigation of both treatment response, as well as delineating factors

predicting favorable outcome. Cognitive abilities prior to training as well as motivation (Guye

et al., 2017; Jaeggi et al., 2014) are currently among our best predictors for beneficial effects.

Our results revealed significant improvement on all included tasks from the training program,

in line with previous reports (Spencer-Smith and Klingberg, 2015; Westerberg et al., 2007).

Our results support that repeated practice, where level of difficulty is adjusted during training,

improves performance on the trained tasks. It is important to note that the current study

investigates possible biomarkers for individual training outcome, and its beyond our aim to

evaluate potential behavioral transfer effects. However, the generalizability of training-effects

following CCT has been debated (Au et al., 2015; Melby-Lervåg and Hulme, 2013; Morrison

and Chein, 2011; Shipstead et al., 2012; von Bastian and Oberauer, 2014). Indeed, recent

large-scale reviews suggest a lack of reliable transfer beyond tasks that share properties with

the trained tasks (Melby-Lervåg et al., 2016; Sala and Gobet, 2018). The lack of transfer

effects is suggested to be caused by the failure of targeting common mechanisms underlying

fluid intelligence such as working memory and attention, and rather targeting task specific

mechanisms through repeated practice (Melby-Lervåg et al., 2016). Despite our promising

behavioral results on trained tasks, the lack of transfer-effects reported in the literature calls

for caution before implementation in a clinical setting.

542   Our results revealed no significant interaction between cognitive improvement and

tDCS stimulation with the exception of one of the trained tasks, which did not reach

significance after removal of an outlier. Although anodal tDCS has shown beneficial effects

on cognitive training compared with sham stimulation in healthy adults (Martin et al., 2013)

and have been associated with steeper learning curves (Ruf et al., 2017), a recent meta-

analysis provided no support of a beneficial effect of tDCS on cognitive training compared to

sham (Medina and Cason, 2017). The current literature investigating the added effect of tDCS

when combined with working memory training in stroke patients is scarce. Single session

anodal tDCS towards the left DLPFC improved recognition accuracy during an N-back task

(Jo et al., 2009), and longitudinal concurrent anodal tDCS and CCT increased accuracy on a

non-trained continuous performance test (CPT) compared to sham (Park et al., 2013).

However, divergence in study protocols and results hampers a direct comparison (daSilva

Filho et al., 2017). Beyond robust improvement on all trained tasks during the course of the

intervention, our results provide no evidence of additional beneficial effects of tDCS

stimulation. Intriguingly, current investigations aim to evaluate how home-based tDCS may

aid patients with mild cognitive impairments (Park et al., 2019). This will potentially provide

novel insights regarding clinical application for tDCS.

Our fMRI results for task activation during the first assessment revealed increased

activation of the DAN, along with deactivation of DMN for passive viewing, load 1 and load

2. Further, contrasting tracking to passive viewing, and load 2 to load 1 revealed that attentive

tracking increased DAN activation in a load-dependent manner, in line with previous reports

based on healthy samples (Alnæs et al., 2014; Dørum et al., 2016).

Importantly, longitudinal studies utilizing fMRI for investigation of within-subject

changes depends on the reliability of brain measurements across time in order to pick up the

relevant signal. Previous investigations suggest favorable reliability in task based compared to

resting-state fMRI (Specht, 2020), indicating higher reliability when the task elicits strong

patterns of activation (Johnstone et al., 2005) increasing contrast to noise ratio (Bennett and

Miller, 2010). Our results are in line with these reports, as results revealed fair (>0.4) to good

(>0.75) reliability across task-associated brain regions following guidelines for ICC

interpretation (Cicchetti, 1994), with poor reliability in non-task associated areas.

Our results revealed no significant interaction between activation during passive

viewing, load 1, load 2, tracking nor load, and time. Although we did not find any predictive

value of fMRI-activation patterns on Cogmed performance improvement, our results revealed

an association between average performance during the Cogmed training period and increased

tracking-related activity in key nodes of the DAN and DMN as well as ACC. DMN is

commonly suppressed during task engagement, with increasing suppression with increased

task demands (Anticevic et al., 2012). Furthermore, cognitive abilities are negatively

correlated with degree of de-activation (Lipp et al., 2012), most likely due to need for higher

effort to resolve the task. In line with these findings our results indicate less suppression of

PCC and precuneus during task engagement, with greater average performance.

Working memory capacity, reflecting both executive control as well as cognitive

flexibility, has been linked to general cognitive abilities (Conway et al., 2003). Furthermore,

higher cognitive abilities have been associated with increased activity in frontal and parietal

cortices (Basten et al., 2015), and cognitive flexibility have been linked to brain activation in

ventrolateral prefrontal cortex, as well as anterior and posterior cingulate (Dajani and Uddin,

2015). As our results indicate that greater cognitive capacity is associated with an increase in

activation during task engagement over time, it may reflect compensation and learning to

more efficiently resolve the task.

The current findings should be interpreted in light of several limitations. Based on the

clinical ratings (NIHSS) at hospital discharge, the patients were sampled from the less severe

part of the stroke severity spectrum. Further, patient drop-out during the intervention was

relatively high, partly due to the labor-intensity of the training. Hence, our sample is biased

towards higher functioning stroke patients, rendering the generalizability of our results to

more severe cases unclear. The lack of a control group for the CCT does not allow us to

disentangle effects of the cognitive training from the effect of time, anticipation, or other

confounders. As our main aim was to identify potential fMRI-based markers for training

outcome, we did not assess possible cognitive and clinical transfer effects, which will be

relevant for evaluating clinical relevance, and further studies are needed to establish the

premises for optimizing the clinical potential for working memory interventions in individual

stroke patients. It has furthermore been reported that potential beneficial effects of tDCS may

emerge at a delayed stage after intervention (Goodwill et al., 2016; Li et al., 2019). As our last

assessment was performed shortly after the final stimulation, any delayed effects may have

been lost. Future follow-up assessments of the same patients may provide relevant

information about potential long-term effects of CCT and tDCS. Potential effects also may be

dependent of stimulation frequency (Boggio et al., 2007; Yun et al., 2015). Increasing

frequency beyond current protocol may have altered our results. Lesion size and location

may influence current flow and cortical excitability (Marquez et al., 2013), and future studies

may test for associations between lesion characteristics and response to tDCS. While the

degree of cognitive impairment at baseline is likely to influence the response to cognitive

training and rehabilitation in general, we found no significant association between baseline

cognitive performance and training gain. However, the lack of pre-stroke neuropsychological

evaluation complicates the assessment of stroke-related cognitive impairment. The group

level task activation patterns in the current study largely mirrored those reported in previous

studies in healthy controls (Alnæs et al., 2015a; Dørum et al., 2016). However, the current

study design did not allow us to make an explicit comparison (D'Esposito et al., 2003). Here

we only considered brain activation in response to task demands. Future studies should also

pursue a promising line of research utilizing imaging indices of structural and functional

connectivity to investigate behavioral correlates of stroke, which may supplement and

increase our understanding of stroke and stroke recovery.

621   In conclusion, we have investigated response to a computerized cognitive training

622 program in 54 stroke survivors in combination with tDCS, as well as the predictive value of

623 neural activation on training outcome, and neural alterations following training. Our results

624 revealed increased performance across all trained tasks, with no additional benefit of tDCS.

625 Brain activation prior to the training was not predictive for training outcome, nor was training

626 gains reflected in altered brain activation. The generalizability of the reported beneficial

627 effects of CCT remains uncertain, and future studies may be able to assess the transfer value

628 of for relevant cognitive and clinical variables in stroke patients.

References

631 Adams, H.P., Bendixen, B.H., Kappelle, L.J., Biller, J., Love, B.B., Gordon, D.L., Marsh,
632 E.E., 1993. Classification of subtype of acute ischemic stroke. Definitions for use in a
633 multicenter clinical trial. TOAST. Trial of Org 10172 in Acute Stroke Treatment. Stroke 24,
634 35-41.
635 Alnæs, D., Kaufmann, T., Richard, G., Duff, E.P., Sneve, M.H., Endestad, T., Nordvik, J.E.,
636 Andreassen, O.A., Smith, S.M., Westlye, L.T., 2015a. Attentional load modulates large-scale
637 functional brain connectivity beyond the core attention networks. NeuroImage 109, 260-272.
638 Alnæs, D., Sneve, M.H., Espeseth, T., Endestad, T., van de Pavert, S.H.P., Laeng, B., 2014.
639 Pupil size signals mental effort deployed during multiple object tracking and predicts brain
640 activity in the dorsal attention network and the locus coeruleus. Journal of Vision 14, 1-1.
641 Alnæs, D., Sneve, M.H., Richard, G., Skåtun, K.C., Kaufmann, T., Nordvik, J.E., Andreassen,
642 O.A., Endestad, T., Laeng, B., Westlye, L.T., 2015b. Functional connectivity indicates
643 differential roles for the intraparietal sulcus and the superior parietal lobule in multiple object
644 tracking. NeuroImage 123, 129-137.
645 Altamura, C., Reinhard, M., Vry, M.-S., Kaller, C.P., Hamzei, F., Vernieri, F., Rossini, P.M.,
646 Hetzel, A., Weiller, C., Saur, D., 2009. The longitudinal changes of BOLD response and
647 cerebral hemodynamics from acute to subacute stroke. A fMRI and TCD study. BMC
648 Neuroscience 10, 151.
649 Andrews, S.C., Hoy, K.E., Enticott, P.G., Daskalakis, Z.J., Fitzgerald, P.B., 2011. Improving
650 working memory: the effect of combining cognitive activity and anodal transcranial direct
651 current stimulation to the left dorsolateral prefrontal cortex. Brain Stimulation 4, 84-89.
652 Anticevic, A., Cole, M.W., Murray, J.D., Corlett, P.R., Wang, X.-J., Krystal, J.H., 2012. The
653 role of default network deactivation in cognition and disease. Trends in Cognitive Sciences
654 16, 584-592.
655 Au, J., Sheehan, E., Tsai, N., Duncan, G.J., Buschkuehl, M., Jaeggi, S.M., 2015. Improving
656 fluid intelligence with training on working memory: a meta-analysis. Psychonomic Bulletin &
657 Review 22, 366-377.
658 Barbay, M., Diouf, M., Roussel, M., Godefroy, O., 2018. Systematic Review and Meta-
659 Analysis of Prevalence in Post-Stroke Neurocognitive Disorders in Hospital-Based Studies.
660 Dementia and Geriatric Cognitive Disorders 46, 322-334.

Basten, U., Hilger, K., Fiebach, C.J., 2015. Where smart brains are different: A quantitative
meta-analysis of functional and structural brain imaging studies on intelligence. Intelligence
51, 10-27.
Bennett, C.M., Miller, M.B., 2010. How reliable are the results from functional magnetic
resonance imaging? Annals of the New York Academy of Sciences 1191, 133-155.
Buitenweg, J.I.V., Van De Ven, R.M., Ridderinkhof, K.R., Murre, J.M.J., 2018. Does
cognitive flexibility training enhance subjective mental functioning in healthy older adults?
Aging, Neuropsychology, and Cognition, 1-23.
Cicchetti, D.V., 1994. Guidelines, criteria, and rules of thumb for evaluating normed and
standardized assessment instruments in psychology. American Psychological Association,
US, pp. 284-290.
Clare Kelly, A.M., Hester, R., Foxe, J.J., Shpaner, M., Garavan, H., 2006. Flexible cognitive
control: Effects of individual differences and brief practice on a complex cognitive task.
NeuroImage 31, 866-886.
Clas, P., Groeschel S Fau - Wilke, M., Wilke, M., 2012. A semi-automatic algorithm for
determining the demyelination load in metachromatic leukodystrophy. Academic Radiology
19.
Constantinidis, C., Klingberg, T., 2016. The neuroscience of working memory capacity and
training. Nature Reviews Neuroscience 17, 438.
Conway, A.R.A., Kane, M.J., Engle, R.W., 2003. Working memory capacity and its relation
to general intelligence. Trends in Cognitive Sciences 7, 547-552.
D'Esposito, M., Deouell, L.Y., Gazzaley, A., 2003. Alterations in the BOLD fMRI signal with
ageing and disease: a challenge for neuroimaging. Nature Reviews Neuroscience 4, 863.
Dajani, D.R., Uddin, L.Q., 2015. Demystifying cognitive flexibility: Implications for clinical
and developmental neuroscience. Trends in Neurosciences 38, 571-578.
daSilva Filho, E.M., Andrade deAlbuquerque, J., Mescouto, K.A., deAbreu Freitas, R.P.,
2017. Effects of transcranial direct current stimulation on the working memory of post-stroke
people: an integrative review. Manual Therapy, Posturology & Rehabilitation Journal=
Revista Manual Therapy 15.
de Haan, B., Clas, P., Juenger, H., Wilke, M., Karnath, H.-O., 2015. Fast semi-automated
lesion demarcation in stroke. NeuroImage. Clinical 9, 69-74.
De Luca, R., Calabrò, R.S., Bramanti, P., 2018a. Cognitive rehabilitation after severe
acquired brain injury: current evidence and future directions. Neuropsychological
Rehabilitation 28, 879-898.
De Luca, R., Leonardi, S., Spadaro, L., Russo, M., Aragona, B., Torrisi, M., Maggio, M.G.,
Bramanti, A., Naro, A., De Cola, M.C., Calabrò, R.S., 2018b. Improving Cognitive Function
in Patients with Stroke: Can Computerized Training Be the Future? Journal of Stroke and
Cerebrovascular Diseases 27, 1055-1060.
Dedoncker, J., Brunoni, A.R., Baeken, C., Vanderhasselt, M.-A., 2016. A Systematic Review
and Meta-Analysis of the Effects of Transcranial Direct Current Stimulation (tDCS) Over the
Dorsolateral Prefrontal Cortex in Healthy and Neuropsychiatric Samples: Influence of
Stimulation Parameters. Brain Stimulation 9, 501-517.
Dørum, E.S., Alnæs, D., Kaufmann, T., Richard, G., Lund, M.J., Tønnesen, S., Sneve, M.H.,
Mathiesen, N.C., Rustan, Ø.G., Gjertsen, Ø., Vatn, S., Fure, B., Andreassen, O.A., Nordvik,
J.E., Westlye, L.T., 2016. Age-related differences in brain network activation and co-
activation during multiple object tracking. Brain and Behavior 6, e00533.
Erickson, K.I., Colcombe, S.J., Wadhwa, R., Bherer, L., Peterson, M.S., Scalf, P.E., Kim,
J.S., Alvarado, M., Kramer, A.F., 2006. Training-Induced Functional Activation Changes in
Dual-Task Processing: An fMRI Study. Cerebral cortex 17, 192-204.
Feigin, V.L., Norrving, B., Mensah, G.A., 2017. Global Burden of Stroke. Circulation
Research 120, 439.

Fregni, F., Boggio Ps Fau - Nitsche, M., Nitsche M Fau - Bermpohl, F., Bermpohl F Fau -
Antal, A., Antal A Fau - Feredoes, E., Feredoes E Fau - Marcolin, M.A., Marcolin Ma Fau -
Rigonatti, S.P., Rigonatti Sp Fau - Silva, M.T.A., Silva Mt Fau - Paulus, W., Paulus W Fau -
Pascual-Leone, A., Pascual-Leone, A., 2005. Anodal transcranial direct current stimulation of
prefrontal cortex enhances working memory. Experimental Brain Research.
Giordano, J., Bikson, M., Kappenman, E.S., Clark, V.P., Coslett, H.B., Hamblin, M.R.,
Hamilton, R., Jankord, R., Kozumbo, W.J., McKinley, R.A., Nitsche, M.A., Reilly, J.P.,
Richardson, J., Wurzman, R., Calabrese, E., 2017. Mechanisms and Effects of Transcranial
Direct Current Stimulation. Dose-Response 15, 1559325816685467.
Goldstein, L.B., Bertels, C., Davis, J.N., 1989. Interrater Reliability of the NIH Stroke Scale.
Archives of Neurology 46, 660-662.
Graham, S., Jiang J Fau - Manning, V., Manning V Fau - Nejad, A.B., Nejad Ab Fau -
Zhisheng, K., Zhisheng K Fau - Salleh, S.R., Salleh Sr Fau - Golay, X., Golay X Fau - Berne,
Y.I., Berne Yi Fau - McKenna, P.J., McKenna, P.J., 2010. IQ-related fMRI differences during
cognitive set shifting. Cerebral cortex 20.
Guye, S., De Simoni, C., von Bastian, C.C., 2017. Do Individual Differences Predict Change
in Cognitive Training Performance? A Latent Growth Curve Modeling Approach. Journal of
Cognitive Enhancement 1, 374-393.
Haut, K.M., Lim, K.O., MacDonald Iii, A., 2010. Prefrontal Cortical Changes Following
Cognitive Training in Patients with Chronic Schizophrenia: Effects of Practice,
Generalization, and Specificity. Neuropsychopharmacology 35, 1850.
Heinzel, S., Lorenz, R.C., Pelz, P., Heinz, A., Walter, H., Kathmann, N., Rapp, M.A., Stelzel,
C., 2016. Neural correlates of training and transfer effects in working memory in older adults.
NeuroImage 134, 236-249.
Hill, A.T., Fitzgerald, P.B., Hoy, K.E., 2016. Effects of Anodal Transcranial Direct Current
Stimulation on Working Memory: A Systematic Review and Meta-Analysis of Findings From
Healthy and Neuropsychiatric Populations. Brain Stimulation 9, 197-208.
Hothorn, T., Bretz, F., Westfall, P., 2008. Simultaneous Inference in General Parametric
Models. Biometrical Journal 50, 346-363.
Jaeggi, S.M., Buschkuehl, M., Shah, P., Jonides, J., 2014. The role of individual differences
in cognitive training and transfer. Memory & Cognition 42, 464-480.
Jenkinson, M., Bannister, P., Brady, M., Smith, S., 2002. Improved Optimization for the
Robust and Accurate Linear Registration and Motion Correction of Brain Images.
NeuroImage 17, 825-841.
Jenkinson, M., Beckmann, C.F., Behrens, T.E.J., Woolrich, M.W., Smith, S.M., 2012. FSL.
NeuroImage 62, 782-790.
Jenkinson, M., Smith, S., 2001. A global optimisation method for robust affine registration of
brain images. Medical Image Analysis 5, 143-156.
Jo, J.M., Kim, Y.-H., Ko, M.-H., Ohn, S.H., Joen, B., Lee, K.H., 2009. Enhancing the
Working Memory of Stroke Patients Using tDCS. American Journal of Physical Medicine &
Rehabilitation 88.
Johnstone, T., Somerville, L.H., Alexander, A.L., Oakes, T.R., Davidson, R.J., Kalin, N.H.,
Whalen, P.J., 2005. Stability of amygdala BOLD response to fearful faces over multiple scan
sessions. NeuroImage 25, 1112-1123.
Kang, N., Summers, J.J., Cauraugh, J.H., 2015. Transcranial direct current stimulation
facilitates motor learning post-stroke: a systematic review and meta-analysis. Journal of
Neurology, Neurosurgery & Psychiatry.
Lawrence, B.J., Gasson, N., Johnson, A.R., Booth, L., Loftus, A.M., 2018. Cognitive Training
and Transcranial Direct Current Stimulation for Mild Cognitive Impairment in Parkinson's
Disease: A Randomized Controlled Trial. Parkinson's Disease 2018.

Leung, I.H.K., Walton, C.C., Hallock, H., Lewis, S.J.G., Valenzuela, M., Lampit, A., 2015.
Cognitive training in Parkinson disease: A systematic review and meta-analysis. Neurology
85, 1843-1851.
Lin, Z.C., Tao, J., Gao, Y.L., Yin, D.Z., Chen, A.Z., Chen, L.D., 2014. Analysis of central
mechanism of cognitive training on cognitive impairment after stroke: Resting-state
functional magnetic resonance imaging study. Journal of International Medical Research 42.
Lipp, I., Benedek, M., Fink, A., Koschutnig, K., Reishofer, G., Bergner, S., Ischebeck, A.,
Ebner, F., Neubauer, A., 2012. Investigating neural efficiency in the visuo-spatial domain: an
FMRI study. PloS one 7, e51316-e51316.
Maffei, L., Picano, E., Andreassi, M.G., Angelucci, A., Baldacci, F., Baroncelli, L.,
Begenisic, T., Bellinvia, P.F., Berardi, N., Biagi, L., Bonaccorsi, J., Bonanni, E., Bonuccelli,
U., Borghini, A., Braschi, C., Broccardi, M., Bruno, R.M., Caleo, M., Carlesi, C., Carnicelli,
L., Cartoni, G., Cecchetti, L., Cenni, M.C., Ceravolo, R., Chico, L., Cintoli, S., Cioni, G.,
Coscia, M., Costa, M., D'Angelo, G., D'Ascanio, P., Nes, M.D., Turco, S.D., Coscio, E.D.,
Galante, M.D., Lascio, N.d., Faita, F., Falorni, I., Faraguna, U., Fenu, A., Fortunato, L.,
Franco, R., Gargani, L., Gargiulo, R., Ghiadoni, L., Giorgi, F.S., Iannarella, R., Iofrida, C.,
Kusmic, C., Limongi, F., Maestri, M., Maffei, M., Maggi, S., Mainardi, M., Mammana, L.,
Marabotti, A., Mariotti, V., Melissari, E., Mercuri, A., Micera, S., Molinaro, S., Narducci, R.,
Navarra, T., Noale, M., Pagni, C., Palumbo, S., Pasquariello, R., Pellegrini, S., Pietrini, P.,
Pizzorusso, T., Poli, A., Pratali, L., Retico, A., Ricciardi, E., Rota, G., Sale, A., Sbrana, S.,
Scabia, G., Scali, M., Scelfo, D., Sicari, R., Siciliano, G., Stea, F., Taddei, S., Tognoni, G.,
Tonacci, A., Tosetti, M., Turchi, S., Volpi, L., 2017. Randomized trial on the effects of a
combined physical/cognitive training in aged MCI subjects: the Train the Brain study.
Scientific Reports 7, 39471.
Mahon, S., Parmar, P., Barker-Collo, S., Krishnamurthi, R., Jones, K., Theadom, A., Feigin,
V., 2017. Determinants, Prevalence, and Trajectory of Long-Term Post-Stroke Cognitive
Impairment: Results from a 4-Year Follow-Up of the ARCOS-IV Study. Neuroepidemiology
49, 129-134.
Marquez, J., van Vliet, P., McElduff, P., Lagopoulos, J., Parsons, M., 2015. Transcranial
Direct Current Stimulation (tDCS): Does it Have Merit in Stroke Rehabilitation? A
Systematic Review. International Journal of Stroke 10, 306-316.
Martin, D.M., Liu, R., Alonzo, A., Green, M., Player, M.J., Sachdev, P., Loo, C.K., 2013.
Can transcranial direct current stimulation enhance outcomes from cognitive training? A
randomized controlled trial in healthy participants. International Journal of
Neuropsychopharmacology 16, 1927-1936.
Medina, J., Cason, S., 2017. No evidential value in samples of transcranial direct current
stimulation (tDCS) studies of cognition and working memory in healthy populations. Cortex
94, 131-141.
Melby-Lervåg, M., Hulme, C., 2013. Is working memory training effective? A meta-analytic
review. Developmental Psychology 49, 270-291.
Melby-Lervåg, M., Redick, T.S., Hulme, C., 2016. Working Memory Training Does Not
Improve Performance on Measures of Intelligence or Other Measures of "Far Transfer":
Evidence From a Meta-Analytic Review. Perspectives on Psychological Science 11, 512-534.
Miró-Padilla, A., Bueichekú, E., Ventura-Campos, N., Flores-Compañ, M.-J., Parcet, M.A.,
Ávila, C., 2018. Long-term brain effects of N-back training: an fMRI study. Brain Imaging
and Behavior.
Morrison, A.B., Chein, J.M., 2011. Does working memory training work? The promise and
challenges of enhancing cognition by training working memory. Psychonomic Bulletin &
Review 18, 46-60.
Motes, M.A., Yezhuvath, U.S., Aslan, S., Spence, J.S., Rypma, B., Chapman, S.B., 2018.
Higher-order cognitive training effects on processing speed–related neural activity: a
randomized trial. Neurobiology of Aging 62, 72-81.

814 Motter, J.N., Pimontel, M.A., Rindskopf, D., Devanand, D.P., Doraiswamy, P.M., Sneed,
815 J.R., 2016. Computerized cognitive training and functional recovery in major depressive
816 disorder: A meta-analysis. Journal of Affective Disorders 189, 184-191.
817 Ørbeck, B., Sundet, K., 2007. WASI (Wechsler Abbreviated Scale of Intelligence) Norsk
818 versjon Manualsupplement. Stockholm: Harcort Asessment Inc.
819 P. Filzmoser, Gschwandtner, M., 2018. mvoutlier: Multivariate Outlier Detection Based on
820 Robust Methods.
821 Park, S.H., Koh, E.J., Choi, H.Y., Ko, M.H., 2013. A double-blind, sham-controlled, pilot
822 study to assess the effects of the concomitant use of transcranial direct current stimulation
823 with the computer assisted cognitive rehabilitation to the prefrontal cortex on cognitive
824 functions in patients with stroke. J Korean Neurosurg Soc. 54.
825 Peers, P.V., Astle, D.E., Duncan, J., Murphy, F., Hampshire, A., Das, T., Manly, T., 2018. A
826 randomised exploratory investigation of the effects of Attention vs Working Memory
827 Training on cognitive performance and everyday functioning following stroke. bioRxiv.
828 Pendlebury, S.T., 2012. Dementia in Patients Hospitalized with Stroke: Rates, Time Course,
829 and Clinico-Pathologic Factors. International Journal of Stroke 7, 570-581.
830 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Team, R.C., 2017. nlme: Linear and
831 Nonlinear Mixed Effects Models - R package version 3.1-131.
832 Prikken, M., Konings, M.J., Lei, W.U., Begemann, M.J.H., Sommer, I.E.C., 2018. The
833 efficacy of computerized cognitive drill and practice training for patients with a
834 schizophrenia-spectrum disorder: A meta-analysis. Schizophrenia Research.
835 Pylyshyn, Z.W., Storm, R.W., 1988. Tracking multiple independent targets: evidence for a
836 parallel tracking mechanism.
837 Ramsay, I.S., MacDonald, I.I.I.A.W., 2015. Brain Correlates of Cognitive Remediation in
838 Schizophrenia: Activation Likelihood Analysis Shows Preliminary Evidence of Neural Target
839 Engagement. Schizophrenia Bulletin 41, 1276-1284.
840 Ruf, S.P., Fallgatter, A.J., Plewnia, C., 2017. Augmentation of working memory training by
841 transcranial direct current stimulation (tDCS). Scientific Reports 7, 876.
842 Sala, G., Gobet, F., 2018. Cognitive Training Does Not Enhance General Cognition. Trends
843 in Cognitive Sciences.
844 Schaapsmeerders, P., Maaijwee Na Fau - van Dijk, E.J., van Dijk Ej Fau - Rutten-Jacobs,
845 L.C.A., Rutten-Jacobs Lc Fau - Arntz, R.M., Arntz Rm Fau - Schoonderwaldt, H.C.,
846 Schoonderwaldt Hc Fau - Dorresteijn, L.D.A., Dorresteijn Ld Fau - Kessels, R.P.C., Kessels
847 Rp Fau - de Leeuw, F.-E., de Leeuw, F.E., 2013. Long-term cognitive impairment after first-
848 ever ischemic stroke in young adults. Stroke 44.
849 Shipstead, Z., Hicks, K.L., Engle, R.W., 2012. Cogmed working memory training: Does the
850 evidence support the claims? Journal of Applied Research in Memory and Cognition 1, 185-
851 193.
852 Smith, S.M., Brady, J.M., 1997. SUSAN—A New Approach to Low Level Image Processing.
853 International Journal of Computer Vision 23, 45-78.
854 Smith, S.M., Jenkinson, M., Woolrich, M.W., Beckmann, C.F., Behrens, T.E.J., Johansen-
855 Berg, H., Bannister, P.R., De Luca, M., Drobnjak, I., Flitney, D.E., Niazy, R.K., Saunders, J.,
856 Vickers, J., Zhang, Y., De Stefano, N., Brady, J.M., Matthews, P.M., 2004. Advances in
857 functional and structural MR image analysis and implementation as FSL. NeuroImage 23,
858 Supplement 1, S208-S219.
859 Smith, S.M., Nichols, T.E., 2009. Threshold-free cluster enhancement: Addressing problems
860 of smoothing, threshold dependence and localisation in cluster inference. NeuroImage 44, 83-
861 98.
862 Specht, K., 2020. Current Challenges in Translational and Clinical fMRI and Future
863 Directions. Frontiers in psychiatry 10, 924-924.

Spencer-Smith, M., Klingberg, T., 2015. Benefits of a Working Memory Training Program
for Inattention in Daily Life: A Systematic Review and Meta-Analysis. PLOS ONE 10,
e0119522.
Strobel, K., Engedal, K., 2008. MMSE-NR, Norsk revidert Mini Mental Status Evaluering.
Book MMSE-NR, Norsk revidert Mini Mental Status Evaluering. Oslo: Nasjonal
kompetansetjeneste i aldring og helse.
Subramaniam, K., Luks, T.L., Garrett, C., Chung, C., Fisher, M., Nagarajan, S., Vinogradov,
S., 2014. Intensive cognitive training in schizophrenia enhances working memory and
associated prefrontal cortical efficiency in a manner that drives long-term functional gains.
NeuroImage 99, 281-292.
Synhaeve, N.E., Arntz Rm Fau - Maaijwee, N.A.M., Maaijwee Na Fau - Rutten-Jacobs,
L.C.A., Rutten-Jacobs Lc Fau - Schoonderwaldt, H.C., Schoonderwaldt Hc Fau - Dorresteijn,
L.D.A., Dorresteijn Ld Fau - de Kort, P.L.M., de Kort Pl Fau - van Dijk, E.J., van Dijk Ej Fau
- de Leeuw, F.-E., de Leeuw, F.E., 2014. Poor long-term functional outcome after stroke
among adults aged 18 to 50 years: Follow-Up of Transient Ischemic Attack and Stroke
Patients and Unelucidated Risk Factor Evaluation (FUTURE) study. Stroke 45.
Team, R.C., 2013. R: A language and environment for statistical
computing
Teasell, R.W., 1992. Long-Term Sequelae of Stroke: How should you handle stroke
complications? Canadian Family Physician 38, 381-388.
Ulrichsen, K.M., Alnæs, D., Kolskår, K.K., Richard, G., Sanders, A.M., Dørum, E.S., Ihle-
Hansen, H., Pedersen, M.L., Tornås, S., Nordvik, J.E., Westlye, L.T., 2019. Dissecting the
cognitive phenotype of post-stroke fatigue using drift diffusion modeling of sustained
attention. bioRxiv 582502.
von Bastian, C.C., Oberauer, K., 2014. Effects and mechanisms of working memory training:
a review. Psychological Research 78, 803-820.
Weicker, J., Villringer, A., Thöne-Otto, A., 2016. Can impaired working memory functioning
be improved by training? A meta-analysis with a special focus on brain injured patients.
Neuropsychology 30, 190-212.
Westerberg, H., Jacobaeus, H., Hirvikoski, T., Clevberger, P., Östensson, M.L., Bartfai, A.,
Klingberg, T., 2007. Computerized working memory training after stroke–A pilot study.
Brain Injury 21, 21-29.
Wolak, M.E., Fairbairn, D.J., Paulsen, Y.R., 2012. Guidelines for Estimating Repeatability.
Methods in Ecology and Evolution 3.
Woolrich, M.W., Ripley, B.D., Brady, M., Smith, S.M., 2001. Temporal Autocorrelation in
Univariate Linear Modeling of FMRI Data. NeuroImage 14, 1370-1386.
Yamasaki, T., Ogawa, A., Osada, T., Jimura, K., Konishi, S., 2018. Within-Subject
Correlation Analysis to Detect Functional Areas Associated With Response Inhibition.
Frontiers in Human Neuroscience 12, 208.
Yang, S., Jiang, C., Ye, H., Tao, J., Huang, J., Gao, Y., Lin, Z., Chen, L., 2014. Effect of
integrated cognitive therapy on hippocampal functional connectivity patterns in stroke
patients with cognitive dysfunction: a resting-state FMRI study. Evidence-based
complementary and alternative medicine : eCAM 2014, 962304-962304.
Ziemus, B., Baumann, O., Luerding, R., Schlosser, R., Schuierer, G., Bogdahn, U., Greenlee,
M.W., 2007. Impaired working-memory after cerebellar infarcts paralleled by changes in
BOLD signal of a cortico-cerebellar circuit. Neuropsychologia 45, 2016-2024.
Zinn, S., Bosworth, H.B., Hoenig, H.M., Swartzwelder, H.S., 2007. Executive Function
Deficits in Acute Stroke. Archives of Physical Medicine and Rehabilitation 88, 173-180.
Alnæs, D., Sneve, M.H., Espeseth, T., Endestad, T., van de Pavert, S.H.P., Laeng, B., 2014.
Pupil size signals mental effort deployed during multiple object tracking and predicts brain
activity in the dorsal attention network and the locus coeruleus. Journal of Vision 14, 1-1.

Boggio, P.S., Nunes, A., Rigonatti, S.P., Nitsche, M.A., Pascual-Leone, A., Fregni, F., 2007.
Repeated sessions of noninvasive brain DC stimulation is associated with motor function
improvement in stroke patients. Restorative neurology and neuroscience 25, 123-129.
Dørum, E.S., Alnæs, D., Kaufmann, T., Richard, G., Lund, M.J., Tønnesen, S., Sneve, M.H.,
Mathiesen, N.C., Rustan, Ø.G., Gjertsen, Ø., Vatn, S., Fure, B., Andreassen, O.A., Nordvik,
J.E., Westlye, L.T., 2016. Age-related differences in brain network activation and co-
activation during multiple object tracking. Brain and Behavior 6, e00533.
Marquez, J., van Vliet, P., McElduff, P., Lagopoulos, J., Parsons, M., 2013. Transcranial
Direct Current Stimulation (tDCS): Does it Have Merit in Stroke Rehabilitation? A
Systematic Review. International Journal of Stroke 10, 306-316.
Olesen, P.J., Westerberg, H., Klingberg, T., 2004. Increased prefrontal and parietal activity
after training of working memory. Nature Neuroscience 7, 75-79.
Park, J., Oh, Y., Chung, K., Kim, K.J., Kim, C.O., Park, J.Y., 2019. Effect of home-based
transcranial direct current stimulation (tDCS) on cognitive function in patients with mild
cognitive impairment: a study protocol for a randomized, double-blind, cross-over study.
Trials 20, 278.
Pylyshyn, Z.W., Strom, R.W., 1988. Tracking multiple independent targets: Evidence for a
parallel tracking mechanism*. Spatial Vision 3, 179-197.
Richard, G., Kolskår, K., Ulrichsen, K., Kaufmann, T., Alnæs, D., Sanders, A.-M., Dørum,
E., Sánchez, J.M., Petersen, A., Ihle-Hansen, H., Nordvik, J.E., Westlye, L., 2019. Reliable
longitudinal brain age prediction in stroke patients: Associations with cognitive function and
response to cognitive training. bioRxiv.
Richard, G., Kolskår, K., Ulrichsen, K.M., Kaufmann, T., Alnæs, D., Sanders, A.-M., Dørum,
E.S., Monereo Sánchez, J., Petersen, A., Ihle-Hansen, H., Nordvik, J.E., Westlye, L.T., 2020.
Brain age prediction in stroke patients: Highly reliable but limited sensitivity to cognitive
performance and response to cognitive training. NeuroImage: Clinical 25, 102159.
Team, R., 2016. RStudio: Integrated Development Environment for R. RStudio, Inc.
Walle, K.M., Nordvik, J.A.-O.X., Espeseth, T., Becker, F., Laeng, B., 2019. Multiple object
tracking and pupillometry reveal deficits in both selective and intensive attention in unilateral
spatial neglect. J Clin Exp Neuropsychol. 41, 270-289.
Winkler, A.M., Ridgway, G.R., Webster, M.A., Smith, S.M., Nichols, T.E., 2014.
Permutation inference for the general linear model. NeuroImage 92, 381-397.
Yun, G.J., Chun, M.H., Kim, B.R., 2015. The Effects of Transcranial Direct-Current
Stimulation on Cognition in Stroke Patients. J Stroke 17, 354-358.