Title: Bats dynamically change echolocation parameters in response to acoustic

playback

Running Title: Adaptations in response to acoustic playback

M. Jerome Beetz (*)[1,2], Manfred Kössl[1], Julio C. Hechavarría[1]

Affiliations

[1]Institut für Zellbiologie und Neurowissenschaft, Goethe-Universität, Frankfurt/M., Germany

[2] current address: Zoology II Emmy-Noether Animal Navigation Group, Biocenter, University of

Würzburg, Germany

* Corresponding author

Mailing address:

M. Jerome Beetz

Email: jerome.beetz@uni-wuerzburg.de

Zoology II, University of Wuerzburg, Am Hubland, 97074, Wuerzburg, Germany, Tel.: +49

9313184528

Keywords *echolocation, active sensing, bioacoustics, signal interference, spatial orientation,*

Jamming Avoidance Response

Summary statement

The frugivorous bat *Carollia perspicillata* dynamically switch between different adaptations when

echolocating in acoustically contaminated environments.

Abstract

Animals extract behaviorally relevant signals from "noisy" environments. To investigate signal

extraction, echolocating provides a rich system testbed. For orientation, bats broadcast calls and assign

each echo to the corresponding call. When orienting in acoustically enriched environments or when

approaching targets, bats change their spectro-temporal call design. Thus, to assess call adjustments

that are exclusively meant to facilitate signal extraction in "noisy" environments, it is necessary to

control for distance-dependent call changes. By swinging bats in a pendulum, we tested the influence

of acoustic playback on the echolocation behavior of *Carollia perspicillata*. This paradigm evokes

reproducible orientation behavior and allows a precise definition of the influence of the acoustic

context. Our results show that bats dynamically switch between different adaptations to cope with

sound-based navigation in acoustically contaminated environments. These dynamics of echolocation

behavior may explain the large variety of adaptations that have been reported in the bat literature.

Introduction

For orientation, echolocating bats emit biosonar calls and listen to their echoes arising from reflections of surrounding objects (Kössl et al., 2014; Moss and Surlykke, 2010; Simmons, 2012). Spectro-temporal parameters of echoes inform the animals about the position and identity of objects nearby (Wohlgemuth et al., 2016b). To gain spatial information, bats must assign incoming echoes to their corresponding calls (Corcoran and Moss, 2017; Suga et al., 1983; Ulanovsky et al., 2004). Call-echo assignments become challenging, however, when biosonar signals from many bats are overlapping (Corcoran and Moss, 2017; Levin et al., 2013; Parsons et al., 2003; Ulanovsky and Moss, 2008). Under these circumstances, bats demonstrate a large repertoire of behavioral adaptations that are thought to represent behavioral strategies to improve signal extraction. These adaptations range from spectro-temporal changes in call design, to changes in call emission patterns (Adams et al., 2017; Amichai et al., 2015; Cvikel et al., 2015; Gillam and McCracken, 2007; Gillam et al., 2007; Habersetzer, 1981; Hage et al., 2013; Hiryu et al., 2010; Ibanez et al., 2004; Jarvis et al., 2013; Luo et al., 2015; Miller and Degn, 1981; Obrist, 1995; Ratcliffe et al., 2004; Roverud and Grinnell, 1985a; Roverud and Grinnell, 1985b; Simmons et al., 1979; Simmons et al., 1978; Takahashi et al., 2014; Tressler and Smotherman, 2009; Ulanovsky et al., 2004).

Our current understanding of why bats show such a large variety of adaptations when echolocating under "noisy" conditions is sparse. For example, it remains unknown whether the adaptations observed are individualistic and/or depend on the environmental context in which bats vocalize (i.e., the distance between the bat and the nearest target).

In this study, we tested the hypothesis that individual bats rely on different combinations of behavioral adaptations to overcome noise and that they can switch adaptation strategies at any timepoint during echolocation. To test this hypothesis and to gain a clearer understanding of echolocation behavior in "noisy" environments, individual bats of the species *Carollia perspicillata* were attached on the mass of a swinging pendulum (Figure 1A). The pendulum offers a behavioral paradigm whereby bats could actively echolocate in controlled scenarios, which could be replicated over several trials (Beetz et al., 2016b; Beetz et al., 2017; Henson et al., 1982; Macias et al., 2016). In

our experiments, during forward swings – which mimicked a bat closing in on a target– the animals

were acoustically stimulated with patterned echolocation calls broadcast from a speaker, which

travelled with and pointing towards the animal (test trial). The call design and emission pattern of test

trials were then compared to those recorded during control trials in which bats were swung in the

absence of playback stimuli. During test trials, we examined whether bats would change different

echolocation parameters, including call duration, call level, call frequency composition, and call

emission pattern.

Materials and Methods

Animals

Experiments were conducted on 10 bats (5 females and 5 males) of the species *Carollia perspicillata*. The bats were bred and kept in a colony at the Institute for Cell Biology and Neuroscience (Goethe-University Frankfurt). The experiments complied with all current German laws on animal experimentation and in accordance with the Declaration of Helsinki. All experimental protocols were approved by the Regierungspräsidium Darmstadt (experimental permit # #FU-1126).

Pendulum paradigm and audio recordings

For controlling behavioral context, bats were positioned on the mass of a pendulum and were repetitively swung towards an acrylic glass wall (50 × 150 cm, Figure 1A) (Beetz et al., 2016b; Beetz et al., 2017; Henson et al., 1982; Macias et al., 2016). During the swing, the bats emitted echolocation sequences that were recorded, together with their echoes, by an ultrasound sensitive microphone (CM16/CMPA, Avisoft Bioacoustics, Germany). The microphone had a sensitivity of 50 mV/Pa and an input-referred self-noise level of 18 dB SPL, as reported by the manufacturer. The frequency response curve was flat (± 3 dB, as specified by the manufacturer) in the range from 30-130 kHz. The microphone travelled with the mass of the pendulum, which was medially positioned above the bat's head. The membrane of the microphone was adjusted as closely as possible to the bat's ears (~ 4 cm). The microphone was connected to a sound acquisition system (Ultra Sound Gate 116Hm mobile recording interface, + Recorder Software, Avisoft Bioacoustics, Germany). To test the influence of acoustic interference on echolocation behavior, bats were swung in the pendulum while they were acoustically stimulated with a playback stimulus (see below). We compared the echolocation behavior recorded in the absence of playback stimuli (control trials) with the one shown in the presence of playback (test trials). Our reasoning was that because the behavioral context was invariant during control and test trials, except for the occurrence of the playback stimulus, we could correlate adaptations in the echolocation behavior with the presence/absence of the playback. Initially, the bats were tested in a control trial followed by test trials where an echolocation call recorded during the forward swing of the control trial was selected to construct an individual-specific

playback stimulus. The playback stimulus consisted of an echolocation call that was presented as

quartets with a call interval of 25 ms and the quartets were repeated with an inter-quartet interval

between 130 and 150 ms. The intensity of the playback stimulus was adjusted to rms values (of single

calls) between 80 and 90 dB SPL for all animals. We reasoned that using an echolocation call of the

tested animal, as a playback stimulus, would be the most effective way of achieving acoustic jamming.

The latter is supported by the fact that subtle inter-individual differences in call design could be

detected by the animals, which reduces signal interference (Yovel et al., 2009). During test trials, the

playback stimulus was presented from an ultrasound speaker (MK 103.1 Microtech Gefell

Microphone Capsule used as speaker) that was flat in the range from 5 to 120 kHz (mean level in

calibration curve 84 ± 3 dB SPL, the speaker's protection cap was replaced with a self-made cap to

prevent energy loss at high frequencies). The speaker was placed pointing towards the bat's head at a

distance of 20 cm. The short distance between speaker and animal and the relatively tight fixation of

the bat's head prevented situations in which the bat could reduce acoustic interference via motor

responses like head "waggling" (Wohlgemuth et al., 2016a). Thus, the bats had to rely mostly on

changes in call design or emission pattern to minimize signal interference. Eight out of ten bats were

tested on two consecutive days, but with different, day-specific, playback stimuli. The latter controlled

for changes of the call design that may occur across days might bias our analysis. An overview of the

call parameters used for constructing playback stimuli is shown in Table 1.

Analyzed echolocation parameters

Since the time pattern of the playback stimuli was kept constant, we could discriminate between

biosonar signals emitted by the bat and the playback stimuli. The call emissions were manually tagged

in the software Avisoft SAS Lab Pro (Avisoft Bioacoustics, Germany). To characterize the

echolocation calls, different call parameters were measured in Avisoft SAS Lab Pro. The present study

focused on call level, call duration, peak frequency at different call time points (start, end, maximum

amplitude, and mean), bandwidth 5 (BW5), BW10, and sweep rate (Figure 1B). Regarding the call

spectra, we considered the peak frequencies (frequencies with the maximum energy at particular time

points of the call or on average of a call), because peak frequencies were likely to be the most salient

spectral information of the echo that would suffer least from reflective attenuation. BW5 and BW10

represents frequency ranges at 5 and 10 dB below the mean peak frequency (Figure 1B). The sweep

rate was calculated by subtracting the initial peak frequency from the terminal peak frequency and by

dividing by the call duration.

The call emission pattern was characterized by measuring the call intervals and the tendency

of grouping the calls. Analysis of the call groups was conducted using custom-written scripts in

Matlab 2014 (MathWorks, USA). Call groups were defined according to two criteria (Beetz et al.,

2018; Kothari et al., 2014). An "island criterion" defined call groups that were isolated in time. An

isolation was fulfilled as soon as the preceding and following call intervals of a call group were 20%

longer than the call intervals within call groups. If the "island criterion" was fulfilled, a second

criterion, the so called "stability criterion", defined the size of the call groups indicated by the number

of calls belonging to a group. The stability criterion was fulfilled if the call intervals within call groups

were stable with a 5% tolerance. Next, we calculated a strobe index for each animal and each

condition (control and test trial). The strobe index represented the relative amount of calls that were

emitted as groups.

Statistics

For statistical analysis, we used the software GraphPad Prism 7 (GraphPad Software, USA; *

$p < 0.05$; ** $p < 0.005$; *** $p < 0.001$; **** $p < 0.0001$). For analyzing distance-dependent changes of

the echolocation behavior in the pendulum, non-parametric Kruskal-Wallis tests and a Dunn`s

multiple comparison post hoc tests were computed. For analyzing individual specific call adaptations

in response to acoustic playback, control and test trials were directly compared from each animal by

performing non-parametric Mann Whitney (in case of non-Gaussian distribution according to

D´Agostino & Pearson normality test; alpha = 0.05) or parametric t-Tests (in case of Gaussian

distribution according to D´Agostino & Pearson normality test; alpha = 0.05). For a comparison of the

echolocation behavior between subsequent trials, non-parametric Kruskal-Wallis tests and Dunn`s

multiple comparison post hoc tests (in case of non-Gaussian distribution according to D´Agostino &

Pearson normality test; alpha = 0.05) or ordinary one-way ANOVA and Tukey´s multiple comparison

post hoc tests (in case of Gaussian distribution according to D´Agostino & Pearson normality test;

alpha = 0.05) were computed.

Results

Pendulum paradigm mimics a natural approach flight

When swinging bats on the mass of a pendulum, they often emit echolocation calls (Beetz et al.,

2016a; Henson et al., 1982; Macias et al., 2016). Thus, a pendulum paradigm allows to describe

echolocation behavior under controlled conditions. This is important to test the influence of acoustic

playbacks on echolocation behavior, independent from changes in the echolocation behavior due to

target distances.

First, we quantified if pendulum-forward swings evoked consistent distance-dependent

adjustments of the echolocation behavior in *C. perspicillata*. Based on 32 forward swings, each

recorded from a different individual, we found that the bats shortened their call duration and inter-call

intervals with decreasing target distance (Figure 2A-2B). In addition, with decreasing target distance,

the bats increased their call intensity, starting peak frequency and peak frequency at the call's

maximum energy (peak freq max; Figure 2C-2E). Since the distance-dependent adjustments in call

duration and call interval are comparable in the pendulum (laboratory condition) as in freely-flying

bats (Thies et al., 1998), we concluded that a forward swing in the pendulum mimics a bat zooming in

on a target in natural conditions.

Fig. 2 Distance-dependent changes of echolocation parameters during pendulum-forward swings
C. perspicillata reduces call durations (**A**) and call-intervals (**B**) with shorter distances to an object. Call intensities (**C**), initial peak frequency (**D**), and maximum peak frequency (**E**) slightly increase with shorter distance to an object. Subfigures on the left represent examples from three animals (M27, F2, F11) and subfigures on the right represent the median and the interquartile range from data of 32 bats in which each trial was normalized to its maximum value. Blue stars indicate significant differences ($p < 0.05$) between subsequent echolocation calls. Red stars indicate significant differences ($p < 0.05$) with calls emitted at echo delays between 23-22 ms (red data point). Kruskal-Wallis test + Dunn`s multiple comparison post hoc test.

Next, we tested the inter-swing variability in the echolocation behavior of nine bats (five females, four males, Supplementary Table 1). Call duration, intensity, starting peak frequency, sweep rate, terminal peak frequency, peak frequency at call's maximum energy, mean peak frequency, bandwidth 5, and bandwidth 10 did not vary over subsequent swings (Supplementary Table 1; $p > 0.05$) indicating that one forward swing reliably represents the echolocation behavior of an individual

bat. Only four animals (female/F9, F10, F11, and male/M12) increased their call intervals across

subsequent trials (Mann-Whitney test for F9 and F10; Kruskal-Wallis test for F11 and M12; $p <$

0.005), which may indicate that the bats habituated to the pendulum and therefore decreased the call

rate.

Individual bats change different call parameters in response to acoustic playback

To quantify adaptations of the echolocation behavior in response to acoustic playback, bats were

swung in the pendulum while presenting an echolocation sequence (playback stimulus). The sequence

was presented through a speaker attached to the pendulum mass, pointing towards the animal's head

(Figure 1A, test trials). One echolocation call from each tested bat served as building block for the

playback stimulus (see methods for details). Thus, for each animal and experimental day, a new

"individualized" playback stimulus was constructed (for stimulus details see methods and Table 1). In

total, the echolocation behavior in the presence of playback stimuli was characterized in ten bats (5

females and 5 males). Echolocation behavior in the presence of playback was compared with the

behavior recorded during an initial control trial in which no acoustic stimulus was played back to the

animals. To minimize habituation to the pendulum paradigm, we decided to have only one control trial

per session (per animal and day). As previously described, call parameters from subsequent control

trials do not vary across swings (except for call intervals, Supple Table 1). Thus, one control trial is

enough to characterize the bat's echolocation behavior in the absence of playback stimuli. Since bats

adjust their call design and emission pattern with the target distance (Figure 2), we pooled the calls

into two groups, namely "long delay calls" and "short delay calls". Echolocation calls that were

broadcasted as the bat was farther than 1 m away from the acrylic glass wall were defined as "long

delay calls". Here, the echoes are delayed by more than 6 ms from the calls. Accordingly, echolocation

calls that were emitted when the bat was closer than 1 m from the acrylic glass wall were defined as

"short delay calls" (echo delays equal to or shorter than 6 ms).

animal (day)	call duration [ms]	intensity rms [dB]	p f start [kHz]	p f end [kHz]	p f centre [kHz]	p f max [kHz]	p f mean [kHz]	BW5 [kHz]	BW10 [kHz]	sweep rate [kHz/ms]
f8 (1)	1,57	80,13	68,8	68,1	67,3	68,1	68,1	4,3	20,5	-0,446
f8 (2)	1,62	83,77	82	82	82,7	82	82	9,5	12,4	0,000
f9 (1)	1,94	87,02	61,5	82	82	82	82,7	27	30	10,567
f9 (2)	1,64	78,8	80,5	90,8	87,1	87,1	87,1	11,7	27,8	6,280
f10 (1)	1,85	85,86	68,1	85,6	82,7	86,4	85,6	16,8	28,5	9,459
f11 (1)	2,34	84,9	74,7	76,9	82	82	82	21,2	29,2	0,940
f11 (2)	2	78,13	69,5	79,8	85,6	86,4	86,4	7,3	13,1	5,150
f12 (1)	1,96	85,6	74,7	79,1	79,1	85,6	79,1	19	24,1	2,245
f12 (2)	2,15	80,25	80,5	82	82,7	87,1	82,7	10,2	27	0,698
m9 (1)	2,04	87,08	79,1	84,2	82,7	71	71	33,6	38,8	2,500
m9 (2)	2,3	83,95	78,3	82,7	82,7	82,7	82,7	5,1	22,7	1,913
m10 (1)	2,21	86,72	71	80,5	80,5	81,2	80,5	5,8	10,2	4,299
m10 (2)	1,62	82,25	67,3	84,9	82	87,1	82,7	9,5	32,9	10,864
m11 (1)	1,68	81,53	70,3	81,2	82	82	82,7	5,1	10,9	6,488
m12 (1)	1,4	83,57	79,1	66,6	61,5	79,1	67,3	23,4	24,9	-8,929
m12 (2)	1,53	77,23	79,8	89,3	87,8	87,8	87,8	4,3	27,8	6,209
m13 (1)	2,64	84,92	72,5	82	82,7	85,6	82,7	7,3	11,7	3,598
m13 (2)	2	76,35	67,3	81,2	80,5	82	80,5	10,9	33,6	6,950

In the presence of the playback stimulus, each individual bat demonstrated different

combinations of adaptations (Table 2 and Supple Table 2). Four bats (F11, F12, M9, M12) increased

the tendency of grouping their calls into call packs (exemplarily shown for F11 in Figure 3A; test trial,

see also population data in Figure 3E). Six bats (F8, F10, F11, M10, M12, and M13) varied their call

intervals. However, only a reduction of the call interval (observed in two bats, F11, M13) could be

interpreted as an adaptation in response to the playback stimulus. Increased call intervals may be

interpreted as habituation to the pendulum paradigm (see also Supple Table 1). Three bats (F8, M9,

M13) increased and another three bats decreased (F9, M11, M12) call intensity during the test trials

(Table 2; example in Figure 3B). Five bats changed their call duration, two shortened (F8, M11), two

lengthened (M9, M10) and one shortened their "short delay calls" and lengthened their "long delay

calls" (F9, Figure 3B; Table 2; Supple Table 2). The adaptation in call duration of F9 indicated that

some bats differentially adapt "long delay calls" and "short delay calls" in response to playbacks.

Changes in call spectra were sometimes prominent (Figure 3C and 3D) but also varied across animals (Table 2; Supplement Table 2). Calls shown in figure 3C and 3D were recorded as the bat had approximately the same distance to the target (~2 m). Seven out of eight bats that changed their calls' sweep rate decreased it when in the presence of playback. This indicated that the call frequency changed more slowly during the test compared to the control trials (Table 2; Supple Table 2). Changes in the sweep rate could be caused by changes of the call's frequency range or by changes in call duration. Because lowering the sweep rate was not associated with lengthening the call, the sweep rate was mainly affected by changes in the frequency range. Seven animals (70%) changed either the BW5 or BW10 of the calls in the test trials. These changes could either be a BW decrease (shown by 40% of the bats tested; F8, F10, M11, M13) or an increase (shown by 30%; F9, F12, M9). Detailed data from three animals (A: F8; B: F9; C: M9) are plotted as boxplots in figure 4A-4C. For reasons of visualization, only call parameters that differed between the test and the control trials are plotted. Data from the remaining animals are presented in Supple figure 1. In conclusion, each animal adapted at least one call parameter in response to the playback stimuli. M11 was the only individual that did not change their call design (short delay calls) during the test trials. Overall, the bats changed different combinations of their call parameters, indicating that there was no common rule as to how to adapt to the playback stimuli.

Fig. 3 Examples of echolocation adaptation behaviors in response to playback stimuli
(**A**) Oscillograms of one control trial (top), the playback stimulus (middle), and one test trial (bottom) from female 11 (F11). Time points of call emissions are indicated by black or red dots above each oscillogram. During the control trial, the bat did not emit echolocation call groups. During the test trial, the bat grouped some calls into doublets (indicated by the number 2) or triplets (indicated by a "3"). Note that the jamming stimulus was recorded in addition to the echolocation calls of the test trial. Thus, oscillogram deflections without a dot represent signals coming from the playback stimulus. (**B**) Oscillogram of one control trial (top), the playback stimulus (middle), and one test trial (bottom) from female 8 (F8). In comparison to the calls emitted during the control trials, the call intensity was increased during the test trials. Numbers above each emitted call indicate the call intensity. (**C-D**) Power spectra (left) and spectrograms (right) of representative calls emitted during the control and test trial for two individuals (F11, M11). To exclude distance-dependent changes in the call design, all four calls were recorded as the bat was ~2 meters away from the acrylic wall. Both bats decreased the bandwidth and mean peak frequency of their calls during the test trials as compared with the calls recorded during the control trials. (**E**) Tendency of emitting grouped calls (strobe index) under control and test conditions in all bats tested (n=10).

Table 2 Changes of the call parameters induced by the presence of playback stimuli. + = higher values for test than for control trials (+ = p < 0.05; ++ p < 0.01; +++ p < 0.001); - = lower values for test than for control trials (- = p < 0.05; -- p < 0.01; --- p < 0.001); F = female; M = male; l = long delay calls; p f = peak frequency; s = short delay calls

Animal ID	F8		F9		F10		F11		F12		M9		M10		M11		M12		M13	
Delay	l	s	l	s	l	s	l	s	l	s	l	s	l	s	l	s	l	s	l	s
Interval	++	+++			+	++	--						++	+			+		--	
Duration	-		+++	---							+		+		--					
Intensity		+++		---							++				--		---	---		++
P F start			-	---					+		---									
Sweep rate		---	---		--	-	-		+		--		--				-			
P F end		+++			+							---	-	---		--				
P F max			---	--	--		-								--		++			
P F mean			---	--			--													
BW5	---		++								+									
BW10	---		+++		---				++		+				--					-

Fig. 4 Individual specific call adaptations in response to playback stimuli
(**A-C**) Boxplots (whiskers represent minimum and maximum values) from three individuals (female 8 = 1st row; female 9 = 2nd row; male 9 = 3rd row), showing call parameters that bats changed in response to the playback stimulus. Calls recorded under control conditions (absence of playback stimulus) are indicated by black boxplots, while white boxplots represent calls recorded under test conditions (presence of playback stimulus). Echolocation calls that are followed by an echo within 6 ms were grouped into "short delay calls". Echoes following a call by more than 6 ms were grouped into "long delay calls". Note that each bat changed their different call parameters under test conditions. MW = Mann-Whitney test; * p < 0.05; ** p < 0.005; *** p < 0.001.

Bats vary adaptation strategies across trials and days

We were interested in assessing if each individual bat prefers the same combination of adaptations or

whether the bats change their strategies across days or even across trials on the same day. However,

before characterizing the temporal dynamics of the adaptations, we quantified the variability of the call

design across subsequent days under controlled conditions (absence of playback stimulus). We tested

ten bats in the absence of playback stimuli for two (F1, M3, M7) or three (F2, F3, F5, F7, M1, M4,

M6) subsequent days (Supple Table 3). Although the bats did not vary their call design across

subsequent control trials on the same day (Supple Table 1), they dramatically varied their call design

across subsequent days (Supple Table 3). Thus, to test if bats change their adaptations in response to

the playback stimulus across days, we recorded an initial control trial on each day and compared the

echolocation behavior from the day-specific control trial with the one recorded during test trials.

Moreover, to perform a trial-by-trial analysis and to gather enough data points for statistical analysis,

we pooled data from long and short delay calls. During the test trials, bats emitted slightly fewer calls

than during control trials (median n of calls: 16.5 control and 13 test; Mann-Whitney test: $p = 0.036$).

By comparing the call parameters from F9 across days (Figure 5; Table 3), it became clear that the

adjustments of call duration (Figure 5C), starting (Figure 5B), maximum (Figure 5F), and mean peak

frequency (Figure 5H), bandwidth 5 (Figure 5G) and bandwidth 10 (Figure 5I) exclusively occurred

on day 1. On day 2, bat F9 mainly changed call intensity (Figure 5A), terminal peak frequency (Figure

5D), and sweep rate (Figure 5J). As already mentioned, increments in call interval (Figure 5E) did not

necessarily represent an adaptation to reduce acoustic interference; possibly, they represented

habituation to the pendulum paradigm across trials/days.

      We observed that echolocation adaptation strategies not only varied across days, but also

across subsequent trials (Table 3). For example, F9 changed the calls' mean peak frequency in three

(trial 3, 4, 5) out of five trials at day 1 (Figure 5H). Changes of other call parameters varied less

dramatically across trials of the same day. In all trials on day 1, F9 decreased its starting (Figure 5B)

and maximum peak frequency (Figure 5F). For detailed data from the remaining nine animals see S2-

S10 figure. Overall, we found in 56 out of 67 test trials (83.6%) statistically significant differences

between the control and test trials (Table 3). In eleven test trials, the bats did not change any call

parameter compared to the control trial.

Fig.5 Bats switch adaptation strategies across trials and days
Call parameters are shown as boxplots (whiskers represent minimum and maximum values) for each trial (8 test trials and 2
control trials) across two days (from one bat). For visualization purposes, each trial is color coded and the control trials are
shown in black. Note that the bat changes some call parameters only at day 1 (e.g., peak freq start; call duration; peak freq
max) and not at day 2. Kruskal-Wallis Test and Dunn's multiple comparison Post hoc test; * $p < 0.05$; ** $p < 0.005$; *** $p <
0.001$, **** $p < 0.0001$.

333 Table 3 Changes of the call parameters across trials. + = higher values for test than for control trials (+ = p < 0.05; ++ p < 0.01; +++ p <
334 0.001); - = lower values for test than for control trials (- = p < 0.05; -- p < 0.01; --- p < 0.001; ----p < 0.0001); F = female; M = male; l = long
335 delay calls; p f = peak frequency; s = short delay calls

Animal ID (calls/trial)	Trial (Day)	Intensity	Duration	Interval	Sweep rate	P F start	P F end	P F max	P F mean	BW5	BW10
F8 (25, 16, 14, 12, 16, 16, 13, 11)	1 (1st)	+	---	+++	-		++			--	---
	2 (1st)							+			
	3 (1st)	++		++						--	----
	4 (2nd)				-		+				
	5 (2nd)										
	6 (2nd)			+							
F9 (28, 15, 10, 15, 17, 23, 18, 34, 16, 16)	1 (1st)					-					
	2 (1st)	-			-	--		--			
	3 (1st)					-		--	-		
	4 (1st)		+++			--		----	---	+	+
	5 (1st)	+				--		-	-		
	6 (2nd)	++			--		+++				
	7 (2nd)		+								
	8 (2nd)	-		+	-		++				
F10 (22, 13, 10, 22, 16, 15, 16)	1 (1st)				--						
	2 (1st)		+		--			--			
	3 (1st)				-						
	4 (1st)				-			-			
	5 (1st)						+				
	6 (1st)			+							
F11 (13, 23, 7, 20, 8, 9, 16, 13, 11, 14, 15, 10, 9, 11)	1 (1st)	+		---				-			
	2 (1st)		-								
	3 (1st)			-			-	--	--		
	4 (1st)			-				-	-		
	5 (2nd)			-						++	
	6 (2nd)										
	7 (2nd)										
	8 (2nd)										
	9 (2nd)										
	10 (2nd)	--									
	11 (2nd)			--							
	12 (2nd)										
F12 (18, 14, 11, 11, 11, 24, 17, 16)	1 (1st)		+								
	2 (1st)			++	-						+
	3 (1st)						-				
	4 (2nd)			----	++++	++	--			++	++
	5 (2nd)			-	+		-			+	+
	6 (2nd)			-	+		-			+	++
M9 (23, 24, 15, 7, 9, 24, 23, 9, 7)	1 (1st)	++	++		-	----					
	2 (1st)			+		-					
	3 (1st)	++++								---	
	4 (1st)				--	----					
	5 (2nd)			--	-	--				+	++
	6 (2nd)									+	+
	7 (2nd)									+	
M10 (17, 7, 14, 18, 12, 12, 7, 7)	1 (1st)						-	--	-		
	2 (1st)						--	----		-	
	3 (2nd)						-				
	4 (2nd)	+++	+++	+							
	5 (2nd)		+	++							
	6 (2nd)		++	+	++	++	----				
M11 (22, 8, 12, 11, 10)	1 (1st)	--					---		--		
	2 (1st)						-				
	3 (1st)								-		
	4 (1st)	---	-								
M12 (20, 10, 7, 11, 13, 11)	1 (1st)						-				
	2 (1st)			++						--	-
	3 (2nd)										
	4 (2nd)				++						
M13 (15, 29, 11, 24, 20, 20, 14, 16, 12, 15)	1 (1st)	++		----	---	+++					
	2 (1st)										
	3 (1st)			-	---	++++					
	4 (2nd)				---	--				--	
	5 (2nd)			-	---	--					
	6 (2nd)										-

7 (2nd)		----	+			+++				-	
8 (2nd)				--	-	+				-	

Bats dynamically switch adaptation strategies within trials

What could have happened during test trials when we could not find an adaptation in the echolocation

behavior? For these trials, was the acoustic interference too weak to evoke adaptations? Alternatively,

might the bats have dynamically changed their adaptations during trials, so that the adaptation would

not be detectable when pooling calls from an entire swing? To assess the latter idea, we compared

parameters of each call from the test trial with the same parameters in the call used to construct the

playback stimulus. The upper color maps, in figure 6A and 6B, exemplarily show the relative

differences between call parameters and the playback parameters for two trials in two different bats

(M9 and F12). The calls are ordered along columns in which the leftmost column represents the call

with the longest echo-delay and the rightmost column represents the call with the shortest echo-delay.

Each line represents the relative difference of a call parameter with respect to a playback parameter.

This result was calculated by subtracting the playback parameters from the call parameters and by

normalizing the difference against its absolute maximal difference for the entire trial. The darker the

red and blue patches are, the more positive and negative were the call parameters in comparison to the

playback stimulus. Based on the trial in figure 6A, the bat initially emitted calls with lower starting

peak frequencies (peak start) and call intensities than the playback stimulus. At an echo delay of ~3 ms

(between the 12^{th} and 13^{th} call, white dashed line in Fig. 6A), the bat abruptly switched the strategy

and increased the maximum and mean peak frequency while decreasing the BW of subsequent calls.

To visualize abrupt changes better, we calculated the differences of the parameters of subsequent calls

and plotted the values in the bottom color maps shown in figure 6A and 6B. We defined an abrupt

change when the considered parameter varied by more than 50% between subsequent calls. For

example, according to figure 6A, the terminal (peak end), maximum peak frequency (peak max), and

sweep rate of call 13, are more than 50% higher than the ones of call 12. This outcome is indicated by

red cells at the corresponding column (white dashed line) in the lower color map of figure 6A.

Fig.6 Bats dynamically change echolocation parameters within trials

(**A-B** upper graphs) Color maps from two representative test trials (M9 in (A) and F12 in (B)), illustrating the differences between calls and playback stimuli in a call-wise manner. Along the x-axis, the calls are ordered according to their emission order during the trial. The echo delay value from some call-echo pairs are indicated in the x-axis. Along the y-axis, normalized call parameter differences are color coded. The differences were normalized to their absolute maximum value at the corresponding parameter for the specific trial. The differences of the following call parameters were considered: call duration, call intensity, peak frequency at the beginning, end and maximum of the call, mean peak frequency of the call, bandwidth 5 (BW5), bandwidth 10 (BW10), and sweep rate. In some trials, a clear transition of the adaptation strategies can be detected (white vertical dashed lines). In some cases, the bats alternate call values, as exemplified for F12 for the terminal and maximum peak frequency indicated by a gray and black arrowhead, respectively. (**A-B** lower graphs) Colormaps illustrating abrupt changes of call parameters across subsequent calls. Abrupt changes occurred when a call parameter between two consecutive calls varied by more than 50% (blue and red cells represent reductions or increases in the corresponding call parameter). Changes of the call parameters that are below 50% were not abrupt enough to be defined as a change (green cells). Transitions between adaptation strategies and alternations between call parameter values can be seen more easily in the lower colormaps. (**C**) Histogram showing the level of parameters that are abruptly changed per call for all investigated calls (n = 889). Note that almost 75% of the calls show at least one abrupt change. (**D**) Pie chart illustrating the distribution of abrupt changes over the call parameters. Abrupt changes occur primarily within the call spectrum and less often for the intensity or duration. (**E**) Mean values of the amount of change per trial, plotted against the call parameter. Spectral parameters are shown to vary more often across trials than non-spectral ones (duration and intensity). (**F**) Pie chart representing the relative distribution of alternations across different call parameters.

Abrupt changes in call design were also visible in other trials, like the one presented in figure

6B. Here, abrupt changes mainly occurred at around 2.5-ms echo delay (white dashed line) by

decreasing the call intensity, starting (peak start), and terminal frequency (peak end) while the

maximum peak frequency (peak max) as well as the call bandwidths (BW5 and BW10) abruptly

increased. By considering all calls (889 calls from 69 trials and 10 animals), about three quarters of the

calls (74.24%) showed abrupt changes in at least one call parameter (Figure 6C). About half of the

calls (50.84%) showed abrupt changes in more than one call parameter. The bats did not focus on a

specific call parameter, but rather changed most of their parameters with equal probability (Figure

6D). Call intensity and call duration were least (7.24%) abruptly changed within the trials.

When taking a closer look on the pattern of call changes over subsequent calls (color maps at

the bottom of figure 6B), it became clear that the bats sometimes changed the call parameters in an

alternating manner. During the second half of the trial, the bat alternated between high and low

terminal (peak end) and maximum peak frequencies (peak max), indicated by gray and black

arrowheads, respectively. Before analyzing the alternations in more detail, we questioned how often

the bats changed a certain call parameter during the trial. The bar plot in figure 6E shows that the bats

changed spectral parameters more often per trial (mean of peak start = 2.85 ± 2.39; mean of peak end

= 2.96 ± 1.59; mean of peak max = 2.8 ± 2.29; mean of BW5 = 2.84 ± 1.75; mean of BW10 = 3.26 ±

2.39; mean of sweep rate = 2.51 ± 1.82) than the call intensity (mean = 1.91 ± 1.57) and the call

duration (mean = 1.73 ± 1.46) ($p < 10^{-5}$ Kruskal-Wallis test). Because spectral parameters varied more

often during the trials, alternations occurred with a higher probability in spectral than in non-spectral

(call intensity and call duration) parameters (Figure 6F). Across the spectral parameters, the

probability of alternations did not differ significantly ($p = 0.91$ Kruskal-Wallis test), indicating that

alternations could equally occur in each of the analyzed call parameters.

Discussion

The present study characterized adaptation strategies of echolocating bats (fruit-eating bat *C. perspicillata*) in the presence of playback stimuli. These playback stimuli potentially interfered with the bat's biosonar signals, making signal extraction for the bat challenging. In their natural environments, adjustments to echolocation behavior not only occur in the presence of an acoustic interferer, but also when bats approach obstacles or transition between different locales. Thus, it was crucial to test for the influence of acoustic interference under an otherwise invariant behavioral context. The pendulum paradigm, indeed, fulfilled these requirements because the behavioral scenario of an approach flight could be repetitively mimicked (Figure 2).

Our results demonstrate that *C. perspicillata* changes its call parameters and emission pattern when echolocating in the presence of playback stimuli. Interestingly, instead of relying on one adaptation strategy, the bats combined different adaptation strategies (Table 2 and Figure 4). To our surprise, the bats switched between different strategies across (Table 3, Figure 5) and even within trials (Figure 6). This flexibility renders the echolocation behavior, in the presence of acoustic interferers, highly dynamic and unique across different individuals and time points. Utilizing such dynamics, the bats can create unique echolocation streams that may be distinguished from interfering signals.

Coping with signal interference

Signal interference is a challenge with which every animal must cope; they must extract behaviorally relevant signals from a noisy background. The greater the similarity between relevant signals and background, the more challenging signal extraction becomes. To facilitate signal extraction, animals show large repertoires of different behavioral adaptations (Corcoran and Moss, 2017; Ulanovsky and Moss, 2008) like orienting their sensory organs towards relevant signals (Eckmeier et al., 2008; Ganguly and Kleinfeld, 2004; Land, 2015; Ribak et al., 2009; Schroeder et al., 2010; Tarsitano and Andrew, 1999; Towal and Hartmann, 2006). For example, bats increase head waggles and the inter-pinna distance when orienting under challenging conditions (Wohlgemuth et al., 2016a). Potentially, this response improves localization of the echo source (Wohlgemuth et al.,

2016a). Additionally, adjustments of the pinna's shape and orientation may increase the directionality

of hearing (Gao et al., 2011). In the present study, head waggles were limited by tightly positioning

the bats on the platform of the pendulum mass. Moreover, by placing the jamming source close to the

animals' head, motor responses would barely facilitate signal extraction under such conditions.

For some behaviors - like communication, electrolocation, or echolocation - animals produce

behaviorally relevant signals, which allow them to directly control the signal's discriminability from

its background. The latter becomes clear when considering the cocktail party problem (Bee and

Micheyl, 2008). In a noisy environment, we can focus on our communication partner by carefully

listening to him/her and improve the signal-to-noise ratio by increasing our voice intensity (Brumm

and Zollinger, 2011; Luo et al., 2015), an adaptation known as the Lombard-effect. Signal extraction

may not only be improved by increasing the signal-to-noise ratio but also by limiting the spectral

overlap between signal and background. This adaptation has originally been described in

electrolocating fish (Bullock et al., 1972; Watanabe and Takeda, 1963). When encountering animals

whose signal frequencies overlap with the fish's own signal frequency, the animals shift the signal

frequencies away from each other. This behavior has been circumscribed as the Jamming Avoidance

Response (JAR) and it reduces the signal interference with signals coming from conspecifics. JAR has

also been demonstrated in different bat species (Gillam and McCracken, 2007; Gillam et al., 2007;

Habersetzer, 1981; Hage et al., 2013; Ibanez et al., 2004; Miller and Degn, 1981; Ratcliffe et al., 2004;

Takahashi et al., 2014; Tressler and Smotherman, 2009; Ulanovsky et al., 2004) and the present study.

However, in contrast to weakly electric fish, which try to occupy an individual specific frequency

band, bats dynamically adjust their call spectra in various situations. Bats adjust their calls when

approaching an obstacle or when transiting between different habitats (Barchi et al., 2013; Falk et al.,

2014; Griffin, 1953; Hiryu et al., 2010; Kalko, 1995; Kalko and Schnitzler, 1989; Knowles et al.,

2015; Kothari et al., 2014; Petrites et al., 2009; Roverud and Grinnell, 1985a; Schnitzler et al., 1987;

Simmons et al., 1978; Surlykke and Moss, 2000; Wheeler et al., 2016). Since frequency adjustments

occur frequently and under various conditions, an adaptation that purely depends on a JAR may not be

sufficient to orient collision-free in the presence of signal interferers. This hypothesis gains support by

recent simulations that have tested for the effectiveness frequency adjustments when navigating in

noisy environments (Mazar and Yovel, 2019). Note that some studies reported that bats do not shift

their frequency in response to acoustic interference (Götze et al., 2016) or that the frequency shifts are

correlated with the object's distance rather with the presence of an acoustic interferer (Cvikel et al.,

2015). Because we compared echolocation calls that were emitted roughly at similar object-distances,

we can exclude that frequency shifts, present in our study, reflect distance-dependent changes of the

call design.

Repertoire of behavioral adaptations in response to interfering signals and their possible neural

correlates

In the present study, most biosonar adjustments occurred in the spectral domain, although others were

also detected in the temporal or energy domain (Figure 5D). Each adaptation may facilitate signal

extraction in noisy environments. For example, decreasing the call bandwidth may reduce the spectral

overlap between the playback stimulus and the biosonar signals. An increase in call bandwidth may

recruit neurons that are not sensitive to the playback stimulus and therefore "selectively" process

frequencies that are not occupied by the interferers. Bats also increase the signal-to-noise ratio by

increasing call intensity (Amichai et al., 2015; Hage et al., 2013; Luo et al., 2015; Simmons, 2017;

Simmons et al., 1978; Takahashi et al., 2014; Tressler and Smotherman, 2009). Unexpectedly, in the

present study, sometimes the bats decreased their call intensity when they echolocated in the presence

of interfering signals. Although this decreases the signal-to-noise ratio, it could still be useful from the

perspective of neuronal processing. Many auditory neurons are more sensitive to low rather than to

high sound levels (Barone et al., 1996; Hechavarría and Kössl, 2014; Park and Pollak, 1993; Suga and

Manabe, 1982; Yang et al., 1992). This attribute makes the neurons highly selective to faint biosonar

signals while being insensitive to intense background stimuli. Some studies have reported that bats

lengthen their calls when flying in noisy environments (Amichai et al., 2015; Simmons, 2017;

Simmons et al., 1979; Simmons et al., 1975; Tressler and Smotherman, 2009). In the present study, we

observed that some bats lengthened, and others shortened their calls. Both adaptations putatively

minimize acoustic interference. Shortening the calls decreases the chance of a temporal overlap

between biosonar signals and the background. Lengthening the calls increases the risk of temporal

overlap, but it could still be useful if only a small portion of the echo needed to be detected to gain enough spatial information.

Furthermore, not only the call design, but also the emission pattern can be adjusted to reduce or even avoid signal interference. Some bat species alternate between two call designs that differ in their frequency spectrum (Obrist, 1995; Roverud and Grinnell, 1985a; Roverud and Grinnell, 1985b). This adaptation allows a higher call rate by emitting a pair of calls before receiving an echo from the first call of the pair (Behr and von Helversen, 2004; Jung et al., 2007). The arising echoes differ in their frequency spectra which makes their discrimination feasible (Hiryu et al., 2010). Alternation of spectral call parameters have also been observed in the present study (Figure 5B, 5F). However, these alternations occurred occasionally and not throughout the entire trial. Thus, the behavioral importance of alternating call parameters in *C. perspicillata* needs to be further assessed.

Some bats reduce their call rate (Adams et al., 2017) and temporally even cease to emit calls (Jarvis et al., 2013). This adaptation may be beneficial if the bats eavesdrop on echolocation signals from conspecifics and use the signals for orientation (Barclay, 1982; Chiu et al., 2008; Leonard and Fenton, 1984; Lin and Abaid, 2015). Although, *C. perspicillata* emitted fewer calls during test compared to control trials, the pendulum paradigm was not designed to test for eavesdropping on the playback stimulus.

Lastly, some bats increase their rate of grouping calls when orienting in cluttered or noisy environments ((Beetz et al., 2018; Beetz et al., 2019; Luo et al., 2015; Roverud and Grinnell, 1985a) and present study). Grouping calls may improve echolocation performance in different ways. First, a defined periodicity of echo arrivals allows echo identification based on prediction (Petrites et al., 2009; Suga et al., 1983; Wheeler et al., 2016; Wohlgemuth et al., 2016a). Second, grouping the calls could create an information redundancy allowing the bats to rely only on a small portion of the call group (Beetz et al., 2018).

Bats show different combinations of adaptations when echolocating in the presence of an acoustic interferer

Instead of adjusting just one echolocation parameter, when echolocating in noisy environments, our

results indicate that bats have a toolbox of different and combinable adaptations to minimize acoustic

interference (Hage et al., 2013; Luo et al., 2015). The dynamics and variability of such strategies are

important factors for explaining the high diversity of behavioral adaptations reported in earlier studies.

We must keep in mind that the discriminability of a signal from the background is dictated by the

difference of the physical parameters between the signal and its background. Essentially, it is

unimportant which physical parameter is adjusted, just so the signal has its own physical identity and

provides a high level of discriminability from its background. For call-echo assignment, it has been

discussed that bats keep an "internal copy" of their broadcasted calls and compare that copy with

received echoes (Simmons, 2012). Neural activity occurring before biosonar production in frontal and

striatal brain regions could contribute to the formation of such "internal copy" (Weineck et al., 2020).

The idea of an "internal copy" is in line with behavioral results showing that correct call-echo

assignment is decreased when spectro-temporal properties of the echo are manipulated (Masters and

Jacobs, 1989; Masters and Raver, 1996; Masters and Raver, 2000) or when echoes are replaced by

noise bursts (Surlykke, 1992). Because of missing behavioral data in *C. perspicillata*, it remains

speculative to what extent the echolocation calls need to differ from the playback stimuli so that bats

can still extract the signals. When comparing different call parameters against the playback stimuli

used in the present study, it becomes clear that some echolocation calls can vastly differ from playback

stimuli (S11 Figure). Although having no detection thresholds from *C. perspicillata*, there are some

behavioral and electrophysiological results from other bat species that use similar call designs as *C.*

perspicillata (for example, *Eptesicus fuscus*: (Chiu et al., 2009; von Stebut and Schmidt, 2001);

Tadarida brasilienisis: (Bartsch and Schmidt, 1993), *Antrozous pallidus* (Fuzessery, 1994)). Based on

these studies, we speculate that *C. perspicillata* can extract signals that differ for one of the following

parameters by more than 10 dB in intensity, by at least 0.7 ms in duration, by more than 5 kHz in the

peak frequency, by more than 12 kHz in bandwidth, and by more than 6 kHz in the sweep rate from

playback stimuli. By considering these thresholds, *C. perspicillata* may be able to extract about 94%

of the calls from the playback stimuli. Only 5.96% of the calls did not reach our hypothetical detection

thresholds for any of the investigated call parameters. Note that the emission pattern could not be

considered for a call-by-call analysis. Thus, it is still probable that the remaining 5.96% of the calls'

echoes could be detected by anticipation of the echo pattern. This ability could be accomplished by

grouping the calls (Figure 3A, (Beetz et al., 2018; Beetz et al., 2019)). In the present study, four out of

ten bats increased the tendency of grouping the calls, which increases the stimulus rate (Figure 3E).

Neurons of the bat's auditory midbrain and cortex can likely extract relevant echolocation information

when the bats face such high call rates (Bartenstein et al., 2014; Beetz et al., 2018; Beetz et al., 2016a;

Beetz et al., 2016b; Beetz et al., 2017; Greiter and Firzlaff, 2017; Macias et al., 2018; Sanderson and

Simmons, 2005).

Acknowledgements We thank Karen Mesce for her helpful comments on the manuscript.

Competing interests The authors declare no competing financial interests.

Funding This work was funded by the German Research Foundation (DFG) (Grant number: KO

987/12-2)

References

Adams, A. M., Davis, K. and Smotherman, M. (2017). Suppression of emission rates
improves sonar performance by flying bats. *Scientific Reports* **7**.
Amichai, E., Blumrosen, G. and Yovel, Y. (2015). Calling louder and longer: how bats use
biosonar under severe acoustic interference from other bats. *Proceedings of the Royal Society B-*
Biological Sciences **282**.
Barchi, J. R., Knowles, J. M. and Simmons, J. A. (2013). Spatial memory and stereotypy of
flight paths by big brown bats in cluttered surroundings. *Journal of Experimental Biology* **216**, 1053-
1063.
Barclay, R. M. R. (1982). Interindividual Use of Echolocation Calls - Eavesdropping by Bats.
Behavioral Ecology and Sociobiology **10**, 271-275.
Barone, P., Clarey, J. C., Irons, W. A. and Imig, T. J. (1996). Cortical synthesis of azimuth-
sensitive single-unit responses with nonmonotonic level tuning: A thalamocortical comparison in the
cat. *Journal of Neurophysiology* **75**, 1206-1220.
Bartenstein, S. K., Gerstenberg, N., Vanderelst, D., Peremans, H. and Firzlaff, U. (2014).
Echo-acoustic flow dynamically modifies the cortical map of target range in bats. *Nature*
Communications **5**, 4668.
Bartsch, E. and Schmidt, S. (1993). Psychophysical Frequency-Modulation Thresholds in a
Fm-Bat, Tadarida-Brasiliensis. *Hearing Research* **67**, 128-138.
Bee, M. A. and Micheyl, C. (2008). The cocktail party problem: What is it? How can it be
solved? And why should animal behaviorists study it? *Journal of Comparative Psychology* **122**, 235-
251.
Beetz, M. J., García-Rosales, F., Kössl, M. and Hechavarría, J. C. (2018). Robustness of
cortical and subcortical processing in the presence of natural masking sounds. *Sci Rep*.
Beetz, M. J., Hechavarria, J. C. and Kossl, M. (2016a). Cortical neurons of bats respond best
to echoes from nearest targets when listening to natural biosonar multi-echo streams. *Scientific*
Reports **6**.
Beetz, M. J., Hechavarria, J. C. and Kossl, M. (2016b). Temporal tuning in the bat auditory
cortex is sharper when studied with natural echolocation sequences. *Scientific Reports* **6**.
Beetz, M. J., Kordes, S., Garcia-Rosales, F., Kossl, M. and Hechavarria, J. C. (2017).
Processing of Natural Echolocation Sequences in the Inferior Colliculus of Seba's Fruit Eating Bat,
Carollia perspicillata. *Eneuro* **4**.
Beetz, M. J., Kossl, M. and Hechavarria, J. C. (2019). Adaptations in the call emission pattern
of frugivorous bats when orienting under challenging conditions. *Journal of Comparative Physiology*
a-Neuroethology Sensory Neural and Behavioral Physiology **205**, 457-467.
Behr, O. and von Helversen, O. (2004). Bat serenades - complex courtship songs of the sac-
winged bat (*Saccopteryx bilineata*). *Behavioral Ecology and Sociobiology* **56**, 106-115.
Brumm, H. and Zollinger, S. A. (2011). The evolution of the Lombard effect: 100 years of
psychoacoustic research. *Behaviour* **148**, 1173-1198.
Bullock, T. H., Hamstra, R. H. and Scheich, H. (1972). Jamming Avoidance Response of High-
Frequency Electric Fish .1. General Features. *Journal of Comparative Physiology* **77**, 1-22.
Chiu, C., Xian, W. and Moss, C. F. (2008). Flying in silence: Echolocating bats cease vocalizing
to avoid sonar jamming. *Proceedings of the National Academy of Sciences of the United States of*
America **105**, 13116-13121.
Chiu, C., Xian, W. and Moss, C. F. (2009). Adaptive echolocation behavior in bats for the
analysis of auditory scenes. *Journal of Experimental Biology* **212**, 1392-1404.
Corcoran, A. J. and Moss, C. F. (2017). Sensing in a noisy world: lessons from auditory
specialists, echolocating bats. *Journal of Experimental Biology* **220**, 4554-4566.

Cvikel, N., Levin, E., Hurme, E., Borissov, I., Boonman, A., Amichai, E. and Yovel, Y. (2015).
On-board recordings reveal no jamming avoidance in wild bats. *Proceedings of the Royal Society B-*
Biological Sciences **282**.
Eckmeier, D., Geurten, B. R. H., Kress, D., Mertes, M., Kern, R., Egelhaaf, M. and Bischof, H.
J. (2008). Gaze Strategy in the Free Flying Zebra Finch (Taeniopygia guttata). *Plos One* **3**.
Falk, B., Jakobsen, L., Surlykke, A. and Moss, C. F. (2014). Bats coordinate sonar and flight
behavior as they forage in open and cluttered environments. *Journal of Experimental Biology* **217**,
4356-4364.
Fuzessery, Z. M. (1994). Response Selectivity for Multiple Dimensions of Frequency Sweeps
in the Pallid Bat Inferior Colliculus. *Journal of Neurophysiology* **72**, 1061-1079.
Ganguly, K. and Kleinfeld, D. (2004). Goal-directed whisking increases phase-locking
between vibrissa movement and electrical activity in primary sensory cortex in rat. *Proceedings of*
the National Academy of Sciences of the United States of America **101**, 12348-12353.
Gao, L., Balakrishnan, S., He, W. K., Yan, Z. and Müller, R. (2011). Ear Deformations Give
Bats a Physical Mechanism for Fast Adaptation of Ultrasonic Beam Patterns. *Physical Review Letters*
107.
Gillam, E. H. and McCracken, G. F. (2007). Variability in the echolocation of Tadarida
brasiliensis: effects of geography and local acoustic environment. *Animal Behaviour* **74**, 277-286.
Gillam, E. H., Ulanovsky, N. and McCracken, G. F. (2007). Rapid jamming avoidance in
biosonar. *Proc Biol Sci* **274**, 651-660.
Götze, S., Koblitz, J. C., Denzinger, A. and Schnitzler, H. U. (2016). No evidence for spectral
jamming avoidance in echolocation behavior of foraging pipistrelle bats. *Scientific Reports* **6**.
Greiter, W. and Firzlaff, U. (2017). Echo-acoustic flow shapes object representation in
spatially complex acoustic scenes. *Journal of Neurophysiology*, jn 00860 2016.
Griffin, D. R. (1953). Bat Sounds under Natural Conditions, with Evidence for Echolocation of
Insect Prey. *Journal of Experimental Zoology* **123**, 435-465.
Habersetzer, J. (1981). Adaptive Echolocation Sounds in the Bat *Rhinopoma-Hardwickei* - a
Field-Study. *J Comp Physiol* **144**, 559-566.
Hage, S. R., Jiang, T. L., Berquist, S. W., Feng, J. and Metzner, W. (2013). Ambient noise
induces independent shifts in call frequency and amplitude within the Lombard effect in echolocating
bats. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 4063-
4068.
Hechavarría, J. C. and Kössl, M. (2014). Footprints of inhibition in the response of cortical
delay-tuned neurons of bats. *Journal of Neurophysiology* **111**, 1703-1716.
Henson, O. W., Pollak, G. D., Kobler, J. B., Henson, M. M. and Goldman, L. J. (1982).
Cochlear Microphonic Potentials Elicited by Biosonar Signals in Flying Bats, *Pteronotus-P-Parnellii*.
Hearing Research **7**, 127-147.
Hiryu, S., Bates, M. E., Simmons, J. A. and Riquimaroux, H. (2010). FM echolocating bats
shift frequencies to avoid broadcast-echo ambiguity in clutter. *Proc Natl Acad Sci USA* **107**, 7048-
7053.
Ibanez, C., Juste, J., Lopez-Wilchis, R. and Nunez-Garduno, A. (2004). Habitat variation and
jamming avoidance in echolocation calls of the sac-winged bat (*Balantiopteryx plicata*). *Journal of*
Mammalogy **85**, 38-42.
Jarvis, J., Jackson, W. and Smotherman, M. (2013). Groups of bats improve sonar efficiency
through mutual suppression of pulse emissions. *Frontiers in Physiology* **4**.
Jung, K., Kalko, E. K. V. and von Helversen, O. (2007). Echolocation calls in Central American
emballonurid bats: signal design and call frequency alternation. *Journal of Zoology* **272**, 125-137.
Kalko, E. K. V. (1995). Insect Pursuit, Prey Capture and Echolocation in Pipistrelle Bats
(Microchiroptera). *Animal Behaviour* **50**, 861-880.
Kalko, E. K. V. and Schnitzler, H. U. (1989). The Echolocation and Hunting Behavior of
Daubenton Bat, *Myotis-Daubentoni*. *Behavioral Ecology and Sociobiology* **24**, 225-238.

Knowles, J. M., Barchi, J. R., Gaudette, J. E. and Simmons, J. A. (2015). Effective biosonar
echo-to-clutter rejection ratio in a complex dynamic scene. *Journal of the Acoustical Society of*
America **138**, 1090-1101.
Kössl, M., Hechavarría, J. C., Voss, C., Macías, S., Mora, E. C. and Vater, M. (2014). Neural
maps for target range in the auditory cortex of echolocating bats. *Current Opinion in Neurobiology*
24, 68-75.
Kothari, N. B., Wohlgemuth, M. J., Hulgard, K., Surlykke, A. and Moss, C. F. (2014). Timing
matters: sonar call groups facilitate target localization in bats. *Frontiers in Physiology* **5**.
Land, M. F. (2015). Eye movements of vertebrates and their relation to eye form and
function. *Journal of Comparative Physiology a-Neuroethology Sensory Neural and Behavioral*
Physiology **201**, 195-214.
Leonard, M. L. and Fenton, M. B. (1984). Echolocation Calls of *Euderma-Maculatum*
(Vespertilionidae) - Use in Orientation and Communication. *Journal of Mammalogy* **65**, 122-126.
Levin, E., Roll, U., Dolev, A., Yom-Tov, Y. and Kronfeld-Shcor, N. (2013). Bats of a Gender
Flock Together: Sexual Segregation in a Subtropical Bat. *Plos One* **8**.
Lin, Y. and Abaid, N. (2015). Modeling perspectives on echolocation strategies inspired by
bats flying in groups. *Journal of Theoretical Biology* **387**, 46-53.
Luo, J. H., Goerlitz, H. R., Brumm, H. and Wiegrebe, L. (2015). Linking the sender to the
receiver: vocal adjustments by bats to maintain signal detection in noise. *Scientific Reports* **5**.
Macias, S., Luo, J. H. and Moss, C. F. (2018). Natural echolocation sequences evoke echo-
delay selectivity in the auditory midbrain of the FM bat, Eptesicus fuscus. *Journal of Neurophysiology*
120, 1323-1339.
Macias, S., Mora, E. C., Hechavarria, J. C. and Kossl, M. (2016). Echo-level compensation and
delay tuning in the auditory cortex of the mustached bat. *European Journal of Neuroscience* **43**, 1647-
1660.
Masters, W. M. and Jacobs, S. C. (1989). Target Detection and Range Resolution by the Big
Brown Bat (Eptesicus-Fuscus) Using Normal and Time-Reversed Model Echoes. *Journal of*
Comparative Physiology a-Sensory Neural and Behavioral Physiology **166**, 65-73.
Masters, W. M. and Raver, K. A. S. (1996). The degradation of distance discrimination in big
brown bats (Eptesicus fuscus) caused by different interference signals. *Journal of Comparative*
Physiology a-Sensory Neural and Behavioral Physiology **179**, 703-713.
Masters, W. M. and Raver, K. A. S. (2000). Range discrimination by big brown bats (Eptesicus
fuscus) using altered model echoes: Implications for signal processing. *Journal of the Acoustical*
Society of America **107**, 625-637.
Mazar, O. and Yovel, Y. (2019). A spectral jamming avoidance response does not help bats
deal with jamming. *bioRxiv*, 2019.12.16.876086.
Miller, L. A. and Degn, H. J. (1981). The Acoustic Behavior of 4 Species of Vespertilionid Bats
Studied in the Field. *Journal of Comparative Physiology* **142**, 67-74.
Moss, C. F. and Surlykke, A. (2010). Probing the natural scene by echolocation in bats.
Frontiers in Behavioral Neuroscience **4**.
Obrist, M. K. (1995). Flexible Bat Echolocation - the Influence of Individual, Habitat and
Conspecifics on Sonar Signal-Design. *Behavioral Ecology and Sociobiology* **36**, 207-219.
Park, T. J. and Pollak, G. D. (1993). Gaba Shapes a Topographic Organization of Response
Latency in the Moustache Bats Inferior Colliculus. *Journal of Neuroscience* **13**, 5172-5187.
Parsons, K. N., Jones, G. and Greenaway, F. (2003). Swarming activity of temperate zone
microchiropteran bats: effects of season, time of night and weather conditions. *Journal of Zoology*
261, 257-264.
Petrites, A. E., Eng, O. S., Mowlds, D. S., Simmons, J. A. and DeLong, C. M. (2009). Interpulse
interval modulation by echolocating big brown bats (*Eptesicus fuscus*) in different densities of
obstacle clutter. *J Comp Physiol A* **195**, 603-617.
Ratcliffe, J. M., ter Hofstede, H. M., Avila-Flores, R., Fenton, M. B., McCracken, G. F.,
Biscardi, S., Blasko, J., Gillam, E., Orprecio, J. and Spanjer, G. (2004). Conspecifics influence call

design in the Brazilian free-tailed bat, Tadarida brasiliensis. *Canadian Journal of Zoology-Revue Canadienne De Zoologie* **82**, 966-971.

Ribak, G., Egge, A. R. and Swallow, J. G. (2009). Saccadic head rotations during walking in the stalk-eyed fly (Cyrtodiopsis dalmanni). *Proceedings of the Royal Society B-Biological Sciences* **276**, 1643-1649.

Roverud, R. C. and Grinnell, A. D. (1985a). Discrimination Performance and Echolocation Signal Integration Requirements for Target Detection and Distance Determination in the CF FM Bat, *Noctilio-Albiventris*. *J Comp Physiol A* **156**, 447-456.

Roverud, R. C. and Grinnell, A. D. (1985b). Echolocation Sound Features Processed to Provide Distance Information in the Cf Fm Bat, Noctilio-Albiventris - Evidence for a Gated Time Window Utilizing Both Cf and Fm Components. *Journal of Comparative Physiology a-Sensory Neural and Behavioral Physiology* **156**, 457-469.

Sanderson, M. I. and Simmons, J. A. (2005). Target representation of naturalistic echolocation sequences in single unit responses from the inferior colliculus of big brown bats. *Journal of the Acoustical Society of America* **118**, 3352-3361.

Schnitzler, H. U., Kalko, E., Miller, L. and Surlykke, A. (1987). The Echolocation and Hunting Behavior of the Bat, Pipistrellus-Kuhli. *Journal of Comparative Physiology a-Sensory Neural and Behavioral Physiology* **161**, 267-274.

Schroeder, C. E., Wilson, D. A., Radman, T., Scharfman, H. and Lakatos, P. (2010). Dynamics of Active Sensing and perceptual selection. *Current Opinion in Neurobiology* **20**, 172-176.

Simmons, J. A. (2012). Bats use a neuronally implemented computational acoustic model to form sonar images. *Current Opinion in Neurobiology* **22**, 311-319.

Simmons, J. A. (2017). Noise interference with echo delay discrimination in bat biosonar. *Journal of the Acoustical Society of America* **142**, 2942-2952.

Simmons, J. A., Fenton, M. B. and Ofarrell, M. J. (1979). Echolocation and Pursuit of Prey by Bats. *Science* **203**, 16-21.

Simmons, J. A., Howell, D. J. and Suga, N. (1975). Information-Content of Bat Sonar Echoes. *American Scientist* **63**, 204-215.

Simmons, J. A., Lavender, W. A., Lavender, B. A., Childs, J. E., Hulebak, K., Rigden, M. R., Sherman, J., Woolman, B. and Ofarrell, M. J. (1978). Echolocation by Free-Tailed Bats (Tadarida). *Journal of Comparative Physiology* **125**, 291-299.

Suga, N. and Manabe, T. (1982). Neural Basis of Amplitude-Spectrum Representation in Auditory-Cortex of the Mustached Bat. *Journal of Neurophysiology* **47**, 225-255.

Suga, N., O'Neill, W. E., Kujirai, K. and Manabe, T. (1983). Specificity of Combination-Sensitive Neurons for Processing of Complex Biosonar Signals in Auditory-Cortex of the Mustached Bat. *Journal of Neurophysiology* **49**, 1573-1626.

Surlykke, A. (1992). Target Ranging and the Role of Time-Frequency Structure of Synthetic Echoes in Big Brown Bats, Eptesicus-Fuscus. *Journal of Comparative Physiology a-Sensory Neural and Behavioral Physiology* **170**, 83-92.

Surlykke, A. and Moss, C. F. (2000). Echolocation behavior of big brown bats, Eptesicus fuscus, in the field and the laboratory. *Journal of the Acoustical Society of America* **108**, 2419-2429.

Takahashi, E., Hyomoto, K., Riquimaroux, H., Watanabe, Y., Ohta, T. and Hiryu, S. (2014). Adaptive changes in echolocation sounds by *Pipistrellus abramus* in response to artificial jamming sounds. *Journal of Experimental Biology* **217**, 2885-2891.

Tarsitano, M. S. and Andrew, R. (1999). Scanning and route selection in the jumping spider Portia labiata. *Animal Behaviour* **58**, 255-265.

Thies, W., Kalko, E. K. V. and Schnitzler, H. U. (1998). The roles of echolocation and olfaction in two Neotropical fruit-eating bats, *Carollia perspicillata* and *C-castanea*, feeding on Piper. *Behavioral Ecology and Sociobiology* **42**, 397-409.

Towal, R. B. and Hartmann, M. J. (2006). Right-left asymmetries in the whisking behavior of rats anticipate head movements. *Journal of Neuroscience* **26**, 8838-8846.

Tressler, J. and Smotherman, M. S. (2009). Context-dependent effects of noise on
echolocation pulse characteristics in free-tailed bats. *J Comp Physiol A* **195**, 923-934.
Ulanovsky, N., Fenton, M. B., Tsoar, A. and Korine, C. (2004). Dynamics of jamming
avoidance in echolocating bats. *Proceedings of the Royal Society B-Biological Sciences* **271**, 1467-
1475.
Ulanovsky, N. and Moss, C. F. (2008). What the bat's voice tells the bat's brain. *Proc Natl*
Acad Sci USA **105**, 8491-8498.
von Stebut, B. and Schmidt, S. (2001). Frequency discrimination threshold at search call
frequencies in the echolocating bat, Eptesicus fuscus. *Journal of Comparative Physiology a-Sensory*
Neural and Behavioral Physiology **187**, 287-291.
Watanabe, A. and Takeda, K. (1963). The Change of Discharge Frequency by A.C. Stimulus in
a Weak Electric Fish. *Journal of Experimental Biology* **40**, 57-+.
Weineck, K., Garcia-Rosales, F. and Hechavarria, J. C. (2020). Neural oscillations in the
fronto-striatal network predict vocal output in bats. *Plos Biology* **18**.
Wheeler, A. R., Fulton, K. A., Gaudette, J. E., Simmons, R. A., Matsuo, I. and Simmons, J. A.
(2016). Echolocating Big Brown Bats, *Eptesicus fuscus*, Modulate Pulse Intervals to Overcome Range
Ambiguity in Cluttered Surroundings. *Frontiers in Behavioral Neuroscience* **10**.
Wohlgemuth, M. J., Kothari, N. B. and Moss, C. F. (2016a). Action Enhances Acoustic Cues
for 3-D Target Localization by Echolocating Bats. *Plos Biology* **14**.
Wohlgemuth, M. J., Luo, J. and Moss, C. F. (2016b). Three-dimensional auditory localization
in the echolocating bat. *Current Opinion in Neurobiology* **41**, 78-86.
Yang, L. C., Pollak, G. D. and Resler, C. (1992). Gabaergic Circuits Sharpen Tuning Curves and
Modify Response Properties in the Moustache Bat Inferior Colliculus. *Journal of Neurophysiology* **68**,
1760-1774.
Yovel, Y., Melcon, M. L., Franz, M. O., Denzinger, A. and Schnitzler, H. U. (2009). The Voice
of Bats: How Greater Mouse-eared Bats Recognize Individuals Based on Their Echolocation Calls. *Plos*
Computational Biology **5**.