Purging of highly deleterious mutations through severe bottlenecks in Alpine ibex

Christine Grossen[1,*], Frederic Guillaume[1], Lukas F. Keller[1,2], Daniel Croll[3,*]

[1]Department of Evolutionary Biology and Environmental Studies, University of Zurich, CH-8057 Zurich, Switzerland

[2]Zoological Museum, University of Zurich, Karl-Schmid-Strasse 4, CH-8006 Zurich, Switzerland

[3]Laboratory of Evolutionary Genetics, Institute of Biology, University of Neuchâtel, CH-2000 Neuchâtel, Switzerland; 0000-0002-2072-380X

[*]Authors for correspondence: christine.grossen@uzh.ch, daniel.croll@unine.ch

Author contributions:

Conception and design of study: CG, LFK, DC

Acquisition and analysis of data: CG

Interpretation of data: CG, FG, LFK, DC

Funding: CG, LFK

Wrote the manuscript with input from the other authors: CG, DC

Keywords: population genomics, deleterious mutations, purging, Alpine ibex

Abstract

Human activity caused dramatic population declines in many wild species. The resulting bottlenecks have a profound impact on the genetic makeup of a species with unknown consequences for health. A key genetic factor for species survival is the evolution of deleterious mutation load, but how bottleneck strength and mutation load interact lacks empirical evidence. Here, we take advantage of the exceptionally well-characterized population bottlenecks of the once nearly extinct Alpine ibex. The species survived one of the most dramatic bottlenecks known for successfully restored species. We analyze 60 complete genomes of six ibex species and the domestic goat. We show that historic bottlenecks rather than the current conservation status predict levels of genome-wide variation. By retracing the recolonization of the Alps by Alpine ibex, we find genomic evidence of concurrent purging of highly deleterious mutations but accumulation of mildly deleterious mutations. This demonstrates how human-induced severe bottlenecks caused both relaxed selection and purging, thus reshaping the landscape of deleterious mutation load. Our findings also highlight that even populations of ~1000 individuals can accumulate mildly deleterious mutations. Hence, conservation efforts should focus on preventing population declines below such levels to ensure long-term survival of species.

Main text

Climate change and pressure from human activities such as hunting caused profound changes in the population and demographic structure of many species [1]. This is because extinction events and subsequent recolonization severely alter the genetic makeup [2]. The demographic changes have important consequences for wildlife management and the conservation of endangered species [2] including raising the risk of genetic disorders [e.g 3-9]. However, nearly all plant and animal populations including humans suffered from temporary reductions in population size – so-called bottlenecks. Bottlenecks increase genetic drift and inbreeding, which leads to a loss of genetic variation, reduces the efficacy of natural selection, and increases the expression of deleterious recessive mutations [10-12]. The expression of recessive mutations under inbreeding creates the potential for selection to act against these mutations. This process known as purging reduces the frequency of deleterious mutations depending on the degree of dominance and the magnitude of the deleterious effects [13]. Because purging depends on levels of inbreeding, bottlenecks tend to purge highly deleterious, recessive mutations unless population sizes are extremely low [13-15]. Bottlenecks also increase genetic drift and reduce the efficacy of selection [16]. This allows mildly deleterious mutations to drift to substantially higher frequencies [4,6,17]. Hence, bottlenecks generate complex dynamics of deleterious mutation frequencies due to the independent effects of purging and reduced selection efficacy [13-15,18,19].

A major gap in our understanding is how reduced selection efficacy and purging jointly determine the mutation load in wild populations. Theoretical predictions are well established [13,15,18,20] but empirical evidence is conflicting [9,21,22] including for humans (see e.g. [23-31]. Previous research used changes in fitness to infer possible purging events, [7,20,32-34], but changes in fitness can result from causes unrelated to purging [12,35]. Direct evidence for purging exists only for isolated mountain gorilla populations that split off larger lowland populations ~20'000 years ago [36]. However, it remains unknown how recent, dramatic bottleneck events on the scale caused by human activity impacts levels of deleterious mutations in the wild. Here, we take advantage of exceptionally well characterized repeated bottlenecks during the reintroduction of the once near-extinct Alpine ibex to retrace the fate of

deleterious mutations. Alpine ibex were reduced to ~100 individuals in the 19th century in a single

population in the Gran Paradiso region of Northern Italy [37]. In less than a century, a census size of ca.

50'000 individuals has been re-established across the Alps. Thus, the population bottleneck of Alpine

ibex is among the most dramatic recorded for any successfully restored species. Most extant

populations experienced two to four, well-recorded bottlenecks leaving strong footprints of low

genetic diversity [38,39].

Figure 1

Figure 1: Genetic diversity and deleterious mutation load of ibex species. A) Geographical distribution and
IUCN conservation status of ibex and wild goat species (LC: Least concern, V: Vulnerable, NT: Near threatened
[40]). Sample sizes: Alpine ibex (*C. ibex*): N=29, Iberian ibex (*C. pyrenaica*): N=4, Bezoar (*C. aegagrus*): N=6,
Siberian ibex (*C. sibirica*): N=2, Markhor (*C. falconeri*): N=1, Nubian ibex (*C. nubiana*): N=2. B) Maximum
likelihood phylogenetic analyses, C) nucleotide diversity, D) proportion of the genome with runs of
homozygosity (ROH) longer than 2.5 Mb and E) percentage of polymorphic sites within species that segregate
highly deleterious mutations. Confidence intervals are based on downsampling to four individuals matching the
sample size of Iberian ibex (100 replicates).

We analyzed 60 genomes covering seven species including the Alpine ibex (*C. ibex*), five additional

wild goats and the domestic goat. Siberian ibex have the largest population size (~200'000), followed

by Alpine and Iberian ibex (both ~50'000), Bezoar (~25'000), Markhor (~6000) and Nubian ibex

(~2500) [40]. We found that Alpine ibex together with Iberian ibex and Markhor (*C. falconeri*) have

exceptionally low genome-wide variation compared to closely related species (Figure 1, Figure S1).

All three species either underwent severe bottlenecks in the past century or are currently threatened

(Figure 1C, Figure S2, Tables S1 and S2). Historic records indicate that Alpine ibex suffered a

bottleneck of ~100 individuals at the end of the 19th century and Iberian ibex a bottleneck of ~1000

individuals [41] (Table S1). In contrast, genome-wide diversity was highest in Siberian ibex (*C.*

sibirica), which have large and relatively well-connected populations [42]. The genomes of some Alpine

ibex, the Markhor and some domestic goat individuals contained more than 20% genome-wide runs

of homozygosity (ROH) (Figure 1D, Figure S3-S5, Table S3) [43]. Overall, there was clear genomic

evidence that the near extinction and recovery of the Alpine ibex resulted in substantial genetic drift

and inbreeding.

We analyzed all *Capra* genomes for evidence of segregating deleterious mutations (Figure S6). We

restricted our analyses to coding sequences with evidence for transcriptional activity in Alpine ibex

organs and high evolutionary conservation (*i.e.* GERP, see methods) [44] yielding a total of 370'853

SNPs (Table S4). We found that across all seven *Capra* species 0.17% of these SNPs carried a highly

deleterious variant with the majority of the highly deleterious variants incurring a stop-gain mutation,

a further 19.1% carried a moderate impact variant and 33.1% carried a low impact variant (Table S4).

We used seven additional mutation impact scoring approaches (see methods). The proportion of

highly deleterious variants segregating within species was inversely correlated with nucleotide

diversity (Figure 1C and E, Figure S7A-D; Pearson, df = 5, r = -0.86, p = 0.012). Hence, species with

the smallest populations or the most severe population size reductions (*i.e.* Alpine ibex, Iberian ibex

and Markhor) showed an accumulation of deleterious mutations relative to closely related species.

Both Alpine and Iberian ibex experienced severe bottlenecks due to overhunting and habitat

fragmentation. We first analyzed evidence for purifying selection using allele frequency spectra. We

focused only on derived sites that were polymorphic in at least one of the two sister species (Figure

2A, S8). We found that frequency distributions of high and moderate impact mutations in Alpine ibex

were downwards shifted compared to modifier (*i.e.* neutral) mutations, which strongly suggests

purifying selection against highly deleterious mutations (Figure 2B, D). Short indels ($\leq$ 10 bp) in

coding sequences revealed a similar downward shift (Figure S9). We found no comparable frequency

shifts in Iberian ibex (Figure 2B). This is consistent with purifying selection acting more efficiently

against highly deleterious mutations in Alpine ibex compared to Iberian ibex. We repeated the

analyses of the site frequency spectra also for three alternative scoring systems (GERP, phyloP and

phastCons) reporting phylogenetic conservation. We found for all three scores an excess in Alpine

ibex of the most deleterious mutation category (Figure S10).

Figure 2

Figure 2: Deleterious mutations segregating in Alpine and Iberian ibex. A) Population sampling locations
of Iberian ibex (left, grey circles) and Alpine ibex (right, colored circles). Each filled circle represents a
population. Circles with a black outline indicate the first three reintroduced populations in Switzerland that were
used for all subsequent population reintroductions of Alpine ibex. Colors associate founder and descendant
populations (see also Figure 3A). (B) Site frequency spectrum for neutral (modifier), mildly (moderate impact)
and highly deleterious (high impact) mutations for Alpine ibex. Alternative allele counts >10 are shown as mean
counts per interval. (C) Site frequency spectrum for Iberian ibex. (D) Site frequency spectrum for Alpine ibex
downsampled to four individuals matching the sample size of Iberian ibex (100 replicates). See Figure S11 for
additional downsampling and jack-knifing analyses of data presented in B-D. E) Rxy analysis contrasting
Iberian with Alpine ibex across the spectrum of impact categories. $Rxy < 1$ indicates a relative frequency deficit
of the corresponding category in Alpine ibex compared to Iberian ibex. Rxy distributions are based on jack-
knifing across chromosomes. All pairwise comparisons were significant except low vs. moderate (Tukey test, p
< 0.0001). F) Individual homozygote counts per impact category for Iberian (light green) and Alpine ibex (dark
green).

To test whether Alpine ibex showed evidence for purging of deleterious mutations, we calculated the

relative number of derived alleles Rxy [29] for the different categories of mutations (Figure 2D). We

used a random set of intergenic SNPs for standardization, which makes Rxy robust against sampling

effects and population substructure [29]. Low and moderate impact mutations (*i.e.* mildly deleterious

mutations) showed a minor excess in Alpine ibex compared to Iberian ibex, indicating a higher load

in Alpine ibex. In contrast, we found that highly deleterious mutations were strongly reduced in

Alpine ibex compared to Iberian ibex (Tukey test, $p < 0.0001$, Figure 2E). Strikingly, the proportion

of SNPs across the genome segregating a highly deleterious mutation is higher in Alpine ibex (Figure

1E), but Rxy shows that highly deleterious mutations have a pronounced downwards allele frequency

shift in Alpine ibex compared to Iberian ibex. Furthermore, the number of homozygous sites with

highly deleterious mutations per individual was significantly lower in Alpine ibex than Iberian ibex (t

test, $p = 0.015$, Figure 2E). Individual allele counts at highly deleterious sites were also significantly

lower in Alpine ibex compared to Iberian ibex (t test, $p = 0.003$). We assessed the robustness of the

Rxy analyses using four additional mutation scoring methods (*i.e.* SIFT, REVEL, CADD and

VEST3). We found that the highest impact category had a consistent deficit in Alpine ibex compared

to Iberian ibex (Figure S12).. Together, this shows that highly deleterious mutations were

substantially purged in Alpine ibex. We also found evidence for the accumulation of mildly

deleterious mutations through genetic drift in Alpine ibex.

Consistent with the fact that all extant Alpine ibex originate from the Gran Paradiso, this population

occupies the center of a principal component analysis (Figure 3A-B, Figure S13A-B; [39]). The first

populations re-established in the Alps were clearly differentiated from the Gran Paradiso source

population and showed reduced nucleotide diversity due to reintroduction bottlenecks [38] (Figure 3A,

C). These initial three reintroduced populations were used to establish additional populations raising

the total number of experienced bottlenecks to 3 or 4. The additional bottlenecks led to further loss of

nucleotide diversity and genetic drift, as indicated by the increasing spread in the principal component

analysis (Figure 3A-C). An exceptional case constitutes the Alpi Marittime population, which was

established through the translocation of 25 Gran Paradiso individuals of which only six successfully

reproduced [45]. As expected from such an extreme bottleneck, Alpi Marittime showed strong genetic

differentiation from all other Alpine ibex populations and highly reduced nucleotide diversity (Figure

3B-C; [46]). To compare the strength of drift experienced by different populations, we estimated long-

term effective population sizes. For this, we used detailed demographic records spanning the near

century since the populations were established [47,48]. We found that nucleotide diversity decreased with

smaller long-term population size (Spearman's rank correlation, rho = 0.93, p = 0.007, Figure 3C). We

found the same trend for the individual number of heterozygous sites per kilobase (Figure S14).

Populations with the lowest harmonic mean population sizes also showed the highest levels of

inbreeding. Genomes from the Gran Paradiso source population generally showed the lowest

proportions of the genome affected by ROH, while reintroduced populations of lowest effective

population size had the highest proportions of the genome affected by ROH (Pearson, df = 19, r = -

0.70, p = 0.0004, Figure 3D and Figure S3).

Figure 3

Figure 3: Genomic consequences of the Alpine ibex recolonization. A) The recolonization history and population pedigree of Alpine ibex. Locations include also zoos and the population Pilatus (pi), which was not sampled for this study. am: Alpi Marittime, gp: Gran Paradiso; ih: Zoo Interlaken Harder; al: Albris; bo: Bire Oɩschinen; br: Brienzer Rothorn; ob: Oberbauenstock; pl: Pleureur; rh: Rheinwald; wh: Weisshorn; pi: Pilatus; pp: Wildpark Peter and Paul. The grey circle represents a population that was founded from more than one population. Figure elements were modified from Biebach and Keller (2009) with permission. B) Principal component analysis of all Alpine ibex individuals included in the study. C) Nucleotide diversity per population. D) Proportion of the genome within runs of homozygosity (ROH) longer than 2.5 Mb. E) *Rxy* analysis contrasting the strongly bottlenecked Alpi Marittime population with all other Alpine ibex populations across the spectrum of impact categories. *Rxy* < 1 indicates a relative frequency deficit of the corresponding category in the Alpi Marittime population. *Rxy* distributions are based on jack-knifing across chromosomes. Circles with a black outline indicate the first three reintroduced populations in Switzerland that were used for all subsequent population reintroductions of Alpine ibex. Colors associate founder and descendant populations.

Bottlenecks affect the landscape of deleterious mutations by randomly increasing or decreasing allele frequencies at individual loci. We find that individuals from populations that underwent stronger bottlenecks carry significantly more homozygotes for nearly neutral and mildly deleterious mutations

(*i.e.* modifier, low and moderate impact mutations; Figure 4A). In contrast, individuals showed no

meaningful difference in the number of homozygotes for highly deleterious (*i.e.* high impact)

mutations across populations. The stability in the number of homozygotes for highly deleterious

mutations through successive bottlenecks despite a step-wise increase in the number of homozygotes

for weaker impact mutations, supports that purging occurred over the course of the Alpine ibex

reintroductions. We repeated the analyses using an alternative scoring of mutations based on the

phylogenetic conservation of the region in which the mutations were found (*i.e.* Genomic

Evolutionary Rate Profiling; GERP). The number of homozygotes for mutations in highly conserved

regions showed a slight upwards trend still indicating purifying but not necessarily purging for this

category of mutations (Figure S15). This suggests that the mutational impact (*e.g.* premature stop

codons) rather than degree of conservation predicts whether purging is likely to occur. This suggests

also that mean fitness should be more directly assessed using *e.g.* simulation datasets (see below).

Because the above findings are contingent on a model where deleterious mutations are recessive, we

also analyzed the total number of derived alleles per individual. We find a consistent but less

pronounced increase in total number of derived alleles per individual for nearly neutral and mildly

deleterious mutations (Figure 4B). In contrast, the total number of derived alleles for highly

deleterious mutations did not correlate with the strength of bottleneck and was lowest in the most

severely bottlenecked Alpi Marittime population (Figure 4B), suggesting that the most deleterious

mutations were purged in this population. The *Rxy* statistics showed a corresponding strong deficit in

the Alpi Marittime population (Figure 3E). We repeated the *Rxy* analyses using four additional

mutation scoring methods (*i.e.* SIFT, REVEL, CADD and VEST3) and found a consistent deficit of

the highest impact category in the Alpi Marittime (Figure S12). Overall, we find evidence for more

purging in the most bottlenecked Alpine ibex population.

Figure 4

Figure 4: Impact of the recolonization on the individual mutation load. (A) Homozygote counts and (B) allele counts per individual for each Alpine ibex population. The schematic between A and B indicates the harmonic mean of the census size of each population, which is inversely correlated with the strength of drift. [*] Estimated numbers. Colors associate founder and descendant populations (see also Figure 3A).

We analyzed the impact of highly deleterious mutations by predicting the protein truncation using homology-based inferences. Focusing on mutations segregating in Alpine ibex, we found that nearly all mutations disrupted conserved protein family (PFAM) domains encoded by the affected genes (Figure 5). This shows that highly deleterious mutations not only are altering the length of open reading frames but that evolutionarily conserved protein domains are affected by the mutations.

Figure 5: Homology-based inference of the impact of highly deleterious mutations. The localization of protein family (PFAM) domains are highlighted in dark. Red dots indicate the relative position of a highly deleterious mutation segregating in Alpine ibex. The frequencies of highly deleterious mutations are summarized for Iberian ibex and three groups of Alpine ibex. Allele frequencies are shown as boxplots per group representing the outcome of downsampling to the smallest group (Alpi Marittime, *n* = 3) based on 100 replicates.

We further analyzed the impact of bottlenecks on different mutation classes using an individual-based

forward simulation model parametrized with the demographic record [49] (Figure 6A). The model

realistically reproduced the reintroduction history and populations were parametrized with the actual

founder size (Figure 6A, S16, Table S1). We used *Rxy* to analyze the evolution of deleterious

mutation frequencies through the reintroduction bottlenecks. The simulations showed a deficit of

highly deleterious mutations, but an increase of mildly deleterious mutations after the reintroduction

bottlenecks (Figure 6B). This is consistent with our empirical evidence for purging during the species

bottleneck (Figure 2E). We computed genetic load defined as the mean individual fitness in females

and found an increase in Alpine ibex following the species bottleneck (Figure S17). Hence, the

accumulation of mildly deleterious mutations was reducing overall fitness despite purging. The

simulations also supported purging at the level of individual populations as found in the extremely

bottlenecked Alpi Marittime population (Figure 3E). The number of derived mildly deleterious

homozygotes increased with the strength of drift experienced by individual populations (Figure 6C,

Figures S18-S21). In contrast, the median number of homozygote counts for high impact mutations

were lower for Alpi Marittime than Gran Paradiso but not statistically significant (Figure 6C, Mann-

Whitney U, $p = 0.39$). The highly deleterious allele counts were significantly lower for Alpi Marittime

compared to Gran Paradiso (Figure S19, $p = 0.001$). We also analyzed Rxy for the simulation data and

found that high impact mutations were indeed relatively less frequent in Alpi Marittime compared to

all other Alpine ibex populations (Figure 6D) and compared to Gran Paradiso (Figure S22). Overall,

the realistically parametrized model recapitulated all major empirical findings of mutation

accumulation and purging across bottlenecks.

Figure 6

Figure 6: Individual-based simulations of the reintroduction history of Alpine ibex. A) The demographic model was parametrized using census data and historical records. Bold numbers show effective population sizes. Numbers not in bold indicate the number of individuals released to found each population. If a population was established from two source populations, the individual numbers are separated by commas. *) Upwards adjusted harmonic means of the census size (historical records were ih=16, Zoo Interlaken Harder, and pp = 20, Wildpark Peter and Paul). The adjustment was necessary to prevent extinction of zoo populations. **) Census numbers were estimated based on historical records of the population but no long-term data census data was available. B) Relative frequency comparison (*Rxy*) of Alpine ibex just before and after the species bottleneck and recolonization. C) Individual homozygote counts per impact category. Boxplots summarize 100 population means across simulation replicates. Colors associate founder and descendant populations (see also Figure 3A). D) *Rxy* analysis contrasting the strongly bottlenecked Alpi Marittime population with all other Alpine ibex populations across the spectrum of impact categories.

Ibex species with recently reduced population sizes accumulated deleterious mutations compared to

closely related species. This accumulation was particularly pronounced in the Iberian ibex that

experienced a severe bottleneck and Alpine ibex that went nearly extinct. We show that even though

Alpine ibex carry an overall higher mutation burden than related species, the strong bottlenecks

imposed by the reintroduction events purged highly deleterious mutations. Importantly, purging was

only effective against highly deleterious mutations. Mildly deleterious mutations actually

accumulated through the reintroductions. Hence, the overall number of deleterious mutations

increased with bottleneck strength. This is consistent with the finding that population-level

inbreeding, which is a strong indicator of past bottlenecks, is correlated with lower population growth

rates in Alpine ibex [50].

Empirical evidence for purging in the wild is scarce [7,20,36]. Here, we show that a few dozen

generations were sufficient to reduce the burden of highly deleterious mutations. Purging may occur

widely in populations undergoing severe bottlenecks contingent on populations surviving the

consequences of inbreeding depression. Failure to purge under extreme bottlenecks can have severely

deleterious consequences for wild populations such as shown for the Isle Royal wolves (Robinson et

al. 2019). Our empirical results from the Alpine ibex reintroduction are in line with theoretical

predictions that populations with an effective size below 100 individuals can accumulate a substantial

burden of mildly deleterious mutations within a relatively short time resulting in increased extinction

risks [5]. The burden of deleterious mutations evident in Iberian ibex supports the notion that even

population sizes of ~1000 still accumulate mildly deleterious mutations. High loads of deleterious

mutations have been shown to increase the extinction risk of a species [5]. Thus, conservation efforts

aimed at keeping effective population sizes above a minimum of 1000 individuals [2] are critical for the

long-term survival of managed species.

Methods

Genomic data acquisition

DNA samples from 29 Alpine ibex, 4 Iberian ibex, 2 Nubian ibex, 2 Siberian ibex and 1 Markhor
individuals were sequenced on an Illumina Hiseq2500 or Hiseq4000 to a depth of 15-38 (median of
17). Table S2 specifies individual sampling locations. Libraries were produced using the TruSeq
DNA Nano kit. Illumina sequencing data of 6 Bezoar and 16 domestic goat (coverage 6x – 14x,
median 12x) were generated by the NextGen Consortium (https://nextgen.epfl.ch). The corresponding
raw data was downloaded from the EBI Short Read Archive: ftp://ftp.sra.ebi.ac.uk/vol1/fastq/.

Read alignment and variant calling

Trimmomatic v.0.36 [51] was used for quality and adapter trimming before reads were mapped to the
domestic goat reference genome (version CHIR1, [52]) using Bowtie2 v.2.2.5 [53]. MarkDuplicates from
Picard (http://broadinstitute.github.io/picard, v.1.130) was used to mark duplicates. Genotype calling
was performed using HaplotypeCaller and GenotypeGVCF (GATK, v.3.6 [54,55]). VariantFiltration of
GATK was used to remove single nucleotide polymorphisms (SNP) if: QD <2.0, FS > 40.0, SOR >
5.0, MQ < 20.0, -3.0 > MQRandkSum > 3.0, -3.0 > ReadPosRankSum > 3.0 and AN < 62 (80% of all
Alpine ibex individuals). Indels up to 10 bp were also retained and filtered using the same filters and
filter parameters, except for not including the filter MQRankSum, because this measure is more likely
to be biased for indels of several base pairs. Filtering parameters were chosen based on genome-wide
quality statistics distributions (see Figures S23 – S40). Variant positions were independently validated
by using the SNP caller Freebayes (v1.0.2-33-gdbb6160 [56]) with the following settings: --no-complex
--use-best-n-alleles 6 --min-base-quality 3 --min-mapping-quality 20 --no-population-priors --hwe-
priors-off.

To ensure high-quality SNPs, we only retained SNPs that were called and passed filtering using
GATK, and that were confirmed by Freebayes. Overall, 97.5 % of all high-quality GATK SNP calls
were confirmed by Freebayes. This percentage was slightly lower for chromosome X (96.7%) and
unplaced scaffolds (95.2%). We tested whether the independent SNP calls of GATK and Freebayes

were concordant and we could validate 99.6% of the biallelic SNPs. We retained genotypes called by

GATK and kept SNPs with a minimum genotyping rate of 90% for all further analysis. The total

number of SNPs detected was 59.5 million among all species. Per species, the number of SNPs

ranged from 21.9 million in the domestic goat (N=16) to 2.0 million in Markhor (N=1, Table S2).

RNA-seq data generation

Tissue samples of a freshly harvested Alpine ibex female were immediately conserved in RNA*later*

(QIAGEN) in the field and stored at -80°C until extraction. The following ten organs were sampled:

retina/uvea, skin, heart, lung, lymph, bladder, ovary, kidney, liver and spleen. RNA was extracted

using the AllPrep DNA/RNA Mini Kit from Qiagen following the manufacturer's protocol.

Homogenization of the samples was performed using a Retsch bead beater (Retsch GmbH) in RLT

plus buffer (Qiagen). RNA was enriched using a PolyA enrichment protocol implemented in the

TruSeq RNA library preparation kit. Illumina sequencing libraries were produced using the Truseq

RNA stranded kit. Sequencing was performed on two lanes of an Illumina Hiseq4000.

Genetic diversity and runs of homozygosity

Genetic diversity measured as individual number of heterozygous sites and nucleotide diversity were

computed using vcftools [57]. Runs of homozygosity were called using BCFtools/RoH [58], an extension

of the software package BCFtools, v.1.3.1. BCFtools/RoH uses a hidden Markov model to detect

segments of autozygosity from next generation sequencing data. Due to the lack of a detailed linkage

map, we used physical distance as a proxy for recombination rates with the option -M and assuming

1.2cM/Mb following sheep recombination rates [59]. Smaller values for -M led to slightly longer ROH

(Figures S3 –S5). Because of small per population sample size, we decided to fix the alternative allele

frequency (option --AF-dflt) to 0.4. Estimates for the population with the largest sample size (Gran

Paradiso, N=7) were very similar if actual population frequencies (option --AF-estimate sp) were used

(Figures S4 and S5). Option --viterbi-training was used to estimate transition probabilities before

running the HMM. Running the analysis without the option --viterbi-training led to less but longer

ROH (Figures S3-S5). ROH were also estimated using PLINK (v1.90b5, https://www.cog-

genomics.org/plink/) with the following settings: --homozyg-window-het 2,

--homozyg-window-missing 5, --homozyg-snp 100, --homozyg-kb 500, --homozyg-density 10, --

homozyg-gap 100, --homozyg-window-threshold .0. ROH estimates based on PLINK were overall

slightly lower but the qualitative trends hold among species and population (Figures S41 and S42).

Identification of high-confidence deleterious mutations

Three lines of evidence were used to identify high-confidence deleterious mutations. First, variants

leading to a functional change are candidates for deleterious mutations. We used snpEff [60] v.4.3 for

the functional annotation of each variant. The annotation file ref_CHIR_1.0_top_level.gff3 was

downloaded from: ftp://ftp.ncbi.nlm.nih.gov/genomes/Capra_hircus/GFF and then converted to gtf

using gffread. Option -V was used to discard any mRNAs with CDS having in-frame stop codons.

SnpEff predicts the effects of genetic variants (e.g. stop-gain variants) and assesses the expected

impact. The following categories were retrieved: high (e.g. stop-gain or frameshift variant), moderate

(e.g missense variant, in-frame deletion), low (e.g. synonymous variant) and modifier (e.g. exon

variant, downstream gene variant). In the case of overlapping transcripts for the same variant, we used

the primary transcript for further analysis. A total of 49.0 % of all detected SNPs were located in

intergenic regions, 43.2 % in introns, 6.5 % down- and upstream of genes. A total of 0.7% of variants

were within CDS, of which ~60% were synonymous and ~40% were missense variants. Overall,

0.002 % were stop-gain mutations.

Protein sequences were annotated using InterProScan v.5.33 by identifying conserved protein family

(PFAM) domains [61].

Second, we assessed the severity of a variant by its phylogenetic conservation score. A non-

synonymous variant is more likely to be deleterious if it occurs in a conserved region of the genome.

We used GERP conservation scores, which are calculated as the number of substitutions observed

minus the number of substitutions expected from the species tree under a neutral model. We

downloaded GERP scores (accessed from http://mendel.stanford.edu/SidowLab/downloads/gerp),

which have been computed for the human reference genome version hg19. The alignment was based

on 35 mammal species but did not include the domestic goat (see https://genome.ucsc.edu/cgi-

bin/hgTrackUi?db=hg19&g=allHg19RS_BW for more information). Exclusion of the focal species

domestic goat is recommended for the computation of conservation scores, as the inclusion of the

reference genome may lead to biases [62]. In order to remap the GERP scores associated to hg19

positions to the domestic goat reference genome positions, we used liftOver

(hgdownload.cse.ucsc.edu, v.287) and the chain file downloaded from hgdownload-

test.cse.ucsc.edu/goldenPath/capHir1.

Third, we ascertained support for gene models annotated in the domestic goat genome with

expression analyses of Alpine ibex tissue samples. We included expression data from 10 organs of an

Alpine ibex female (see RNA-seq data section above) to assess expression levels of each gene model.

Quality filtering of the raw data was performed using Trimmomatic [51] v.0.36. Hisat2 [63] v.2.0.5 was

used to map the reads of each organ to the domestic goat reference genome. The mapping was run

with option --rna-strandness RF (stranded library) and supported by including a file with known splice

sites (option --known-splicesite-infile). The input file was produced using the script

hisat2_extract_splice_sites.py (part of hisat2 package) from the same gtf file as the one used for the

snpEff analyis (see above). For each organ, featureCounts [64] (subread-1.5.1) was used to count reads

per each exon using the following options: -s 2 (reverse stranded) –f (count reads at the exon level), –

O (assign reads to all their overlapping features), –C (excluding read pairs mapping to different

chromosomes or the same chromosome but on a different strand). The R package edgeR [65] was used

to calculate FPKM (Fragments Per Kilobase Of Exon Per Million Fragments Mapped) per each gene

and organ. For variant sites that were included in more than one exon, the highest FPKM value was

used. We found that 16'013 out of 17'998 genes showed transcriptional activity of at least one exon

(FPKM > 0.3). Overall 166'973 out of 178'504 exons showed evidence for transcription. In a total of

1928 genes, one or more exons showed no evidence for transcription. Retained SNPs were found

among 118'756 exons and 17'685 genes. Overall 611'711 out of 677'578 SNPs were located in genes

with evidence for transcription.

Deleterious mutations are assumed to be overwhelmingly derived mutations. We used all ibex species

except Alpine and Iberian ibex as an outgroup to define the derived state. For each biallelic site,

which was observed in alternative state in Alpine ibex or Iberian ibex, the alternative state was

defined as derived if its frequency was zero in all other species (a total of 44'730 autosomal SNPs).

For loci with more than two alleles, the derived state was defined as unknown. For comparisons

among all species, we only used the following criteria to select SNPs (370'853 biallelic SNPs

retained): transcriptional activity (FPKM > 0.3 in at least one organ) and GERP > -2. The minimum

GERP score cut-off was set to retain only high-quality chromosomal regions following previously

established practice e.g. [24]. We further followed the following categorizations adopted for human

populations to identify moderate to highly deleterious mutations e.g. [24]: -2 to +2 is considered neutral

or near neutral, 2 to 4 considered as moderate, 4 to 6 as large and >6 as extreme effects. We also

required a minimal distance to the next SNP of 3bp to avoid confounding effects of potential multi-

nucleotide polymorphisms (MNPs).

Population genetic analyses

Site frequency spectra (SFS) were calculated using the R packages *plyr* and *dplyr*. SFS analyses were

performed for SnpEff and GERP categories and two additional conservation scores: phyloP [66] and

phastCons [67]. We chose a cutoff of 1 to distinguish conserved from less conserved sites. In the case of

phyloP, sites with a score above 1 were defined as conserved. For phastCons, sites with a score equal

to 1 (the maximum observed value) were considered as conserved.

For individual counts of derived alleles or homozygotes, we used all biallelic sites polymorphic either

in Alpine or Iberian ibex (or both) for which the derived state was known with a maximal missing rate

per locus of 10%. We retained all sites matching these criteria for any downstream analyses even if a

particular site was not polymorphic in any given population. The effective rate of missing data per

locus was between 0.03-0.07% (Figure S43) and no correlation was found between missing rate per

population and counts. We found no qualitative differences if we included only loci with a 100%

genotyping rate (Figure S44).

We calculated the relative number of derived alleles Rxy [29] for the different categories of mutations.

Rxy compares the number of derived alleles found at sites within a specific category. Following[9,29],

we used a random set of 65592 intergenic SNPs for standardization, which makes Rxy robust against

sampling effects and population substructure. We performed the Rxy analysis for the four SnpEff

categories as well as four additional mutation scoring methods: SIFT [68], REVEL [69], CADD [70] and

VEST3 [71] . Human scores mapped to hg38 chromosomal positions were retrieved using the web

interface of the Variant Effect Predictor (VEP) by ensemble[72]. The scores were mapped to

chromosomal positions in the domestic goat reference genome using liftOver

(http://hgdownload.cse.ucsc.edu, v.287) with the chain file accessed from http://hgdownload-

test.cse.ucsc.edu/goldenPath/capHir1. As we applied these scores outside of humans, we did not have

pathological evidence underpinning score cut-offs for deleteriousness. We conservatively used score

cut-offs proposed as best-practices by the tool developers (Revel: 0.75, CADD: 20, SIFT: 0.05) or

used a very conservative percentile (99%, 0.91 VEST3).

Individual-based simulations with Nemo

Individual-based forward simulations were run using the software Nemo [49] v.2.3.51. A customized

version of aNEMOne [73] was used to prepare input files for parameter exploration. The sim.ini file for

the final set of parameters run in 100 replicates is available as Supplementary File 1. All populations

relevant for the founding of the populations under study were included in the model. See Figure 6A

for the simulated demography, which was modeled with the actual founder numbers (assuming a sex-

ratio of 1:1), while the translocations were simplified into four phases (data from [48], DRYAD entry

doi:10.5061/dryad.274b1 and [47]). The harmonic mean of the population census from the founding up

to the final sampling year (2007) was used to define the population carrying capacity. Mating was

assumed to be random and fecundity (mean number of offspring per female) set to five. The selection

coefficients of 5000 biallelic loci subject to selection were drawn from a gamma distribution with a

mean of 0.01 and a shape parameter of 0.3 resulting in $s < 1\%$ for 99.2% of all loci [74] (Figure S45).

Based on empirical evidence, we assumed a negative relationship between h and s [75]. We used the

exponential equation $h = \exp(-51*s)/2$ with a mean h set to 0.37 following [76]. We assumed hard

selection acting at the offspring level. In addition to the 5000 loci under selection, we simulated 500

neutral loci. Recombination rates among each neutral or deleterious locus was set to 0.5. This

corresponds to an unlinked state. Initial allele frequencies for the burn-in were set to $\mu / h * s =$

0.0014 (corresponding to the expected mean frequency at mutation-selection balance [77]). Mutation

rate μ was set to 5e-05 and deleterious mutations were allowed to back-mutate at a rate of 5e-07.

A burn-in of 3000 generations was run with one population ($N = 1000$) representing the entire species

allowing to reach a quasi-equilibrium. N was reduced to $N = 500$ for five generations before a brief,

two generation bottleneck of $N = 80$. At generation 3007, the population recovered to $N = 1000$ and

three generations later the reintroduction was started with the founding of the two zoos Interlaken

Harder (ih) and Peter and Paul (pp). The founding of new populations was modeled by migration of

offspring into an empty patch.

The zoo ih (Interlaken Harder) and several populations did not survive all replicates of the

simulations. Extinction rates were as follows: ih (Zoo Interlaken Harder) 84%, bo (Bire Öschinen)

3%, wh (Weisshorn) 3%, ob (Oberbauenstock) 9%, am (Alpi Marittime) 14% and pil (Pilatus) 2%.

The high extinction rate of the zoo Interlaken Harder did not affect the outcome of the simulations.

The extinctions were a result of the strong reduction in population size during the founding and

occurred always after the founding (see also Figure S16). The extinctions of the reintroduced

populations did not affect the estimates of derived allele counts but reduced sample sizes and, hence,

affected the variance of estimators. Genetic load information was retrieved using the *delet* statistics

option and is defined in Nemo as the mean realized fitness ($L = 1 - W_{mean}/W_{max}$, where W_{max} is the

maximum number of surviving offspring per female computed from her deleterious mutations).

Data availability

Raw whole-genome sequencing data produced for this project was deposited at the NCBI Short Read

Archive under the Accession nos. SAMN10736122–SAMN10736160 (BioProject PRJNA514886).

Raw RNA sequencing data produced for this project was deposited at the NCBI Short Read Archive

under the Accession nos. SAMN10839218-SAMN10839227 (BioProject PRJNA517635).

Acknowledgments

We thank the following organizations and colleagues who contributed samples to this project. Iris Biebach, the Swiss hunting authorities of the cantons of Bern, Nidwalden, Obwalden, Uri, Graubünden and Wallis; the Gran Paradiso National Park (Alice Brambilla) and the Alpi Marittime National Park (Laura Martinelli), Sebastien Regnaut and Richard Kock, Zoological Society of London, Christian Siegenthaler, Ruedi Kunz and Samer Angelone-Alasaad. We are thankful to Glauco Camenisch and Kasia Sluzek, who provided access to Alpine ibex RNAseq datasets. We thank Laurent Excoffier, Stephan Peischl, Kimberly Gilbert, Heidi Lischer, Stefan Wyder, Thomas Wicker, Alan Brelsford, Jessica Purcell, Sarah P. Otto, Andreas Wagner, Sam Yeaman and Nicolas Perrin for helpful advice and comments on previous versions of the manuscript. We are grateful for drawings by Nadine Coline of the Zoological Museum of Zürich. This work was supported by the University of Zurich through a University Research Priority Program "Evolution in Action" pilot project grant and the Swiss Federal Office for the Environment. DC and CG were supported by the Swiss National Science Foundation (grant 31003A_173265 and 31003A_182343, respectively). This study makes use of data generated by the NextGen Consortium, which was supported by grant agreement number 244356 of the European Union's Seventh Framework Programme (FP7/2010-2014).

References:

1. Pimm, S. L. *et al.* The biodiversity of species and their rates of extinction, distribution, and protection. *Science* **344,** 1246752–1246752 (2014).
2. Frankham, R., Bradshaw, C. J. A. & Brook, B. W. Genetics in conservation management: Revised recommendations for the 50/500 rules, Red List criteria and population viability analyses. *Biol Conserv* **170,** 56–63 (2014).
3. Cruz, F., Vilà, C. & Webster, M. T. The legacy of domestication: accumulation of deleterious mutations in the dog genome. *Mol Biol Evol* **25,** 2331–2336 (2008).
4. Renaut, S. & Rieseberg, L. H. The accumulation of deleterious mutations as a consequence of domestication and improvement in sunflowers and other *compositae* crops. *Mol Biol Evol* **32,** 2273–2283 (2015).
5. Lynch, M., Conery, J. & Burger, R. Mutation accumulation and the extinction of small populations. *Am Nat* **146,** 489–518 (1995).
6. Marsden, C. D. *et al.* Bottlenecks and selective sweeps during domestication have increased deleterious genetic variation in dogs. *PNAS* **113,** 152–157 (2016).
7. Robinson, J. A., Brown, C., Kim, B. Y., Lohmueller, K. E. & Wayne, R. K. Purging of strongly deleterious mutations explains long-term persistence and absence of inbreeding depression in Island Foxes. *Curr Biol* 1–13 (2018). doi:10.1016/j.cub.2018.08.066
8. Laenen, B. *et al.* Demography and mating system shape the genome-wide impact of purifying selection in Arabis alpina. *PNAS* **115,** 816–821 (2018).
9. Xue, Y. Mountain gorilla genomes reveal the impact of long-term population decline and inbreeding. *Science* **348,** 239–242 (2015).
10. Charlesworth, B. Effective population size and patterns of molecular evolution and variation. *Nat Rev Genet* **10,** 195–205 (2009).
11. Kimura, M. Stochastic processes and distribution of gene frequencies under natural selection. *Cold Spring Harb. Symp. Quant. Biol.* **20,** 33–53 (1955).
12. Keller, L. F. & Waller, D. Inbreeding effects in wild populations. *Trends Ecol Evol* **17,** 230–241 (2002).
13. Glémin, S. How are deleterious mutations purged? Drift versus nonrandom mating. *Evolution* **57,** 2678–2687 (2003).
14. Garcia-Dorado, A. Understanding and predicting the fitness decline of shrunk populations: inbreeding, purging, mutation, and standard selection. *Genetics* **190,** 1461–1476 (2012).
15. Kirkpatrick, M. & Jarne, P. The effects of a bottleneck on inbreeding depression and the genetic load. *Am Nat* **155,** 154–167 (2000).
16. Crow, J. F. in *Mathematical Topics in Population Genetics* (ed. Kojima, K. I.) 128–177 (Springer-Verlag, 1970).
17. Robinson, J. A. *et al.* Genomic flatlining in the endangered island fox. *Curr Biol* **26,** 1183–1189 (2016).
18. Bataillon, T. & Kirkpatrick, M. Inbreeding depression due to mildly deleterious mutations in finite populations: size does matter. *Genet Res* **75,** 75–81 (2000).
19. Willi, Y., Fracassetti, M., Zoller, S. & Van Buskirk, J. Accumulation of mutational load at the edges of a species range. *Mol Biol Evol* **22,** 140 (2018).
20. Hedrick, P. W. & Garcia-Dorado, A. Understanding inbreeding depression, purging, and genetic rescue. *Trends Ecol Evol* **31,** 940–952 (2016).
21. Rogers, R. L. & Slatkin, M. Excess of genomic defects in a woolly mammoth on Wrangel island. *PLoS Genet* **13,** e1006601 (2017).
22. Robinson, J. A., Brown, C., Kim, B. Y., Lohmueller, K. E. & Wayne, R. K. Purging of strongly deleterious mutations explains long-term persistence and absence of inbreeding depression in island foxes. *Curr Biol* 1–13 (2018). doi:10.1016/j.cub.2018.08.066
23. Simons, Y. B., Turchin, M. C., Pritchard, J. K. & Sella, G. The deleterious mutation load is insensitive to recent population history. *Nat Genet* **46,** 220–224 (2014).
24. Henn, B. M. *et al.* Distance from sub-Saharan Africa predicts mutational load in diverse human genomes. *PNAS* **113,** E440–E449 (2015).

25.   Henn, B. M., Botigué, L. R., Bustamante, C. D., Clark, A. G. & Gravel, S. Estimating the
mutation load in human genomes. *Nature* **16,** 333–343 (2015).
26.   Lopez, M. *et al.* The demographic history and mutational load of African hunter-gatherers
and farmers. *Nature Ecology & Evolution* **2,** 1–13 (2018).
27.   Scott, E. M. *et al.* Characterization of Greater Middle Eastern genetic variation for enhanced
disease gene discovery. *Nat Genet* **48,** 1071–1076 (2016).
28.   Peischl, S. *et al.* Relaxed selection during a recent human expansion. *Genetics* **208,** 763–777
(2018).
29.   Do, R. *et al.* No evidence that selection has been less effective at removing deleterious
mutations in Europeans than in Africans. *Nature* **47,** 126–131 (2015).
30.   Lohmueller, K. E. The distribution of deleterious genetic variation in human populations.
Curr Opin Genet Dev **29,** 139–146 (2014).
31.   Fu, W., Gittelman, R. M., Bamshad, M. J. & Akey, J. M. Characteristics of Neutral and
Deleterious Protein-Coding Variation among Individuals and Populations. *Am J Hum Genet*
95, 421–436 (2014).
32.   Laws, R. J. & Jamieson, I. G. Is lack of evidence of inbreeding depression in a threatened
New Zealand robin indicative of reduced genetic load? *Animal Conservation* **14,** 47–55
(2011).
33.   Kennedy, E. S., Grueber, C. E., Duncan, R. P. & Jamieson, I. G. Severe inbreeding
depression and no evidence of purging in an extremely inbred wild species--the Chatham
Island black robin. *Evolution* **68,** 987–995 (2014).
34.   Crnokrak, P. & Barrett, S. Perspective: Purging the genetic load: A review of the
experimental evidence. *Evolution* **56,** 2347–2358 (2002).
35.   Kalinowski, S. T., Hedrick, P. W. & Miller, P. S. Inbreeding depression in the Speke's
gazelle captive breeding program. *Conserv Biol* **14,** 1375–1384 (2000).
36.   Xue, Y. *et al.* Mountain gorilla genomes reveal the impact of long-term population decline
and inbreeding. *Science* **348,** 242–245 (2015).
37.   Grodinsky, C. & Stuwe, M. The reintroduction of the Alpine ibex to the Swiss Alps.
Smithsonian **18,** 68–77 (1987).
38.   Biebach, I. & Keller, L. F. A strong genetic footprint of the re-introduction history of Alpine
ibex (*Capra ibex ibex*). *Mol Ecol* **18,** 5046–5058 (2009).
39.   Grossen, C., Biebach, I., Angelone-Alasaad, S., Keller, L. F. & Croll, D. Population
genomics analyses of European ibex species show lower diversity and higher inbreeding in
reintroduced populations. *Evol Appl* **11,** 123–139 (2018).
40.   IUCN 2018. *httpwww.iucnredlist.org* Available at: (Accessed: 10 November 2018)
41.   Couturier, M. *Le Bouquetin des Alpes.* (1962).
42.   Reading, R. & Shank, C. *Capra sibirica:* The IUCN Red List of Threatened Species 2008.
doi:10.2305/IUCN.UK.2008.RLTS.T42398A10695735.en
43.   McQuillan, R. *et al.* Runs of homozygosity in European populations. *The American Journal*
of Human Genetics **83,** 359–372 (2008).
44.   Cooper, G. M. *et al.* Distribution and intensity of constraint in mammalian genomic
sequence. *Genome Res* **15,** 901–913 (2005).
45.   Terrier, G. & Rossi, P. Le bouquetin (*Capra ibex ibex*) dans les alpes maritimes franco-
italiennes: occupation de l'éspace, clonisation et régulation naturelles. *Traveaux Scientifiques*
du Parc National de la Vanoise **XVIII,** 271–288 (1994).
46.   Maudet, C. *et al.* Microsatellite DNA and recent statistical methods in wildlife conservation
management: applications in Alpine ibex *Capra ibex (ibex). Mol Ecol* **11,** 421–436 (2002).
47.   Biebach, I. & Keller, L. F. Inbreeding in reintroduced populations: the effects of early
reintroduction history and contemporary processes. *Conserv Genet* **11,** 527–538 (2010).
48.   Aeschbacher, S., Futschik, A. & Beaumont, M. A. Approximate Bayesian computation for
modular inference problems with many parameters: the example of migration rates. *Mol Ecol*
22, 987–1002 (2013).
49.   Guillaume, F. & Rougemont, J. Nemo: an evolutionary and population genetics
programming framework. *Bioinformatics* **22,** 2556–2557 (2006).

50. Bozzuto, C., Biebach, I., Muff, S., Ives, A. R. & Keller, L. F. Inbreeding reduces long-term
growth of Alpine ibex populations. *Nature Ecology & Evolution* in press

51. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30,** 2114–2120 (2014).

52. Dong, Y. *et al.* Sequencing and automated whole-genome optical mapping of the genome of
a domestic goat (*Capra hircus*). *Nat Biotechnol* **31,** 135–141 (2012).

53. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nature Methods*
9, 357–359 (2012).

54. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing
next-generation DNA sequencing data. *Genome Res* **20,** 1297–1303 (2010).

55. DePristo, M. A. *et al.* A framework for variation discovery and genotyping using next-
generation DNA sequencing data. *Nat Genet* **43,** 491–498 (2011).

56. Garrison, E. & Marth, G. *Garrison: Haplotype-based variant detection from short-read*
sequencing. arXiv 1207.3907 (2012)

57. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27,** 2156–2158
(2011).

58. Narasimhan, V. *et al.* BCFtools/RoH: a hidden Markov model approach for detecting
autozygosity from next-generation sequencing data. *Bioinformatics* **32,** 1749–1751 (2016).

59. Dumont, B. L.Payseur, Bret A. Evolution of the genomic rate of recombination in mammals.
Evolution **62,** 276–294 (2008).

60. Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w(1118);
iso-2; iso-3. *fly* **6,** 80–92 (2012).

61. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics*
30, 1236–1240 (2014).

62. Fu, W., Gittelman, R. M., Bamshad, M. J. & Akey, J. M. Characteristics of neutral and
deleterious protein-coding variationamong individuals and populations. *Am J Hum Genet* **95,**
421–436 (2014).

63. Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory
requirements. *Nature Methods* **12,** 357–360 (2015).

64. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30,** 923–930 (2014).

65. Robinson, M. D., McCarthy, D. J., Bioinformatics, G. S.2010. edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.

66. Pollard, K. S., Hubisz, M. J., Rosenbloom, K. R. & Siepel, A. Detection of nonneutral
substitution rates on mammalian phylogenies. *Genome Res* **20,** 110–121 (2010).

67. Siepel, A. *et al.* Evolutionarily conserved elements in vertebrate, insect, worm, and yeast
genomes. *Genome Res* **15,** 1034–1050 (2005).

68. Ng, P. C. SIFT: predicting amino acid changes that affect protein function. *Nucleic Acids*
Research **31,** 3812–3814 (2003).

69. Ioannidis, N. M. *et al.* REVEL: An Ensemble Method for Predicting the Pathogenicity of
Rare Missense Variants. *Am J Hum Genet* **99,** 877–885 (2016).

70. Kircher, M. *et al.* A general framework for estimating the relative pathogenicity of human
genetic variants. *Nat Genet* **46,** 310–315 (2014).

71. Carter, H., Douville, C., Stenson, P. D., Cooper, D. N. & Karchin, R. Identifying Mendelian
disease genes with the Variant Effect Scoring Tool. *Bmc Genomics* **14,** S3 (2013).

72. McLaren, W. *et al.* Deriving the consequences of genomic variants with the Ensembl API
and SNP Effect Predictor. *Bioinformatics* **26,** 2069–2070 (2010).

73. kjgilbert/aNEMOne. (2017).

74. Keightley, P. D. The distribution of mutation effects on viability in Drosophila melanogaster.
Genetics **138,** 1315–1322 (1994).

75. Agrawal, A. F. & Whitlock, M. C. Inferences about the distribution of dominance drawn
from yeast gene knockout data. *Genetics* **187,** 553–566 (2011).

76. Gilbert, K. J. *et al.* Local Adaptation Interacts with Expansion Load during Range
Expansion: Maladaptation Reduces Expansion Load. *Am Nat* **189,** 368–380 (2017).

77.   Crow, J. F. & Kimura, M. *An Introduction to Population Genetics Theory*. (New Jersey:
685   Blackburn Press, 1970).
686

Figure 1

Figure 2

Figure 3

Figure 4

Relative position in protein

Figure 6