Shifting from fear to safety through deconditioning-update: a novel

approach to attenuate fear memories

Bruno Popik[1], Felippe E. Amorim[2], Olavo B. Amaral[2] and Lucas de Oliveira Alvares[1*]

[1]Neurobiology of Memory Lab, Biophysics Department, Biosciences Institute, Federal

University of Rio Grande do Sul, Porto Alegre, Brazil. [2]Institute of Medical

Biochemistry Leopoldo de Meis, Federal University of Rio de Janeiro, Rio de Janeiro,

Brazil,

Corresponding author: Lucas de Oliveira Alvares, Laboratório de Neurobiologia da

Memória, Departamento de Biofísica, Instituto de Biociências, Universidade Federal do

Rio Grande do Sul (UFRGS), Av. Bento Gonçalves 9500, Prédio 43422, Sala 210, CEP

91.501-970, Porto Alegre, Rio Grande do Sul, Brasil. E-mail: lucas.alvares@ufrgs.br

Abstract

Aversive memories are at the heart of psychiatric disorders such as phobias and post-traumatic stress disorder (PTSD). Here, we present a new behavioral approach in rats that robustly attenuates aversive memories. This method consists of "deconditioning" animals previously trained to associate a tone with a strong footshock by replacing it with a much weaker one during memory retrieval. Our results indicate that deconditioning-update is more effective than traditional extinction in reducing fear responses; moreover, such effects are long lasting and resistant to renewal and spontaneous recovery. Remarkably, this strategy overcame important boundary conditions for memory updating, such as remote or very strong traumatic memories. The same beneficial effect was found in other types of fear-related memories. Deconditioning was mediated by L-type voltage-gated calcium channels and is consistent with computational accounts of mismatch-induced memory updating. Our results suggest that shifting from fear to safety through deconditioning-update is a promising approach to attenuate traumatic memories.

Introduction

Memory is a dynamic process that allows for adaptation to the demands of a

continuously changing environment. The ability to update old memories in accordance

with new experiences is crucial for maintaining their relevance over time. Particularly, it

has been shown that after retrieval (or reactivation), memories may undergo a cycle of

labilization and restabilization commonly known as reconsolidation (Nader et al., 2000).

The labile state induced by this process can thus allow changes in memory strength

and/or content (De Oliveira Alvares et al., 2013). This has been most extensively

studied in aversive conditioning paradigms in rodents and humans and is of potential

relevance to the management of psychiatric disorders involving dysfunctional memories

(Monfils & Holmes, 2018).

Repeated exposure to a conditioned stimulus (CS) in the absence of an aversive

unconditioned stimulus (US) also leads to a progressive reduction in fear responses,

commonly known as extinction. However, extinction is thought not to erase the original

memory; instead, it induces new learning that transiently inhibits fear expression

(Bouton, 2002). Therefore, the fear memory typically reemerges with the passage of

time (spontaneous recovery), exposure to the US (reinstatement), or when the CS is

presented independently of the extinction context (renewal) (Rescorla and Heth, 1975,

Archbold et al., 2010, Bouton et al., 2012). Thus, behavioral strategies that can weaken

traumatic memories and reduce memory recovery can be relevant for improving the

effectiveness of extinction.

Reconsolidation has been described in several experimental paradigms and species,

from invertebrates to humans, suggesting that it might a fundamental property of

memories (Nader and Einarsson, 2010). Beyond its biological role in memory updating,

it opens a window of opportunity as a potential therapeutic strategy to modify

pathological memories. Several studies in the last decades have attempted to

pharmacologically disrupt the reconsolidation of traumatic memories (Beckers and

Kindt, 2017), as at least in some situations, this strategy can be less susceptible to

spontaneous or induced recovery than traditional extinction (Duvarci and Nader, 2004,

Monfils et al., 2009). However, most treatments that interfere with memory

reconsolidation are toxic and cannot be readily administered to humans. Thus, in spite

of almost two decades of research on memory reconsolidation, the evidence for the

efficacy of reconsolidation-blocking treatments in clinical settings is still limited

(Monfils and Holmes, 2018).

Research on reconsolidation-extinction boundaries suggests that the transition from one

process to the other depends on the degree of mismatch between the original memory

and the reactivation experience. Many studies have suggested that some degree of

mismatch or prediction error is necessary for reconsolidation to occur (Lee, 2009,

Fernandez et al., 2016); however, if prediction error is too high, extinction may occur

instead (Suzuki et al., 2004, Sevenster et al., 2014). Computational models suggest that,

in models in which prediction error is low, memory updating/reconsolidation

mechanisms are preferentially engaged, as the new experience is recognized as a new

instance of the former one; however, as mismatch increases, the chances of it being

attributed to a new latent cause (Gershman et al., 2017) or forming a new attractor in a

neural network (Osan et al., 2011) increases.

This framework suggests that lowering the degree of mismatch between learning and

reexposure might conceivably potentiate memory updating during extinction. This has

been the rationale behind so-called retrieval-extinction procedures (Kredlow et al.,

2016) and has also been explored in short-term extinction protocols (Gershman et al.,

2013). In this work, we propose a novel approach for long-term attenuation of traumatic

memories that we term "deconditioning-update". This strategy consists in weakening

fear memories by updating the aversive information, substituting the original US by a

mildly aversive stimulus during the plastic state induced by reactivation.

107   **Results**

In order to approach our hypothesis, male Wistar rats were trained in auditory fear

conditioning, where they received 5 conditioning trial tones (CS) that co-terminated

with a 0.5-mA, 1-s footshock (US). On days 3 to 6 (reactivation), animals received 3

CSs during a 400-s daily session in a different context. In the no-footshock group, they

were unpaired, while in the deconditioning-update group, each tone co-terminated with

a 0.1-mA footshock. A third group remained in the homecage (control group). On day

7, both groups were tested in the reactivation context with 3 unpaired CSs. On day 8,

animals were tested in the training context for renewal, and, on day 28, they were

retested for spontaneous recovery (**Fig.1A**).

Figure 1

Figure 1 - Weakening fear memory through deconditioning-update training. (**A**)

Experimental design: male rats were fear-conditioned with 5 tone-shock pairings

(context A; 5 CS + US, 0.5mA). 48 h later, the no-footshock and footshock

(deconditioning-update*)* groups were exposed to 4 daily reactivation sessions (context

B). After this, animals underwent test (context B), renewal (context A) and spontaneous

recovery (context B) sessions. Black circles represent context A, while white rectangles

represent context B. (**B**) Freezing levels during reactivation sessions. Rats exposed to

weak footshocks during reactivation sessions showed a significant reduction in freezing

responses, maintained during the test (**C**), renewal (**E**) and spontaneous recovery (**D**)

sessions. (**F**) Experimental design: female rats were fear-conditioned (context A;

5CS+US, 0.5mA). 48 hours later, the no-footshock and footshock groups were exposed

to 4 daily reactivation sessions (context B). After this, all groups underwent test,

renewal, and spontaneous recovery sessions. Animals were reconditioned (context A;

3CS+US, 0.5mA) on the next day and retested 24 h later. (**G**) Freezing levels during

memory reactivation. Rats exposed to weak footshocks showed a significant reduction

in freezing responses, maintained during the test (**H**), renewal (**I**), spontaneous recovery

(**J**) and retraining test (**K**) sessions. Bars represent mean ± SEM. Statistical comparisons

were performed using two-way repeated-measures ANOVA followed by a Bonferroni

post-hoc (reactivation sessions) or one-way ANOVA followed by Tukey post-hoc (test,

renewal, spontaneous recovery, and retraining test). * $p < 0.05$; ** $p < 0.005$; *** $p <$

0.0005; **** $p < 0.0001$. For full statistics, see **Table S1**.

Over the course of reactivation sessions, both groups presented a decrease in freezing,

but this was more marked in the deconditioning-update group (**Fig 1B**), which presented

a decrease in freezing of around 70% in comparison to the no-footshock group and 80%

in comparison to the homecage control group in the test session (**Fig 1C**). Animals in

the deconditioning-update group also had lower freezing responses in the renewal (**Fig**

1D) and spontaneous recovery (**Fig. 1E**) tests, although it should be noted that memory

recovery was not observed in the no-footshock group either, perhaps due to a ceiling

effect caused by insufficient extinction (for complete statistics, see **Table 1**). When

performing the same experiment, but with each tone co-terminating with a 0.3-mA

footshock in the reactivations, no fear reduction was shown (**Fig. S1**); on the contrary,

the footshock group presented higher levels of freezing than the no-footshock group

throughout the extinction sessions, as well as in the test. When using a single

reactivation session with 0.1mA, no difference was found in the test in comparison to

the no-footshock group (**Fig. S2**), suggesting that deconditioning-update requires

several sessions to take place.

One could argue that the exposure to weak footshocks could simply lead to habituation

and consequent devaluation of the US, without altering the conditioned association

itself (Rescorla, 1973; Rescorla and Heth, 1975). In order to test this alternative

interpretation, rats were conditioned as described above and the same 3 0.1 mA weak

footshocks were given in another context without being paired with the tones (**Fig.**

S3A). In this case, no fear reduction was found in comparison to homecage controls.

(**Fig. S3B**). To further rule out the devaluation hypothesis, another set of animals was

submitted to reinstatement after deconditioning, in order to test whether restoring the

original footshock valence outside of the extinction context might lead to memory

recovery. We found that the deconditioning group expressed a lower fear level when

compared with the no-footshock group in the test even after reinstatement (**Fig. S3E**),

suggesting that the deconditioning-update effect is not due to US devaluation, but rather

to updating of the CS-US association.

Next, we asked whether the deconditioning-update effect would also be efficient in

reducing fear in female rats. As observed in males, females from the deconditioning-

update group showed lower freezing levels throughout the reactivation sessions, as well

as in the test session, when compared with the control group (**Fig 1H**). The same pattern

was maintained in the renewal and spontaneous recovery tests (**Fig 1I** and **1J**,

respectively). As a further way to test for persistence of the original memory, we

performed a retraining session in the original context with 3 0.5-mA CS-US pairings 24

h after the spontaneous recovery test to assess savings. The deconditioning-update

group had lower freezing compared with the other groups in a subsequent test session in

the extinction context, suggesting that our protocol also lowers the rate of re-acquisition

of an aversive memory (**Fig. 1K**).

Boundary conditions such as training intensity and memory age have been reported to

prevent fear memories from being modified. Protocols with high training intensity make

memory less prone to interference by pharmacological agents in the reconsolidation

window (Frankland et al., 2006, Suzuki et al., 2004, Wang et al., 2009), while older

memories are also less susceptible to modification than recent ones (Milekic and

Alberini, 2002, Frankland et al., 2006, Suzuki et al., 2004, Bustos et al., 2009). Thus,

our next experiments investigated whether deconditioning-update could attenuate fear

expression in these cases. First, we trained the animals in the same protocol described

above, but starting reactivations 40 days after conditioning (**Fig. 2A**). Once more, the

deconditioning-update group showed lower freezing expression throughout the

reactivation sessions (**Fig. 2B**). 24 h after the last reactivation, both the footshock and

no-footshock groups presented a comparable decrease in freezing behavior when

compared to controls (**Fig. 2C**). However, in the renewal and spontaneous recovery

tests, fear expression reemerged in the no-footshock group, while the deconditioning-

update group maintained its low freezing levels (**Fig. 2D** and **2E**, respectively; **Table**

2).

Figure 2

Figure 2 - Deconditioning-update weakens both remote and strong fear memories.

(**A**) Experimental design for remote memory: rats were fear-conditioned with 5 tone-

shock pairings (context A; 5 CS + US, 0.5mA). Starting 40 days later, the no-footshock

and footshock (deconditioning-update) groups were exposed to daily reactivation

sessions (context B). Then, all groups underwent test (context B), renewal (context A)

and spontaneous recovery (context B) sessions. Black circles represent context A, while

white rectangles represent context B. (**B**) Freezing levels during reactivation sessions.

Rats exposed to weak footshocks during reactivation sessions showed similar freezing

levels to no-footshock animals during the test session (**C**) and lower freezing levels at

the renewal (**D**) and spontaneous recovery (**E**) ones. (**F**) Experimental design for strong

training (5CS+US, 1mA). (**G**) Freezing levels during reactivation sessions. Rats

exposed to weak footshocks during reactivation sessions showed a significant reduction

in freezing responses that was maintained during the test (**H**), renewal (**I**) and

spontaneous recovery (**J**) sessions. Bars represent mean ± SEM. Statistical comparisons

are performed using two-way repeated-measures ANOVA followed by a Bonferroni

post-hoc (reactivation sessions) or one-way ANOVA followed by a Tukey post-hoc

(test and spontaneous recovery). * $p < 0.05$; ** $p < 0.005$; *** $p < 0.0005$; **** $p <$

0.0001. For full statistics, see **Table S2**.

Next, we tested whether a stronger fear memory would constrain the deconditioning-

update effect by training animals with 5 CS-US pairings using 1mA shocks, while

maintaining the rest of the protocol unchanged. In spite of the stronger shock intensity

in the training session, the deconditioning-update group presented reduced freezing

levels in reactivation sessions when compared to the no-footshock group (**Fig. 2G**).

These lower freezing levels were maintained in the test, renewal and spontaneous

recovery sessions, in which robust recovery was observed in no-footshock animals, but

not in the deconditioning-update group (**Fig. 2H**, **2I** and **2J**, respectively; **Table 2**).

Similar results were observed in females in a slightly modified protocol with 3

reactivation sessions (**Fig. S4**). These experiments suggest that the deconditioning-

update induces a plastic state, allowing the fear memory trace to be altered even in

boundary conditions that usually constrain memory updating (Pedraza et al., 2018).

In order to investigate whether the deconditioning-update approach is effective in

attenuating other types of aversive memories, we trained animals in different fear-

related tasks. First, we used a contextual fear conditioning paradigm, which is known to

involve a set of brain regions that is partially distinct from auditory conditioning and

includes the prefrontal cortex and hippocampus (Maren et al., 2013). Animals were

placed in the conditioning chamber for 3 min before receiving two 0.5-mA, 2-s

footshocks separated by a 30-s interval. On days 3 to 6, rats were reexposed to the same

context for 4 min, with those in the deconditioning-update group receiving weak

footshocks (0.1mA, 2 s) 3 min after being placed in the chamber (**Fig. 3A**). The

deconditioning-update group had lower fear expression during reactivations (**Fig. 3B**)

and maintained these lower freezing levels compared with the other groups in the test

(**Fig. 3C**). The same pattern was observed in the spontaneous recovery test, performed

20 days later, suggesting that the decrease in freezing caused by deconditioning-update

is long-lasting (**Fig. 3D**; **Table 3**).

Another set of animals underwent the step-through inhibitory avoidance task, in which

training consists of applying 4 0.5mA, 1-s footshocks with 5-s intervals when the

animal enters the dark compartment of a shuttle box. During reactivations, animals were

placed in the dark compartment for 30 s, with those in the deconditioning-update group

receiving 2 0.1-mA shocks. In the test session, the animals were put in the light

compartment, and the latency to enter the dark compartment was measured (**Fig. 3E**).

The deconditioning-update group showed a much shorter latency to reach the dark

chamber (**Fig. 3F**) and spent more time in the dark compartment over the 10-min

session compared to the other groups (**Fig 3G**; **Table 3**), suggesting that memory was

more robustly updated in this group during reactivation. Taken together, these results

suggest that the deconditioning-update strategy is effective in weakening distinct types

of fear-related memories.

Figure 3

Figure 3 - Deconditioning-update weakens fear memory in different behavioral

tasks. (**A**) Experimental design in contextual fear conditioning: rats were fear-

conditioned with 5 contextual-shock pairings (4 min context + 4 US, 0.5mA). Starting

48 h later, the no-footshock and footshock groups were exposed to daily reactivation

sessions. 24 h after the last reactivation, all groups were tested; 20 days later, they were

tested for spontaneous recovery. (**B**) Freezing levels during reactivation sessions. Rats

exposed to weak footshocks during reactivation sessions showed a significant reduction

in freezing responses maintained during the test (**C**) and spontaneous recovery (**D**)

sessions. (**E**) Experimental design in inhibitory avoidance: rats were placed in the

lighted compartment and received footshocks (4 US, 0.5mA) upon entering the dark

one. Starting 48 h later, the no-footshock and footshock groups were exposed to daily

30-s reactivation sessions in the dark compartment; 24 h after the last reactivation, all

groups were tested. Rats exposed to weak footshocks during reactivation sessions

showed lower latencies to cross to the dark compartment (**F**) and spent more time in it

during the test (**G**). Bars represent mean ± SEM or median with interquartile range (in F

and G). Statistical comparisons for contextual fear conditioning are performed using

two-way repeated-measures ANOVA followed by a Bonferroni post-hoc (reactivation

sessions) or one-way ANOVA followed by a Tukey post-hoc (test, renewal, and

spontaneous recovery). For inhibitory avoidance, a Kruskal-Wallis test followed by a

Dunn post-hoc was performed. $*p < 0.05$; $**p < 0.005$; $***p < 0.0005$; $****p < 0.0001$.

For full statistics, see **Table S3**.

To address whether the deconditioning-update effect would also be observed within a

single, long-lasting extinction session, we trained animals and exposed them 48 h later

to a session containing 12 CSs, with the deconditioning-update group receiving a 0.1-

mA footshock at the end of each tone. Fear reduction was limited and largely similar

across groups in the extinction session and in the subsequent tests (**Fig. S5**), albeit with

slightly lower renewal in the deconditioning-update group, suggesting that the pairing

of the CS with a weak footshock is not as effective in accelerating single-session

extinction. In order to further explore this possibility, we subjected animals to a 24-tone

single-session extinction protocol (**Fig. 4A**). This led to robust fear reduction both

within the extinction session (**Fig 4B**) and in a subsequent test (**Fig 4C**) in the no-

footshock group, while freezing remained largely unchanged in the deconditioning-

update group. However, fear memory reemerged in the renewal and spontaneous

recovery test among no-footshock animals (**Fig 4D** and **4E**; **Table 4**), as typically

occurs with extinction learning. These results show that the presence of a weak shock at

the end of every CS not only does not enhance extinction occurring over a single

behavioral session, but can actually impair it in the short-term.

Other studies have shown that activation of L-type voltage-gated Ca^{++} channels

(LVGCC) is necessary both for destabilizing a reactivated memory during

reconsolidation (Suzuki et al., 2008, Lee and Flavell, 2014, Crestani et al., 2015,

Haubrich et al., 2015) and for some forms of extinction (Cain et al., 2002). Thus, we

used the LVGCC antagonist nimodipine as a pharmacological tool to investigate

whether deconditioning-update involved memory destabilization mechanisms (Flavell et

al., 2011, Sierra et al., 2013, Crestani et al., 2015, Haubrich et al., 2015). Animals were

trained and divided into four groups: no-footshock + vehicle, no-footshock +

nimodipine, deconditioning-update + vehicle and deconditioning-update + nimodipine,

with nimodipine or vehicle administered systemically 30 min before each reactivation

(**Fig 4F**). Nimodipine attenuated freezing decrease in both behavioral protocols,

suggesting that the drug impaired both deconditioning-update and regular extinction

(**Fig. 4G** and **4H**). In the renewal and spontaneous recovery sessions, high freezing

levels reemerged both in the no-footshock vehicle group and in nimodipine-treated

animals, while the deconditioning-update vehicle group maintained a lower freezing

level (**Fig. 4I** and **4J**; **Tables 4**). This suggests that deconditioning-update is mediated

by memory destabilization processes requiring Ca^{++} influx through LVGCCs.

Figure 4

Figure 4 - Deconditioning-update is based on memory destabilization mechanisms.

(**A**) Experimental design: rats were fear-conditioned with 5 tone-shock pairings (context

A; 5CS+US, 0.5mA). 48 h later, the no-footshock and footshock groups underwent a

single extinction session (context B, 24 CSs), followed by test (context B), renewal

(context A) and spontaneous recovery (context B) sessions. (**B**) Freezing levels during

extinction. Weak footshocks impaired extinction within the session and in the test

session (**C**), but not in renewal (**D**) or spontaneous recovery (**E**). (**F**) Experimental

design: rats were fear-conditioned (context A; 5CS+US, 0.5mA). 48 h later, all animals

underwent daily reactivation sessions (context B), receiving nimodipine (16 mg/kg, i.p.)

or vehicle 30 min before each one. They then underwent test (context B), renewal

(context A) and spontaneous recovery (context B) sessions. Nimodipine prevented

freezing decrease across reactivation sessions in both groups (**G**). Freezing was similar

between groups in the test session (**H**), but was lower in the vehicle-footshock group in

the renewal (**I**) and spontaneous recovery (**J**) sessions. Bars represent mean ± SEM.

Statistical comparisons are performed using two-way repeated-measures ANOVA

followed by Bonferroni post-hoc (extinction), one-way ANOVA followed by Tukey

post-hoc (test, renewal, and spontaneous recovery following extinction), three-way

repeated-measures ANOVA followed by Bonferroni post-hoc (reactivation sessions

with nimodipine/vehicle) and two-way ANOVA followed by Bonferroni post-hoc (test,

renewal, and spontaneous recovery following nimopidine/vehicle). * $p < 0.05$; ** $p <$

0.005; *** $p < 0.0005$; **** $p < 0.0001$ in between-group comparisons. For full

statistics, see **Table S4**,

One explanation for our findings is that pairing the CS with a weak footshock could

lead to a smaller degree of prediction error during reexposure sessions, biasing them

towards memory updating as opposed to new learning. Computational models using

neural networks (Osan et al., 2011) or Bayesian approaches (Gershman et al., 2017)

have explored how different degrees of mismatch between stored memories and new

experiences can lead to these two outcomes, suggesting lower degrees of mismatch or

prediction error could lead to greater destabilization of stored memories. With this in

mind, we used an adaptation of one of these models (Osan et al., 2011) to explore if this

framework could account for our main results – i.e. accelerated extinction over multiple

sessions and lower memory recovery when mismatch is reduced during reexposure.

This fully connected Hopfield-like network (Hopfield, 1984) of 100 neurons is capable

of storing patterns using Hebbian learning rules and retrieving them according to the

inputs presented, which in our simulations included neurons representing tone and

context information, as well as shock/non-shock information (**Fig. 5A**). Additionally,

the network also updates synaptic weights according to mismatch between a cue input

and the retrieved network pattern (Osan et al., 2011).

As shown in Fig. **5B,** a lower degree of mismatch in the reexposure pattern, caused by

weak activation of shock-related neurons (as would be expected from deconditioning-

update), can indeed lead to a greater reduction in subsequent fear behavior than a "pure

extinction" pattern (sharing only the tone information with the original memory). This

effect was sensitive to blockade of the mismatch-induced degradation term – the model

equivalent of blocking memory destabilization mechanisms such as LVGCCs with a

drug like nimodipine (**Fig. 5C**). This occurs because the "minor footshock" pattern

leads to recovery of the original memory as the outcome of the initial sessions (**Fig 5D**),

causing mismatch-induced updating of the weights supporting these memories and

weakening of the mutual connections between shock and context neurons (**Fig. 5E**). On

the other hand, no-footshock extinction leads to immediate formation of a new attractor,

leaving the connection weights related to the shock memory unaltered throughout the

extinction process.

Figure 5 – Lower mismatch accelerates extinction and decreases renewal in a neural network model. (**A**) Cue inputs presented to the network during training (shock memory), reexposure (with or without footshock) and test sessions (consisting of the tone and either context B (test) or A (renewal)). Color scale shows the cue received by each of the 100 neurons (**B**) Extinction over multiple sessions using the no-footshock (red bars) or footshock (blue bars) cue. Bars represent freezing, expressed as the activity ratio between shock neurons and the sum of shock and non-shock neurons in response to the test cue, at reexposure days 1 to 4. After each test, memory is updated according to the activity reached in response to the full reexposure pattern. (**C**) Effect of LVGCC

blockade (i.e. setting MID to 0). Removing the degradation term blocks deconditioning-

update, but not regular extinction. **(D)** Network activity in retrieval tests during tone

presentations (e.g. cued with the tone alone) and at the end of reexposure (e.g. cued with

the full reactivation pattern), as well as on test and retrieval sessions. Lower mismatch

(i.e. weak footshock) leads to retrieval of the original pattern on the first days, leading to

memory updating through mismatch-induced degradation and lower retrieval on

subsequent tests. **(E)** Mean synaptic weights between different neuronal groups after

training and at the end of each extinction session. Heat map represents the connection

from neuronal populations in the Y axis to those in the X axis in the no-footshock and

footshock groups. Deconditioning-update leads to weakening of connections between

context and shock neurons and of their inhibitory connections to other neurons. On no-

footshock extinction, an extinction memory is formed with sparing of the shock

representation.

Because of this, deconditioning-update was sensitive to blockade of mismatch-induced

degradation, while no-footshock extinction – which depends basically on new learning

– was not (Fig. **5C**). Unlike in the model, however, nimodipine did affect regular

extinction in the experimental results. This suggests that the mechanistic distinction

between deconditioning-update and classic extinction is not so clear-cut, and that these

two processes might share common mechanisms, as has been proposed for

reconsolidation and extinction (Almeida-Corrêa and Amaral, 2014). Capturing these

subtleties, however, likely requires model implementations that are more complex than

this simple adaptation of the classic Hopfield formulation.

Discussion

Taken as a whole, our findings demonstrate that presenting a tone followed by a weak

footshock (deconditioning-update) was more effective in reducing fear memory than

presenting a tone in the absence of shock (extinction training). Attenuation of fear

responses following deconditioning-update was robust, long-lasting, and less sensitive

to renewal and spontaneous recovery. Remarkably, this strategy was also effective in

reducing fear within boundary conditions in which memories have been described to be

less sensitive to modification (e.g., very strong training protocols and remote

memories). The same effect was found in other types of fear-related memory

(contextual fear conditioning and inhibitory avoidance tasks), and was sensitive to

pharmacological blockade of LVGCCs. We suggest that the CS–US association is

weakened during the deconditioning-update approach, leading to lower fear expression.

Considering that (i) fear reduction is long-lasting, without spontaneous recovery,

renewal and savings, (ii) weak-footshock pairings do not have the same effect in single-

session extinction protocols or when stronger footshocks are applied, and (iii) this effect

is prevented by LVGCC antagonism, we suggest that deconditioning-update is mediated

by the memory destabilization effects commonly associated with reconsolidation.

These results are in line with predictions from a neural network model previously built

to account for transitions between reconsolidation and extinction with increasing

reexposure time in contextual fear conditioning (Osan et al., 2011). In this model,

modification of existing connection weights is triggered by a protein synthesis-

independent set of processes when there is mismatch between the memory pattern

retrieved by the network (based on previously learned experiences) and the currently

experienced sensory state, as when the prediction of a strong footshock is offset by a

mild one. However, if mismatch is too extensive, as during extinction in the absence of

a footshock, a new attractor is formed in the network at the first reexposure session

already, preventing mismatch-induced weakening of the original memory.

Although we have not tested this explicitly, our results also appear compatible with the

Bayesian inference framework proposed by Gershman et al. (2017), in which the

probability that an experience is attributed to a new latent cause increases in proportion

to the degree of prediction error generated by previous experience. This can explain

why a lower degree of mismatch, such as that caused by reactivation followed by a mild

footshock, can lead to a greater probability of memory updating and decrease the

recovery of fear. Our results are also in accordance with studies showing that greater

mismatch between the training and reactivation sessions, as caused by increasing

reexposure time (Suzuki et al., 2004) or number of non-reinforced CSs (Lee et al., 2006,

Sevenster et al., 2014), can lead to transitions from reconsolidation to extinction, as

assessed by the effects of pharmacological agents affecting these processes.

In this sense, the rationale for the greater effectiveness of deconditioning-update in

reducing fear might be similar to that observed in so-called retrieval extinction

paradigms (Monfils et al., 2009, Kredlow et al., 2016). In this case, the first retrieval

trial, usually consisting of a single CS, is thought to induce retrieval-induced

destabilization of the original memory, as the prediction error generated by this

reexposure is not sufficient to immediately form an extinction memory. Nevertheless,

following this initial retrieval session with a standard extinction procedure leads to

lower fear recovery than when extinction is performed without it, although this effect

has been observed rather inconsistently across studies (Auber et al., 2013, Kredlow et

al., 2016).

Interestingly, deconditioning-update only occurred when reactivation sessions were

spaced across multiple days. When a massed extinction procedure with 24 non-

reinforced CSs was performed within a single extinction session, on the other hand, a

mild footshock at the end of the tones actually prevented within-session extinction, and

increased freezing in a test performed on the following day. This seems to reinforce the

notion that computations linking the degree of prediction error with memory

destabilization occur only at the end of reexposure (Perez-Cuesta et al., 2007, Osan et

al., 2011). It is also in line with the idea that within- and between-session extinction are

distinct processes, with different dynamics and molecular requirements (Plendl and

Wotjak, 2010, Almeida-Corrêa et al., 2015). An interesting challenge for future

theoretical models would be to study whether and how within-session extinction relates

mechanistically to mismatch-induced updating and between-session extinction.

In contrast to our work, Gershman et al. (2013) did find an effect of reducing mismatch

at the start of a single 24-CS extinction session by pairing some of the initial tones with

full-strength footshocks. That said, their effect was only observed in spontaneous

recovery sessions and post-reinstatement sessions, and not within the extinction session

itself. Moreover, the approach to induce lower degrees of mismatch in their experiment

– i.e. gradually reducing the frequency of regular footshocks – was different from ours,

in which this was achieved by providing a low-intensity shock at the end of every tone.

Studying whether both approaches could be combined – by gradually decreasing

footshock intensity, for example – could be an interesting topic for further investigation

of the deconditioning-update effect.

Psychiatric disorders associated with pathological memories are prevalent, cause

important social and economic burden, and approaches to translate basic knowledge of

fear conditioning for potentiation of exposure therapy have met limited success so far in

clinical settings (Monfils and Holmes, 2018). We believe that exploring the principles

of deconditioning-update – i.e. the notion that there is an ideal window of prediction

error to potentiate reexposure effects – is a promising therapeutic avenue that could be

explored in more depth in the setting of trauma-focused psychotherapy. The high

efficacy, long-lasting effects, and drug-free nature of this approach make it particularly

appealing for translation to human memory-related disorders, such as trauma, phobias

and drug abuse.

496   **Methods**

A total of 323 male and female Wistar rats (2-3 months old, weighing between 300 and

400 g) from CREALat the Federal University of Rio Grande do Sul (UFRGS) were

used for all the experiments. Only one animal was excluded (in the experiment in Fig.

1A) due to health conditions. They were housed in plexiglass boxes, with 4 animals per

cage, with block randomization using the cage as subgroup to ensure that each cage

contained at least one animal per experimental group. Sample sizes ranged from 6 to 10,

yielding statistical power of at least 90% to detect an absolute difference of 30% in

freezing time (e.g. from 60 to 30%) with a standard deviation of 15% at $\alpha = 0.05$ in a 2-

tailed t test.

Animals were kept on 12/12h light/dark cycle under controlled temperature (21ºC ±2),

with regular chow and water available *ad libitum* and humidity of approximately 65%.

All the procedures followed the Brazilian ethical guidelines for animal research set by

the National Council for the Control of Experimental Animal Research (CONCEA).

Methods and results are reported according to the revised ARRIVE guidelines (Percie

du Sert et al., 2019).

Auditory fear conditioning

Apparatus: The conditioning chamber (context A) consisted of an illuminated plexiglass

box ($33 \times 22 \times 22$ cm), with a floor grid of parallel 0.1 cm caliber stainless steel bars

spaced 1 cm apart. All context chambers had the same dimensions, but context A had

black walls, whereas context B was vertically striped in black and white. Context C

consisted of white and brown lateral walls and a transparent front wall.

Training session: Rats were habituated for 2 days in context B (10 min each), and 24 h

later were placed in context A, where they received 5 conditioning trials consisting of a

30-s presentation of a 5-kHz, 75-dB tone (CS) that co-terminated with a 0.5 mA (or

1mA in Fig.2F-H), 1-s footshock (US) (5 tone-footshock pairings). The first CS was

presented 2 min into the session, with an interpairing interval of 1 min. One minute after

the final pairing, rats were returned to their home cages.

Reactivation sessions: In daily sessions taking place in context B and starting 48 h after

training (or 40 days later in Fig.2A-E), animals in the no-footshock group received 3

unpaired CSs, while the footshock group (deconditioning-update) received 3 CSs that

co-terminated with a 0.1 (or 0.3-mA in Fig.S1), 1-s shock. The percentage of time

freezing during each tone presentation was quantified and the mean freezing percentage

for the 3 tones was used as a measure of fear. The first CS was presented 2 min into the

session, with an interpairing interval of 1 min. One minute after the final pairing, rats

were returned to their home cages. Most experiments used 4 reactivation sessions,

except for those in Fig. S2 (1 session) and Fig. S4 (3 sessions). In the devaluation

experiment (Fig S3A-B), the protocol was the same, except that the 0.1mA footshocks

were presented in context C without being paired with the tone.

Test session: 24 h after the last reactivation session, both groups were presented with 3

unpaired CSs in context B. The percentage of time freezing during each tone

presentation was quantified, and the average for the 3 tones was used as a fear measure.

The first CS was presented 2 min into the session, with an interpairing interval of 1 min.

One minute after the final pairing, rats were returned to their home cages.

Renewal: 24 h after the test session, animals were placed in context A, where they

received 3 unpaired CSs. The percentage of time freezing during each tone presentation

was quantified, and the average percentage was used as a measure of fear recovery. The

first CS was presented 2 min into the session, with an interpairing interval of 1 min.

One minute after the final pairing, rats were returned to their home cages.

Spontaneous Recovery: 20 days after the renewal session, animals were placed in

context B and received 3 unpaired CSs. The percentage of time freezing during each

tone presentation was quantified and the average was used as a measure of fear

recovery. The first CS was presented 2 min into the session, with an interpairing

interval of 1 min. One minute after the final pairing, rats were returned to their home

cages.

Reinstatement: 24 h after the test session, animals were exposed in to context C for 5

sec, where they received 2-sec 0.7mA footshocks. 24 h later, they were tested for

reinstatement in context B.

Retraining: 24 h after the spontaneous recovery test, rats were submitted to a training

procedure identical to the one described above, except that footshock intensity was

lower (0.3mA). They were then tested 24 h later to assess savings.

Single-session extinction: Animals were placed in context B 48 h after training, where

they received either 12 or 24 CSs depending on the protocol. In the no-footshock group

these tones were unpaired, while the footshock group received tones that co-terminated

with a 0.1 mA footshock. The first CS was presented 2 min into the session, with an

interpairing interval of 1 min. One minute after the final pairing, rats were returned to

their home cages.

Contextual Fear Conditioning

Apparatus: The conditioning chamber consisted of an illuminated plexiglass box (33 ×

22 × 22 cm grid of parallel 0.1 cm caliber stainless steel bars spaced 1 cm apart).

Training session: In the training session, rats were placed in the conditioning chamber

for 3 min before receiving two 2-s, 0.5-mA footshocks separated by a 30-s interval; they

were kept in the conditioning context for an additional 30 s before returning to their

home cage.

Reactivation session: 48 h after the training session, animals were reexposed to the

same conditioning chamber for 4 daily 4-min sessions. Rats from the footshock group

received two pairs of 0.1-mA, 2-s shocks after 180 and 210 s, while the no-footshock

group did not receive any shocks.

Test session and spontaneous recovery: 24 h after the last reactivation session, animals

were re-exposed to the same conditioning chamber for a 4-min test session and the

percentage of time freezing was quantified. 20 days later, the procedure was repeated to

assess spontaneous recovery.

Step-through inhibitory avoidance

Apparatus: The apparatus consists of an automated box (Insight Ltda., Brazil) with two

compartments, a dark one and a lighted one, each measuring 33 X 22 X 22 cm. The

floor consisted of a grid of metal bars with 1-mm diameter placed 1 cm from each other

Training session: Animals were placed in the lighted compartment. When they entered

the dark compartment, the door was closed and the animals received 4 0.5-mA, 1-s

footshocks, with intervals of 5 s between them. They were removed from the box 10 s

after the last footshock.

Reactivation sessions: In daily sessions starting 48 h after training, animals in the

footshock and no-footshock groups were placed in the dark compartment for 30 s, with

no access to the lighted compartment. The footshock group received 2 0.1-mA shocks

(at 25 s and 30 s) while the no-footshock group did not receive any shocks.

Test session: All animals were placed in the lighted compartment and left free to

explore the box. The latency to enter into the dark compartment for the first time and

the time spent in each compartment were counted over a 10-min session and used as

measures of memory.

Behavioral Assessment

Freezing behavior was used as a memory index in the fear conditioning tasks, being

registered with a stopwatch in real time by an experienced observer that was blinded to

the experimental group. Freezing was defined as total cessation of all movements except

those required for respiration.

Open Field

Exploratory activity and anxiety-like behavior were assessed in the open field test in

order to exclude non-specific effects of nimodipine. The apparatus consisted of a

circular arena (90-cm diameter) with 50-cm high walls. The floor was subdivided into

12 quadrants and 3 concentric zones (periphery, intermediary and center). Animals were

exposed to the apparatus for 5 minutes, during which the time spent on the periphery

(thigmotaxis) and the number of crossings between quadrants were measured.

Nimodipine (16 mg/kg) or vehicle was measured intraperitonally 30 minutes before the

test.

Drugs

Nimodipine (Sigma), an antagonist of the L-type voltage-gated calcium channels

(LVGCCs) was dissolved in sterile isotonic saline solution with 8% dimethylsulfoxide

to a concentration of 16 mg/mL. Nimodipine or its vehicle was injected

intraperitoneally at a volume of 1 mL/kg (16 mg/kg) 30 min before memory reactivation

or open field sessions.

Statistical Analysis

Data are expressed as mean ± SEM, always using the animal as the experimental unit.

The statistical tests used and their results are detailed for every experiment in **Tables**

S1-S10; they include two-tailed Student's *t* test; one-way, two-way or three-way

analysis of variance (ANOVA), followed by Tukey's or Bonferroni's post hoc test, when

necessary; and Kruskal-Wallis test, followed by Dunn's post hoc. Values of $p < 0.05$

were considered statistically significant. Unit-level data for all figures is provided as

Supplementary Data.

Computational simulations

Model Network: In order to propose a mechanistic explanation for the experimental

results, we used an adaptation of the attractor network model described in Osan et al

(2011). This Hopfield-like network is capable of storing and retrieving memories using

Hebbian learning rules dependent on neuronal activity, which in turn depends on the

inputs presented to a fully connected network of 100 neurons. In this network, the

activity of each neuron i is determined by

$$\tau \frac{du_i}{d_t} = -u_i + \frac{1}{2}\left(1 + \tanh\left(\sum_{j=1}^{N} w_{ij}u_j + I_i\right)\right)$$

where τ is the neural time constant and u_i represents the level of activation of neuron i

which can vary continuously from 0 to 1 – unlike in the original Hopfield continuous

activity model (Hopfield, 1984), in which activity varies from -1 to 1.

As a fully connected neural network, every neuron i is connected with every neuron j.

For the learning process, the network needs to reinforce the connections between

neurons that fire together, while creating inhibition when presynaptic neuron i is active

and postsynaptic j is silent. Changes in the synaptic weight matrix $W = (w_{ij})$ are

determined by the equation

$$\Delta W = -\gamma W + HLP + MID$$

where Hebbian learning plasticity (HLP) and mismatch-induced degradation are two

independent learning rules (see below) and $0 \leq \gamma \leq 1$ is a time-dependent synaptic decay

factor.

Learning occurs by presenting an input I_i to the circuit, corresponding to sensory

information provided by the environment and/or internal cues, which lead to changes in

the plastic connections between neurons. The cue has a one-to-one topology to the

memory network, with every neuron receiving a cue input that can be either excitatory

or inhibitory. Modifications on the synaptic weight matrix follow the HLP rule,

corresponding to the Hebbian formulation implemented in classic Hopfield networks

and described as

$$HLP = S(u^T * u) - S((1 - u)^T * u)$$

where vector u = (u$_1$,u$_2$,…,u$_N$) is the steady state of the network after input I_i

presentation, while S is a factor representing requirements for Hebbian plasticity, such

as protein synthesis, receptor activation, intracellular signaling and other mechanisms.

When the cue input leads to the retrieval of a previously stored memory, this can lead to

mismatch between the cue input and the retrieved attractor if the two are not the same.

This leads to concomitant activation of the MID learning rule, corresponding to a

memory-updating system akin to that involved in memory destabilization during

reconsolidation and defined by

$$MID = D(m^T * u)$$

where D is a factor representing requirements for memory destabilization (such as

protein degradation and LVGCCs), $m = I_{norm} - u$ is the mismatch vector defined and I_{norm}

is a normalized cue vector. The *MID* term leads to weakening of connections

responsible for the mismatch in order to update the existing memory.

Learning, retrieval and reactivation

Non-overlapping neuron clusters in the network were chosen to represent the training or

extinction contexts (6 neurons each), tone (2 neurons), aversive stimulus/shock (10

neurons) or safety/absence of shock (10 neurons) (**Fig. 5A**). Initially, a pattern

698 representation of a memory unrelated to fear conditioning was presented as a cue to the

699 network. This was followed by a training pattern activating neurons representing

700 context A, tone and shock while inhibiting the remaining ones.

702 Retrieval was evaluated at every training or reactivation session through presentation of

703 a cue activating neurons corresponding to context B and tone, with no input to the

704 remaining neurons. This corresponds to the period in which freezing is assessed (e.g.

705 during the tone itself, shown as 'tone' in **Fig. 5D**), and was modeled with the same cue

706 irrespectively of the presence of shock at the end of reactivation. For the renewal test,

707 the retrieval cue activated neurons representing context A and tone. To quantify

708 memory retrieval, we used the mean activity of neurons representing shock and absence

709 of shock, which was converted to a 'freezing percentage' by dividing the total activity

710 of shock neurons by the total activity of both groups – thus, 100% freezing corresponds

711 to full activation of shock neurons and no activation of non-shock neurons.

713 At the end of each reactivation session, the network underwent a new learning round

714 with a pattern that varied according to the experimental group. To model standard

715 extinction over multiple retrieval sessions, we activated the non-shock neuron cluster

716 along with the neurons representing the extinction context and tone, while inhibiting the

717 remaining ones ('no-footshock reactivation' in **Fig. 5A**). For deconditioning-update, we

718 assumed an intermediate representation between the learning and extinction patterns

719 ('footshock reactivation' in **Fig. 5A**). Synaptic weights were updated according to the

720 activation pattern reached in response to these cues (shown as 'session end' in **Fig. 5D**)

721 Unlike in the original model, no synaptic decay was assumed (i.e. γ was set to 0) and

722 learning strength (as defined by S) was assumed to be smaller during reactivation

sessions in both groups due to the lower intensity of the stimuli, thus allowing

extinction to occur over multiple sessions.

After each learning or reactivation session, the mean synaptic weight between each

cluster of neurons (tone, contexts, shock and non-shock) was calculated by taking the

average of the connections between all presynaptic neurons of a subpopulation and all

postsynaptic neurons of the other subpopulation. This was used to create the synaptic

weight matrix between clusters shown in **Fig. 5E**.

Model parameters

All simulations were performed in MATLAB R2018a (Mathworks) using $N = 100$; $\tau =$

1; $\gamma = 0$; $s_0 = 1$. For training sessions, we set $S = 0.8$, while in reactivation sessions we

used $S = 0.25$. D was set to 0.95 for all sessions, except for reactivations using

nimodipine, in which $D = 0$. Each unit i during learning received an input I_i varying

between -5 and 5. In the deconditioning update group, the aversive shock cluster

received an input I_i of -2.31, while non-shock neurons received 2.31 (corresponding to

t=6 in the transformation used by Osan et al. (2011) to create intermediate patterns). In

the retrieval cue, each targeted neuron had an input $I_j = 1.5$ and 0 for other neurons.

One hundred simulations of each experiment were performed, with different initial

conditions determined by Gaussian noise in the initial weight matrices (with a normal

distribution on [-0.05, 0.05]) and in the neuronal activation at the start of every session

(with a normal distribution on [0, 0.1]). For each simulation, 100 retrieval trials were

run in each session to determine freezing percentage. All results are displayed as the

mean ± S.E.M of these 100 simulations. Matlab code to perform all simulations and

generate figures 5B, 5C and 5E is presented as **Supplementary Code.**

751   **Acknowledgements**

753   This work was supported by the Brazilian government agencies CAPES and CNPq

754   (Universal 2018 - 405100/2018-3), who had no role in the design, analysis or reporting

755   of the study. The authors acknowledge Isabel Cristina Marques Scarello for her kind

756   technical assistance.

758   **Author contributions**

760   L.O.A and B.P. designed this study. B.P. conducted the behavioral, pharmacological

761   and performed statistical analyses. O.B.A. planned and F.E.A. conducted the

762   computational simulations. L.O.A and O.B.A wrote the manuscript. L.O.A supervised

763   the project.

765   **Competing interests**

767   The authors declare no competing interests

769   **References**

Almeida-Corrêa S, Amaral OB (2014) Memory labilization in reconsolidation and
extinction-evidence for a common plasticity system? J Physiol Paris 108:292-
306.

Almeida-Corrêa S, Moulin TC, Carneiro CF, Goncalves MM, Junqueira LS, Amaral
OB (2015) Calcineurin inhibition blocks within-, but not between-session fear
extinction in mice. Learn Mem 22:159-169.
Archbold GE, Bouton ME, Nader K (2010) Evidence for the persistence of contextual
fear memories following immediate extinction. Eur J Neurosci 31:1303-1311.
Auber A, Tedesco V, Jones CE, Monfils MH, Chiamulera C (2013) Post-retrieval
extinction as reconsolidation interference: methodological issues or boundary
conditions? Psychopharmacology (Berl) 226:631-647.
Beckers T, Kindt M (2017) Memory Reconsolidation Interference as an Emerging
Treatment for Emotional Disorders: Strengths, Limitations, Challenges, and
Opportunities. Annu Rev Clin Psychol 13:99-121.
Bouton ME (2002) Context, ambiguity, and unlearning: sources of relapse after
behavioral extinction. Biol Psychiatry 52:976-986.
Bouton ME, Winterbauer NE, Todd TP (2012) Relapse processes after the extinction of
instrumental learning: renewal, resurgence, and reacquisition. Behav Processes
90:130-141.
Bustos SG, Maldonado H, Molina VA (2009) Disruptive effect of midazolam on fear
memory reconsolidation: decisive influence of reactivation time span and
memory age. Neuropsychopharmacology 34:446-457.
Cain CK, Blouin AM, Barad M (2002) L-type voltage-gated calcium channels are
required for extinction, but not for acquisition or expression, of conditional fear
in mice. J Neurosci 22:9113-9121.
Crestani AP, Zacouteguy Boos F, Haubrich J, Ordonez Sierra R, Santana F, Molina JM,
Cassini Lde F, Alvares Lde O, Quillfeldt JA (2015) Memory reconsolidation
may be disrupted by a distractor stimulus presented during reactivation. Sci Rep
5:13633.
De Oliveira Alvares L, Crestani AP, Cassini LF, Haubrich J, Santana F, Quillfeldt JA
(2013) Reactivation enables memory updating, precision-keeping and
strengthening: exploring the possible biological roles of reconsolidation.
Neuroscience 244:42-48.
Duvarci S, Nader K (2004) Characterization of fear memory reconsolidation. J Neurosci
24:9269-9275.
Fernandez RS, Bavassi L, Kaczer L, Forcato C, Pedreira ME (2016) Interference
conditions of the reconsolidation process in humans: The role of valence and
different memory systems. Front Hum Neurosci 10:641.
Flavell CR, Barber DJ, Lee JL (2011) Behavioural memory reconsolidation of food and
fear memories. Nat Commun 2:504.
Frankland PW, Ding HK, Takahashi E, Suzuki A, Kida S, Silva AJ (2006) Stability of
recent and remote contextual fear memory. Learn Mem 13:451-457.
Gershman SJ, Jones CE, Norman KA, Monfils MH, Niv Y (2013) Gradual extinction
prevents the return of fear: implications for the discovery of state. Front Behav
Neurosci 7:164.
Gershman SJ, Monfils MH, Norman KA, Niv Y (2017) The computational nature of
memory modification. Elife 6.
Haubrich J, Crestani AP, Cassini LF, Santana F, Sierra RO, Alvares Lde O, Quillfeldt
JA (2015) Reconsolidation allows fear memory to be updated to a less aversive
level through the incorporation of appetitive information.
Neuropsychopharmacology 40:315-326.

Hopfield JJ (1984) Neurons with graded response have collective computational properties like those of two-state neurons. Proc Natl Acad Sci USA 81:3088-3092.

Kredlow MA, Unger LD, Otto MW (2016) Harnessing reconsolidation to weaken fear and appetitive memories: A meta-analysis of post-retrieval extinction effects. Psychol Bull 142:314-336.

Lee JL (2009) Reconsolidation: maintaining memory relevance. Trends Neurosci 32:413-420.

Lee JL, Flavell CR (2014) Inhibition and enhancement of contextual fear memory destabilization. Front Behav Neurosci 8:144.

Lee JL, Milton AL, Everitt BJ (2006) Reconsolidation and extinction of conditioned fear: inhibition and potentiation. J Neurosci 26:10051-10056.

Maren S, Phan KL, Liberzon I (2013) The contextual brain: implications for fear conditioning, extinction and psychopathology. Nat Rev Neurosci 14:417-428.

Milekic MH, Alberini CM (2002) Temporally graded requirement for protein synthesis following memory reactivation. Neuron 36:521-525.

Monfils MH, Cowansage KK, Klann E, LeDoux JE (2009) Extinction-reconsolidation boundaries: key to persistent attenuation of fear memories. Science 324:951-955.

Monfils MH, Holmes EA (2018) Memory boundaries: opening a window inspired by reconsolidation to treat anxiety, trauma-related, and addiction disorders. Lancet Psychiatry 5:1032-1042.

Nader K, Einarsson EO (2010) Memory reconsolidation: an update. Ann N Y Acad Sci 1191:27-41.

Nader K, Schafe GE, Le Doux JE (2000) Fear memories require protein synthesis in the amygdala for reconsolidation after retrieval. Nature 406:722-726.

Osan R, Tort AB, Amaral OB (2011) A mismatch-based model for memory reconsolidation and extinction in attractor networks. PLoS One 6:e23113.

Pedraza LK, Sierra RO, Lotz FN, de Oliveira Alvares L (2018) Periodical reactivation under the effect of caffeine attenuates fear memory expression in rats. Sci Rep 8:7260.

Percie du Sert N, Hurst V, Ahluwalia A, et al. (2019). The ARRIVE guidelines 2019: updated guidelines for reporting animal research. bioRxiv, doi: 10.1101/703181

Perez-Cuesta LM, Hepp Y, Pedreira ME, Maldonado H (2007) Memory is not extinguished along with CS presentation but within a few seconds after CS-offset. Learn Mem 14:101-108.

Plendl W, Wotjak CT (2010) Dissociation of within- and between-session extinction of conditioned fear. J Neurosci 30:4990-4998.

Rescorla RA, Heth CD (1975) Reinstatement of fear to an extinguished conditioned stimulus. J Exp Psychol Anim Behav Process 1:88-96.

Rescorla R (1973). Effects of US habituation following conditioning. J Comp Physiol Psychol 82:137–143.

Sevenster D, Beckers T, Kindt M (2014) Prediction error demarcates the transition from retrieval, to reconsolidation, to new learning. Learn Mem 21:580-584.

Sierra RO, Cassini LF, Santana F, Crestani AP, Duran JM, Haubrich J, de Oliveira Alvares L, Quillfeldt JA (2013) Reconsolidation may incorporate state-dependency into previously consolidated memories. Learn Mem 20:379-387.

Suzuki A, Josselyn SA, Frankland PW, Masushige S, Silva AJ, Kida S (2004) Memory reconsolidation and extinction have distinct temporal and biochemical signatures. J Neurosci 24:4787-4795.

Suzuki A, Mukawa T, Tsukagoshi A, Frankland PW, Kida S (2008) Activation of
LVGCCs and CB1 receptors required for destabilization of reactivated
contextual fear memories. Learn Mem 15:426-433.
Wang SH, de Oliveira Alvares L, Nader K (2009) Cellular and systems mechanisms of
memory strength as a constraint on auditory fear reconsolidation. Nat Neurosci
12:905-912.

878