Shifting from fear to safety through deconditioning-update: a novel approach to attenuate fear memories

Bruno Popik[1], Felippe Espinelli Amorim[2], Olavo Boher Amaral[2] and Lucas de Oliveira Alvares[1*]

[1]Neurobiology of Memory Lab, Biophysics Department, Biosciences Institute, Federal University of Rio Grande do Sul, Porto Alegre, Brazil. [2]Institute of Medical Biochemistry Leopoldo de Meis, Federal University of Rio de Janeiro, Rio de Janeiro, Brazil,

Corresponding Author: Lucas de Oliveira Alvares, Laboratório de Neurobiologia da Memória, Departamento de Biofísica, Instituto de Biociências, Universidade Federal do Rio Grande do Sul (UFRGS), Av. Bento Gonçalves 9500, Prédio 43422, Sala 210, CEP 91.501-970, Porto Alegre, Rio Grande do Sul, Brasil. E-mail: lucas.alvares@ufrgs.br

Abstract

Traumatic memories are at the heart of psychiatric disorders such as PTSD. Here, we present a new behavioral approach in rats that robustly attenuates aversive memories in a long-lasting way. This method consists of "deconditioning" animals previously trained to associate a tone with a strong footshock by replacing it with a much weaker one during memory retrieval. Our results indicate that deconditioning-update is more effective than traditional extinction in reducing fear responses; moreover, such effects are long-lasting and resistant to renewal and spontaneous recovery. Remarkably, this strategy overcame important boundary conditions for memory updating, such as remote or very strong traumatic memories. The same beneficial effect was found in other types of fear-related memories. Deconditioning was mediated by LVGCCs and is consistent with computational accounts of mismatch-induced memory updating. Our results suggest that shifting from fear to a non-fear memory through deconditioning-update is a promising approach to remove pathological memories.

Introduction

Exposure to strong aversive experiences can lead to the formation of enduring traumatic memories, which can trigger many debilitating psychiatric disorders including posttraumatic stress disorder (PTSD), phobias and depression [1]. These pathological memories have a high prevalence and cause tremendous social and economic burden worldwide [1]. Exposure-based therapies and pharmacological treatments are the most common interventions to attenuate these pathological memories [2], but a significant proportion of patients are not responsive to these approaches, with only 20–30% showing full remission [3,4].

During exposure-based therapies, traumatic reminders (conditioned stimuli, CS) are repeatedly presented in a safe environment in the absence of trauma (unconditioned stimulus, US), leading to a progressive reduction in fear expression reminiscent of extinction of fear conditioning. However, extinction does not erase the original memory; instead, it induces new learning that transiently inhibits fear expression [5]. Therefore, the fear memory typically reemerges with the passage of time (spontaneous recovery, SR), exposure to the US (reinstatement), or when the CS is presented outside of the extinction context (renewal) [6-8]. Thus, behavioral strategies that weaken traumatic memories and reduce memory recovery can be relevant for improving the effectiveness of therapies and alleviate the burden of patients.

Memory is a dynamic process that allows for adaptation to the demands of a continuously changing environment. The ability to update old memories in accordance

with new experiences is crucial for maintaining their relevance over time. Particularly, it has been shown that after retrieval (or reactivation), memories may undergo a cycle of labilization and restabilization commonly known as reconsolidation [9]. The labile state induced by this process can thus allow changes in memory strength and/or content [10].

Reconsolidation has been described in several experimental paradigms and species, from invertebrates to humans, suggesting that it might a fundamental property of memories [11]. Beyond its biological role in memory updating, this process also opens a window of opportunity as a potential therapeutic strategy to modify pathological memories. Several studies in the last decades have attempted to pharmacologically disrupt the reconsolidation of traumatic memories [12]. One of the advantages of interfering with reconsolidation mechanisms is that fear reduction through this strategy, at least in some situations, can be less susceptible to spontaneous or induced recovery than traditional extinction [13,14]. However, most treatments that interfere with memory reconsolidation are toxic and cannot be readily administered to humans. Moreover, in spite of almost two decades of research on memory reconsolidation, the evidence for the efficacy of reconsolidation-blocking treatments in clinical settings is still limited [15].

Research on reconsolidation-extinction boundaries suggests that the transition from one process to the other depends on the degree of mismatch between the original memory and the reactivation experience. Many studies have suggested that some degree of mismatch or prediction error is necessary for reconsolidation to occur [16,17]; however, if prediction error is too high, extinction may occur instead [18,19]. Computational models suggest that, in models in which prediction error is low, memory updating/reconsolidation mechanisms are preferentially engaged, as the new experience

is recognized as a new instance of the former one; however, as mismatch increases, the chances of it being attributed to a new latent cause [20] or forming a new attractor in a neural network [21] increases.

This framework suggests that lowering the degree of mismatch between learning and reexposure might conceivably potentiate memory updating during extinction. This has been the rationale behind so-called retrieval-extinction procedures [22] and has also been explored in short-term extinction protocols [23]. In this work, we propose a novel approach for long-term attenuation of traumatic memories that we term "deconditioning-update". This strategy consists in weakening fear memories by updating the aversive information, substituting the original CS by a mildly aversive stimulus during the plastic state induced by reactivation.

Results

In order to approach our hypothesis, male Wistar rats were trained in auditory fear conditioning, where they received 5 conditioning trial tones (CS) that co-terminated with a 0.5-mA, 2-s foot-shock (US). On days 3 to 5 (reactivation), animals received 3 CSs during a 400-s daily session in a different context. In the no-footshock group, they were unpaired, while in the deconditioning-update group, each tone co-terminated with a 0.1-mA footshock. A third group remained in the homecage (control group). On day 6, both groups were tested in the reactivation context with 3 unpaired CSs. On day 7, animals were tested in the training context for renewal, and, on day 27, they were retested for spontaneous recovery (**Fig.1A**).

Over the course of reactivation sessions, both groups presented a decrease in freezing, but this was more marked in the deconditioning-update group (**Fig 1B**). In the test session, it presented a decrease in freezing of around 70% in comparison to the no-footshock group, and 80% in comparison to the homecage control group (**Fig 1C**). Animals in the deconditioning-update group also had lower freezing responses in the renewal (**Fig 1D**) and spontaneous recovery (**Fig. 1E**) tests, although it should be noted that memory recovery was not observed in the no-footshock group either, perhaps due to a ceiling effect caused by insufficient extinction (for complete statistics, see **Table S1**). When performing the same experiment, but using a single reactivation session, no difference was found in the test (**Fig. S1**), suggesting that deconditioning-update requires several sessions to take place.

Next, we asked whether the deconditioning-update effect would also be efficient in reducing fear in female rats. As observed in males, females from the deconditioning-update group showed lower freezing levels throughout the reactivation sessions, as well as in the test session, when compared with the control group (**Fig 1H**). The same pattern was maintained in the renewal and spontaneous recovery tests (**Fig 1I** and **1J**, respectively). As a further way to test for persistence of the original memory, we performed a retraining session in the original context with 3 0.5-mA CS-US pairings 24h after the spontaneous recovery test to assess savings. The deconditioning-update group had lower freezing compared with the other groups in a subsequent test session in the extinction context, suggesting that our protocol also lowers the rate of re-acquisition of an aversive memory (**Fig. 1K**).

Figure 1 - Weakening fear memory through deconditioning-update training. (**A**) Experimental design: male rats were fear-conditioned with five tone-shock pairings (context B; 5CS+US, 0.5mA). 48h later, the no-footshock and footshock (deconditioning-update*)* groups were exposed to 4 daily reactivation sessions (context A). After this, animals underwent test (context A), renewal (context B) and spontaneous recovery (context A) sessions. Grey rectangles represent context A, while black circles represent context B. (**B**) Freezing levels during reactivation sessions. Rats exposed to weak footshocks during reactivation sessions showed a significant reduction in freezing responses, maintained during the test (**C**), renewal (**D**) and spontaneous recovery (**E**) sessions. For statistics, see **Table S1** (**F**) Experimental design: female rats were fear-conditioned (context B; 5CS+US, 0.5mA). 48 hours later, the no-footshock and footshock groups were exposed to 4 daily reactivation sessions (context A). After this, all groups underwent test, renewal, and spontaneous recovery sessions. Animals were reconditioned (context B; 3CS+US, 0.5mA) on the next day and retested 24h later. (**G**) Freezing levels during memory reactivation. Female rats exposed to weak footshocks showed a significant reduction in freezing responses, maintained during the test (**H**), renewal (**I**), spontaneous recovery (**J**) and retraining test (**K**) sessions. Bars represent mean ± SEM. Statistical comparisons were performed using a two-way repeated-measures ANOVA followed by a Bonferroni post-hoc (reactivation sessions) or one-way ANOVA followed by Tukey post hoc (test, renewal, spontaneous recovery, and retraining test). * $p < 0.05$; ** $p < 0.005$; *** $p < 0.0005$; **** $p < 0.0001$.

Boundary conditions such as training intensity and memory age have been reported to prevent fear memories from being modified. Protocols with high training intensity make memory less prone to interference by pharmacological agents in the reconsolidation window [24-27], while older memories are also less susceptible to modification than recent ones [26-28]. Thus, our next experiments investigated whether deconditioning-update could attenuate fear expression in these cases.

First, we trained the animals in the same protocol described above, but starting reactivations 40 days after conditioning (**Fig. 2A**). Once more, the deconditioning-

update group showed lower freezing expression throughout the reactivation sessions (**Fig. 2B**). 24 h after the last reactivation, both reactivated groups presented a comparable decrease in freezing behavior when compared with controls (**Fig. 2C**). However, in the renewal test, as well as in the spontaneous recovery test, fear expression reemerged in the no-footshock group, while the deconditioning-update group maintained its low freezing levels (**Fig. 2D** and **2E**, respectively; **Table S2**).

Next, we tested whether a stronger fear memory would constrain the deconditioning-update effect by training animals with 5 CS-US pairings using 1mA shocks, while maintaining the rest of the protocol unchanged. In spite of the stronger shock intensity, the deconditioning-updating group presented reduced freezing levels in reactivation sessions when compared to the no-footshock group (**Fig. 2G**). These lower freezing levels were maintained in the test, renewal and spontaneous recovery sessions, in which robust recovery was observed in no-footshock animals, but not in the deconditioning-update group (**Fig. 2H**, **2I** and **2J**, respectively; **Table S2**). Similar results were observed in females in a slightly modified protocol with 3 reactivation sessions (**Fig. S2**). These experiments suggest that the deconditioning-update induces a plastic state, allowing the fear memory trace to be altered even in boundary conditions that usually constrain memory updating [29].

Figure 2 - Deconditioning-update weakens both remote and strong fear memories. (**A**) Experimental design for remote memory: rats were fear-conditioned with five tone-shock pairings (context B; 5CS+US, 0.5mA). Starting 40 day later, the no-footshock and footshock (deconditioning-update) groups were exposed to daily reactivation sessions (context A). Then, all groups underwent test (context A), renewal (context B) and spontaneous recovery (context A) sessions. Grey rectangles represent context A and black circles represent context B. (**B**) Freezing levels during reactivation sessions. Rats exposed to weak footshocks during reactivation sessions showed similar freezing levels to no-footshock animals during the test session (C) and lower freezing leves at the renewal (**D**) and spontaneous recovery (**E**) ones. For statistics, see **Table S2** (**F**) Experimental design for strong training (5CS+US, 1mA). (**G**) Freezing levels during reactivation sessions. Rats exposed to weak footshocks during reactivation sessions showed a significant reduction in freezing responses that was maintained during the test (**H**), renewal (**I**) and spontaneous recovery (**J**) sessions. Bars represent mean ± SEM. Statistical comparisons are performed using two-way repeated-measures ANOVA followed by a Bonferroni post-hoc (reactivation sessions) or one-way ANOVA followed by a Tukey post-hoc (test and spontaneous recovery). * $p < 0.05$; ** $p < 0.005$; *** $p < 0.0005$; **** $p < 0.0001$.

In order to investigate whether the deconditioning-update approach is effective in attenuating other types of aversive memories, we also trained animals in different fear-related tasks. First, we used a contextual fear conditioning paradigm, which is known to involve a partially distinct set of brain regions, including the prefrontal cortex and hippocampus [30]. Animals were placed in the conditioning chamber for 3 min before receiving two 0.5-mA, 2-s footshocks separated by a 30 s interval. On days 3 to 6, rats were reexposed to the same context for 4 min, with those in the deconditioning-update group receiving weak footshocks (0.1mA, 2 s) 3 min after being placed in the chamber (**Fig. 3A**). The deconditioning-update group had lower fear expression during reactivations (**Fig. 3B**) and maintained these lower freezing levels compared with the other groups in the test (**Fig. 3C**). The same pattern was observed in the spontaneous

recovery test, performed 20 days later, suggesting that the decrease in freezing caused by deconditioning-update is long-lasting (**Fig. 3D**; **Table S3**).

A further set of animals underwent the step-through inhibitory avoidance task, in which training consists in applying 4 0.5mA, 1-s footshocks with 5-s intervals when the animal enters the dark compartment of a shuttle box. During reactivations, animals were placed in the dark compartment for 30 s, with those in the deconditioning-update group receiving 2 0.1-mA shocks. In the test session, the animals were put in the light compartment, and the latency to enter the dark compartment was measured (**Fig. 3E**). The deconditioning-update group showed a much shorter latency to reach the dark chamber (**Fig. 3F**) and spent more time in the dark compartment over the 10-minute session compared to the other groups (**Fig 3G**; **Table S3**), suggesting that memory was more robustly updated in this group during reactivation. Taken together, these results suggest that the deconditioning-update strategy is effective in weakening distinct types of fear-related memories.

Figure 3 - Deconditioning-update weakens fear memory in different behavioral tasks. (**A**) Experimental design in contextual fear conditioning: rats were fear-conditioned with five contextual-shock pairings (4 minutes CS + 4 US, 0.5mA). Starting 48h later, the no-footshock and footshock groups were exposed to daily reactivation sessions. 24h after the last reactivation, all groups were tested; 20 days later, they were tested for spontaneous recovery. (**B**) Freezing levels during reactivation sessions. Rats exposed to weak footshocks during reactivation sessions showed a significant reduction in freezing responses maintained during the test (**C**) and spontaneous recovery (**D**) sessions. For statistics, see **Table S3** (**E**) Experimental design in inhibitory avoidance: rats were placed in the light compartment and received footshocks (4 US, 0.5mA) upon

entering the dark one. Starting 48h later, the non-foot shock and footshock groups were exposed to daily 30-s reactivation sessions in the dark compartment; 24h after the last reactivation, all groups were tested. Rats exposed to weak footshocks during reactivation sessions showed lower latencies to cross to the dark compartment (**F**) and spent more time in it (**G**). Bars represent mean ± SEM or median with interquartile range (in F and G). Statistical comparisons for contextual fear conditioning are performed using two-way repeated-measures ANOVA followed by a Bonferroni post-hoc (reactivation sessions) or one-way ANOVA followed by a Tukey post-hoc (test, renewal, and spontaneous recovery). For inhibitory avoidance, a Kruskal-Wallis test followed by a Dunn post-hoc was performed. $*p < 0.05$; $**p < 0.005$; $***p < 0.0005$; $****p < 0.0001$.

To address whether the deconditioning-update effects would also be observed within a single, long-lasting extinction session, we trained animals and exposed them 48h later to a session containing 12 CSs, with the deconditioning-update group receiving a 0.1-mA footshock at the end of each tone. Fear reduction was limited and largely similar across groups in the extinction session and in the subsequent tests (**Fig. S3**), albeit with slightly lower renewal in the deconditioning-update group, suggesting that the pairing of the CS with a weak footshock is not as effective in accelerating single-session extinction. In order to further explore this possibility, we subjected animals to a 24-tone single-session extinction protocol (**Fig. 4A**). This led to robust fear reduction both within the extinction session (**Fig 4B**) and in a subsequent test (**Fig 4C**) in the no-footshock group, while freezing remained largely unchanged in the deconditioning-update group. However, fear memory reemerged in the renewal and spontaneous recovery test among no-footshock animals (**Fig 4D** and **4E**; **Table S4**), as typically occurs with extinction learning. These results show that the presence of a weak shock at the end of every CS not only does not enhance extinction occurring over a single behavioral session, but can actually impair it in the short-term.

Other studies have shown that activation of LVGCCs is necessary both for destabilizing a reactivated memory during reconsolidation [26,31-33] and for some forms of extinction (Cain et al., 2002). Thus, we used the LVGCC antagonist nimodipine as a pharmacological tool to investigate whether deconditioning-update involved memory destabilization mechanisms [32-35]. Animals were trained and divided into four groups: no-footshock + vehicle, no-footshock + nimodipine, deconditioning-update + vehicle and deconditioning-update + nimodipine, with nimodipine or vehicle administered systemically 30 min before each reactivation (**Fig 4F)**. Nimodipine attenuated freezing decrease in both behavioral protocols, suggesting that the drug impaired both deconditioning-update and regular extinction (**Fig. 4G** and **4H**). In the renewal and spontaneous recovery, high freezing levels reemerged both in the no-footshock vehicle group and in nimodipine-treated animals, while the deconditioning-update vehicle group maintained a lower freezing level (**Fig. 4I** and **4J**; **Tables S4**). This suggests that the deconditioning-update approach is mediated by memory destabilization processes requiring Ca^{++} influx through LVGCCs.

Figure 4 - Deconditioning-update is based on memory destabilization mechanisms. (**A**) Experimental design: rats were fear-conditioned with 5 tone-shock pairings (context B; 5CS+US, 0.5mA). 48h later, the no-footshock and foot shock groups underwent a single extinction session (context A, 24 CSs), followed by test (context A), renewal (context B) and spontaneous recovery (context A) sessions. (**B**) Freezing levels during extinction. Weak footshocks impaired extinction within the session and in the test session (**C**), but not in renewal (**D**) or spontaneous recovery (**E**). For statistics, see **Table S4**. (**F**) Experimental design: rats were fear-conditioned (context B; 5CS+US, 0.5mA). 48 hours later, all animals underwent daily reactivation sessions (context A), receiving nimodipine (16 mg/kg, i.p.) or vehicle 30 min before each one. They then underwent test (context A), renewal (context B) and spontaneous recovery (context A) sessions. Nimodipine prevented freezing decrease across reactivation sessions in both groups (**G**). Freezing was similar between groups in the test session (**H**), but was lower in the vehicle-footshock group in the renewal (**I**) and spontaneous recovery (**J**) sessions. Bars represent mean ± SEM. Statistical comparisons are performed using two-way repeated-measures ANOVA followed by Bonferroni post-hoc (extinction), one-way ANOVA followed by Tukey post-hoc (test, renewal, and spontaneous recovery following extinction), three-way repeated-measures ANOVA followed by Bonferroni post-hoc (reactivation sessions with nimodipine/vehicle) and two-way ANOVA followed by Bonferroni post-hoc (test, renewal, and spontaneous recovery following nimopidine/vehicle). * $p < 0.05$; ** $p < 0.005$; *** $p < 0.0005$; **** $p < 0.0001$ in between-group comparisons.

One explanation for our findings is that pairing the CS with a weak footshock could lead to a smaller degree of prediction error during reexposure sessions, biasing them towards memory updating as opposed to new learning. Computational models using neural networks [21] or Bayesian approaches [20] have explored how different degrees of mismatch between stored memories and new experiences can lead to these two outcomes, suggesting lower degrees of mismatch or prediction error could lead to greater destabilization of stored memories. With this in mind, we used an adaptation of one of these models [21] to explore if its framework could account for our main results –

i.e. accelerated extinction over multiple sessions and lower memory recovery when mismatch is reduced during reexposure. This fully connected Hopfield-like network (Hopfield et al., 1984). of 100 neurons is capable of storing and retrieving memories using Hebbian learning rules dependent on the inputs presented, which included neurons representing cue tone and context information, as well as shock/non-shock information (**Fig. 5A**). Additionally, the network also updates synaptic weights according to mismatch between a cue input and the retrieved network pattern.

As shown in Fig. **5B,** a lower degree of mismatch in the reexposure pattern, caused by the partial activation of shock-related neurons (as would be expected from deconditioning-update), can indeed lead to a greater reduction in subsequent fear behavior than a "pure extinction" pattern (sharing only the tone information with the original memory). This effect was sensitive to blockade of the mismatch-induced degradation term – the model equivalent of blocking memory destabilization mechanisms such as LVGCCs with a drug like nimodipine (**Fig. 5C**). This occurs because the "minor footshock" pattern leads to recovery of the original memory in the initial sessions (**Fig 5D**), causing mismatch-induced updating of the weights supporting these memories and weakening of the mutual connections between shock and context neurons **(Fig. 5E)**. On the other hand, no-footshock extinction leads to immediate formation of a new attractor, leaving the connection weights related to the shock memory unaltered throughout the extinction process.

Figure 5 – Lower mismatch accelerates extinction and decreases renewal in a neural network model. (A) Cue inputs presented to the network during training (shock memory), reexposure (with or without footshock) and test sessions (consisting of the tone and either context B (test) or A (renewal)). **(B)** Extinction over multiple sessions using the no-footshock (red) or footshock (blue) cue. Bars represent freezing, expressed as the activity ratio between shock neurons and the sum of shock and non-shock neurons in response to the test cue. After the cue, memory is updated according to the corresponding reexposure pattern in days 1-4. **(C)** Effect of LVGCC blockade (i.e. setting MID to 0). Removing the degradation term blocks deconditioning-update (blue

bars), but not regular extinction. **(D)** Network activity in retrieval tests during tone presentations (e.g. cued tests) and at the end of reexposure, as well as on test and retrieval sessions. Lower mismatch (i.e. weak footshock) leads to retrieval of the original pattern on the first days, leading to memory updating through mismatch-induced degradation and lower retrieval on subsequent tests. **(E)** Mean synaptic weights between different neuronal groups after training and at the end of each extinction session. Heat map represents the connection from the groups in the Y axis to those in the X axis in the no-footshock and footshock groups. Deconditioning-update leads to weakening of connections between context and shock neurons and of their inhibitory connections to other neurons. On no-footshock extinction, an extinction memory is formed with sparing of the shock representation.

Discussion

Taken as a whole, our findings demonstrate that presenting a tone followed by a weak footshock (deconditioning-update) was more effective in reducing fear memory than presenting a tone in the absence of shock (extinction training). Attenuation of fear responses following deconditioning-update was robust, long-lasting, and less sensitive to renewal and spontaneous recovery. Remarkably, this strategy was also effective in reducing fear within boundary conditions in which memories have been described to be less sensitive to modification (e.g., very strong training protocols and remote memories). The same beneficial effect was found in other types of fear-related memory (contextual fear conditioning and inhibitory avoidance tasks), and was sensitive to pharmacological blockade of LVGCCs. We suggest that the CS–US association is weakened during the deconditioning-update approach, leading to lower fear expression. Considering that (i) fear reduction is long-lasting, without spontaneous recovery, renewal and savings, (ii weak-footshock pairings do not have the same effect in single-session extinction protocols, and (iii) this effect is prevented by LVGCC antagonism,

we suggest that deconditioning-update is mediated by the memory destabilization effects commonly associated with reconsolidation.

These results are in line with predictions from a neural network model previously built to account for transitions between reconsolidation and extinction with increasing reexposure time in contextual fear conditioning [21]. In this model, modification of existing connection weights is triggered by a protein synthesis-independent set of processes when there is mismatch between the memory pattern retrieved by the network (based on previously learned experiences) and the currently experienced sensory state, as when the prediction of a strong footshock is offset by a mild one. However, if mismatch is too extensive, as during extinction in the absence of a footshock, a new attractor is formed in the network at the first reexposure session already, preventing mismatch-induced weakening of the original memory.

Although we have not tested this explicitly, our results also seem to be compatible with the Bayesian inference framework proposed by Gershman et al. [20], in which the probability that an experience is attributed to a new latent cause increases in proportion to the degree of prediction error generated by previous experience. This can explain why a lower degree of mismatch, such as that caused by reactivation followed by a mild footshock, can lead to a greater probability of memory updating and decrease the recovery of fear. Our results are also in accordance with studies showing that greater mismatch between the training and reactivation sessions, as caused by increasing reexposure time [18] or number of non-reinforced CSs [19,37], can lead to transitions from reconsolidation to extinction, as assessed by the effects of pharmacological agents affecting these processes.

In this sense, the rationale for the greater effectiveness of deconditioning-update in reducing fear might be similar to that observed in so-called retrieval extinction paradigms [14,22]. In this case, the first retrieval trial, usually consisting of a single CS, is thought to induce retrieval-induced destabilization of the original memory, as the prediction error generated by this reexposure is not sufficient to immediately form an extinction memory. Nevertheless, following this initial retrieval session with a standard extinction procedure leads to lower fear recovery than when extinction is performed without it, although this effect has been observed rather inconsistently across studies [22,38].

Interestingly, deconditioning-update only occurred when reactivation sessions were spaced across multiple days. When a massed extinction procedure with 24 non-reinforced CSs was performed within a single extinction session, on the other hand, a mild footshock at the end of the tones actually prevented within-session extinction, and increased freezing in a test performed on the following day. This seems to reinforce the notion that computations linking the degree of prediction error with memory destabilization occur only at the end of reexposure [21,39]. It is also in line with the idea that within- and between-session extinction are distinct processes, with different dynamics and molecular requirements [40,41]. An interesting challenge for future theoretical models would be to study whether and how within-session extinction relates mechanistically to mismatch-induced updating and between-session extinction.

In contrast to our work, Gershman et al. [23] did find an effect of reducing mismatch at the start of a single 24-CS extinction session by pairing some of the initial tones with full-

strength footshocks. That said, their effect was only observed in spontaneous recovery sessions and post-reinstatement sessions, and not within the extinction session itself. Moreover, the approach to induce lower degrees of mismatch in their experiment – i.e. gradually reducing the frequency of regular footshocks – was different from ours, in which this was achieved by providing a low-intensity shock at the end of every tone. Studying whether both approaches could be combined – by gradually decreasing footshock intensity, for example – could be an interesting topic for further investigation of the deconditioning-update effect.

Psychiatric disorders associated with pathological memories are prevalent, cause important social and economic burden, and approaches to translate basic knowledge of fear conditioning for potentiation of exposure therapy have met limited success so far in clinical settings [15]. We believe that exploring the principles of deconditioning-update – i.e. the notion that there is an ideal window of prediction error to potentiate reexposure effects – is a promising therapeutic avenue that could be explored in more depth in the setting of trauma-focused psychotherapy. The high efficacy, long-lasting effects, and drug-free nature of this approach make it particularly appealing for translation to human memory-related disorders, such as trauma, phobias and drug abuse.

Methods

Male and female Wistar rats (2-3 months old, weighing approximately 350 g) from the Biophysics Department at the Federal University of Rio Grande do Sul (UFRGS) were used for all the experiments. They were housed in plexiglass boxes, with 4 animals per cage. Animals were kept on 12/12h light/dark cycle under controlled temperature (21ºC

±2), with food and water available *ad libitum*. All the procedures followed the Brazilian ethical guidelines for animal research.

<u>Auditory fear conditioning</u>

Apparatus: The conditioning chamber (context B) consisted of an illuminated plexiglass box (33 × 22 × 22 cm grid of parallel 0.1 cm caliber stainless steel bars spaced 1 cm apart). The context A has black walls whereas the context B is vertically striped in black and white.

Training session: Rats were habituated for 2 days in context A (10 min each), and 24 h later, they were placed in a different context (context B), where they received 5 conditioning trials consisting of a 30-s presentation of a 5-kHz, 75-dB tone (CS) that co-terminated with a 0.5-mA, 1-s footshock (US) (five tone-footshock pairings). The first CS was presented 2 min into the session, with an interpairing interval of 1 min. One minute after the final pairing, rats were returned to their home cages.

Reactivation sessions: In daily sessions taking place in context A and starting 48 h after training, animals of the no-footshock group received 3 unpaired CSs, while the footshock group (deconditioning-update) received 3 CSs that co-terminated with a 0.1-mA, 1-s shock. The percentage of time freezing during each tone presentation was quantified and the mean freezing percentage for the 3 tones was used as a measure of fear. The first CS was presented 2 min into the session, with an interpairing interval was 1 min. One minute after the final pairing, rats were returned to their home cages.

Test session: 24 h after the last reactivation session, both groups were presented with 3 unpaired CSs in context A. The percentage of time freezing during each tone presentation was quantified, and the average for the 3 tones was used as a fear measure. The first CS was presented 2 min into the session, with an interpairing interval of 1 min. One minute after the final pairing, rats were returned to their home cages.

Renewal: 24h after the test session, animals were placed in context B, where they received 3 unpaired CSs. The percentage of time freezing during each tone presentation was quantified and the average percentage was used as a measure of fear recovery. The first CS was presented 2 min into the session, with an interpairing interval of 1 min. One minute after the final pairing, rats were returned to their home cages.

Spontaneous Recovery: 20 days after the renewal session, animals were placed in context A and received 3 unpaired CSs. The percentage of time freezing during each tone presentation was quantified and the average was used as a measure of fear recovery. The first CS was presented 2 min into the session, with an interpairing interval of 1 min. One minute after the final pairing, rats were returned to their home cages.

Single-session extinction: Animals were placed in context A 48h after training, where they received either 12 or 24 CSs depending on the protocol. In the no-footshock group these tones were unpaired, while the footshock group received tones that co-terminated with a 0.1 mA footshock. The first CS was presented 2 min into the session, with an interpairing interval was 1 min. One minute after the final pairing, rats were returned to their home cages.

Step-through inhibitory avoidance

Apparatus: *The* apparatus consists of an automated box (Insight Ltda., Brazil) with two compartments, a dark one and a lighted one, each measuring 33 X 22 X 22 cm. The floor consisted of a grid of metal bars with 1-mm diameter placed 1 cm from each other.

Training session: Animals were placed in the lighted compartment. When they entered the dark compartment, the door was closed and the animals received 4 0.5-mA, 1-s footshocks, with intervals of 5 s between them. They were removed from the box 10 s after the last footshock.

Reactivation sessions: In daily sessions starting 48 h after training, the animals in the footshock group and in the no-footshock group were placed in the dark compartment for 30 s, with no access to the light compartment. The footshock group received 2 0.1-mA shocks (sec 25 and 30) while the no-footshock group did not receive any shocks.

Test session: All animals were placed in the lighted compartment and left free to explore the box. The latency to enter into the dark compartment for the first time and the time spent in each compartment were counted over a 10-min session and used as measures of memory.

Contextual Fear Conditioning

Apparatus: The conditioning chamber consisted of an illuminated plexiglass box (33 × 22 × 22 cm grid of parallel 0.1 cm caliber stainless steel bars spaced 1 cm apart).

Training session: In the training session, rats were placed in the conditioning chamber for 3 min before receiving two 2-s, 0.5-mA footshocks separated by a 30s interval; they were kept in the conditioning context for an additional 30s before returning to their home cage.

Reactivation session: 48 h after the training session, animals were reexposed to the same conditioning chamber for 4 daily 4-min sessions. Rats from the footshock group received two pairs of 0.1-mA, 2-s shock after 3 min, and again 30 s later, while the no-footshock group did not receive any shocks.

Test session and Spontaneous Recovery: 24 h after the last reactivation session, animals were re-exposed to the same conditioning chamber for a 4-min test session and the percentage of time freezing was quantified. 20 days later, the procedure was repeated to assess spontaneous recovery.

Behavioral Measurements

Freezing behavior was used as a memory index in the fear conditioning tasks, being registered with a stopwatch in real time by an experienced observer that was blinded to the experimental group. Freezing was defined as total cessation of all movements except those required for respiration.

Open Field

Exploratory activity and anxiety-like behavior were assessed in the open field test in order to exclude non-specific effects of nimodipine. The apparatus consisted of a circular arena (90-cm diameter) with 50-cm high walls. The floor was subdivided into 12 quadrants and 3 concentric zones (periphery, intermediary and center). Animals were exposed to the apparatus for 5 minutes, during which the time spent on the periphery (thigmotaxis behavior) and the number of crossings between quadrants were measured. Nimodipine (16 mg/kg) or vehicle was measured intraperitonally 30 minutes before the test.

Drugs

Nimodipine (Sigma), an antagonist of the L-type voltage-gated calcium channels (LVGCCs) was dissolved in sterile isotonic saline solution with 8% dimethylsulfoxide to a concentration of 16 mg/mL. Nimodipine or its vehicle was injected intraperitoneally at a volume of 1 mL/kg (16 mg/kg) 30 min before memory reactivation sessions.

Statistical Analysis

Data were expressed as mean ± SEM. The statistical analyses were performed using two-tailed Student's t test; one-way, two-way or three-way analysis of variance, followed by Tukey's or Bonferroni's post hoc test, when necessary; Kruskal-Wallis test, followed by Dunn's post hoc. Values of $p < 0.05$ were considered statistically

significant. All experiments were randomized, performed double-blindly and without any form of P-hacking.

Computational simulations

Model network: In order to propose a mechanistic explanation for the experimental results, we used an adaptation of the attractor network model described in Osan et al (2011). This Hopfield-like network is capable of storing and retrieving memories using Hebbian learning rules dependent on the inputs presented to a fully connected network of 100 neurons. In this network, the activity of each neuron i is determined by

$$\tau \frac{du_i}{d_t} = -u_i + \frac{1}{2}\left(1 + \tanh\left(\sum_{j=1}^{N} w_{ij}u_j + I_i\right)\right)$$

where τ is the neural time constant and u_i represents the level of activation of neuron i which can vary continuously from 0 to 1 – unlike in the original Hopfield continuous activity model (Hopfield, 1984), in which activity varies from -1 to 1.

As a fully connected neural network, every neuron i is connected with every neuron j. For the learning process, the network needs to reinforce the connections between neurons that fire together, while creating inhibition when presynaptic neuron i is active and postsynaptic j is silent. Changes in the synaptic weight matrix $W = (w_{ij})$ are determined by the equation

$$\Delta W = -\gamma W + HLP + MID$$

where Hebbian learning plasticity (HLP) and mismatch-induced degradation (MID) are two independent learning rules (see below) and $0 \leq \gamma \leq 1$ is a time-dependent synaptic decay factor.

Learning occurs by presenting an input I_i to the circuit, corresponding to sensory information provided by the environment and/or internal cues, which lead to changes in the plastic connections between neurons. The cue has a one-to-one topology to the memory network, with every neuron receiving a cue input that can be either excitatory or inhibitory. Modifications on the synaptic weight matrix follow the HLP rule, corresponding to the Hebbian formulation implemented in classic Hopfield networks and described as:

$$HLP = S(u^T * u) - S((1 - u)^T * u)$$

where the vector u=$(u_1, u_2, \ldots, u_N)$ is the steady state of the network after input I_i presentation, while S is a factor representing requirements for Hebbian plasticity, such as protein synthesis, receptor activation, intracellular signaling and other mechanisms.

When the cue input leads to the retrieval of a previously stored memory, this can lead to mismatch between the cue input and the retrieved attractor if the two are not the same. This leads to concomitant activation of the MID learning rule, corresponding to a memory-updating system akin to that involved in memory destabilization during reconsolidation and defined by

$$MID = D(m^T * u)$$

where D is a factor representing requirements for memory destabilization (such as protein degradation and LVGCCs), $m = I_{norm} - u$ is the mismatch vector defined and I_{norm} is a normalized cue vector. The *MID* term leads to weakening of connections responsible for the mismatch in order to update the existing memory.

Learning and reactivation protocol: Non-overlapping neuron clusters in the network were chosen to represent the training or extinction contexts (6 neurons each), tone (2

neurons), aversive stimulus/shock (10 neurons) or safety/absence of shock (10 neurons) (**Fig. 5A**). Initially, a pattern representation of a memory unrelated to fear conditioning was presented as a cue to the network. This was followed by a training pattern activating neurons representing context A, tone and shock while inhibiting the remaining ones. To model standard extinction over multiple retrieval sessions, we activated the non-shock neuron cluster along with the neurons representing the extinction context and tone, while inhibiting the remaining ones. For deconditioning-update, we assumed an intermediate representation between the learning and extinction patterns. Unlike in the original model, no synaptic decay was assumed (i.e. γ was set to 0) and learning strength (as defined by S) was assumed to be smaller during reactivation sessions in both groups due to the lower intensity of the stimuli, thus allowing extinction to occur over multiple sessions.

Retrieval was evaluated after every training or reactivation session through presentation of a cue activating neurons corresponding to the extinction context and tone, with no input to the remaining neurons. For the renewal test, the retrieval cue activated neurons representing the training context and tone. To quantify memory retrieval, we used the mean activity of neurons representing shock and absence of shock, which was converted to a 'freezing percentage' by dividing the total activity of shock neurons by the total activity of both groups – thus, 100% freezing corresponds to full activation of shock neurons and no activation of non-shock neurons.

After each learning or reactivation session, the mean synaptic weight between each cluster of neurons (tone, contexts, shock and non-shock) was calculated by taking the average of the connections between all presynaptic neurons of a subpopulation and all

postsynaptic neurons of the other subpopulation. This was used to create the synaptic weight matrix between clusters shown in **Fig. 5E**.

Model parameters: All simulations used the following parameters: $N = 100$; $\tau = 1$; $\gamma = 0$; $s_0 = 1$. For training sessions, we set $S = 0.8$, while in reactivation sessions we used $S = 0.25$. D was set to 0.95 for all sessions, except for reactivations using nimodipine, in which $D = 0$. Each unit i during learning received an input I_i varying between -5 and 5. In the deconditioning update group, the aversive shock cluster received an input I_i of -2.31, while non-shock neurons received 2.31 (corresponding to t=6 in the transformation used by Osan et al. (2011) to create intermediate patterns). In the retrieval cue, each targeted neuron had an input $I_j = 1.5$ and 0 for other neurons. 100 simulations of each experiment were performed, with different initial conditions determined by Gaussian noise in the initial weight matrices (with a normal distribution on [-0.05, 0.05]) and in the neuronal activation at the start of every session (with a normal distribution on [0, 0.1]). For each simulation, 100 retrieval trials were run in each session to determine freezing percentage. All results are displayed as the mean ± S.E.M of these 100 simulations.

Acknowledgements

This work was supported by the Brazilian government agencies CAPES and CNPq (Universal 2018 - 405100/2018-3). The authors acknowledge Isabel Cristina Marques Scarello for her kind technical assistance.

Author contributions

L.O.A and B.P. designed this study. B.P. conducted the behavioral, pharmacological and performed statistical analyses. F.E.A. conducted the computational simulations. L.O.A and O.B.A wrote the manuscript. L.O.A supervised the project.

Competing interests

The authors declare no competing interests

References

Sareen, J. Posttraumatic stress disorder in adults: impact, comorbidity, risk factors, and treatment. *Can J Psychiatry* **59**, 460-467, doi:10.1177/070674371405900902 (2014).

Jonas, D. E. *et al.*, Psychological and pharmacological treabtments for adults with posttraumatic streess disorder (PTSD) Rockville (MD): Agency for Healthcare Research and Quality (US) doi:NBK137702 [bookaccession] (2013).

Berger, W. *et al.* Pharmacologic alternatives to antidepressants in posttraumatic stress disorder: a systematic review. *Prog Neuropsychopharmacol Biol Psychiatry* **33**, 169-180, doi: 10.1016/j.pnpbp.2008.12.004 (2009).

Vervliet, B., Craske, M. G. & Hermans, D. Fear extinction and relapse: state of the art. *Annu Rev Clin Psychol* **9**, 215-248, doi:10.1146/annurev-clinpsy-050212-185542 (2013).

Bouton, M. E. Context, ambiguity, and unlearning: sources of relapse after behavioral extinction. *Biol Psychiatry* **52**, 976-986, doi: 10.1016/S0006-3223(02)01546-9 (2002).

Archbold, G. E., Bouton, M. E. & Nader, K. Evidence for the persistence of contextual fear memories following immediate extinction. *Eur J Neurosci* **31**, 1303-1311, doi: 10.1111/j.1460-9568.2010.07161.x (2010).

Bouton, M. E., Winterbauer, N. E. & Todd, T. P. Relapse processes after the extinction of instrumental learning: renewal, resurgence, and reacquisition. *Behav Processes* **90**, 130-141, doi: 10.1016/j.beproc.2012.03.004 (2012).

Rescorla, R. A. & Heth, C. D. Reinstatement of fear to an extinguished conditioned stimulus. *J Exp Psychol Anim Behav Process* **1**, 88-96 (1975).

Nader, K., Schafe, G. E. & Le Doux, J. E. Fear memories require protein synthesis in the amygdala for reconsolidation after retrieval. *Nature* **406**, 722-726, doi:10.1038/35021052 (2000).

De Oliveira Alvares, L. *et al.* Reactivation enables memory updating, precision-keeping and strengthening: exploring the possible biological roles of reconsolidation. *Neuroscience* **244**, 42-48, doi: 10.1016/j.neuroscience.2013.04.005 (2013).

Nader, K. & Einarsson, E. O. Memory reconsolidation: an update. *Ann N Y Acad Sci* **1191**, 27-41, doi: 10.1111/j.1749-6632.2010.05443.x (2010).

Beckers, T. & Kindt, M. Memory Reconsolidation Interference as an Emerging Treatment for Emotional Disorders: Strengths, Limitations, Challenges, and Opportunities. *Annu Rev Clin Psychol* **13**, 99-121, doi:10.1146/annurev-clinpsy-032816-045209 (2017).

Duvarci, S. & Nader, K. Characterization of fear memory reconsolidation. *J Neurosci* **24**, 9269-9275, doi:10.1523/JNEUROSCI.2971-04.2004 (2004).

Monfils, M. H., Cowansage, K. K., Klann, E. & LeDoux, J. E. Extinction-reconsolidation boundaries: key to persistent attenuation of fear memories. *Science* **324**, 951-955, doi: 10.1126/science.1167975 (2009).

Monfils, M. H. & Holmes, E. A. Memory boundaries: opening a window inspired by reconsolidation to treat anxiety, trauma-related, and addiction disorders. *Lancet Psychiatry* **5**, 1032-1042, doi:10.1016/S2215-0366(18)30270-0 (2018).

Lee, J. L. Reconsolidation: maintaining memory relevance. *Trends Neurosci* **32**, 413-420, doi:10.1016/j.tins.2009.05.002 (2009).

Fernandez, R. S., Bavassi, L., Kaczer, L., Forcato, C. & Pedreira, M. E. Interference Conditions of the Reconsolidation Process in Humans: The Role of Valence and Different Memory Systems. *Front Hum Neurosci* **10**, 641, doi:10.3389/fnhum.2016.00641 (2016).

Suzuki, A. *et al.* Memory reconsolidation and extinction have distinct temporal and biochemical signatures. *J Neurosci* **24**, 4787-4795, doi:10.1523/JNEUROSCI.5491-03.2004 (2004).

Sevenster, D., Beckers, T. & Kindt, M. Prediction error demarcates the transition from retrieval, to reconsolidation, to new learning. *Learn Mem* **21**, 580-584, doi:10.1101/lm.035493.114 (2014).

Gershman, S. J., Monfils, M. H., Norman, K. A. & Niv, Y. The computational nature of memory modification. *Elife* **6**, doi:10.7554/eLife.23763 (2017).

Osan, R., Tort, A. B. & Amaral, O. B. A mismatch-based model for memory reconsolidation and extinction in attractor networks. *PLoS One* **6**, e23113, doi:10.1371/journal.pone.0023113 (2011).

Kredlow, M. A., Unger, L. D. & Otto, M. W. Harnessing reconsolidation to weaken fear and appetitive memories: A meta-analysis of post-retrieval extinction effects. *Psychol Bull* **142**, 314-336, doi:10.1037/bul0000034 (2016).

Gershman, S. J., Jones, C. E., Norman, K. A., Monfils, M. H. & Niv, Y. Gradual extinction prevents the return of fear: implications for the discovery of state. *Front Behav Neurosci* **7**, 164, doi:10.3389/fnbeh.2013.00164 (2013).

Wang, S. H., de Oliveira Alvares, L. & Nader, K. Cellular and systems mechanisms of memory strength as a constraint on auditory fear reconsolidation. *Nat Neurosci* **12**, 905-912, doi:10.1038/nn.2350 (2009).

Milekic, M. H. & Alberini, C. M. Temporally graded requirement for protein synthesis following memory reactivation. *Neuron* **36**, 521-525, doi:10.1016/S0896-6273(02)00976-5 (2002).

Suzuki, A., Mukawa, T., Tsukagoshi, A., Frankland, P. W. & Kida, S. Activation of LVGCCs and CB1 receptors required for destabilization of reactivated contextual fear memories. *Learn Mem* **15**, 426-433, doi:10.1101/lm.888808 (2008).

Frankland, P. W. *et al.* Stability of recent and remote contextual fear memory. *Learn Mem* **13**, 451-457, doi: 10.1101/lm.183406 (2006).

Bustos, S. G., Maldonado, H. & Molina, V. A. Disruptive effect of midazolam on fear memory reconsolidation: decisive influence of reactivation time span and memory age. *Neuropsychopharmacology* **34**, 446-457, doi:10.1038/npp.2008.75 (2009).

Pedraza, L. K., Sierra, R. O., Lotz, F. N. & de Oliveira Alvares, L. Periodical reactivation under the effect of caffeine attenuates fear memory expression in rats. *Sci Rep* **8**, 7260, doi:10.1038/s41598-018-25648-6 (2018).

Maren, S., Phan, K. L. & Liberzon, I. The contextual brain: implications for fear conditioning, extinction and psychopathology. *Nat Rev Neurosci* **14**, 417-428, doi:10.1038/nrn3492 (2013).

Lee, J. L. & Flavell, C. R. Inhibition and enhancement of contextual fear memory destabilization. *Front Behav Neurosci* **8**, 144, doi:10.3389/fnbeh.2014.00144 (2014).

Crestani, A. P. *et al.* Memory reconsolidation may be disrupted by a distractor stimulus presented during reactivation. *Sci Rep* **5**, 13633, doi: 10.1038/srep13633 (2015).

Haubrich, J. *et al.* Reconsolidation allows fear memory to be updated to a less aversive level through the incorporation of appetitive information. *Neuropsychopharmacology* **40**, 315-326, doi: 10.1038/npp.2014.174 (2015).

Sierra, R. O. *et al.* Reconsolidation may incorporate state-dependency into previously consolidated memories. *Learn Mem* **20**, 379-387, doi:10.1101/lm.030023.112 (2013).

Flavell, C. R., Barber, D. J. & Lee, J. L. Behavioural memory reconsolidation of food and fear memories. *Nat Commun* **2**, 504, doi:10.1038/ncomms1515 (2011).

Almeida-Correa, S. & Amaral, O. B. Memory labilization in reconsolidation and extinction--evidence for a common plasticity system? *J Physiol Paris* **108**, 292-306, doi:10.1016/j.jphysparis.2014.08.006 (2014).

Lee, J. L., Milton, A. L. & Everitt, B. J. Reconsolidation and extinction of conditioned fear: inhibition and potentiation. *J Neurosci* **26**, 10051-10056, doi:10.1523/JNEUROSCI.2466-06.2006 (2006).

Auber, A., Tedesco, V., Jones, C. E., Monfils, M. H. & Chiamulera, C. Post-retrieval extinction as reconsolidation interference: methodological issues or boundary conditions? *Psychopharmacology (Berl)* **226**, 631-647, doi:10.1007/s00213-013-3004-1 (2013).

Perez-Cuesta, L. M., Hepp, Y., Pedreira, M. E. & Maldonado, H. Memory is not extinguished along with CS presentation but within a few seconds after CS-offset. *Learn Mem* **14**, 101-108, doi:10.1101/lm.413507 (2007).

Plendl, W. & Wotjak, C. T. Dissociation of within- and between-session extinction of conditioned fear. *J Neurosci* **30**, 4990-4998, doi:10.1523/JNEUROSCI.6038-09.2010 (2010).

Almeida-Correa, S. *et al.* Calcineurin inhibition blocks within-, but not between-session fear extinction in mice. *Learn Mem* **22**, 159-169, doi:10.1101/lm.037770.114 (2015).