1   **Acquired FGFR and FGF alterations confer resistance to estrogen receptor (ER)**
2   **targeted therapy in ER+ metastatic breast cancer**
3

4   Pingping Mao[1,2,3,4,£], Ofir Cohen[1,2,3,4,£], Kailey J. Kowalski[1,2,3,4], Justin G. Kusiel[1,2,3,4],
5   Jorge E. Buendia-Buendia[1,2,3,4], Pedro Exman[2], Seth A. Wander[1,2,3,4,5], Adrienne G.
6   Waks[1,2,3,4,5], Jon Chung[6], Vincent A. Miller[6], Federica Piccioni[4], David E. Root[4], Eric P.
7   Winer[2,3,5], Nancy U. Lin[2,3,5], Nikhil Wagle[1,2,3,4,5]
8
1. Center for Cancer Precision Medicine, Dana-Farber Cancer Institute, Boston, MA
2. Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA
3. Harvard Medical School, Boston, MA
4. Broad Institute of MIT and Harvard, Cambridge, MA
5. Department of Medicine, Brigham and Women's Hospital, Boston, MA
6. Foundation Medicine, Inc. Cambridge, MA
15
16 Running title: FGFR/FGF alterations confer resistance to endocrine therapy
17
18 Key words: Drug resistance, endocrine therapy, ER+ breast cancer, FGF, FGFR
19
*Address correspondence to:
Nikhil Wagle, MD
Department of Medical Oncology
Dana-Farber Cancer Institute
450 Brookline Ave, Dana 820A
Boston, MA 02215
Phone: 617-632-6419
e-mail: nikhil_wagle@dfci.harvard.edu
28
29 [£]These authors contributed equally

30
31 Word count (excluding abstract, references and figure legends): 6182
32 Total number of figures: 7
33

Competing Financial Interest Statement
S.A.W. is a consultant for Foundation Medicine and Eli Lilly, and a consultant and equity
holder for InfiniteMD. J.C. is an employee/stockholder in Foundation Medicine, a wholly
owned subsidiary of Roche. V.A.M. is an employee of Foundation Medicine/Roche,
officer and board member of Revolution Medicines, and stockholder of Roche and
Revolution Medicines. N.U.L is a consultant for Puma Biotechnology and Daichii, and
received research funding from Pfizer, Genentech, Novartis and Seattle Genetics. E.P.W.
is a consultant for InfiniteMD, Genentech, and Eli Lilly. N.W. was previously a
stockholder and consultant for Foundation Medicine; has been a consultant/advisor for
Novartis and Eli Lilly; and has received sponsored research support from Novartis and
Puma Biotechnology. None of these entities had any role in the conceptualization, design,
data collection, analysis, decision to publish, or preparation of the manuscript

Abstract

Beyond acquired mutations in the estrogen receptor (ER), mechanisms of resistance to ER-directed therapies in ER+ breast cancer have not been clearly defined. We conducted a genome-scale functional screen spanning 10,135 genes to investigate genes whose overexpression confer resistance to selective estrogen receptor degraders. Pathway analysis of candidate resistance genes demonstrated that the FGFR, ERBB, insulin receptor, and MAPK pathways represented key modalities of resistance. In parallel, we performed whole exome sequencing in paired pre-treatment and post-resistance biopsies from 60 patients with ER+ metastatic breast cancer who had developed resistance to ER-targeted therapy. The FGFR pathway was altered via *FGFR1*, *FGFR2,* or *FGF3/FGF4* amplifications or *FGFR2* mutations in 24 (40%) of the post-resistance biopsies. In 12 of the 24 post-resistance tumors exhibiting FGFR/FGF alterations, these alterations were not detected in the corresponding pre-treatment tumors, suggesting that they were acquired or enriched under the selective pressure of ER-directed therapy. *In vitro* experiments in ER+ breast cancer cells confirmed that FGFR/FGF alterations led to fulvestrant resistance as well as cross-resistance to the CDK4/6 inhibitor palbociclib, through activation of the MAPK pathway. The resistance phenotypes were reversed by FGFR inhibitors and, to a lesser extent, MEK inhibitors, suggesting potential treatment strategies.

Significance

A genome-scale overexpression screen revealed a broad spectrum of resistance mechanisms against SERDs, which can provide a resource for researchers studying resistance to ER-directed therapies as well as the biology of estrogen receptor

dependencies in ER+ breast cancer. We demonstrate that activating FGFR/FGF

alterations are a mechanism of acquired resistance to ER-directed therapies and CDK4/6

inhibitors in ER+ metastatic breast cancer and can be overcome by combination therapy

targeting both the ER and the FGFR pathway. The detection of targetable, clonally

acquired genetic alterations in metastatic tumor biopsies highlights the value of serial

tumor testing to dissect mechanisms of resistance in human breast cancer and its potential

application in directing clinical management.

Introduction

Approximately 70% of breast cancers express the estrogen receptor (ER), and estrogen

signaling drives breast cancer cell growth and progression [1]. Endocrine therapies are

commonly used to treat ER+ breast cancer and work by reducing estrogen levels or

targeting the estrogen receptor through functional inhibition or degradation. Although

these endocrine therapies, including tamoxifen, aromatase inhibitors (AI), and the

selective estrogen receptor degrader (SERD) fulvestrant have improved survival for ER+

breast cancer patients, within the metastatic setting resistance to endocrine therapies is

nearly universal and remains a key challenge in reducing breast cancer morbidity and

mortality [2].

Although various resistance mechanisms have been proposed for tamoxifen and

aromatase inhibitor resistance, including loss or modification in ER expression (*ESR1*

activating mutations and *ESR1* fusions) [3-7], and regulation of alternative signal

transduction pathways (PI3K/AKT/mTOR and EGFR/ERBB2/MAPK) [8-10],

mechanisms of resistance to SERDs remain understudied. Mechanisms of endocrine

resistance identified in patients include acquired mutations in the estrogen receptor itself

[4-7], acquired activating mutations in *ERBB2* (HER2) [11, 12], loss of function of *NF1*

[13], and other alterations in MAPK pathway genes [14]. Additional mechanisms remain

to be identified.

Gain-of-function screens have played a pivotal role in identification of resistance

mechanisms to targeted therapies in various cancer types [15-17]. In breast cancer,

several functional screen studies identified *IGF1R*, *KRAS* and *ESR1* as mechanisms of

resistance to tamoxifen and/or estrogen deprivation [18-20]. However, genome-scale

functional screens for SERD resistance have not been reported.

We conducted a genome-scale gain-of-function screen in ER+ breast cancer cells

spanning 17,255 overexpressed lentiviral open reading frames (ORFs) to investigate

genes whose overexpression was sufficient to confer resistance to the SERDs fulvestrant

and GDC-0810 [21]. In parallel, we sought to identify endocrine resistance mechanisms

of clinical significance through genomic profiling of paired pre-treatment and post-

treatment tumor samples from 60 patients with ER+ metastatic breast cancer who

developed resistance to endocrine therapy.

The intersection of top candidate resistance mechanisms from both approaches converged

on the fibroblast growth factor receptor (FGFR) pathway. Here, we demonstrate that

acquired FGFR/FGF alterations identified in patients with resistant metastatic breast

cancer cause resistance to a variety of ER-directed therapies as well as to CDK4/6

inhibitors, and that this can be overcome by combination therapy targeting both the ER

and the FGFR pathway.

Results

A Genome-Scale Gain-of-Function Screen for Resistance to Selective Estrogen

Receptor Degraders

To identify the spectrum of genes whose overexpression confers resistance to SERDs *in*

vitro, we expressed 17,255 human open reading frames (ORFs), corresponding to 10,135

distinct human genes, in ER+ T47D breast cancer cells in the presence of fulvestrant or

GDC-0810. T47D cells were infected with the pooled lentiviral ORF library hORFeome

[22]. Fulvestrant, GDC-0810, or vehicle control (DMSO) was added following infection

and selection. ORF representation was assessed by sequencing after 21 days of drug

exposure. Genes that confer drug resistance will be enriched under drug selection,

indicated by a positive log fold change (LFC) for ORF representation before and after

DMSO/drug selection.

Using a Z score >3 as a criterion to identify resistance candidates, we identified 64 genes

(93 ORFs) that conferred resistance to fulvestrant and 57 genes (83 ORFs) that conferred

resistance to GDC-0810 (Fig.1A and Supplemental Table.1). 37 genes (55 ORFs)

conferred resistance to both drugs, a degree of overlap which was anticipated given the

mechanistic similarities between fulvestrant and GDC-0810. The LFC and corresponding

Z score for each ORF in fulvestrant and GDC-0810 treatment arms were highly

correlated, with a correlation coefficient of 0.77 (Fig.1A).

To confirm these results, we conducted a secondary screen using a smaller pooled library

consisting of 570 ORFs to validate candidates nominated by the primary screen. The

secondary screen was performed in both T47D and MCF7 cell lines with a similar screen

process as the primary screen. Top resistance genes found in the primary screen were

again enriched in the secondary screen, including *FGF* genes, *FOXR1*, *AKT* genes, *PIM*

genes and several *GPCR* genes (Supplemental Data Fig.S1). Many top ranked resistance

genes (*CSF1R, FGF3, FGF6, FOXR1* and *PIM2*) were shared between T47D and MCF7

cells (Supplemental Data Fig.S2). However, distinct resistance genes were also observed

in each cell line, suggesting some resistance mechanisms may be cell context-dependent.

Functional categories of candidate resistance genes include serine/threonine kinases

(*PIK3CA*, *AKT1/2/3*, *PIM1/2/3*), receptor tyrosine kinases (*EGFR*, *ERBB2, PDGFRB*),

growth factors (*FGF3/6/10/22*), cell cycle regulatory proteins (*CCND1*, *CCND2,*

CCND3, *CDK6*) and G-protein coupled receptors (*GPCR*) (Fig.1B). As further

validation, we overexpressed 13 ORFs belonging to these categories individually in

T47D cells and they all conferred resistance to fulvestrant (Fig.1C and Supplemental

Data Fig.S3A). Most of the 13 ORFs also conferred resistance to GDC-0810

(Supplemental Data Fig.S3B).

Gene Set Enrichment Analysis (GSEA) of the candidate resistance genes demonstrated

enrichment in 4 functional pathways: FGFR signaling, ERBB signaling, insulin receptor

signaling, and the MAPK pathway (Fig.1D, Supplemental Data Fig.S4 and Supplemental

Table.2). Consistent with this, we recently demonstrated that acquired *ERBB2* activating

mutations activate the MAPK pathway and cause endocrine resistance in patients with

ER+ metastatic breast cancer [11]. Several recent studies have also shown that alterations

in MAPK pathway genes are enriched in endocrine-resistant tumors [13, 14]. We sought

to further examine the role of FGFR and FGF genes in resistance to SERDs in MBC.

Identification of acquired FGFR and FGF alterations in metastatic biopsies from

patients with resistant ER+ MBC

To examine the potential role of the FGFR and FGF alterations in the development of

endocrine resistance clinically, we analyzed whole exome sequencing (WES) data from

paired pre-treatment and post-treatment metastatic tumor biopsies or cell free DNA from

60 patients with ER+ metastatic breast cancer who had received at least one endocrine

therapy (tamoxifen, AI, SERDs) for more than 120 days between the two biopsies [23].

Amongst the 60 post-treatment samples, we found *FGFR1* amplifications in 15% (9/60),

FGFR2 amplifications in 5% (3/60), *FGFR2* activating mutations in 3.3% (2/60), and

FGF3/FGF4 amplifications (which are adjacent genes and therefore co-amplified) in

28.3% (17/60) – for a total of 40% (24/60) of the cohort with at least one alteration in one

of these four genes (Fig.2A). Overall, the prevalence of *FGFR1*, *FGFR2*, and

FGF3/FGF4 alterations in the resistant metastatic setting seen here is increased

compared to what was observed in previously published cohorts of primary ER+ breast

cancer, such as The Cancer Genome Atlas (TCGA) [24] (Supplemental Table.3). The

incidence of *FGFR2* alterations (6.7%), in particular, is markedly increased compared to

primary treatment-naive breast cancer, in which the incidence is less than 2% in TCGA

(Supplemental Table.3).

To determine if this enrichment of FGFR/FGF alterations in the metastatic setting was

due to acquisition/selection under the selective pressure of endocrine therapy, we

compared the WES from the paired pre-treatment and post-treamtent samples for the 24

patients that exhibited FGFR/FGF alterations in their post-treatment samples. These 24

pairs of samples included 23 tumor biopsies and one cell-free DNA sample at the pre-

treatment timepoint, and 22 tumor biopsies, and two cell-free DNA samples at the post-

treatment timepoint. We performed an evolutionary analysis to evaluate clonal structure

and dynamics, including changes in mutations and copy number. The evolutionary

inference and clonal dynamics of mutations was based on changes in the estimated

fraction of tumor cells harboring each genomic alteration (the cancer cell fraction, CCF)

as previously shown for acquired HER2 mutations [11]. The evolutionary inference of

copy number changes was based on measuring differences in copy number amplitudes

between pre-treatment and post-treatment samples, while accounting for differences in

cancer cell fraction ("purity") in the sample and correcting for differences in ploidy. The

resultant purity-corrected values provide an estimate of "copy number above ploidy"

(CNAP) (see Methods).

For this analysis, we define "acquired" alterations as alterations with higher

representation in the post-treatment sample as compared to the pre-treatment sample. For

single nucleotide variants (SNVs), this means that the mutation had a substantially higher

CCF in the post-treatment sample compared to the pre-treatment sample (including lack

of detection in the pre-treatment sample despite having sufficient power to detect the

mutation). For copy number changes, this means that there was a substantial increase in

the overall copy-number in the post-treatment sample, suggesting a clonal expansion of

the amplification. Although we use the term "acquired", we recognize that when the

mutation is not detected in the pre-treatment sample, we cannot distinguish between pre-

existing alteration that was selected for and clonally enriched versus *de novo* alterations

that developed during the treatment.

In 12 of the 24 patients with FGFR or FGF alterations (50%), the alterations were

acquired in the post-treatment sample as compared to the pre-treatment sample (Fig. 2A,

marked in red). Five out of nine *FGFR1* amplifications were acquired (55.6%), while all

four *FGFR2* alterations were acquired (100%), including one patient (Pt 0300350) with

acquisition of both an *FGFR2* mutation and amplification. *FGF3/FGF4* amplifications

were acquired in 4 of 17 tumors (23.5%), including one case in which an *FGFR1*

amplification was co-acquired. The concurrent acquisition may suggest that the

evolutionary selection of both the ligand and receptor provided additional fitness in this

tumor. Among the other 12 patients, the alterations in eight patients were shared in both

pre-treatment and post-treatment samples (Fig 2A, marked in black), and evolutionary

status of alterations in the remaining four patients was inconclusive (Fig 2A, marked in

grey). The increase in copy number (corrected for tumor purity and ploidy, Supplemental

Table.4) from pre-treatment to post-treatment for *FGFR1*, *FGFR2*, and the *FGF3/FGF4*

amplicon in all 12 patients is depicted in Fig.2B.

Two of the acquired alterations found in these 12 patients were SNVs in the *FGFR2*

gene: M538I (chr10:123258070C>T, GRCh37, also denoted as M537I, depending on the

isoform) and N550K (chr10:123258034A>T, GRCh37, also denoted as N549K,

depending on the isoform). N550K is the most common FGFR2 mutation in breast cancer

while M538I was previously identified in lung cancer but has not yet been characterized

in breast cancer [25]. Figure 2C illustrates the change in the estimated fraction of tumor

cells harboring each genomic alteration (CCF) from the pre-treatment biopsy to the

resistant biopsy. In both patients, the *FGFR2* mutations were either not detected in the

primary tumor, despite sufficient power to detect mutations at this locus (N550K in Pt

0300350) or detected by a single read, inferred in a small fraction (CCF of 2%) of the

pre-treatment tumor (M538I in Pt 0300348). In both patients the activating *FGFR2*

mutations in the post-treatment biopsies were clonally acquired (CCF of 100%). Pt

0300350 was also found to have an acquired *FGFR2* amplification while Pt 0300348 was

found to have gained low-level amplification in *FGFR2* post treatment (Fig.2B).

Notably, the acquired alterations in *FGFR1*, *FGFR2*, and *FGF3/FGF4* were largely

mutually exclusive with acquired *ESR1* mutations. *ESR1* mutations are the most common

mechanism described for acquired endocrine resistance [26]. Although the overall rate of

acquired *ESR1* mutation in this cohort is 22% (13/60), among the 12 cases of acquired

FGFR and FGF alterations, only one patient also has an acquired ESR1 mutation (Figure

2A).

FGF3 and *FGF4* reside in genomic proximity to *FGF19* and *CCND1* and these four

genes are often co-amplified. However, here, in 3 out of the 4 cases with acquired

FGF3/FGF4 amplification, *FGF3/FGF4* copies were gained without co-acquisition of

CCND1 amplification, suggesting that this acquisition can occur as an independent

genomic event. Similarly, 2 out of the 4 cases with acquired *FGF3/FGF4* did not have

co-acquisition of *FGF19* amplification. The relationship of acquisitions of *FGF3*, *FGF4*,

FGF19, and *CCND1* are depicted in Supplemental Data Fig.S5 and further details about

these amplicons are described in Supplemental Methods. Further exploration of the full

genomic contexts and other concurrent genetic alterations for the 12 patients with

acquired FGFR/FGF alterations are shown in Supplemental Data Fig.S6 and

Supplemental Table.5-6.

Figure 3 depicts clinical vignettes for six of the patients with acquired *FGFR1*, *FGFR2*,

and/or *FGF3/FGF4* alterations in their post-treatment biopsies. All patients were treated

with ER-directed therapy before acquiring FGF or FGFR alterations, including tamoxifen

(3 patients), AIs (6 patients), and fulvestrant (3 patients). Vignettes for the other six

patients with acquired *FGFR1*, *FGFR2*, and/or *FGF3/FGF4* alterations in their post-

treatment biopsies are shown in Supplemental Data Fig.S7. Detailed clinicopathological

features and therapy details for all 12 patients are found in Supplemental Table.7.

In addition to these 12 patients in our cohort, we identified several additional patients

with acquired *FGFR1* and *FGFR2* activating mutations following the development of

resistance to endocrine therapy (Supplemental Data Fig.S8 and Supplemental Table.8).

FM patient 1 acquired a clonal *FGFR1* N546K mutation (a known activating mutation

paralogous to *FGFR2* N550K) following treatment with an AI. FM patient 2 acquired a

subclonal *FGFR2* N550K mutation after treatment with tamoxifen, AI and fulvestrant.

FM patient 3 acquired a subclonal *FGFR2* K660N mutation, another activating mutation

in the kinase domain [27], after treatment with tamoxifen.

In summary, we observed acquired alterations in FGFR or FGF in 20% (12/60) of

patients with endocrine resistant ER+ MBC – comparable to the known frequency of

acquired mutations in *ESR1* – highlighting the important role of the FGFR pathway in

acquired resistance to ER-directed therapies.

Active FGFR signaling leads to resistance to SERDs through activation of the MAP

kinase pathway

To further investigate how FGFR/FGF genes may confer resistance to ER-directed

therapy, we treated T47D cells with FGF3, FGF6, FGF10 or FGF22 ligand. Each of these

ligands resulted in resistance to fulvestrant (Fig.4A). This effect was reversed by

PD173074, a pan-FGFR inhibitor (Fig.4A). The addition of FGF ligand enhanced

phosphorylation of ERK and AKT, which was reversed by PD173074 (Fig.4B). The

effect of FGF ligands on downstream effectors was enhanced when FGFR1 was

simultaneously overexpressed in T47D, which has relatively low expression of FGFR1

[28] (Supplemental Data Fig.S9). FGF3, FGF6, FGF10 and FGF22 also reduced

fulvestrant sensitivity in MCF7 cells (Supplemental Data Fig. S10 A-B). Similar results

have been shown previously for FGF2, which was reported to activate MEK-ERK to

drive fulvestrant resistance in ER+ breast cancer cells [29].

We next overexpressed FGFR1, FGFR2, or GFP in T47D cells through lentiviral

transduction and examined the impact on susceptibility to SERDs. Overexpression of

FGFR1 or FGFR2 alone did not affect sensitivity to fulvestrant or GDC-0810. However,

with the addition of FGF2 ligand, both FGFR1 and FGFR2 rendered cells highly resistant

to the two SERDs (Fig. 4C and Supplemental Data Fig. S11). In comparison, FGF2

ligand alone reduced sensitivity to SERDs in control cells expressing GFP to a much

lesser extent than in the FGFR1 or FGFR2 expressing cells, suggesting the potent

resistance phenotype requires both FGF ligand and receptor. This requirement for the

presence of both FGF ligand and receptor for maximal resistance phenotype may also

explain why only FGFs but not FGFR1 or FGFR2 scored in the resistance screen (Fig.1

A-B). The resistance phenotype resulting from FGFR1 and FGFR2 overexpression was

completely reversed by the addition of PD173074 (Fig. 4C). Similar results were

obtained in MCF7 cells (Supplemental Data Fig.S12 A-B).

FGFR1 and FGFR2 overexpression (in the presence of FGF2 ligand) induced more

potent phosphorylation of AKT and ERK than the GFP control (Fig.4D). These results

are consistent with previous findings that FGFR1 activation led to MAPK activation and

fulvestrant resistance [28]. Activation of downstream effectors p-ERK and p-AKT by

FGFR1/2 overexpression was reversed to baseline levels with PD173074 (Fig. 4D and

Supplemental Data Fig. S12C). Examination of a larger number of kinases using kinase

antibody arrays demonstrated that AKT, ERK and RSK (downstream effector of ERK)

were the only kinases of those tested to exhibit increased phosphorylation following

FGFR1/2 overexpression and FGF2 stimulation (Supplemental Data Fig.S13).

Collectively, these findings suggest that FGFR1 and FGFR2 cause SERD resistance

through the activation of MAPK and/or PI3K/AKT pathways.

We examined the sensitivity of cells overexpressing FGFR1 or FGFR2 to several

inhibitors of downstream effectors: the MEK inhibitor trametinib, the AKT inhibitor

AZD5363, and the mTOR inhibitor everolimus. FGFR1 or FGFR2 overexpression in the

presence of FGF2 led to hypersensitivity to trametinib (Supplemental Data Fig.S14A). In

contrast, FGFR1 or FGFR2 overexpression in the presence of FGF2 reduced sensitivity

to AKT and mTOR inhibitors (Supplemental Data Fig.S14A).

We attempted to reverse FGFR-induced resistance to fulvestrant by inhibiting the MAPK

pathway. Treatment of FGFR1 overexpressing cells with trametinib partially resensitized

cells to fulvestrant, and treatment of FGFR2 overexpressing cells with trametinib fully

resensitized the cells to fulvestrant (Fig.4E-F). Treatment with the mTOR inhibitor

everolimus also partially blocked resistance conferred by FGFR1 or FGFR2

overexpression (Supplemental Data Fig.S14B). We also performed long-term colony

formation assays and observed similar results (Supplemental Data Fig.S14C).

Together, these results suggest that the MAPK pathway is the primary downstream

effector of FGFR activation resulting in endocrine resistance. This is consistent with the

pathway analysis of resistance genes in our initial overexpression screen, which

demonstrated enrichment in MAPK pathway genes (Figure 1D), as well as our prior

findings of MAPK activation through acquired HER2 mutations in endocrine

resistance—adding further support to the idea that the MAPK pathway may be a common

node of endocrine resistance [11, 14].

FGFR activation confers cross-resistance to CDK4/6 inhibitors

Since the combination of endocrine therapy and CDK4/6 inhibitors is now a standard of

care treatment for patients with ER+ metastatic breast cancer, we also examined the

effect of FGFR signaling on sensitivity to the combination of fulvestrant and the CDK4/6

inhibitor palbociclib. In T47D cells, FGFR1 and FGFR2 overexpression in the presence

of FGF2 also conferred resistance to combination treatment of fulvestrant and palbociclib

(Fig. 5A). The resistance phenotype was again abrogated by PD173074 (Fig. 5A).

Resistance to fulvestrant and palbociclib was also partially reversed by trametinib (Fig.

5B and Supplemental Data Fig. S14D), further providing the support for the role of

MAPK pathway activation in FGFR-mediated drug resistance. The reversal of resistance

by trametinib was accompanied by reduced ERK phosphorylation (Fig. 5C). Similar

results were achieved in MCF7 cells, although everolimus was more effective than

trametinib in reversing the resistance phenotype by FGFR1 or FGFR2 overexpression

(Supplemental Data Fig.S15 and Fig.S16).

In the presence of fulvestrant and palbociclib, FGFR1 or FGFR2 overexpression was

accompanied by increased p-Rb and CCND1 levels, both of which were partially

reversed by trametinib (Fig.5C). CCND1 knockdown in cells overexpressing GFP,

FGFR1 or FGFR2 impaired cell proliferation similarly across all three cell lines without

affecting the IC50 of fulvestrant (Supplemental Data Fig.S17). This suggests that the

proliferation advantage provided by active FGFR signaling is partially dependent on

CCND1. This is consistent with prior results suggesting that CCND1 was involved in

FGF2-mediated drug resistance [29].

Clinical evidence also supports the finding that FGFR alterations can cause resistance to

CDK4/6 inhibitors. Following the acquisition of *FGFR2* N550K (along with *FGFR2*

amplification), Pt 0300350 did not respond to the combination of letrozole and

palbociclib (Fig.3), suggesting that *FGFR2* alterations may lead to intrinsic resistance to

the combination of endocrine therapy and CDK4/6 inhibitors. Another patient with an

FGFR2 N550K mutation (FM Patient 2) also did not respond to the combination of

fulvestrant and palbociclib (Supplemental Data Fig.S8). Collectively, this suggests

targeting the FGFR pathway may also be a viable strategy to overcome FGFR/FGF-

mediated resistance to SERDs and CDK4/6 inhibitors.

FGFR2 mutations found in patients are activating and can be targeted with

irreversible FGFR inhibitors

We identified 3 acquired mutations in the kinase domain of FGFR2 in patients who

developed resistance to endocrine therapy. Two of these, FGFR2 N550K and K660N, are

known activating FGFR2 mutations that have been previously identified in breast cancer

[25, 27]. FGFR2 N550K is part of the molecular brake at the kinase hinge region, which

allows the receptor to adopt an active conformation more easily (Fig.6A). N550K is a

recurring hotspot mutation reported to confer resistance to several FGFR inhibitors

including PD173074 and dovitinib [30]. FGFR2 K660N is located in a conserved region

in the tyrosine kinase domain and has been confirmed to increase kinase activity [27, 30].

The third mutation, FGFR2 M538I, has not been previously reported in breast cancer.

Based on its location, M538I appears to stabilize the active kinase confirmation by

strengthening the hydrophobic spine of the FGFR2 kinase [30] (Fig.6A).

We expressed all three FGFR2 kinase domain mutants in T47D cells through lentiviral

transduction, as well as wildtype (WT) FGFR2 and GFP as negative controls. All three

mutants elicited higher kinase activity than WT FGFR2 constitutively, demonstrated by

levels of p-FRS2, a direct substrate for FGFR2 (Fig.6B). MAPK and AKT signaling were

also increased, as indicated by increased p-ERK and p-AKT levels, respectively (Fig.6B).

The addition of FGF2 ligand further enhanced downstream signaling for all FGFR2

mutants, and the enhanced signaling was blocked by PD173074 for FGFR2 M538I and

K660N, but not for N550K (Fig.6B).

FGFR2 mutants were also expressed under a tetracycline responsive promoter in T47D

cells grown in low doses of doxycycline to determine the functionality at lower

expression levels. At lower levels of expression, FGFR2 N550K still led to increased

levels of p-ERK and p-AKT, independent of FGF2 ligand stimulation. FGFR2 M538I

and K660N also resulted in higher p-ERK and p-AKT levels in the presence of FGF2 as

compared to FGFR2 WT (Fig.6C). FGFR2 mutants rendered cells more sensitive to

trametinib than did GFP or WT FGFR2 (Fig.6D), further supporting the finding that

FGFR signaling requires the MAPK pathway in this context. Taken together, these results

indicate that all three FGFR2 mutations acquired in breast cancer patients are functionally

active – FGFR2 N550K is constitutively active while FGFR2 M538I and K660N may be

more ligand-dependent at low levels of expression.

All 3 FGFR2 mutants led to modest resistance to fulvestrant (Fig.6E), which was

enhanced in the presence of FGF2 ligand. PD173074 resensitized cells overexpressing

FGFR2 M538I and FGFR2 K660N as well as WT FGFR to fulvestrant, but not cells

overexpressing FGFR2 N550K (Fig.6E). Consistent results were observed when cells

were treated with the combination of fulvestrant and palbociclib (Fig.6F). Similar results

were obtained in MCF7 cells (Supplemental Data Fig.S18).

FGFR2 M538I and N550K have been shown to confer resistance to multi-kinase inhibitor

dovitinib in BaF3 cells, and N550K is also resistant to PD173074 [30]. Because of the

differential responses of these FGFR2 mutants to PD173074, we tested the ability of

additional FGFR inhibitors to resensitize cells expressing these mutants to fulvestrant.

FIIN-2 and FIIN-3 are two irreversible covalent pan-FGFR inhibitors that target a

cysteine conserved in FGFR1-4 and have exquisite selectivity for some FGFR2 mutations

including M538I and K659N [31]. In addition, FGFR2 N550K was previously shown to

respond to the selective FGFR inhibitor AZD4547 [32].

Both FIIN-2 and FIIN-3 were more effective in inhibiting the downstream signaling (p-

FRS2, p-ERK and p-AKT) induced by FGFR2 N550K as compared to PD173074 and

AZD4547 (Fig.7A). Furthermore, FIIN compounds reduced the level of downstream

effectors back to baseline levels, with FIIN-3 being more potent than FIIN-2

(Supplemental Data Fig.S19). T47D cells stably overexpressing FGFR2 mutant were

exquisitely sensitive to FIIN-2 and FIIN-3 as compared to cells expressing GFP or

FGFR2 WT (Fig.7B). The resistance to fulvestrant induced by both WT FGFR2 and all

mutant FGFR2 was completely blocked by FIIN-2 (1 μM) and FIIN-3 (100 nM)

(Fig.7C).

While resistance to WT FGFR1/2 and FGFR2 M538I and FGFR2 K660N can be

reversed by multiple FGFR inhibitors, for some mutants like FGFR2 N550K, only the

irreversible pan-FGFR inhibitors successfully resensitized cells to fulvestrant,

highlighting the fact that specific resistance mutations might require different strategies

to overcome or preempt endocrine resistance.

Discussion

In this study, we used a genome-scale gain-of-function screen to identify potential

mechanisms of resistance to selective estrogen receptor degraders. We nominated several

different candidate resistance genes and pathways, particularly genes in the ERBB

pathway, FGFR pathway, insulin receptor signaling and the MAPK pathway. Consistent

with this finding, genomic profiling of paired pre-treatment and post-treatment tumor

samples from 60 patients with ER+ metastatic breast cancer who developed resistance to

endocrine therapy identified acquired alterations in *FGFR1*, *FGFR2*, and *FGF3/FGF4* in

20% of patients. Experimental studies confirmed that these alterations confer resistance

to endocrine therapy as well as CDK4/6 inhibitors, through activation of MAPK pathway,

and demonstrated that this resistance can be reversed by FGFR inhibitors. Taken

together, our results suggest that activating FGFR pathway alterations are a distinct

mechanism of acquired resistance to multiple forms of ER-directed therapy in MBC that

can be overcome by FGFR inhibitors.

Our genome-scale screen provided a comprehensive view into the resistance mechanisms

to SERDs. Similar resistance genes were nominated for fulvestrant and GDC-0810,

thereby confirming the two drugs have similar mechanism of action. Of note, two ESR1

ORFs conferred resistance specifically to GDC-0810 but not fulvestrant, possibly due to

GDC-0810 having a less potent effect on ER degradation than fulvestrant [21]. Among

the resistance mechanisms shared by fulvestrant and GDC-0810, many are frequently

altered in ER+ MBC, such as *CCNDs/CDK6*, *KRAS/MAPK*, *EGFR/ERBB2* and

PIK3CA/AKTs/PIMs, and agents targeting those alterations are under clinical

development to be combined with endocrine therapy [14, 33]. We also identified

potential resistance mechanisms that are not characterized to the same extent, such as G

protein-coupled receptors, Wnt pathway (*FZD10, RSPO1, RSPO3*) and Src family

kinases (*YES1, FYN, FGR*), providing clues as to the potential crosstalk between these

pathways and ER signaling [34-36] and suggesting that breast cancer patients harboring

functional alterations in these pathways may develop resistance to SERDs. Notably, some

of the resistance genes were also nominated in gain-of-function screens designed to

identify resistance mechanisms for MAPK pathway inhibitors in melanoma (*AXL, CRK,*

CRKL, FGR, GPCR genes) [15] and PI3K inhibitors in ER+ breast cancer (*AKT1, AKT2,*

CRKL, FGF3, FGF10, PIM genes) [17]. This may reveal multi-drug resistance

mechanisms and thus guide clinical drug combinations to overcome resistance. We have

provided the full genome-scale screen data as a resource to the community of researchers

interested in resistance to ER-directed therapies as well as the biology of estrogen

receptor dependencies in ER+ breast cancer.

Our ultimate goal is to identify resistance mechanisms that are clinically relevant and can

be therapeutically targeted. By comparing paired pre-treatment and post-treatment

tumors, our evolutionary analyses identified acquired *FGFR1* and *FGFR2*, and

FGF3/FGF4 alterations in 12 out of 60 post-treatment samples, further highlighting a

potential role for the FGFR pathway in driving drug resistance and disease progression.

Most notably, all four alterations in FGFR2 in our cohort were found to be acquired after

the development of resistance to endocrine therapy. Our overall findings are consistent

with two recent cohort studies which noted some patients with acquired *FGFR1* and

FGFR2 alterations following treatment of endocrine therapy [37, 38], and provides a

mechanistic explanation for these acquisitions.

This analysis was enabled by a novel method we developed to compare the magnitude of

amplification in matched pre- and post-treatment samples while considering key

confounders to allow for more reliable assignment of copy gain or loss. Since matched

tumor samples of the same patient are highly variable in the cancer cell fraction (purity)

and often variable in ploidy (with genome duplication taking place in the metastatic

setting), we computed the purity-corrected copy number above ploidy and set a relatively

stringent threshold of changes in CNAP to define acquired amplification (See Methods),

as cancer clones bearing amplifications with high focality and magnitude in FGFR/FGF

genes are more likely to induce dependency on FGFR pathway and result in endocrine

resistance.

Our genomic analysis has some limitations and caveats. The observed alterations may not

exclusively result from endocrine therapy as some patients received other therapies

between the two collected biopsies. Moreover, tumors with FGFR/FGF alterations also

harbor alterations in other cancer genes, which may contribute to drug resistance as well

(Supplemental Data Fig.S6 and Supplemental Table.5). Despite these caveats, with the

evidence from unbiased screens, genomic evidence in relevant patient samples, and

confirmatory experimental models, the FGFR pathway clearly emerges as a clinically

important resistance mechanism for SERDs and CDK4/6 inhibitors.

Strategies to target the FGFR pathway in breast cancer patients with FGFR alterations are

currently being assessed in clinical trials. FGFR inhibitors currently under clinical

development in breast cancer include non-selective tyrosine kinase inhibitors (dovitinib

and lucitanib), FGFR1-3 selective inhibitors (AZD4547 and BGJ398), and others,

although clinical trials have achieved mixed results to date [39-44]. The combination of

FGFR inhibitors and endocrine therapy is also being clinically investigated. For example,

the combination of dovitinib and fulvestrant showed promising clinical activity [39]. As

FGFR pathway activation also results in resistance to CDK4/6 inhibitors, a triple

combination with the addition of CDK4/6 inhibitors may also be considered. One

challenge for the use of FGFR inhibitors is to identify reliable biomarkers. Our results

suggest focal and high level amplifications, clonal activating mutations or high

expression levels of FGFR and FGF genes, particularly in the metastatic setting, may be

used to guide the clinical use of FGFR inhibitors. Activating alterations in *FGFR2*, which

are rare in primary treatment naïve breast cancer but appear to be clonally acquired in a

subset of patients with resistant ER+ MBC, may be a particularly good biomarker for the

development of FGFR inhibitors.

Our work also highlights that the effective clinical use of FGFR inhibitors needs to

consider the variable drug sensitivity of different FGFR2 mutations, which were acquired

in some patients following endocrine therapy. The two irreversible pan-FGFR kinase

inhibitors, FIIN-2 and FIIN-3, had superior efficacy in targeting all FGFR2 mutants

including N550K when compared to other FGFR inhibitors, although *in vivo* efficacy,

off-target effects and toxicity of FIIN compounds still warrant further investigation.

Alterations in FGFR1 and FGFR2 activated the MAPK pathway, and MEK inhibition

was able to overcome the resistance conferred by FGFR pathway to some degree. We

previously demonstrated acquired ERBB2 mutations resulted in elevated MAPK

transcriptional signature [11]. Furthermore, increased frequency of alterations in MAPK

pathway genes was found in tumors post hormonal therapy, including *EGFR, ERBB2* and

NF1 [14]. The fact that multiple mechanisms of resistance to ER-directed therapies

and/or CDK4/6 inhibitors activate the MAPK pathway suggests that this may be an

important node of resistance in ER+ MBC. Thus, combining endocrine therapy and

CDK4/6 inhibitors with agents that target MAPK pathway, such as MEK inhibitors

and/or SHP2 inhibitors [45, 46], may be a unifying strategy to overcome or prevent

resistance resulting from multiple genetic aberrations that lead to resistance in ER+

MBC.

In summary, the integration of a functional genomic screen and genomic analysis of pre-

and post-treatment biopsies revealed the FGFR pathway as an important resistance

mechanism for endocrine therapy and CDK4/6 inhibitors in ER+ breast cancer. With the

increasing use of SERDs and CDK4/6 inhibitors in the clinic, we anticipate that the

prevalence of FGFR/FGF alterations might increase in the future. Targeting the FGFR

pathway with FGFR inhibitors or agents that target downstream MAPK signaling may

improve clinical outcomes in patients with aberrations in FGFR/FGF genes. Furthermore,

our study highlights the need to sequence metastatic biopsy or blood biopsies at the time

of resistance to identify patients with these alterations who may benefit from targeting the

FGFR pathway.

Methods

Cell culture

293T, T47D and MCF7 cells were purchased from American Type Culture Collection

(ATCC) and were cultured as described in the Supplemental Methods.

Genome-Scale Gain-of-Function Screen

The pooled lentiviral ORF library hORFeome [22] consists of 17,255 barcoded human

open reading frames (ORFs), corresponding to 10,135 distinct human genes with at least

99% nucleotide and protein match. These ORFs were cloned into pLX317 vector and

pooled together for transfection into 293T cells to make pooled lentivirus (with 2nd

generation packaging plasmids). In 6-well plates, pooled lentivirus was infected in cells

to achieve ~50% infection rate and ensure ~1000 infected cells per ORF for 17255.

Media was supplemented with 4 μg/mL polybrene (Thermo Fisher Scientific #

TR1003G) to boost transfection efficiency. After infection, cells were pooled and

selected with 1.5 μg/mL puromycin for 5 days. Upon completion of selection, cells were

plated for three different drug conditions: DMSO, 100 nM fulvestrant, 1 μM GDC-0810.

There were three replicates for each condition screened. A subset of cells was saved for

sequencing as early time point (ETP) samples to confirm ORF representation. The dose

for each drug was chosen for the two drugs to achieve potent anti-proliferation effect that

could be rescued with ESR1 mutant Y537N and Y537S. Infected cells were passaged

upon confluency and maintained in DMSO or drugs for 21 days to allow sufficient time

for cells carrying resistance to be enriched from the population. At the end of the time

course, cells were harvested for isolating genomic DNA as late time point samples (LTP).

All genomic DNA samples were amplified with PCR primers flanking the ORF region

and sequenced. The ORF representation at the final harvesting (LTP) is compared to the

representation of ORFs in cells collected before drug addition (ETP). Cells carrying

ORFs that are driving resistance will grow and gradually enrich the population and

therefore, will be over-represented in the sequencing data for the final passage compared

to the early time point. An ORF with significant enrichment (a Z score >3) is defined as a

resistance candidate gene. A secondary validation screen was performed as described in

the Supplemental Methods.

Patients and Tumor Samples

Prior to any study procedures, all patients provided written informed consent for research

biopsies and whole exome sequencing of tumor and normal DNA, as approved by the

Dana-Farber/Harvard Cancer Center Institutional Review Board (DF/HCC Protocol 05-

246). Metastatic core biopsies were obtained from patients and samples were

immediately snap frozen in OCT and stored in -80°C. Archival FFPE blocks of primary

tumor samples were also obtained. A blood sample was obtained during the course of

treatment, and whole blood was stored at -80°C until DNA extraction from peripheral

blood mononuclear cells (for germline DNA) was performed. In a few instances, cell free

DNA was obtained from plasma for circulating tumor DNA analysis, as previously

described[47].

WES and data analysis

DNA was extracted from primary tumors, metastatic tumors, plasma, and peripheral

blood mononuclear cells (for germline DNA) from all patients and whole exome

sequencing was performed, as detailed in the Supplemental Methods. Sequencing data

were analyzed using tools to identify somatic point mutations and small

insertions/deletions (indels), and copy number changes using established algorithms (see

Supplemental Methods).

To better measure segment-specific copy-number, we subtracted the genome ploidy for

each sample to compute copy number above ploidy (CNAP). CNAP of at least 3 are

considered as amplifications (AMP), CNAP below 3 are considered low amplification

and ignored in our analysis). CNAP of at least 6 are considered high amplifications

(HighAMP), and CNAP of at least 9 and fewer than 100 genes [48] is considered very

high focal amplification (FocalAMP).

The evolutionary classification of amplifications accounts for the magnitude of the

observed copy-number difference between the pre-treatment and the post-treatment

samples. If the difference between the CNAP of the post-treatment and the CNAP of the

pre-treatment is smaller than 50%, the amplification is defined as "Shared". If the CNAP

of the post-treatment is larger than the CNAP by more than 50% and the lower pre-

treatment CNAP is not at "FocalAMP" level, the evolutionary classification is

"Acquired". If CNAP of the post-treatment is smaller by at least 50%, comparing to the

pre-treatment sample and the lower post-treatment CNAP is not at "FocalAMP" level, the

evolutionary classification is "Loss". Otherwise, the evolutionary classification of

amplifications is defined as "Indeterminate".

Generation of plasmids and engineered cells

T47D or MCF7 cells were infected with lentivirus to derive stable cell lines

overexpressing wildtype (WT) or mutant ORFs. All WT ORFs were obtained from the

Broad Institute. Mutant ORFs (FGFR2 M538I, N550K and K660N) were made using

QuickChange II site-directed mutagenesis kit (Agilent Technologies #200523). Most

stable cells lines express ORFs in pLX317 vector and were selected with puromycin (Life

Technologies #A1113803). Stable cell lines expressing CCND1 and PIM1 in pLX304

vector were selected with blasticidin (Life Technologies #A1113903).

For inducible cell lines, WT or Mutant FGFR2 IIIb ORFs were cloned into pInducer20

Tetracycline-inducible lentiviral vector (Addgene #44012) using gateway cloning

technology (Invitrogen #11791019). Lentivirus was then generated and infected to

establish Tet-inducible cells lines cultured in 10% Tet-system approved FBS (Clontech

#631106) and 500 µg/mL G418 (Life Technologies #10131035). Doxycycline (Clontech

#631311) was used to induce ORF expression.

Kill curves and CellTiter-Glo viability assay

Cells were plated in 96-well tissue culture ViewPlate (Perkin Elmer # 6005181) on Day 1

and treated with drug on Day 2. Media with or without drugs was refreshed on Day 5. On

Day 8, cells were equilibrated to room temperature, media was removed, and cells were

lysed in a mixture of 50 µL media and 50 µL CellTiter-Glo 2.0 reagent (Promega #

G9243) per well. Plates were then incubated on an orbital shaker for 2 mins. Following

another 10 mins of incubation at room temperature to stabilize signal, luminescence was

recorded to measure cell viability on Infinite M200 Pro microplate reader (Tecan).

Colony formation assay

2,000-30,000 cells were plated in 6-well plates on Day 1 and treated on Day 2. Media

was refreshed every 3-4 days until crystal violet staining. On the day of staining, cells

were fixed with ice-cold 100% methanol for 10 minutes and then incubated with 0.5%

crystal violet solution (Sigma Aldrich #C6158) in 25% methanol at room temperature for

10 minutes.

Western blotting

Western blotting was performed as described in supplemental methods.

Statistical analysis

Statistical analyses related to drug response curve were performed with student t-test in Graphpad Prism.

Acknowledgements

We thank Qaren Quartey, Christian Kapstad and Gabriela Johnson for technical assistance; Karla Helvie, Laura Dellostritto, Lori Marini, Nelly Oliver, Shreevidya Periyasamy, Colin Mackichan, Max Lloyd, and Mahmoud Charif for assistance with patient sample collection and annotation; and Flora Luo, Tinghu Zhang and Nathanael Gray for providing reagents. We thank Jorge Gómez Tejeda Zañudo for helpful discussions and comments on the manuscript. We are grateful to all the patients who volunteered for our tumor biopsy protocol and generously provided the tissue analyzed in this study.

Grant support

This work was supported by the Department of Defense W81XWH-13-1-0032 (N.W.), AACR Landon Foundation 13-60-27-WAGL (N.W.), NCI Breast Cancer SPORE at DF/HCC #P50CA168504 (N.W., N.U.L and E.P.W), Susan G. Komen CCR15333343 (N.W.), The V Foundation (N.W.), The Breast Cancer Alliance (N.W.), The Cancer

Author contributions

P.M., O.C. and N.W. conceived and designed the study; P.M. K.K. and J.K. performed

experiments; O.C. and J.B. performed the computational analyses; P.E., S.A.W., A.G.W.,

and N.W. performed clinical data abstraction and annotation; J.C., V.M., and M.C.

provided the clinical and genomic data for the 3 patients in the Foundation Medicine

cohort; F.P and D.R. helped with the design and execution of the ORF screen; E.P.W.,

N.U.L., and N.W. oversaw patient enrollment and sample collection on the metastatic

biopsy protocol; P.M., O.C. and N.W. wrote the manuscript with input from all authors;

N.W. supervised the study.

References
712

713   1.   Chen WY, Colditz GA: **Risk factors and hormone-receptor status:**
714   **epidemiology, risk-prediction models and treatment implications for breast**
715   **cancer**. *Nature clinical practice Oncology* 2007, **4**(7):415-423.

716   2.   Osborne CK, Schiff R: **Mechanisms of endocrine resistance in breast cancer**.
717   *Annual review of medicine* 2011, **62**:233-247.

718   3.   Lei JT, Gou X, Ellis MJ: **ESR1 fusions drive endocrine therapy resistance and**
719   **metastasis in breast cancer**. *Molecular & cellular oncology* 2018,
720   **5**(6):e1526005.

721   4.   Robinson DR, Wu YM, Vats P, Su F, Lonigro RJ, Cao X, Kalyana-Sundaram S,
722   Wang R, Ning Y, Hodges L *et al*: **Activating ESR1 mutations in hormone-**
723   **resistant metastatic breast cancer**. *Nature genetics* 2013, **45**(12):1446-1451.

724   5.   Toy W, Shen Y, Won H, Green B, Sakr RA, Will M, Li Z, Gala K, Fanning S,
725   King TA *et al*: **ESR1 ligand-binding domain mutations in hormone-resistant**
726   **breast cancer**. *Nature genetics* 2013, **45**(12):1439-1445.

727   6.   Merenbakh-Lamin K, Ben-Baruch N, Yeheskel A, Dvir A, Soussan-Gutman L,
728   Jeselsohn R, Yelensky R, Brown M, Miller VA, Sarid D *et al*: **D538G mutation**
729   **in estrogen receptor-alpha: A novel mechanism for acquired endocrine**
730   **resistance in breast cancer**. *Cancer Res* 2013, **73**(23):6856-6864.

731   7.   Jeselsohn R, Yelensky R, Buchwalter G, Frampton G, Meric-Bernstam F,
732   Gonzalez-Angulo AM, Ferrer-Lozano J, Perez-Fidalgo JA, Cristofanilli M,
733   Gomez H *et al*: **Emergence of constitutively active estrogen receptor-alpha**
734   **mutations in pretreated advanced estrogen receptor-positive breast cancer**.
735   *Clin Cancer Res* 2014, **20**(7):1757-1767.

736   8.   Garcia-Becerra R, Santos N, Diaz L, Camacho J: **Mechanisms of resistance to**
737   **endocrine therapy in breast cancer: focus on signaling pathways, miRNAs**
738   **and genetically based resistance**. *International journal of molecular sciences*
739   2012, **14**(1):108-145.

740   9.   Miller TW, Perez-Torres M, Narasanna A, Guix M, Stal O, Perez-Tenorio G,
741   Gonzalez-Angulo AM, Hennessy BT, Mills GB, Kennedy JP *et al*: **Loss of**
742   **Phosphatase and Tensin homologue deleted on chromosome 10 engages**
743   **ErbB3 and insulin-like growth factor-I receptor signaling to promote**
744   **antiestrogen resistance in breast cancer**. *Cancer Res* 2009, **69**(10):4192-4201.

745   10.   Massarweh S, Osborne CK, Jiang S, Wakeling AE, Rimawi M, Mohsin SK,
746   Hilsenbeck S, Schiff R: **Mechanisms of tumor regression and resistance to**
747   **estrogen deprivation and fulvestrant in a model of estrogen receptor-positive,**
748   **HER-2/neu-positive breast cancer**. *Cancer Res* 2006, **66**(16):8266-8273.

749   11.   Nayar U, Cohen O, Kapstad C, Cuoco MS, Waks AG, Wander SA, Painter C,
750   Freeman S, Persky NS, Marini L *et al*: **Acquired HER2 mutations in ER(+)**
751   **metastatic breast cancer confer resistance to estrogen receptor-directed**
752   **therapies**. *Nature genetics* 2018.

753   12.   Croessmann S, Formisano L, Kinch LN, Gonzalez-Ericsson PI, Sudhan DR, Nagy
754   RJ, Mathew A, Bernicker EH, Cristofanilli M, He J *et al*: **Combined Blockade of**
755   **Activating ERBB2 Mutations and ER Results in Synthetic Lethality of**
756   **ER+/HER2 Mutant Breast Cancer**. *Clin Cancer Res* 2019, **25**(1):277-289.

757   13.   Sokol ES, Feng YX, Jin DX, Basudan A, Lee AV, Atkinson JM, Chen J, Stephens
758   PJ, Frampton GM, Gupta PB *et al*: **Loss of function of NF1 is a mechanism of**
759   **acquired resistance to endocrine therapy in lobular breast cancer**. *Ann Oncol*
760   2019, **30**(1):115-123.

761   14.   Razavi P, Chang MT, Xu G, Bandlamudi C, Ross DS, Vasan N, Cai Y, Bielski
762   CM, Donoghue MTA, Jonsson P *et al*: **The Genomic Landscape of Endocrine-**
763   **Resistant Advanced Breast Cancers**. *Cancer cell* 2018, **34**(3):427-438 e426.

764   15.   Johannessen CM, Johnson LA, Piccioni F, Townes A, Frederick DT, Donahue
765   MK, Narayan R, Flaherty KT, Wargo JA, Root DE *et al*: **A melanocyte lineage**
766   **program confers resistance to MAP kinase pathway inhibition**. *Nature* 2013,
767   **504**(7478):138-142.

768   16.   Wilson FH, Johannessen CM, Piccioni F, Tamayo P, Kim JW, Van Allen EM,
769   Corsello SM, Capelletti M, Calles A, Butaney M *et al*: **A functional landscape of**
770   **resistance to ALK inhibition in lung cancer**. *Cancer cell* 2015, **27**(3):397-408.

771   17.   Le X, Antony R, Razavi P, Treacy DJ, Luo F, Ghandi M, Castel P, Scaltriti M,
772   Baselga J, Garraway LA: **Systematic Functional Characterization of**
773   **Resistance to PI3K Inhibition in Breast Cancer**. *Cancer discovery* 2016,
774   **6**(10):1134-1147.

775   18.   Gonzalez-Malerva L, Park J, Zou L, Hu Y, Moradpour Z, Pearlberg J, Sawyer J,
776   Stevens H, Harlow E, LaBaer J: **High-throughput ectopic expression screen for**
777   **tamoxifen resistance identifies an atypical kinase that blocks autophagy**.
778   *Proceedings of the National Academy of Sciences of the United States of America*
779   2011, **108**(5):2058-2063.

780   19.   Fox EM, Miller TW, Balko JM, Kuba MG, Sanchez V, Smith RA, Liu S,
781   Gonzalez-Angulo AM, Mills GB, Ye F *et al*: **A kinome-wide screen identifies**
782   **the insulin/IGF-I receptor pathway as a mechanism of escape from hormone**
783   **dependence in breast cancer**. *Cancer research* 2011, **71**(21):6773-6784.

784   20.   Mendes-Pereira AM, Sims D, Dexter T, Fenwick K, Assiotis I, Kozarewa I,
785   Mitsopoulos C, Hakas J, Zvelebil M, Lord CJ *et al*: **Genome-wide functional**
786   **screen identifies a compendium of genes affecting sensitivity to tamoxifen**.

Proceedings of the National Academy of Sciences of the United States of America
2012, **109**(8):2730-2735.

21.   Joseph JD, Darimont B, Zhou W, Arrazate A, Young A, Ingalla E, Walter K,
Blake RA, Nonomiya J, Guan Z *et al*: **The selective estrogen receptor**
downregulator GDC-0810 is efficacious in diverse models of ER+ breast
cancer. *eLife* 2016, **5**.

22.   Yang X, Boehm JS, Yang X, Salehi-Ashtiani K, Hao T, Shen Y, Lubonja R,
Thomas SR, Alkan O, Bhimdi T *et al*: **A public genome-scale lentiviral**
expression library of human ORFs. *Nature methods* 2011, **8**(8):659-661.

23.   Cohen O, Kim D, Oh C, Waks A, Oliver N, Helvie K, Marini L, Rotem A, Lloyd
M, Stover D *et al*: **Abstract S1-01: Whole exome and transcriptome**
sequencing of resistant ER+ metastatic breast cancer. *Cancer Research* 2017,
77(4 Supplement):S1-01.

24.   Cancer Genome Atlas N: **Comprehensive molecular portraits of human breast**
tumours. *Nature* 2012, **490**(7418):61-70.

25.   Gallo LH, Nelson KN, Meyer AN, Donoghue DJ: **Functions of Fibroblast**
Growth Factor Receptors in cancer defined by novel translocations and
mutations. *Cytokine Growth Factor Rev* 2015, **26**(4):425-449.

26.   Jeselsohn R, Buchwalter G, De Angelis C, Brown M, Schiff R: **ESR1 mutations-**
a mechanism for acquired endocrine resistance in breast cancer. *Nat Rev Clin*
Oncol 2015, **12**(10):573-583.

27.   Reintjes N, Li Y, Becker A, Rohmann E, Schmutzler R, Wollnik B: **Activating**
somatic FGFR2 mutations in breast cancer. *PloS one* 2013, **8**(3):e60264.

28.   Turner N, Pearson A, Sharpe R, Lambros M, Geyer F, Lopez-Garcia MA,
Natrajan R, Marchio C, Iorns E, Mackay A *et al*: **FGFR1 amplification drives**
endocrine therapy resistance and is a therapeutic target in breast cancer.
Cancer Res 2010, **70**(5):2085-2094.

29.   Shee K, Yang W, Hinds JW, Hampsch RA, Varn FS, Traphagen NA, Patel K,
Cheng C, Jenkins NP, Kettenbach AN *et al*: **Therapeutically targeting tumor**
microenvironment-mediated drug resistance in estrogen receptor-positive
breast cancer. *The Journal of experimental medicine* 2018, **215**(3):895-910.

30.   Byron SA, Chen H, Wortmann A, Loch D, Gartside MG, Dehkhoda F, Blais SP,
Neubert TA, Mohammadi M, Pollock PM: **The N550K/H mutations in FGFR2**
confer differential resistance to PD173074, dovitinib, and ponatinib ATP-
competitive inhibitors. *Neoplasia* 2013, **15**(8):975-988.

31.   Tan L, Wang J, Tanizaki J, Huang Z, Aref AR, Rusan M, Zhu SJ, Zhang Y, Ercan
D, Liao RG *et al*: **Development of covalent inhibitors that can overcome**

resistance to first-generation FGFR kinase inhibitors. *Proc Natl Acad Sci U S*
A 2014, **111**(45):E4869-4877.

32.   Yu M, Bardia A, Aceto N, Bersani F, Madden MW, Donaldson MC, Desai R,
Zhu H, Comaills V, Zheng Z *et al*: **Cancer therapy. Ex vivo culture of**
circulating breast tumor cells for individualized testing of drug susceptibility.
Science 2014, **345**(6193):216-220.

33.   AlFakeeh A, Brezden-Masley C: **Overcoming endocrine resistance in hormone**
receptor-positive breast cancer. *Curr Oncol* 2018, **25**(Suppl 1):S18-S27.

34.   Loh YN, Hedditch EL, Baker LA, Jary E, Ward RL, Ford CE: **The Wnt**
signalling pathway is upregulated in an in vitro model of acquired tamoxifen
resistant breast cancer. *BMC Cancer* 2013, **13**:174.

35.   Won HS, Lee KM, Oh JE, Nam EM, Lee KE: **Inhibition of beta-Catenin to**
Overcome Endocrine Resistance in Tamoxifen-Resistant Breast Cancer Cell
Line. *PloS one* 2016, **11**(5):e0155983.

36.   Vallabhaneni S, Nair BC, Cortez V, Challa R, Chakravarty D, Tekmal RR,
Vadlamudi RK: **Significance of ER-Src axis in hormonal therapy resistance**.
Breast cancer research and treatment 2011, **130**(2):377-385.

37.   Yates LR, Knappskog S, Wedge D, Farmery JHR, Gonzalez S, Martincorena I,
Alexandrov LB, Van Loo P, Haugland HK, Lilleng PK *et al*: **Genomic Evolution**
of Breast Cancer Metastasis and Relapse. *Cancer cell* 2017, **32**(2):169-184
e167.

38.   O'Leary B, Cutts RJ, Liu Y, Hrebien S, Huang X, Fenwick K, Andre F, Loibl S,
Loi S, Garcia-Murillas I *et al*: **The Genetic Landscape and Clonal Evolution of**
Breast Cancer Resistance to Palbociclib plus Fulvestrant in the PALOMA-3
Trial. *Cancer discovery* 2018, **8**(11):1390-1403.

39.   Musolino A, Campone M, Neven P, Denduluri N, Barrios CH, Cortes J,
Blackwell K, Soliman H, Kahan Z, Bonnefoi H *et al*: **Phase II, randomized,**
placebo-controlled study of dovitinib in combination with fulvestrant in
postmenopausal patients with HR(+), HER2(-) breast cancer that had
progressed during or after prior endocrine therapy. *Breast Cancer Res* 2017,
19(1):18.

40.   Andre F, Bachelot T, Campone M, Dalenc F, Perez-Garcia JM, Hurvitz SA,
Turner N, Rugo H, Smith JW, Deudon S *et al*: **Targeting FGFR with dovitinib**
(TKI258): preclinical and clinical data in breast cancer. *Clin Cancer Res*
2013, **19**(13):3693-3702.

41.   Smyth EC, Turner NC, Pearson A, Peckitt C, Chau I, Watkins DJ, Starling N, Rao
S, Gillbanks A, Kilgour E *et al*: **Phase II study of AZD4547 in FGFR amplified**

tumours: Gastroesophageal cancer (GC) cohort pharmacodynamic and
biomarker results. *Journal of Clinical Oncology* 2016, **34**(4_suppl):154-154.

42.   Babina IS, Turner NC: **Advances and challenges in targeting FGFR signalling**
in cancer. *Nat Rev Cancer* 2017, **17**(5):318-332.

43.   Chae YK, Ranganath K, Hammerman PS, Vaklavas C, Mohindra N, Kalyan A,
Matsangou M, Costa R, Carneiro B, Villaflor VM *et al*: **Inhibition of the**
fibroblast growth factor receptor (FGFR) pathway: the current landscape
and barriers to clinical application. *Oncotarget* 2017, **8**(9):16052-16074.

44.   Nogova L, Sequist LV, Perez Garcia JM, Andre F, Delord JP, Hidalgo M,
Schellens JH, Cassier PA, Camidge DR, Schuler M *et al*: **Evaluation of BGJ398,**
a Fibroblast Growth Factor Receptor 1-3 Kinase Inhibitor, in Patients With
Advanced Solid Tumors Harboring Genetic Alterations in Fibroblast
Growth Factor Receptors: Results of a Global Phase I, Dose-Escalation and
Dose-Expansion Study. *Journal of clinical oncology : official journal of the*
American Society of Clinical Oncology 2017, **35**(2):157-165.

45.   Matalkah F, Martin E, Zhao H, Agazie YM: **SHP2 acts both upstream and**
downstream of multiple receptor tyrosine kinases to promote basal-like and
triple-negative breast cancer. *Breast Cancer Res* 2016, **18**(1):2.

46.   Chen YN, LaMarche MJ, Chan HM, Fekkes P, Garcia-Fortanet J, Acker MG,
Antonakos B, Chen CH, Chen Z, Cooke VG *et al*: **Allosteric inhibition of SHP2**
phosphatase inhibits cancers driven by receptor tyrosine kinases. *Nature*
2016, **535**(7610):148-152.

47.   Adalsteinsson VA, Ha G, Freeman SS, Choudhury AD, Stover DG, Parsons HA,
Gydush G, Reed SC, Rotem D, Rhoades J *et al*: **Scalable whole-exome**
sequencing of cell-free DNA reveals high concordance with metastatic
tumors. *Nature communications* 2017, **8**(1):1324.

48.   Ulz P, Belic J, Graf R, Auer M, Lafer I, Fischereder K, Webersinke G, Pummer
K, Augustin H, Pichler M *et al*: **Whole-genome plasma sequencing reveals**
focal amplifications as a driving force in metastatic prostate cancer. *Nature*
communications 2016, **7**:12008.
891

Figure 1. A genome-scale gain-of-function screen identified resistance genes to fulvestrant and GDC-0810. 17,255 human open reading frames (ORFs), corresponding to 10,135 distinct genes, were expressed in ER+ T47D breast cancer cells in the presence of fulvestrant or GDC-0810. Fulvestrant, GDC-0810, or vehicle control (DMSO) was added following infection and selection. ORF representation was assessed by sequencing after 21 days of drug exposure. Genes that confer drug resistance will be enriched under drug selection, indicated by a positive log fold change (LFC) for ORF representation before and after DMSO/drug selection. A, The average Z score for LFC of each ORF was plotted for both the fulvestrant (X-axis) and GDC-0810 (Y-axis) arms. The average Z score was calculated from three replicates in each condition. The ORFs with a Z score > 3 in both drug arms are highlighted and labeled with gene ID. The shape of each data point represents the total number of ORFs for that gene in the library. B, Heatmap of top ORF hits with a Z score > 3 in fulvestrant or GDC-0810 arm. The Z score in the DMSO arm is also presented. ORF hits are grouped by their molecular function according to Uniprot annotation. Information on the complete list of ORFs can be found in Supplemental Table.1. C, Individual ORFs were overexpressed in T47D cells and validated to confer resistance to fulvestrant by drug response curves. *KRAS* G12D ORF was used for overexpression in T47D cells while other selected ORFs are wildtype. Cell viability was measured by CellTiter-Glo and all the data points were normalized to growth under DMSO condition. D, Gene set enrichment analysis was performed for the gene list ranked by LFC in the fulvestrant arm. For genes with multiple ORFs, the ORF with highest LFC was selected. 1000 permutations were performed for the analysis. NES, normalized enrichment score. The full list of nominated pathways is shown in Supplemental Table.2.

Figure 2

Figure 2. Identification of acquired FGFR and FGF alterations in metastatic biopsies from patients with resistant ER+ MBC. A, Evolutionary status of *ESR1*, *FGFR1*, *FGFR2*, and *FGF3/FGF4* alterations is presented by comparing the pre-treatment and post-treatment mutational status for each patient (red = acquired, blue = lost, black = shared, grey = indeterminate). Clinical and pathology tracks depict the site of biopsy for both matched samples, and the duration between the pre-treatment and post-treatment biopsies. B, Copy number alterations for *FGFR1*, *FGFR2* and *FGF3/FGF4* in pre- and post-treatment tumor samples are shown with copy number above ploidy (CNAP) depicted to illustrate the magnitude of the acquired amplification in each case. To better measure segment-specific copy number, we subtracted the genome ploidy for each sample to compute CNAP. The purity and ploidy for tumor samples are shown in Supplemental Table.4. C, Clonal evolution analysis showing the overall clonal structure and acquisition for FGFR2 mutations observed in two patients. In the pre-treatment biopsies, *FGFR2* M538I (*ID 0300348*) and *FGFR2* N550K (*ID 0300350*) were with cancer cell fraction (CCF) of 2% (single read) and 0% (unobserved), respectively, while being observed as clonal mutations in the post-treatment sample with a CCF of 1. The phylogenetic relationships among clones are reconstructed for each patient starting from the normal cell (white circle) connected to the ancestral cancer cells (grey trunk). The phylogenetic divergence to the pre-treatment clones (and subclones) is depicted with blue edges, and phylogenetic divergence to the metastatic clones (and subclones) is in red. Selected mutations in cancer genes are marked on the corresponding branches of the cancer phylogeny.

Figure 3

Figur...
patien...
therap...
infor...
the asterisk indicates the time that metastatic disease was diagnosed. The complete clinicopathologic information for each patient is provided in
Supplemental Table.7. IDC: invasive ductal carcinoma, IDLC: invasive ductal-lobular carcinoma; CNAP: copy number above ploidy; yo: years old;
Bx: biopsy; PR: progesterone receptor; wt: wildtype.

Figure 4

Figure 4. Active FGFR signaling leads to resistance to SERDs through activation of MAPK pathway. A, FGF ligands lead to resistance to fulvestrant, which was blocked by FGFR inhibitor PD173074. Recombinant FGF ligands were added into media every three days at the concentration of 100 ng/mL with or without 1 μM PD173014. T47D cells were treated with heparin (1 μg/mL) that facilitates the binding between FGF ligand and receptor, and sensitivity to 100 nM fulvestrant over six days was normalized to DMSO control. * p-value < 0.01. B, FGF ligands increased ERK and AKT phosphorylation, which was blocked by PD173074. T47D cells were treated as indicated for one hour before protein harvest for western blot. C, FGFR1 or FGFR2 overexpression leads to resistance to fulvestrant, which was blocked by PD173074. GFP, FGFR1 or FGFR2 was overexpressed in T47D cells to establish stable T47D_GFP, T47D_FGFR1 and T47D_FGFR2 cells. The fulvestrant sensitivity of various cell lines were determined in the presence or absence of 10 ng/mL FGF2 and 1 μM PD173074 over six days of drug treatment. D, FGFR1 and FGFR2 induced phosphorylation of ERK and AKT in the presence of FGF2, which was blocked by PD173074. Cells were treated with indicated conditions for one hour before protein harvest. E and F, Trametinib abrogated the resistance to fulvestrant conferred by FGFR1 or FGFR2. Cells were treated with different conditions as indicated: 10 ng/mL FGF2; 100 nM fulvestrant; 500 nM trametinib. CellTiter-Glo assay was performed to measure cell viability after six days for all dose response curves (E). 2,000 cells were plated on Day 1 and treated on Day 2, and colony formation assay was performed after three weeks of drug treatment (F). * p-value < 0.01, n.s. not significant.

Figure 5

Figure 5. **Active FGFR signaling leads to resistance to the combination of SERDs and CDK4/6 inhibitors.** A, T47D_GFP, T47D_FGFR1 and T47D_FGFR2 cells were treated with the combination of fulvestrant and palbociclib at various doses for six days. The percentage of cell survival shown was normalized to DMSO control for each condition. The combination treatment was carried out under three conditions: 1, control; 2, with 10 ng/mL FGF2; 3, with 10 ng/mL FGF2 and 1 μM PD173074 (FGFRi), with media refreshed every three days. B, MEK inhibitor trametinib abrogated the resistance to palbociclib alone (Top panel) or the combination of fulvestrant and palbociclib (Bottom panel) conferred by FGFR1 or FGFR2. Cells were treated with different conditions as indicated for six days before CellTiter-Glo assay. Concentration of drugs used: 10 ng/mL FGF2; 100 nM fulvestrant (Fulv); 1 μM palbociclib (Palbo); 500 nM trametinib. * p-value < 0.01. n.s. not significant. C, Trametinib blocked ERK phosphorylation and reduced CCND1 and p-Rb levels. Cells were treated as indicated daily for two days before protein harvest and western blot.

Figure 6

Figure 6. FGFR2 M538I, N5550K and K660N were activating mutations and conferred resistance to fulvestrant and/or palbociclib. A, Crystal structure of activated FGFR2 protein with mutations shown. FGFR2 is in complex with ATP analog (in yellow) and substrate peptide (PDB ID: 2PVF). B, Stable cell lines overexpressing FGFR2 wildtype (WT), M538I, N550K, and K660N were treated with 10 ng/mL FGF2 and/or 1 μM PD173014 for one hour before protein harvest. C, Tetracycline-inducible cell lines that express FGFR2 WT, M538I, N550K and K660N were established and treated with 100 ng/mL doxycycline (Dox) to induce gene expression. Before protein harvest, cells were treated with doxycycline 24 hours before FGF2 stimulation for 3 hours followed by another 3 hours of FGF2 stimulation. Cells were treated with heparin (1 μg/mL) that facilitates the binding between FGF2 and FGFR2. Protein lysates from T47D cells expressing GFP and T47D cells with constitutive overexpression of FGFR2 were used as control. D, Trametinib sensitivity of T47D cell lines expressing GFP or different FGFR2 constructs was examined. E and F, Stable cell lines constitutively overexpressing GFP or FGFR2 constructs (as previously described) were examined for sensitivity to fulvestrant (E) or combination of fulvestrant and palbociclib (F) with or without the treatment of FGF2 and/or PD173074. * p-value < 0.01, calculated as compared to GFP group in all conditions.

Figure 7

Figure 7. Activating FGFR2 mutations were targeted by irreversible kinase inhibitors FIIN-2 and FIIN-3. A, T47D cells overexpressing FGFR2 N550K cells were treated as indicated for three days and retreated for three hours before protein harvest and western blot. B, Stable cell lines overexpressing FGFR2 mutants (M538I, N550K and K660N) exhibited higher sensitivity to FGFR inhibitors AZD4547, FIIN-2 and FIIN-3, as compared to T47D cells overexpressing GFP or wildtype FGFR2. C, All stable cells lines expressing GFP or FGFR2 constructs were treated with fulvestrant under the following conditions: control, 10 ng/mL FGF2, 10 ng/mL FGF2 with 1 μM AZD4547, 10 ng/mL FGF2 with 1 μM FIIN-2, or 10 ng/mL FGF2 with 100 nM FIIN-3. Drug response curves were determined by CellTiter-Glo.