Can hot temperatures limit disease transmission? A test of mechanisms in a zooplankton-

fungus system

Marta S. Shocket[1][*][†] (mshocket@stanford.edu), Alexandra Magnante[1], Meghan A. Duffy[2], Carla

E. Cáceres[3], and Spencer R. Hall[1]

[1] Department of Biology, Indiana University, Bloomington, IN 47405 USA

[2] Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI

48109 USA

[3] School of Integrative Biology, University of Illinois at Urbana-Champaign, Urbana, IL 61801

* Corresponding Author: Marta S. Shocket, 650-723-5923, mshocket@ucla.edu

† Present Address: Department of Ecology and Evolutionary Biology, University of California,

Los Angeles, CA, USA

Keywords: *Daphnia dentifera*, fungal pathogen, host-parasite interaction, infectious disease,

Metschnikowia bicuspidata, temperature, thermal ecology, transmission rate, upper thermal

constraints, upper thermal limits

Author Contributions: SRH, CEC, and MAD designed and obtained funding for the field

survey. SRH and MSS collected field data. MSS and SRH designed the laboratory studies; MSS

and AM conducted them. MSS wrote the first draft of the manuscript, and all authors revised and

approved the final version.

Data Accessibility: Upon acceptance all data and code will be stored in Dryad Data Repository.

ABSTRACT (316 / 350 WORDS)

1. Thermal ecology theory predicts that transmission of infectious diseases should respond unimodally to temperature, i.e., be maximized at intermediate temperatures and constrained at extreme low and high temperatures. However, empirical evidence linking hot temperatures to decreased transmission in nature remains limited.

2. We tested the hypothesis that hot temperatures constrain transmission in a zooplankton-fungus (*Daphnia dentifera-Metschnikowia bicuspidata*) disease system where autumnal epidemics typically start after lakes cool from their peak summer temperatures. This pattern suggested that maximally hot summer temperatures could be inhibiting disease spread.

3. Using a series of lab experiments, we examined the effects of high temperatures on five mechanistic components of transmission. We found that (1) high temperatures increased exposure to parasites by speeding up foraging rate but (2) did not alter infection success post-exposure. (3) High temperatures lowered parasite production (due to faster host death and an inferred delay in parasite growth). (4) Parasites made in hot conditions were less infectious to the next host (instilling a parasite 'rearing' or 'trans-host' effect of temperature during the prior infection). (5) High temperatures in the free-living stage also reduce parasite infectivity, either by killing or harming parasites.

4. We then assembled the five mechanisms into an index of disease spread. The resulting unimodal thermal response was most strongly driven by the rearing effect. Transmission peaked at intermediate-hot temperatures (25-26°C) and then decreased at maximally hot temperatures (30-32°C). However, transmission at these maximally hot temperatures only trended slightly lower than the baseline control (20°C), which easily sustains epidemics

in laboratory conditions and in nature. Overall, we conclude that while exposure to hot

epilimnetic temperatures does somewhat constrain disease, we lack evidence that this

effect fully explains the lack of summer epidemics in this natural system. This work

demonstrates the importance of experimentally testing hypothesized mechanisms of

thermal constraints on disease transmission. Furthermore, it cautions against drawing

conclusions based on field patterns and theory alone.

INTRODUCTION

How do high temperatures affect the spread of infectious diseases? In the current prevailing view, warming from climate change will shift the geographic range of diseases: some new areas will become warm enough to support disease, whereas others that previously sustained disease will become too hot (Altizer, Ostfeld, Johnson, Kutz, & Harvell, 2013; Lafferty, 2009; Lafferty & Mordecai, 2016). This hypothesis stems from a principle of thermal biology: most biological traits have unimodal reaction norms, where performance peaks at intermediate temperatures and declines to zero at cooler and warmer temperatures (Dell, Pawar, & Savage, 2011). Thus, once temperatures exceed the thermal optima of traits driving transmission, disease should decline. Many models predict upper thermal constraints on diseases, e.g., helminthic ungulate parasites (Molnár, Kutz, Hoar, & Dobson, 2013), a rhizocephalan crab parasite (Gehman, Hall, & Byers, 2018), a microsporidian *Daphnia* parasite (Kirk et al., 2018), schistosomiasis (Mangal, Paterson, & Fenton, 2008), and mosquito-borne pathogens (Mordecai et al., *in press*). Additionally, there is evidence for upper thermal constraints on disease in natural populations of the crab parasite (Gehman et al., 2018), mosquito-borne pathogens (e.g., Shocket et al., *in review*) and fungi infecting grasshoppers (Carruthers, Larkin, & Firstencel, 1992), amphibians (Berger et al., 2004; Raffel, Michel, Sites, & Rohr, 2010), and bats (Langwig et al., 2015). However, temperature often co-varies with other seasonal environmental factors, so causally linking temperature to observed patterns of disease is challenging (Altizer et al., 2006; Pascual & Dobson, 2005). Thus, the generality of upper thermal constraints excluding disease remains unclear.

Conceptually, upper thermal constraints act like fever, taking advantage of a common thermal mismatch between hosts and parasites. Because hosts can often endure hotter

environments than their parasites, many animals increase their body temperature when infected

(see citations below). In ectotherms, fever arises from behavioral thermoregulation (microhabitat

selection) and is widespread, occurring in vertebrates (including amphibians, reptiles, and fish:

Rakus, Ronsmans, & Vanderplasschen, 2017), snails (Zbikowska, Wrotek, Cichy, & Kozak,

2013), and insects (including bees, flies, grasshoppers, mosquitoes, and beetles: Stahlschmidt &

Adamo, 2013; Thomas & Blanford, 2003). Behavioral fever can impair parasite performance,

enhancing clearance or reducing virulence of infection. An analogous process can occur within

ectothermic hosts inhabiting high ambient temperatures (regardless of infection status)—in

essence, an environmental fever. High ambient temperatures can also harm parasites with free-

living stages outside of hosts. Mechanistically linking high temperatures to reduced disease

requires examining thermal effects on components of the transmission process (McCallum et al.

2017). We use the term 'transmission (process)' to broadly refer to the full parasite life cycle,

including infective propagule production and propagule survival in the environment; we also use

'transmission rate' narrowly defined as the rate of new infections (i.e., the parameter 'β'

calculated from infection prevalence and densities of hosts and parasites; McCallum et al. 2017).

Here, we use a series of experiments to evaluate mechanisms for potential upper thermal

constraints on transmission in a planktonic-fungal disease system. Autumnal epidemics start

once lake waters cool below summer maxima (Fig. 1A). These delayed starts could reflect hot

temperatures inhibiting disease if they push any of five transmission components past their

thermal optima (Fig. 1B). First, hot temperatures could slow host feeding and lower

consumption-based exposure to parasites. Second, hot temperatures could reduce parasite

infectivity inside hosts, lowering the probability of successful infection (via effects on hosts

and/or parasites). Third, hot temperatures could decrease the quantity of parasite propagules

[spores] produced by an infection. This decrease could stem from slower host growth rate (since

parasite production often scales with host growth: Hall, Knight, et al., 2009; Hall, Simonis,

Nisbet, Tessier, & Cáceres, 2009), slower parasite growth independent from host growth, or

enhanced mortality of infected hosts (truncating production time; Auld, Hall, Housley Ochs,

Sebastian, & Duffy, 2014; Civitello, Forys, Johnson, & Hall, 2012). Fourth, hot temperatures

could lower the quality of parasite spores released from dead hosts into the environment

(Shocket, Vergara, et al., 2018). Finally, these free-living spores could be harmed or killed by

hot temperatures. Thus, high temperatures could constrain this fungal disease at multiple stages

of the transmission process.

108   **STUDY SYSTEM**

The hosts (*Daphnia dentifera*) are zooplankton grazers in freshwater temperate lakes

across the Midwestern United States; the fungal parasite *Metschnikowia biscupidata* causes

epidemics in some host populations, with prevalence reaching up to 60% (Penczykowski, Hall,

Civitello, & Duffy, 2014). Hosts become infected when they filter-feed on algae and

inadvertently consume fungal spores (Hall et al., 2007). The spores pierce the host's gut wall,

entering its body cavity. Inside, fungal conidia replicate in the hemolymph before maturing into

new spores (Stewart Merrill & Cáceres, 2018). Following host death, spores are released into the

water for new hosts to consume (Ebert, 2005).

The seasonality of epidemics motivated a focus on high temperatures. Epidemics

typically begin in late summer or early fall (August–October) and wane in late fall or early

winter (November–December; Penczykowski, Hall, et al., 2014). During this time, lake water

temperature declines (Shocket, Strauss, et al., 2018). Many traits that influence disease spread

(host demographic traits, transmission rate, and spore production) change plastically with

temperature (Hall, Tessier, Duffy, Huebner, & Cáceres, 2006; Shocket, Strauss, et al., 2018).

Transmission increases with constant temperatures up to 26°C, and hosts cannot be cultured in

constant temperatures above 27°C (Shocket, Strauss, et al., 2018). However, organisms can

withstand otherwise lethal temperatures in fluctuating environments (Niehaus, Angilletta, Sears,

Franklin, & Wilson, 2012). For instance, in our stratified lakes, hosts experience temperatures

exceeding 27°C in summer (typical maxima 29–32°C; Fig. 1A): they migrate between the colder,

deeper hypolimnion during day (to avoid mortality from visually-oriented fish predators) and the

warmer, upper epilimnion at night (to take advantage of greater algal resources and faster growth

in warmer temperatures) (Hall, Duffy, Tessier, & Cáceres, 2005; Lampert, 1989). Epidemics

often begin as lakes start cooling from maximum summer temperatures (Fig. 1A). This pattern

suggested that high temperatures could constrain disease spread, as predicted by theory (Lafferty,

2009; Lafferty & Mordecai, 2016).

METHODS

Field Survey

Field survey data generated the motivating pattern (Fig. 1A: the relationship between

epidemic start date and epilimnetic temperature). We surveyed 10–28 lakes in Indiana (Greene

and Sullivan Counties) weekly (2009–2011) or bi-weekly (2013–2014) from August to

December. For each visit, we collected a zooplankton sample (13 cm diameter net with 153 μm

mesh) and measured lake water temperature data at 0.5–1 meter intervals with a Hydrolab

multiprobe (Hach Environmental). For each sample, we visually diagnosed 400+ live hosts with

a dissecting scope (20-50X magnification). An epidemic 'started' when infection prevalence first

exceeded 1% for two consecutive sampling visits (Shocket, Strauss, et al., 2018). We calculated

the epilimnetic temperature by fitting a spline to temperature across water depth, and averaging

from the water surface to the depth where the temperature gradient first exceeded 1°C m^{-1} (i.e.,

the thermocline; see Hite et al. 2016 Appendix S2).

General Approach

We measured how high temperatures influence five components of the transmission

process with laboratory assays (Table 1). Then we combined them into a synthetic index of

disease spread: 'transmission potential' (Auld et al., 2014). For mechanisms involving the host or

host-parasite interaction (mechanisms 1-3: foraging rate [f], spore infectivity from within-host

processes [u], and spore yield [σ]), we used fluctuating temperatures to expose hosts to high

temperatures for part of the day (they cannot survive constant temperatures >27°C). Hosts were

kept on a 16:8 light:dark cycle. All hosts experienced the same 20°C temperature for 8 hours,

and then 20, 26, or 32°C for 16 hours ('maximum temperature'). For mechanisms 4-5 (rearing

effect on spore quality [ρ] and free-living spore effect [φ]), we conducted common garden

infection assays, exposing uniform hosts at constant 20°C to spores from different treatments.

Thus, variation in transmission rate can be attributed to differences in spore infectivity.

Temperatures varied slightly among experiments (25 or 26°C, 30 or 32°C) based on incubator

availability. For calculating transmission potential, we treat temperature categorically and pool

these treatments.

Due to time and incubator constraints, we were unable to replicate experiments across

multiple incubators. Thus, our temperature treatments are 'pseudo-replicated' in that all

replicates for a treatment were conducted in the same incubator at the same time. Accordingly,

our results may be influenced by random incubator effects.

Mechanisms 1 & 2: Foraging rate (*f*) and spore infectivity from within-host processes (*u*)

170   We measured foraging rate of hosts by comparing the fluorescence of ungrazed and

grazed algae (Penczykowski, Lemanski, et al., 2014; Sarnelle & Wilson, 2008). We added

estimates of foraging rate at 30°C to those at 20 and 25°C presented elsewhere using the same

methods (Shocket, Vergara, et al., 2018). In both experiments, we measured foraging rate across

a gradient of host body size (Kooijman, 2009) to index foraging at a common size among

experiments (1.5 mm). We used maximum likelihood estimation (MLE) to fit size-dependent

functions of foraging with the 'bbmle' package (Bolker & R Development Core Team, 2017) in

R (R Core Team, 2017). See Appendix for details.

178   We measured how high temperature impacts transmission rate (β) and spore infectivity

from within-host processes (*u*) with an infection assay ('β + *u* measurement assay'). For

successful infection, the fungus must break through the host gut barrier and then replicate and

develop within the host hemolymph. High temperatures could inhibit the parasite during either

process. Thus, we factorially manipulated the maximum temperature (20 and 32°C) during

parasite exposure and infection establishment (for four exposure/establishment treatments: 20/20,

20/32, 32/20, and 32/32°C) to reveal if high temperatures interfere at either step (similar to Allen

& Little, 2011). Hosts were exposed individually in their 'exposure temperature' for 24 hours,

then moved to their 'infection establishment temperature.' Later, hosts were visually diagnosed

for infection. Transmission rate was calculated from proportion infected (see Appendix). We

calculated spore infectivity from within-host processes (*u*) for each treatment by dividing

transmission rate (β) by foraging rate (*f*) at the exposure temperature ($u=\beta/f$).

Mechanism 3: Spore yield (σ) and related host and parasite traits

We measured how high temperatures impact final spore yield (σ) of infected hosts that died from their infection. This trait estimates spore release into the environment. We pooled spore yields from the $\beta + u$ measurement assay (above; treatments = 20/20 and 32/32°C) and the within-host parasite growth assay (below; treatments = 20, 26, and 32°C) since they did not differ statistically (20°C: p=0.65; 32°C p=0.93). We tested for differences between temperatures by fitting a suite of models via MLE: in each model spore yield was normally distributed and temperature treatments could exhibit the same or different means and standard deviations. We compared models using AIC and calculated p-values with likelihood ratio tests.

To distinguish between three possible mechanisms driving the thermal response of spore yield, we quantified related host and parasite traits. First, we measured host growth rate (g_h) with a juvenile growth rate assay (Lampert & Trubetskova, 1996)(see Appendix), since spore yield often scales with g_h (e.g., with different host food resources: Hall, Knight, et al., 2009; Hall, Simonis, et al., 2009). We compared treatments with t-tests. Second, we measured parasite growth (i.e., number of mature spores within hosts over time) using a sacrifice series ('within-host parasite growth assay;' see Appendix), since spore yield could decline if the number of parasites increases more slowly, independently of host condition (Thomas & Blanford, 2003). We fit and bootstrapped linear models of 'spore load' over time to estimate parasite growth rate (g_P, the model slope). 'Spore load' estimates included spores in living (i.e., sacrificed) hosts, unlike 'spore yield,' which was calculated only from dead hosts that were killed by the parasite. Spore yield is directly relevant for the epidemiology of the system, while spore load measures an underlying process (parasite growth rate per day, g_p) that contributes to spore yield. Spore load

increased linearly over the full time series at 26 and 32ºC. Spore load plateaued after day 19 at

20ºC, so we truncated the time series to estimate the linear slope for only that portion. Finally,

we calculated death rate (d) of infected hosts (see Appendix), since spore yield can decline with

shorter host lifespan (Auld et al., 2014; Civitello et al., 2012). We compared treatments with

randomization tests.

Mechanisms 4 & 5: rearing (ρ) and free-living spore (φ) effects on infectivity

We measured how high temperatures modify spore infectivity prior to encountering hosts

via a rearing effect on baseline spore quality (ρ) and harm to free-living spores (φ). We

conducted infection assays on 'common garden' groups of hosts at 20°C using different spore

treatments (i.e., from different spore rearing temperatures for ρ and from different spore

incubation temperatures for φ). Thus, variation in transmission rate reflects differences in spore

infectivity. To measure ρ, we conducted two experiments, one with spores produced in the $\beta + u$

measurement assay (20/20 and 32/32°C treatments) and another with spores produced in the

within-host parasite growth assay (20, 26, and 32°C treatments). To measure φ, we used spores

incubated at three temperatures (20, 25, and 30°C) for two durations (1-day and 7-days) in

constant, non-fluctuating temperatures (spores do not migrate between stratified water layers).

One-day incubations were stored at 4°C for the first six days (standard procedure for spore

storage). We estimated transmission rates (β) from the prevalence data (see Appendix).

Both mechanisms influence transmission by modifying spore infectivity (already

estimated from within-host processes as u, mechanism 1). Thus, in order to incorporate these

mechanisms into a synthetic metric for disease spread (transmission potential, see below), we

calculated unit-less rearing (ρ) and free-living (φ) effects standardized to infectivity at 20°C.

Specifically, we calculated the parameters by dividing the estimates for transmission rate (β) at

26 and 32°C by that at 20°C. Accordingly, values of $\rho<1$ or $\varphi<1$ mean spores are less infectious

due to rearing or free-living effects than at 20°C, respectively; conversely, values >1 mean

spores are more infectious than at 20°C. To calculate confidence intervals at 20°C, we divided a

bootstrapped distribution of transmission rates by a randomly-shuffled version of itself.

Additionally, because harm to free-living spores occurs over time as spores are removed by

hosts, we used a simple model to estimate time-weighted transmission rates for φ. We assumed

that spore infectivity declined linearly over the 7-day assay, and that hosts consume spores at a

constant foraging rate (resulting in an exponential decay in spores remaining over time). Thus,

we weighted the estimated transmission rate on each day by the proportion of spores consumed

by hosts on that day (see Appendix for detailed methods and a sensitivity analysis of the model).

Transmission potential

We calculated an index of disease spread to synthesize the effects of all five mechanisms.

We defined transmission potential as the product of all five parameters ($f\,u\,\sigma\,\rho\,\varphi$). We generated

confidence intervals using bootstrapped parameter distributions. To visualize the contribution of

each parameter, we calculated transmission potential for each of the five possible four-parameter

combinations, holding the fifth parameter constant at its 20°C point estimate. These values reveal

how each parameter affects the magnitude and uncertainty of transmission potential (i.e., a type

of sensitivity analysis).

Additional Statistical Analyses

For all parameters, we bootstrapped 95% confidence intervals (data sampled within

groups, with replacement; 10,000 samples). For parameters derived from transmission rates (β, u,

ρ, and φ), we used randomization tests to compare temperature treatments, since a single value is

calculated from all individuals (treatment labels shuffled among host individuals, without

replacement; 10,000 samples). For f and transmission potential (for which traditional statistical

tests were not available), we used the bootstrapped distributions to compare treatments.

Specifically, we calculated the cumulative probability density of the best estimate from one

treatment according to the bootstrapped distribution of the other. These 'PD-values' are

analogous to p-values. We considered treatments significantly different if $PD<0.025$. See

Appendix for details and a complete list of statistical tests and results.

269   **RESULTS**

270   **Mechanisms 1 & 2: Foraging rate (*f*) and spore infectivity from within-host processes (*u*)**

271   Contrary to our predictions, high temperature did not lower transmission rate (β) during

either step (exposure or infection establishment; Fig. 2A). Instead, transmission rate was higher

when hosts were exposed at 32°C than at 20°C (20°C infection establishment: $p=0.0013$; 32°C

infection establishment: $p<0.0001$). Temperature during infection establishment exerted no effect

on transmission rate (20°C exposure: $p=0.10$; 32°C exposure: $p=0.31$). When exposure and

establishment temperatures were equal (as in nature; the 20/20 and 32/32°C treatments here),

transmission rate was higher at 32°C than at 20°C ($p=0.0068$). Thus, even at maximal

epilimnetic temperatures, the impacts of higher temperatures on transmission rate promoted

rather than inhibited disease.

280   The thermal response of transmission rate was mechanistically driven by foraging rate of

hosts (*f*), not spore infectivity from within-host processes (*u*). Foraging rate increased from 20 to

25°C (*PD*=0; see *Methods* and Appendix for a description of PD values, which are analogous but

not identical to *p*-values) and then plateaued at 30°C (*PD*=0.11; Fig. 2B). Thus, hosts encounter

more spores at 25 and 30°C than at 20°C. After we accounted for predicted host-parasite contact,

spore infectivity was fairly insensitive to high temperatures (Fig. 2C). Temperature during

infection establishment did not impact spore infectivity (20°C exposure: *p*=0.10; 32°C exposure:

p=0.31). Exposure temperature increased spore infectivity (20°C infection establishment:

p=0.034; 32°C infection establishment: *p*=0.0052), but in the opposite direction of the

hypothesized mechanism (hotter temperature increased infectivity). When exposure and infection

establishment temperatures were equal (as in nature), spore infectivity did not differ (*p*=0.37).

Thus, high temperatures increased the foraging rate of hosts, elevating host contact with spores,

while spore infectivity barely changed. These changes in parasite exposure led to more

transmission at high temperatures.

Mechanism 3: Spore yield (σ) and other measures of host and parasite growth

Final spore yield (σ) in hosts that died from infection was lower at 32°C than at 20 and

26°C (Fig. 3A; best-fitting model had two means, see Table S5 for model AIC scores and Akaike

weights). This pattern was not explained by host condition estimated via growth rate. Host

growth rate (g_h, Fig. 3B) always increased with temperature (20 versus 26°C: $p=4.7 \times 10^{-6}$; 26 vs.

32°C: *p*=0.00038). Instead, the pattern was explained by a combination of host death rates and

delays in spore maturation. Infected hosts died more quickly at 26°C than 20°C (*p*<0.0001), and

death rate trended higher from 26 to 32°C (*p*=0.063; Fig. 3C). Meanwhile, growth rate of mature

parasite spores (g_p, time series in Fig. 3D, linear slopes [growth rate] in Fig. 3E) did not change

with temperature (*PD*>0.15). However, temperature did affect the timing of initial spore

production within hosts (i.e., intercepts of linear model). At the earliest point in the sacrifice

series (day 8), spore load was highest at 26°C, intermediate at 32°C, and nearly zero at 20°C

(Fig. 3D). Given thermally insensitive daily growth rates of parasites (g_P; Fig. 3E), these head-

starts were maintained over time (Fig. 3D). This effect on early spore production, coupled with

host death rate (Fig 3C), explains the spore yield pattern. Final spore yield was lower at 32 than

26°C because there were fewer spores initially (on day 8) and hosts died more quickly (less time

to produce spores). At 20°C, spore production started even later, but the delay was compensated

for by much longer lifespans of infected hosts (lower death rate, d; Fig. 3C).

Mechanisms 4 & 5: rearing (ρ) and free-living spore (φ) effects on infectivity

Spore infectivity (measured as transmission rate) responded unimodally to temperature in

the previous infection (rearing effect on spore quality, ρ; Fig. 4A). Infectivity increased

significantly for spores made at 20 versus 26°C for one of two spore sources (p=0.0083 for

spores from $\beta + u$ measurement assay [square, Fig 4A]; p=0.092 for spores from within-host

growth assay [diamond]). Infectivity then declined for spores made at 26 versus 32°C (p=0.0001

for both spore sources). Infectivity was significantly lower for spores made at 32 versus 20°C for

one of two spore sources (p=0.16 for spores from $\beta + u$ measurement assay [square]; p=0.026 for

spores from within-host growth assay [diamond]). The parameter ρ (Fig. 4C) shows the rearing

effect pooled for both spore sources and normalized by transmission rate at 20°C (used for

calculating transmission potential).

The thermal environment of free-living spores also impacted their infectivity (φ; Fig

4B,D). Spore infectivity decreased with higher incubation temperatures after 7 days (20 versus

25°C: p=0.0031; 25 versus 30°C: p<0.0001; diamonds on Fig. 4B). However, spore infectivity

did not change after 1-day incubations (flat line in Fig. 4B [squares]; 20 versus 25°C: p=0.65, 25

versus 30°C: p=0.64). All 1-day incubations used stored (refrigerated) spores. They had lower

infectivity than the 7-day incubation at 20°C, likely because storage at 4°C also lowers spore

infectivity (1 versus 7-day incubations at 20°C: p<0.0001) (Duffy & Hunsberger, 2019). The

parameter φ (Fig. 4D) shows the free-living spore effect assuming that spores lose infectivity

gradually over seven days as they are consumed by hosts (see *Methods* and Appendix) and

normalized by transmission rate at 20°C (used for calculating transmission potential).

Transmission potential ($f\,u\,\sigma\,\rho\,\varphi$)

Transmission potential, the product of all five mechanisms ($f\,u\,\sigma\,\rho\,\varphi$), responded

unimodally to high temperatures. This metric first increased from 20 to 25/26°C (PD=0.017);

then, it decreased from 25/26 to 30/32°C (PD=0.0001; Fig. 5A, 'full transmission potential').

Transmission potential at 30/32°C trended (non-significantly) lower than at 20°C (PD=0.11).

Thus, high temperatures do not constrain disease enough via these five mechanisms to explain

the absence of summer epidemics.

The initial increase in transmission potential from 20 to 25/26°C was driven most

strongly by host foraging (f, mechanism 1) and the rearing effect on spore quality (ρ, mechanism

4): holding either trait constant removes the significant difference between temperatures (Fig. 5B

and 5E, respectively). The subsequent drop in transmission potential from 25/26 to 30/32°C was

driven most strongly by the rearing effect (ρ): holding it constant again removes the significant

difference (Fig. 5E). Harm to free-living spores (φ, mechanism 5) also contributes somewhat

(Fig. 5F vs. Fig. 5A), though not enough to affect the statistical significance. Additionally, the

thermal response of host foraging (f) is key for maintaining transmission at high temperatures:

without increased exposure to spores, the remaining mechanisms would significantly reduce

transmission at 30/32°C compared to 20°C (Fig 5B). Spore infectivity from within-host

processes (u, mechanism 2) and spore yield (σ, mechanism 3) had no effect (Fig. 5C vs. Fig. 5A)

and very little effect (Fig. 5D vs. Fig. 5A) on transmission potential, respectively.

356   **DISCUSSION**

357   We investigated upper thermal constraints on fungal epidemics in a *Daphnia* zooplankton

host. The seasonality of the autumnal epidemics suggested that hot conditions might constrain

disease: epidemics usually start after lakes cool from maximal summer temperatures in the

epilimnion (29-32°C). We tested five potential thermal constraints on transmission. First,

foraging (exposure) rate of hosts (f) increased at high temperatures (Fig 2B), while, second, high

temperatures did not affect the infectivity of spores from within-host processes (u; Fig 2C). Thus,

high temperatures increased transmission rate, β (where $\beta=uf$; Fig 2A). Third, spore yield (σ)

declined slightly at 32°C (Fig 3A). Fourth, a rearing effect on spore quality driven by

temperature during the previous infection (ρ) emerged: spores made at 32°C were less infectious

than those made at 26°C and sometimes 20°C (for one of two spore sources, Fig 4A). Finally,

harm to free-living spores (φ) lowered infectivity as temperature increased (Fig 4B). Overall,

transmission potential is much lower at 32°C than 26°C, but still similar to at 20°C (Fig 5A), a

temperature that easily supports epidemics in both nature (Shocket, Strauss, et al., 2018) and

laboratory environments (Civitello et al., 2012; Shocket, Strauss, et al., 2018). Thus, maximally

high temperatures do constrain disease, but not sufficiently to explain the absence of summer

epidemics on their own.

373   Contrary to our initial hypothesis, high temperatures *increased* transmission rate (Fig

2A). In principle, high temperatures can lower infection success if pathogens tolerate heat less

well than hosts (Thomas & Blanford, 2003). For instance, many ectothermic hosts behaviorally

induce fever to reduce the negative costs of infection (Rakus et al., 2017; Stahlschmidt &

Adamo, 2013). Further, fungi are particularly sensitive to high temperatures compared to other

pathogen taxa (Robert & Casadevall, 2009) and fungal pathogens are often limited by high

temperatures (Berger et al., 2004; Carruthers et al., 1992; Langwig et al., 2015; Raffel et al.,

2010; Thomas & Blanford, 2003). However, high temperatures did not interfere with this

fungus's success at either stage of transmission: the day of exposure, when most spores penetrate

the host's gut, or infection establishment, when the fungus replicates and develops within the

host (Stewart Merrill & Cáceres, 2018). Instead, high temperatures elevated host foraging rate

(Fig 2B), which increases exposure to parasites, thereby increasing transmission rate (Hall et al.,

2007). In lakes, the thermal response of foraging (exposure) drives variation in the size of

epidemics, which occur in autumn: epidemics that start earlier in warmer conditions grow larger

than those starting later and colder (Shocket, Strauss, et al., 2018). This foraging-controlled

exposure to parasites is a potentially general mechanism: higher temperatures also increase

outbreak size for armyworms that consume virus particles on leaves (Elderd & Reilly, 2014).

However, transmission plateaued with temperature for another ingested *Daphnia* pathogen (Vale,

Stjernman, & Little, 2008).

Spore yield (σ) declined at the highest temperature (32°C; Fig. 3). Although the effect on

transmission potential was minimal (Fig 5D), the results for related traits provide mechanistic

insights into host-parasite interactions. Parasite burdens often decline at temperatures near the

thermal maxima of the host and/or parasite, e.g., for nematodes in slugs (Wilson, Digweed,

Brown, Ivanonva, & Hapca, 2015), trematodes in snails (Paull, Lafonte, & Johnson, 2012),

bacteria in *Daphnia* (Vale et al., 2008) and fruit flies (Lazzaro, Flores, Lorigan, & Yourth, 2008),

and powdery mildew in plants (Laine, 2007). In theory, reduced parasite production at hot

temperatures could arise from several mechanisms. First, parasite production could decline if

host growth slows, since spore yield often scales with host growth, at least along resource

gradients (Hall, Knight, et al., 2009; Hall, Simonis, et al., 2009). However, here host growth rate

(g_h) increased with temperature while spore yield was flat and then decreased (Fig 3B).

Therefore, spore production was decoupled from host growth rate (i.e., the link between host

growth and parasite production that occurs for resources did not occur for temperature). Second,

the parasite itself could grow more slowly at high temperatures. For example, high temperatures

slow bacterial growth inside fruit flies (Lazzaro et al., 2008), fungal growth in grasshoppers

(Springate & Thomas, 2005), and fungal growth on warm-adapted (but not cold-adapted)

amphibians (Cohen et al., 2017). In contrast, here parasite growth rate (g_p) did not respond to

temperature (slope in Fig 3D; Fig 3E).

Instead, the decline in spore production at high temperatures arose from a combination of

host death rate and the timing of initial spore production. Temperature determined spore load on

day 8 (the earliest sampling time in the assay; Fig. 3D). Based on that information (and the

constant parasite growth rates, Fig 3E), we infer that spore production began earliest at 26°C,

followed by 32°C, and then 20°C. These head starts were maintained over time and explain the

spore yield pattern when combined with death rate of infected hosts (Fig. 3C). In general, shorter

lifespan of infected hosts decreases time for spore production, thereby depressing spore yield

(Auld et al., 2014; Civitello et al., 2012). Here, spore yield was lower at 32 than 26°C because

spore production started later and hosts died more quickly. At 20°C, spore production started

even later, but longer host lifespan compensated for this delay (i.e., the fungus had longer to

grow within hosts). Do similar patterns exist in other systems? Unfortunately, few studies focus

on traits underlying thermal responses of parasite load. Hence questions remain: How often does

temperature change the timing versus the rate of parasite production? How often does

temperature decouple positive relationships between host growth and parasite production? The

answers matter because spore yield can influence epidemic size for obligate killer parasites (like

the fungus here: Civitello et al., 2015). Thus, developing a general framework from data across

host-parasite systems remains a fruitful area for future research.

High temperatures reduced transmission potential via two effects on spore infectivity that

act outside the focal host. First, a rearing effect on spore quality (ρ) driven by temperature of

spore production in the previous host elevated (26°C) and then lowered (32°C) spore infectivity

(compared to 20°C). Rearing effects on parasite performance can arise with variation in

resources consumed by hosts (Cornet, Bichet, Larcombe, Faivre, & Sorci, 2014; Little, Birch,

Vale, & Tseng, 2007; Tseng, 2006), temperature experienced by hosts (Shocket, Vergara, et al.,

2018), or host genotype (Searle et al., 2015). These understudied rearing effects may drive

performance of parasites to an unappreciated extent (Shocket, Vergara, et al., 2018). Second,

harm to free-living spores (φ, including spore mortality) also inhibited infection at high

temperatures. After seven days in 30°C, spores lost 92% of their initial infectivity. This

constraint may arise in other systems: for example, high temperatures elevate mortality in free-

living helminths of Arctic ungulates (Molnár et al., 2013). However, in the planktonic system

here, the 7-day result likely exaggerates the thermal constraint. While difficult to quantify,

physical sinking, consumption (Civitello, Pearsall, Duffy, & Hall, 2013; Penczykowski, Hall, et

al., 2014; Shocket, Vergara, et al., 2018; Strauss, Civitello, Cáceres, & Hall, 2015) and damage

from radiation (Overholt et al., 2012) likely remove most spores before 7 days. To acknowledge

this mortality, we weighted this component of infectivity (φ) using a model of spore longevity.

Assuming this modeled weighting reflects reality in lakes, the free-living effect lacks enough

strength to inhibit epidemics during summer, even when combined with the other mechanisms

(see Appendix for sensitivity analysis of the time-weighting model). However, more realistic

dynamical models and better resolved trait data for the free-living spore effect could change the

estimates for how high temperatures affect transmission.

Although the impact of temperature on these five mechanisms does not explain the lack

of epidemics during summer, other co-varying environmental factors could combine with

thermal effects to sufficiently inhibit transmission. Such factors include damage to free-living

spores by solar radiation (Overholt et al., 2012), consumption of spores by resistant zooplankton

species that are more abundant earlier in the year (Penczykowski, Hall, et al., 2014), and low

spore production due to poor quality of host food resources (Hall, Knight, et al., 2009). These

mechanisms could contribute to the observed field pattern, and interact with the thermal effects

examined here. Furthermore, climate change could disrupt covariation among drivers. For

example, high temperatures may persist later in the year when damaging solar radiation is less

intense. Incorporating these other factors may help explain the current field pattern and improve

predictions for how climate change will impact epidemics. These predictions should also

explicitly account for the effects of temperature variation and extremes, which have distinct

impacts on organismal performance (Dowd, King, & Denny, 2015). Here, we employed a

relevant form of thermal variation, mimicking migratory behavior of hosts in stratified lakes, but

did not isolate effects of thermal variation. Future efforts could estimate this effect to better

predict how climate change will impact the host, the parasite, and their interaction.

The current prevailing view argues that hot temperatures should constrain disease

transmission in nature (Altizer et al., 2013; Lafferty, 2009; Lafferty & Mordecai, 2016). This

constraint arises when unimodal thermal reaction norms depress key traits that drive disease

spread. However, such constraints have been rigorously tested in only a handful of systems.

Here, we hypothesized that high summer temperatures limit transmission of a zooplankton-

fungus disease system with autumnal epidemics (i.e., during cooler conditions). High

temperatures constrained disease transmission enough to produce a unimodal thermal response.

This response arose primarily through a rearing effect on spore quality and due to harm to free-

living spores. However, the thermal mechanisms estimated here were not sufficient explain the

lack of summer epidemics. Hence, we draw two major lessons. First, we need to continue to

rigorously evaluate multiple mechanisms of thermal constraints on components of disease

transmission. Second, our example cautions against drawing conclusions about constraints on

disease from warming based on field patterns and theory alone.

479   **ACKNOWLEDGMENTS**

K. Boatman (2009 and 2010), Z. Brown and K. Malins (2011), O. Schmidt (2013), A.

Bowling (2014), and P. Orlando and J. Walsman (2015) conducted field sampling. MSS was

supported by NSF GRFP. This work was supported by NSF DEB 0841679, 0841817, 1120316,

1120804, 1353749, and 1354407.

Table 1: The experiments (and spore sources) used to test the five mechanistic components of

disease transmission.

Mechanism	Experiment(s)	Spore source(s)
1. Foraging rate (f, Fig. 2B)	Foraging rate assay	NA
2. Within-host spore infectivity (u, Fig. 2C)	$\beta + u$ measurement assay Foraging rate assay	General lab stock
3. Spore yield (σ, Fig. 3A)	$\beta + u$ measurement assay Within-host parasite growth assay	General lab stock
4. Rearing effect on infectivity (ρ, Fig 4A,C)	Common garden infection assay #1	$\beta + u$ measurement assay Within-host parasite growth assay
5. Free-living spore effect on infectivity (φ, Fig 4B,D)	Common garden infection assay #2	General lab stock incubated at different temperatures

488
Figure 1: Motivating field pattern and mechanistic components of transmission. (A) Fungal
epidemics usually start (dark grey bars) after lakes have cooled from the maximum summer
temperature (light grey bars). Epidemics never started when the epilimnion (upper, warmer
layer) was hotter than 30°C, suggesting an upper thermal constraint. Data summarize 74
epidemics from 20 lakes in Indiana (USA) sampled from 2009-2015. (B) High temperature could
limit transmission via five mechanisms. 1-2) Hosts become infected at transmission rate β, which
can be divided into 1) host foraging rate (f), i.e., exposure to spores, and 2) spore infectivity, as
determined by within-host processes (u). 3) Parasite spores are produced at spore yield (σ). 4) A
rearing effect from temperature during the previous infection (ρ) determines initial spore
infectivity. 5) Harm to free-living spores (φ) might also impact their infectivity. The product of
all five components ($f\,u\,\sigma\,\rho\,\varphi$) determines 'transmission potential'.

Figure 2: High temperature impacts on transmission rate (*β*), foraging rate (*f*, mechanism 1) and spore infectivity from current within-host processes (*u*, mechanism 2). In A and C, the effect of high temperature during parasite exposure and infection establishment (20°C infection establishment = white circles, solid line; 32°C infection establishment = dark grey circles, dotted line). (A) Transmission rate (*β*) increased when hosts were exposed at 32°C and did not change with infection establishment temperature. For constant temperatures, transmission is higher at 32°C than at 20°C. (B) Foraging (exposure) rate of hosts (*f*) is higher at 26°C (light grey) and 32°C (dark grey) than at 20°C (white). (C) Spore infectivity (*u*=*β*/*f*) increased when hosts were exposed at 32°C for both infection establishment temperatures. However, for constant temperatures, infectivity did not differ between 20 and 32°C. Error bars show 95% CIs. Letters indicate significant differences.

Figure 3: High temperature impacts on spore yield (σ, mechanism 3) and possible underlying traits. (A) Final spore yield at host death (σ) was lower at 32°C (dark grey) than at 20°C (white) or 26°C (light grey). (B) Host growth rate (g_h) increased with temperature for all treatments. (C) Death rate of infected hosts (d) increased from 20 to 26°C and trended higher from 26 to 32°C. (D) Spore load within hosts through time at 32°C (dotted line), 26°C (dashed line), and 20°C (solid line), fit with linear models. 26°C points are shifted over for visual clarity of error bars. (E) Parasite growth rate (g_p, slopes of lines in panel D) did not change with temperature. Hence, declining σ stems from higher death rate of infected hosts and low initial parasite growth, not slower growth rates of hosts or parasites. (A-C,E) Error bars show 95% CIs. (D) Error bars show SE; square points are single hosts. Letters indicate significant differences.

536
Figure 4: High temperature impacts on a rearing effect (ρ, mechanism 4) and harm to free-
living spores (φ, mechanism 5). Variation in transmission rate from common garden infection
assays reflects differences in spore infectivity. (A) Spores came from the $\beta + u$ measurement
assay (Fig. 2; squares) and the within host parasite growth assay ('WHPG'; Fig. 3; diamonds).
Spore infectivity increased with rearing temperature from 20°C (white) to 26°C (light grey; $\beta + u$
only) and decreased with rearing temperatures from 26°C to 32°C (dark grey, both spore
sources). Spore infectivity was lower at 32°C than at 20°C (WHPG spores only). (B) Spore
infectivity decreased when free-living spores were incubated in high temperatures for 7 days but
not for 1 day. Storage at 4°C for 6 days (for all 1-day incubations) also lowered spore infectivity
relative to the 7-day incubation at 20°C. (C-D) Parameter values (transmission rate scaled by
values at 20°C) for (C) rearing effect, ρ, and (D) free-living effect, φ. Phi values also based on
time-weighted model (see text for details). Error bars show 95% CIs. Letters indicate significant
differences.

Figure 5: High temperature impacts on transmission potential. (A) Transmission potential
(*fuσρφ*) responds unimodally, increasing from 20°C (white) to 25/26°C (light grey) and
decreasing from 25/26°C to 30/32°C (dark grey). (B-F) Transmission potential with each
mechanism held constant to show sensitivity to each parameter: (B) foraging rate (*f*), (C) spore
infectivity from within-host effects (*u*), (D) spore yield (*σ*), (E) rearing effect (*ρ*), and (F) harm
free-living spores (*φ*). The rearing effect (E) has the largest impact on transmission potential
(hence, without it, the response of transmission potential is flat with temperature). Error bars
show 95% CIs. Letters indicate significant differences. Y-axis is ln-transformed.

559   **WORKS CITED**

Allen, D. E., & Little, T. J. (2011). Dissecting the effect of a heterogeneous environment on the
interaction between host and parasite fitness traits. *Evolutionary Ecology*, *25*(2), 499–508.
doi:10.1007/s10682-010-9413-2

Altizer, S., Dobson, A., Hosseini, P., Hudson, P., Pascual, M., & Rohani, P. (2006). Seasonality
and the dynamics of infectious diseases. *Ecology Letters*, *9*(4), 467–84. doi:10.1111/j.1461-
0248.2005.00879.x

Altizer, S., Ostfeld, R. S., Johnson, P. T. J., Kutz, S., & Harvell, C. D. (2013). Climate change
and infectious diseases: from evidence to a predictive framework. *Science*, *341*(6145), 514–
9. doi:10.1126/science.1239401

Auld, S. K., Hall, S. R., Housley Ochs, J., Sebastian, M., & Duffy, M. A. (2014). Predators and
patterns of within-host growth can mediate both among-host competition and evolution of
transmission potential of parasites. *American Naturalist*, *184*(S1), S77-90.
doi:10.1086/676927

Berger, L., Speare, R., Hines, H. B., Marantelli, G., Hyatt, A. D., McDonald, K. R., … Tyler, M.
J. (2004). Effect of season and temperature on mortality in amphibians due to
chytridiomycosis. *Australian Veterinary Journal*, *82*(7), 434–439. doi:10.1111/j.1751-
0813.2004.tb11137.x

Bolker, B. M., & R Development Core Team. (2017). bbmle: Tools for General Maximum
Likelihood Estimation.

Carruthers, R. I., Larkin, T. S., & Firstencel, H. (1992). Influence of thermal ecology on the
mycosis of a rangeland grasshopper. *Ecology*, *73*(1), 190–204. doi:10.2307/1938731

Civitello, D. J., Forys, P., Johnson, A. P., & Hall, S. R. (2012). Chronic contamination decreases
disease spread: a Daphnia-fungus-copper case study. *Proceedings of the Royal Society B:*
Biological Sciences, *279*(May), 3146–3153. doi:10.1098/rspb.2012.0684

Civitello, D. J., Pearsall, S., Duffy, M. A., & Hall, S. R. (2013). Parasite consumption and host
interference can inhibit disease spread in dense populations. *Ecology Letters*, *16*(5), 626–34.
doi:10.1111/ele.12089

Civitello, D. J., Penczykowski, R. M., Smith, A. N., Shocket, M. S., Duffy, M. A., & Hall, S. R.
(2015). Resources, key traits and the size of fungal epidemics in Daphnia populations.
Journal of Animal Ecology, *84*, 1010–1017. doi:10.1111/1365-2656.12363

Cohen, J. M., Venesky, M. D., Sauer, E. L., Civitello, D. J., McMahon, T. A., Roznik, E. A., &
Rohr, J. R. (2017). The thermal mismatch hypothesis explains host susceptibility to an
emerging infectious disease. *Ecology Letters*, *20*(2), 184–193. doi:10.1111/ele.12720

Cornet, S., Bichet, C., Larcombe, S., Faivre, B., & Sorci, G. (2014). Impact of host nutritional
status on infection dynamics and parasite virulence in a bird-malaria system. *Journal of*
Animal Ecology, *83*(1), 256–265. doi:10.1111/1365-2656.12113

Dell, A. I., Pawar, S., & Savage, V. M. (2011). Systematic variation in the temperature
dependence of physiological and ecological traits. *Proceedings of the National Academy of*
Sciences, *108*(26), 10591–10596.

Dowd, W. W., King, F. A., & Denny, M. W. (2015). Thermal variation, thermal extremes and
the physiological performance of individuals. *Journal of Experimental Biology*, *218*(12),
1956–1967. doi:10.1242/jeb.114926

Duffy, M. A., & Hunsberger, K. K. (2019). Infectivity is influenced by parasite spore age and
exposure to freezing: do shallow waters provide Daphnia a refuge from some parasites?
Journal of Plankton Research, *41*(1), 12–16. doi:10.1093/plankt/fby046

Ebert, D. (2005). *Ecology, epidemiology, and evolution of parasitism in* Daphnia. Bethesda:
National Library of Medicine (USA), Center for Biotechnology Information.

Elderd, B. D., & Reilly, J. R. (2014). Warmer temperatures increase disease transmission and
outbreak intensity in a host-pathogen system. *Journal of Animal Ecology*, *83*(4), 838–849.
doi:10.1111/1365-2656.12180

Gehman, A.-L. M., Hall, R. J., & Byers, J. E. (2018). Host and parasite thermal ecology jointly
determine the effect of climate warming on epidemic dynamics. *Proceedings of the*
National Academy of Sciences, *115*(4), 744–749. doi:10.1073/pnas.1705067115

Hall, S. R., Duffy, M. A., Tessier, A. J., & Cáceres, C. E. (2005). Spatial heterogeneity of
daphniid parasitism within lakes. *Oecologia*, *143*, 635–644. doi:10.1007/s00442-005-0005-
8

Hall, S. R., Knight, C. J., Becker, C. R., Duffy, M. A., Tessier, A. J., & Cáceres, C. E. (2009).
Quality matters: resource quality for hosts and the timing of epidemics. *Ecology Letters*,
12(2), 118–128. doi:10.1111/j.1461-0248.2008.01264.x

Hall, S. R., Simonis, J. L., Nisbet, R. M., Tessier, A. J., & Cáceres, C. E. (2009). Resource
Ecology of Virulence in a Planktonic Host-Parasite System: An Explanation Using
Dynamic Energy Budgets. *The American Naturalist*, *174*(2), 149–162. doi:10.1086/600086

Hall, S. R., Sivars-Becker, L., Becker, C., Duffy, M. A., Tessier, A. J., & Cáceres, C. E. (2007).
Eating yourself sick : transmission of disease as a function of foraging ecology. *Ecology*
Letters, *10*, 207–218. doi:10.1111/j.1461-0248.2006.01011.x

Hall, S. R., Tessier, A. J., Duffy, M. A., Huebner, M., & Cáceres, C. E. (2006). Warmer does not
have to mean sicker: temperature and predators can jointly drive timing of epidemics.
Ecology, *87*(7), 1684–95.

Hite, J. L., Penczykowski, R. M., Shocket, M. S., Strauss, A. T., Orlando, P. A., Duffy, M. A., …
Hall, S. R. (2016). Parasites destabilize host populations by shifting stage-structured
interactions. *Ecology*, *97*(2), 439–449. doi:10.1890/15-1065.1

Kirk, D., Jones, N., Peacock, S., Phillips, J., Molnár, P. K., Krkošek, M., & Luijckx, P. (2018).
Empirical evidence that metabolic theory describes the temperature dependency of within-
host parasite dynamics. *PLOS Biology*, *16*(2), e2004608. doi:10.1371/journal.pbio.2004608

Kooijman, S. A. L. M. (2009). *Dynamic Energy Budget Theory for Metabolic Organisation* (3rd
ed.). New York, New York: Cambridge University Press.

Lafferty, K. D. (2009). The ecology of climate change and infectious diseases. *Ecology*, *90*(4),
888–900. doi:10.1890/09-1656.1

Lafferty, K. D., & Mordecai, E. A. (2016). The rise and fall of infectious disease in a warmer
world. *F1000Research*, *5*.

Laine, A. L. (2007). Pathogen fitness components and genotypes differ in their sensitivity to
nutrient and temperature variation in a wild plant-pathogen association. *Journal of
Evolutionary Biology*, *20*(6), 2371–2378. doi:10.1111/j.1420-9101.2007.01406.x

Lampert, W. (1989). The adaptive significance of diel vertical migration of zooplankton.
Functional Ecology, *3*(1), 21–27.

Lampert, W., & Trubetskova, I. (1996). Juvenile growth rate as a measure of fitness in Daphnia.
Functional Ecology, *10*(5), 631–635.

Langwig, K. E., Frick, W. F., Reynolds, R., Parise, K. L., Drees, K. P., Hoyt, J. R., … Kilpatrick,
A. M. (2015). Host and pathogen ecology drive the seasonal dynamics of a fungal disease,
white-nose syndrome. *Proceedings of the Royal Society B: Biological Sciences*, *282*(1799),
20142335. doi:10.1098/rspb.2014.2335

Lazzaro, B. P., Flores, H. A., Lorigan, J. G., & Yourth, C. P. (2008). Genotype-by-environment
interactions and adaptation to local temperature affect immunity and fecundity in
Drosophila melanogaster. *PLoS Pathogens*, *4*(3), 1–9. doi:10.1371/journal.ppat.1000025

Little, T., Birch, J., Vale, P., & Tseng, M. (2007). Parasite transgenerational effects on infection.
Evolutionary Ecology Research, *9*(3), 459–469.

Mangal, T. D., Paterson, S., & Fenton, A. (2008). Predicting the impact of long-term temperature
changes on the epidemiology and control of schistosomiasis: a mechanistic model. *PloS
One*, *3*(1), e1438. doi:10.1371/journal.pone.0001438

McCallum, H., Fenton, A., Hudson, P. J., Lee, B., Levick, B., Norman, R., … Lello, J. (2017).
Breaking beta: deconstructing the parasite transmission function. *Philosophical
Transactions of the Royal Society B: Biological Sciences*, *372*, 20160084.
doi:10.1098/rstb.2016.0084

Molnár, P. K., Kutz, S. J., Hoar, B. M., & Dobson, A. P. (2013). Metabolic approaches to
understanding climate change impacts on seasonal host-macroparasite dynamics. *Ecology
Letters*, *16*(1), 9–21. doi:10.1111/ele.12022

Mordecai, E. A., Caldwell, J. M., Grossman, M. K., Lippi, C. A., Johnson, L. R., Neira, M., …
Villena, O. (n.d.). Thermal biology of mosquito-borne disease. *Ecology Letters*.

Niehaus, A. C., Angilletta, M. J., Sears, M. W., Franklin, C. E., & Wilson, R. S. (2012).
Predicting the physiological performance of ectotherms in fluctuating thermal
environments. *The Journal of Experimental Biology*, *215*(Pt 4), 694–701.

671   doi:10.1242/jeb.058032

672 Overholt, E. P., Hall, S. R., Williamson, C. E., Meikle, C. K., Duffy, M. A., & Cáceres, C. E.
673   (2012). Solar radiation decreases parasitism in Daphnia. *Ecology Letters*, *15*(1), 47–54.
674   doi:10.1111/j.1461-0248.2011.01707.x

675 Pascual, M., & Dobson, A. (2005). Seasonal patterns of infectious diseases. *PLoS Medicine*,
676   *2*(1), 0018–0020. doi:10.1371/journal.pmed.0020005

677 Paull, S. H., Lafonte, B. E., & Johnson, P. T. J. (2012). Temperature-driven shifts in a host-
678   parasite interaction drive nonlinear changes in disease risk. *Global Change Biology*, *18*(12),
679   3558–3567. doi:10.1111/gcb.12018

680 Penczykowski, R. M., Hall, S. R., Civitello, D. J., & Duffy, M. A. (2014). Habitat structure and
681   ecological drivers of disease. *Limnology and Oceanography*, *59*(2), 340–348.
682   doi:10.4319/lo.2014.59.2.0340

683 Penczykowski, R. M., Lemanski, B. C. P., Sieg, R. D., Hall, S. R., Housley Ochs, J., Kubanek,
684   J., & Duffy, M. A. (2014). Poor resource quality lowers transmission potential by changing
685   foraging behaviour. *Functional Ecology*, *28*(5), 1245–1255. doi:10.1111/1365-2435.12238

686 R Core Team. (2017). *R: A language and for statistical computing*. Vienna, Austria: R
687   Foundation for Statistical Computing.

688 Raffel, T. R., Michel, P. J., Sites, E. W., & Rohr, J. R. (2010). What Drives Chytrid Infections in
689   Newt Populations? Associations with Substrate, Temperature, and Shade. *EcoHealth*, *7*(4),
690   526–536. doi:10.1007/s10393-010-0358-2

691 Rakus, K., Ronsmans, M., & Vanderplasschen, A. (2017). Behavioral fever in ectothermic
692   vertebrates. *Developmental & Comparative Immunology*, *66*, 84–91.
693   doi:https://doi.org/10.1016/j.dci.2016.06.027

694 Robert, V. A., & Casadevall, A. (2009). Vertebrate Endothermy Restricts Most Fungi as
695   Potential Pathogens. *Journal of Infectious Diseases*, *200*(10), 1623–1626.
696   doi:10.1086/644642

697 Sarnelle, O., & Wilson, A. E. (2008). Type III functional response in *Daphnia*. *Ecology*, *89*(6),
698   1723–1732. doi:10.1890/07-0935.1

699 Searle, C. L., Ochs, J. H., Cáceres, C. E., Chiang, S. L., Gerardo, N. M., Hall, S. R., & Duffy, M.
700   A. (2015). Plasticity, not genetic variation, drives infection success of a fungal parasite.
701   *Parasitology*, *142*(06), 839–848. doi:10.1017/S0031182015000013

702 Shocket, M. S., Strauss, A. T., Hite, J. L., Šljivar, M., Civitello, D. J., Duffy, M. A., … Hall, S.
703   R. (2018). Temperature Drives Epidemics in a Zooplankton-Fungus Disease System: A
704   Trait-Driven Approach Points to Transmission via Host Foraging. *The American Naturalist*,
705   *191*(4), 435–451. doi:10.1086/696096

706 Shocket, M. S., Vergara, D., Sickbert, A. J., Walsman, J. M., Strauss, A. T., Hite, J. L., … Hall,
707   S. R. (2018). Parasite rearing and infection temperatures jointly influence disease

transmission and shape seasonality of epidemics. *Ecology*. doi:10.1002/ecy.2430

Shocket, M. S., Verwillow, A. B., Numazu, M. G., Slamani, H., Cohen, J. M., El Moustaid, F.,
… Mordecai, E. A. (n.d.). Transmission of West Nile virus and other temperate mosquito-
borne viruses occurs at lower environmental temperatures than tropical mosquito-borne
diseases.

Springate, S., & Thomas, M. B. (2005). Thermal biology of the meadow grasshopper,
Chorthippus parallelus, and the implications for resistance to disease. *Ecological*
Entomology, *30*(6), 724–732. doi:10.1111/j.0307-6946.2005.00743.x

Stahlschmidt, Z. R., & Adamo, S. A. (2013). Context dependency and generality of fever in
insects. *Naturwissenschaften*, *100*(7), 691–696. doi:10.1007/s00114-013-1057-y

Stewart Merrill, T. E., & Cáceres, C. E. (2018). Within-host complexity of a plankton-parasite
interaction. *Ecology*.

Strauss, A. T., Civitello, D. J., Cáceres, C. E., & Hall, S. R. (2015). Success, failure and
ambiguity of the dilution effect among competitors. *Ecology Letters*, *18*(9), 916–926.
doi:10.1111/ele.12468

Thomas, M. B., & Blanford, S. (2003). Thermal biology in insect-parasite interactions. *Trends in*
Ecology & Evolution, *18*(7), 344–350. doi:10.1016/s0169-5347(03)00069-7

Tseng, M. (2006). Interactions between the parasite's previous and current environment mediate
the outcome of parasite infection. *The American Naturalist*, *168*(4), 565–571.
doi:10.1086/507997

Vale, P. F., Stjernman, M., & Little, T. J. (2008). Temperature-dependent costs of parasitism and
maintenance of polymorphism under genotype-by-environment interactions. *Journal of*
Evolutionary Biology, *21*(5), 1418–1427. doi:10.1111/j.1420-9101.2008.01555.x

Wilson, M. J., Digweed, A. J., Brown, J., Ivanonva, E. S., & Hapca, S. H. (2015). Invasive slug
pests and their parasites—temperature responses and potential implications of climate
change. *Biology and Fertility of Soils*, *51*(6), 739–748. doi:10.1007/s00374-015-1022-3

Zbikowska, E., Wrotek, S., Cichy, A., & Kozak, W. (2013). Thermal preferences of wintering
snails Planorbarius corneus (L.) exposed to lipopolysaccharide and zymosan. *Journal of*
Invertebrate Pathology, *112*(1), 57–61. doi:10.1016/j.jip.2012.08.011

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of this article.

Appendix S1: Methods, Figures, and Tables

Figure S1: Components of a simple model used to estimate parameter φ.

Figure S2: Sensitivity analysis for spore consumption model parameter (c) affecting damage to
free-living spores (φ) and transmission potential.

Table S1: Raw results from the $\beta + u$ measurement assay.

Table S2: Sample sizes for estimating spore yield (σ) and related traits.

Table S3: Raw results from the rearing effect (ρ) and free-living spore effect (φ) assays.

Table S4: p-values from randomization tests.

Table S5: p-values and ΔAIC from model selection.

Table S6: PD (probability density) values for traits.

Table S7: PD (probability density) values for transmission potential 'sensitivity analysis'
calculations.