1 **Environmental drivers define contrasting microbial habitats, diversity, and community**
2 **structure in Lake Baikal, Siberia**
3

4 Paul Wilburn[1,2,3], Kirill Shchapov[4], Edward C. Theriot[5], Elena Litchman[1,2,3]
5
[1]W.K. Kellogg Biological Station, Michigan State University, USA
[2]Department of Integrative Biology, Michigan State University, USA
[3]Ecology, Evolutionary Biology, and Behavior Graduate Program, Michigan State University,
USA
[4]Large Lakes Observatory, University of Minnesota, Duluth, USA
[5]University of Texas at Austin, USA

ABSTRACT

Understanding how microbial communities respond to environmental change requires knowing

the main drivers of their structure, diversity and potential resilience. Lake Baikal is the world's

most ancient, deep, voluminous, and biodiverse lake, holding 20 percent of unfrozen fresh water

that is undergoing rapid warming. Despite its global importance, little is known about Baikal's

bacterioplankton communities and their drivers. In this extensive survey, we show that

temperature, along with stratification, nutrients, and dissolved oxygen, but not geographic

distance, define major microbial habitats and community similarity. Mixed layer and deep water

communities exhibited contrasting patterns of richness, diversity and evenness, and comprised

different cohesive modules in the whole Baikal OTU co-occurrence network. The network's

small-world properties indicated likely resistance to perturbations but sensitivity to abundance

changes of central, most connected OTUs. Previous studies showed Baikal water temperature

rising by over 1.2°C since 1946, and this trend is predicted to accelerate. Because temperature

emerged as the most significant predictor of the mixed layer community structure, we

hypothesize that it is most likely to drive future community changes. Understanding how

temperature and other abiotic factors structure microbial communities in this and other rapidly

changing ecosystems will allow better predictions of ecosystem responses to anthropogenic

stressors.

<u>INTRODUCTION</u>

The ecological importance of microorganisms in aquatic systems has been recognized at least

since the appearance of "ooze" in Lindeman's trophic energy transfer diagram [1]. Their central

place in material and energy fluxes is now recognized for nearly all nutrient cycles, with greater

relative importance of prokaryotic organisms in more oligotrophic systems [2]. The advent of

next-generation sequencing, combined with environmental monitoring, enabled new discoveries

of microbial diversity and function in various aquatic habitats. However, the environmental

drivers of microbial community diversity, community structure, function, and stability remain

poorly characterized in many aquatic ecosystems, including the world's most ancient (25 My)

lake Baikal – a UNESCO heritage site and known hotspot for endemism of its biota. Baikal is the

world's deepest (1643 m) and most voluminous lake, holding about 20% of world's surface

unfrozen freshwater [3]. Of the approximately 2600 plant and animal species in the lake, two-

thirds are endemic, including the dominant primary producers, phytoplankton, zooplankton

grazers, benthic and pelagic fish and the top predator – world's only freshwater seal [3, 4].

Hampton *et al.* [4] showed that Baikal water temperatures have risen by 1.2 °C over 60 years

in a high-resolution time series, contributing to an increase in the abundance of non-endemic

zooplankton and algal species, with potential consequences for nutrient cycling, food web

structure [3] and microbial communities. Moreover, Lake Baikal region is predicted to warm by

3-4°C in the next century [5], with ongoing changes likely to continue and even accelerate.

Because the biota of Lake Baikal, including microbial communities, are adapted to cold

temperatures, it may be especially vulnerable to warming. Additionally, other changing

environmental factors may also alter the lake's microbial communities.

Despite the remarkable size of this freshwater ecosystem, its uniqueness and global

significance as a natural evolution experiment and the single largest reservoir of unfrozen

freshwater, the microbial communities in the lake are just beginning to be characterized.

Historically, culture-dependent efforts focused on sewage runoff monitoring, with one

amplicon effort using a taxonomically targeted approach [6]. Culture-independent efforts have

explored oil seeps [7, 8], sediments, invertebrate gut [9], diatom-associated [10] and pelagic

communities, with the latter on Sanger [11, 12] and 454 [13–15] platforms. Kurilkina and

colleagues sampled a one station depth profile in September and June [14]. In both seasons, the

main taxonomic groups included Proteobacteria, Actinobacteria, Bacteroidetes, Firmicutes,

Chloroflexi, Acidobacteria, Cyanobacteria, while Caldiserica was substantially represented in

September only.

Recent culture-independent studies emphasized analyses of metagenome-assembled genomes

(MAGs), where contigs assembled from metagenomic shotgun sequences are binned into draft

genomes. While these approaches offer additional insights into genes and function, they have

been limited in scope to 1-5 samples per study. This lack of replication inherently limits the

ability to connect genomic data with environmental covariates, as well as investigate co-

occurrence patterns between taxa.

An amplicon-based survey with good spatial coverage was recently reported by Mikhailov *et*

al. (2019), where the authors examined the possible biotic and abiotic influences on prokaryotic

and eukaryotic communities. The survey data were from a June cruise, thus capturing the non-

stratified homothermic period between spring and summer phytoplankton blooms. The study

found correlation between overall community structure and environmental variables in

multivariate space, as well as a few interesting cross-kingdom co-occurrences.

Here we present a comprehensive report of microbial communities during summer

stratification in Lake Baikal. Our survey and analyses of microbial plankton span all three basins,

from open waters to shallow bays, and multiple depths, including above and below thermocline,

down to 500 m. We reveal major community composition trends that correlate with continuous

environmental gradients in a spatial context. Multiple linear regression and model averaging

reveals which specific environmental covariates are associated with the two most fundamental

measures of diversity: richness and evenness.

We then identify strongest environmental covariates to community composition in a

multivariate framework. Finally, we use co-occurrence network analyses to reveal clusters of

OTUs with contrasting environmental associations and identify clades and taxa most responsible

for maintaining the structure of each network cluster. We conclude by suggesting that

temperature may be the main driver of the microbial community changes in Baikal in response to

global environmental change.

MATERIALS AND METHODS

Our survey was guided by the recorded natural history of Baikal [3, 16, 17] that divides the

lake into eight distinct regions (Fig. 1a, S1). Among them, are the open Baikal basins, Chivyrkuy

Bay, Proval Bay, and Selenga river plume. Chivyrkuy Bay was sampled extensively to capture

transition from the shallow innermost bay (9 m depth) to open waters. Proval Bay and Selenga

plume stations represented the two most eutrophic areas in Baikal. In total, we collected samples

from 24 stations, of which 10 were sampled at various depths for a total of 46 samples.

Temperature and dissolved oxygen profiles were measured with a YSI Instruments sonde (YSI,

Inc., Yellow Springs, OH, USA). Whole water was taken using a Van Dorn sampler (Wildco,

Inc., Yulee, Florida, USA), and 5 L were filtered onto 3 μm and 0.22 μm mixed nitrocellulose

acetate membranes (EMD Millipore, Billerica, MA, USA) and stored at −20°C in RNAlater

(Life Technologies, Grand Island, NY, USA) to capture particle-attached (3 μm filter) and free-

living (0.22 μm filter) fractions of microorganisms. Genomic DNA was extracted using the Mo-

Bio PowerSoil Kit (Mo-Bio Laboratories, Carlsbad, CA), following manufacturer's protocol.

Then, the V4 region of the 16S rRNA gene was amplified using 515F–806R dual-index primers

sequenced on a MiSeq platform (250PE), as described previously [18]. Raw sequences and

metadata needed to reproduce our results are available in GenBank under BioProject

PRJNA530071.

Sequences were processed generally following the *de novo* clustering mothur SOP. Statistical

analyses were performed in the R (3.2.2) environment, unless noted otherwise. Multiple

regression search and model selection were done with the glmulti package [19]. Distance

matrices, ordinations, and correlations with environmental variables were calculated with vegan

[20] and phyloseq [21] packages. We constructed the co-occurrence network using sparCC [22],

following recommended best practices by Berry and Widder [23]. Next, we identified network

modules with the optimum modularity algorithm in the iGraph package [24]. We followed the

WGCNA package workflow [25] to generate eigenvectors for each module and correlate the first

eigenvector (first principal component, PC1) with environmental variables to generate plots and

a heatmap (Fig. 5).

RESULTS AND DISCUSSION

Taxonomic diversity. At a depth of 28059 sequences per sample, we detected 38457 OTUs in

the combined 3.0 μm and the 0.22 μm fractions. Non-singletons (6099 in 3.0 μm and 3346 in

0.22 μm) were classified into 61 phyla, dominated by typical freshwater Actinobacteria,

Bacteroidetes, Cyanobacteria, Proteobacteria and Verrucomicrobia (Fig. 1).

Microbial community composition differed from some previous studies of Lake Baikal,

which explored pelagic communities [11–15]. In a recent effort, Kurilkina et al. [14] sampled

one station depth profile in September and June to reveal the main taxonomic groups as

Proteobacteria, Actinobacteria, Bacteroidetes, Firmicutes, Chloroflexi, Acidobacteria, and

Cyanobacteria in both seasons, and Caldiserica in September only. Our study did not detect

Caldiserica and revealed only a very minor presence of Chloroflexi, while showing a substantial

Verrucomicrobia presence (absent in Kurilkina *et al.*) in every sampled region (Fig. 1).

Interestingly, in the most recent amplicon survey of the lake [15], Mikhailov and colleagues

report Verrucomicrobia relative abundance up to 59% in integrated water column samples taken

across 0-25 m. The differences may be sensitive to the choice of amplicon primer sets, as

Kurilkina *et al.*, Mikhailov *et al.*, and present study used three highly cited but ultimately

different sets.

Our study revealed compositional differences between the 0.22 μm (free-living) and 3 μm

(mostly particle-attached) fractions. Actinobacteria were significantly enriched in the 0.22 μm

fraction (Table S1), similar to results from lakes in Michigan, USA [26]. Proteobacteria were

only marginally more prevalent in the 0.22 μm fraction. We also found enrichment of

Bacteroidetes and Cyanobacteria on the 3 μm fraction, the latter owing to filamentous taxa.

Indeed, the top three most differentially abundant taxa identified with permutation-based

analyses [27] were all classified as the Nostocales genus *Dolichospermum* and contributed up to

90% of Cyanobacteria in the 3 μm fraction. The high abundance of a nitrogen-fixing

cyanobacterium confirms the significance of N limitation in the lake (O'Donnell et al. 2017).

Figure 1: (A) Lake Baikal sampling locations. Stations in outside of Chivyrkuy Bay, Proval
Bay, and Selenga Plume were sampled at multiple depths. (B) Taxonomic composition of the
free-living (0.22 μm, top) and particle-attached (3 μm, bottom) size fractions across surface
samples. While both fractions were mostly dominated by common freshwater phyla,
Actinobacteria were enriched in the 3 μm fraction.
152
Free-living (0.22 μm) surface samples in the open Baikal were dominated by

betaproteobacterium *Limnohabitans* sp. Littoral zones were more variable, represented by

multiple OTUs classified as *Limnohabitans*, *Synechococcus* and *Actinobacteria* acI. Surface

samples of the shallow Proval Bay and the Selenga river plume represented the extreme end of

the eutrophic gradient and were dominated by Actinobacteria acI and Verrucomicrobia

Chthoniobacter. A representative from this genus is reportedly incapable of growth on amino

acids or organic acids other than pyruvate [28]. Indeed, organic acids are among the most

oxidized organic compounds that can be seen in a lake and would be found in areas where labile

carbon is exhausted. *Chtinobacter*, therefore was found where expected - in eutrophic areas with

ample labile carbon. Together with Bacteroidetes *Sediminibacterium* and *Cytophagia*, and

Betaproteobacteria *Limonhabitans* and *Polynucleobacter*, the six OTUs make up ~46% of each

Proval Bay and Selenga River plume's relative abundance. The deepest samples in this study,

collected at 500 m and 300 m, were dominated by Actinobacteria acI and acIV clades as well as

ammonia oxidizing Group 1a Crenarchaeota *Nitrosopumilus*. The latter has been found in

various environments. However, its type species is ubiquitous specifically in oligotrophic oceans,

albeit in the surface layer and not at 500 and 300m [29].

Community Richness and Diversity. Free-living community diversity showed opposing trends

in the upper mixed layer (ML) and deeper waters (DW). Overall, richness and evenness strongly

correlated with temperature in the ML and with depth in DW. For each sample, we calculated the

effective number of species (ENS), a measure of diversity [30–32]. Hill [30] identified ENS as

the number of equally-common species that yields a given value of a diversity index, such as the

Shannon H'. Jost *et al.* [31] have argued that because ENS scales linearly with richness of

equally-common species, it is the preferred metric for quantitative analyses (see Supplementary

Methods). We show that the ENS increased with depth in DW (Fig. 2A) and was driven by an

increase in evenness (Fig. 2C) with no accompanying trend in OTU richness (Fig. 2B). In the

ML, ENS showed a modest increase at the surface in samples collected at 0 m (Fig. 2B). Unlike

in DW, higher ML diversity was generated by higher OTU richness (Fig. 2B), while evenness

remained unaffected. Mixed layer diversity was positively correlated with temperature (Fig. 2D),

along with OTU richness (Fig. 2E). Deep water diversity and richness were marked by high

overall variability, and richness showed a marginally significant increase in samples from cooler

water. Evenness was not directly correlated with temperature in either layer (Fig. 2F).

Furthermore, in mixed layer, richness had a greater response (slope) to temperature than

diversity did, suggesting that samples at higher temperatures and shallower depths, while

supporting the greatest richness, were dominated by a few successful OTUs. This could be due to

variable and high resource conditions leading to coexistence of more taxa, including the

persistence of rare taxa.

Figure 2: Diversity and evenness trends across depth and temperature in the upper mixed layer
and deep waters. Effective Number of Species (ENS) was driven by richness in mixed layer and
by evenness in deep waters. Effect size from Pearson correlation are above the panels.

Deep waters offer the opposite story. While richness had a significant but weak relationship

with depth, ENS and Pielou evenness were very strongly positively correlated with depth

(R^2=0.79, p=2.2x10^{-8}; R^2=0.70, p=7.6x10^{-7}), indicating that a substantial increase in diversity in

the deep hypolimnion was not driven by richness but by an even community structure. These

trends reveal an intriguing picture: warmer shallower part of mixed layer and deeper parts of the

deep water layer both support more diverse (higher ENS) communities. But ML diversity, while

rich in OTUs, is highly uneven, whereas the deep water communities achieve diversity by

increasing evenness. The trends were similar for the 3 μm size fraction (Fig. S7). To better

resolve underlying drivers that underpin these contrasting community features, we modeled ENS

diversity, richness, and evenness with measured environmental variables using iterative multiple

regression and model averaging [19].

Figure 3: Model-averaged importance of environmental predictors for ENS diversity (top), OTU
richness (middle) and Pielou Evenness (bottom) in the upper mixed layer (left) and the deeper
waters (right). Tested predictors included total dissolved silica (TDS), total and dissolved
phosphorus (TP, TDP), and total and dissolved nitrogen (TN, TDN).

Diversity trends in the ML were different, compared with deep waters. Diversity in the mixed

layer correlated the most with temperature and total dissolved silica (TDS) (Fig 3A). These

covariates were also significant for the OTU richness, while no measured environmental variable

was a reliable predictor of ML evenness (Fig. 3B). Depth was not an important predictor for any

ML community feature (Fig. 3C), possibly because depth variation was not large. These results

highlighted the importance of temperature and silica in predicting the ML diversity, by elevating

OTU richness. In the deep waters, total phosphorus (TP) and depth correlated with ENS (Fig.

3D). Furthermore, only TP was a significant predictor for the deep water OTU richness, while

only depth was for evenness (Fig. 3E, F). Temperature, when controlling for other covariates,

was not an important community feature predictor in deep water. The importance of temperature

in ML and depth in DW was not unexpected, considering that the surveyed mixed layer spanned

hundreds of kilometers across all three Baikal's basins and multiple bays, where temperature

ranged from a median 9.5°C in open waters to 21°C in Proval Bay (Fig. S1), while deep waters

were characterized by a large range of depths from 10 to 500 m and a relatively constant

temperature environment. Additionally, nutrients appeared to associate specifically with OTU

richness. Dissolved silica, combined with light availability in ML, is known to promote diatom

growth. Diatom exudates, acting as resources, could create additional niches, enabling

coexistence of a greater number of OTUs. In deep waters, TP could promote OTU richness by

supporting higher phosphorus demands often attributed to fast-growing copiotrophic organisms

[33]. Logue and colleagues [34] found TP alone to be significantly correlated with OTU richness

in a survey of Swedish lakes. Our results suggest that eutrophication leads primarily to an

increase in the overall number of OTUs, but also to dominance of those with opportunistic

lifestyles.

Multivariate trends. Ordination of the Bray-Curtis dissimilarity matrix of the 0.22 μm

samples revealed clear grouping, which we designated into four significant clusters (Fig. 4A,

permANOVA p<0.001). The first cluster (eutrophic) comprised samples from shallow and warm

areas, specifically inner Chivyrkuy and Proval bays. The second cluster (transition) included

samples from the Chivyrkuy Bay to open lake Baikal gradient, the main tributary Selenga River

plume, and the two samples collected in Barguzin Bay – the deepest and most open of the three

examined bays. The third cluster (open) contained the bulk of our samples collected in open

waters that largely separated in ordination space along a depth gradient. The last cluster (deep)

comprised the three deep water samples from 300 and 500 m. *Post hoc* pairwise comparisons of

cluster centroids revealed significant differences between each cluster pair (FDR corrected

p<0.05). Thus, the clusters broadly captured the different habitats of lake Baikal, separated by

depth and the transitions between open lake waters and bays. Given the confounding effects of

abiotic forcing and spatial autocorrelation, we used reciprocal causal modeling [35, 36] to test

whether selection by abiotic environment or spatial dispersal/distance better explained

community composition trends in mixed layer.

Figure 4: Ordination of the Bray-Curtis dissimilarity matrix for all 0.22 μm fraction samples (A)
and the mixed layer only samples (B). (C) Principal component analysis of the mixed layer
samples in environmental space using all measured covariates. In all panels, colors reflect major
recognized regions of lake Baikal (15, 22). In (A) and (B), numbers below bubbles indicate
sample depth (m), and arrows show correlation of environmental covariates with the layout of
sample points in ordination space. Each arrow length = R^2; p-values were obtained using
permutations. In (A), ellipses were drawn to aid visualization.

Environmental conditions, and not the distance, played a dominant role in structuring free-

living mixed layer communities. Reciprocal causal effects models test the hypotheses for

significant correlation between the community dissimilarity matrix and each of the geographic

distance and environmental distance matrices, while controlling for the other. We found that,

when controlling for environment, geographic distance had no correlation with community

dissimilarity. In contrast, when controlling for geographic distance, community structure showed

a significant correlation with the environment (R^2=0.33, $p<1\text{x}10^{-6}$). Our results are consistent

with other studies in freshwater [37, 38] and marine systems [39], which usually note stronger

effects of environment on species composition, compared with dispersal, suggesting that

dispersal is likely not limiting community assembly at this scale.

Temperature, depth, and total dissolved nitrogen (TDN) were the strongest environmental

covariates with multivariate dissimilarity trends among free-living communities (Fig. 4). The

temperature vector (Fig 4A, R^2=0.71, $p=1\text{x}10^{-4}$) indicated the greatest temperature variability

along the open waters to bays gradient. We also confirmed the presence of the depth covariate

among samples collected in the open waters (R^2=0.52, $p=1\text{x}10^{-4}$). Interestingly, open water

community dissimilarities also revealed correlation with TDN in approximately the same

direction as depth. Because depth can confound the effects of various environmental factors, we

further considered an ordination of samples just from the mixed layer (Fig. 4B). Among those

samples, temperature was still the strongest predictor of community dissimilarity (R^2=0.56,

$p=2\text{x}10^{-4}$), and TDS was also significant (R^2=0.36, $p=0.012$). Notably, these were also the only

two model-averaged important predictors of OTU richness and ENS (Fig. 2A, B). However, TN

and TDP were also significant predictors of community dissimilarity (Fig. 4B), although they did

not show an association with alpha diversity metrics (Fig. 2A, B, C). As expected, depth was a

not a significant predictor of community dissimilarity in the mixed layer. Lastly, we investigated

whether the same environmental variables that predicted differences in microbial community

structure also accounted for major abiotic differences between sampled sites. For this, we

constructed a PCA of mixed layer samples using all measured environmental factors (Fig. 4C).

The first two principal components captured 89.8% of the variation. TDS and light had the

highest scores of all environmental variables. Furthermore, they were almost parallel to PC1 and

PC2, respectively, suggesting that the two were responsible for explaining most of the measured

abiotic differences between sampled ML sites. While TDS was a significant environmental

covariate in ordination of biotic community structure, light was not (Fig. 4B). Similarly, light

was not a significant predictor of biotic alpha diversity metrics either (Fig 2). TDP was the least

important predictor of abiotic differences in ML (Fig. 4C); however, it was the second most

significant predictor of biotic dissimilarity in ML (Fig. 4C). These features highlight that

changes in microbial community structure were not correlated with simply the most variable

environmental factors, and that significant associations of particular environmental factors

warrant attention in further studies.

OTU co-occurrence networks. To gain more insight into bacterial community structure in Lake

Baikal and its dependence on the abiotic drivers and to identify OTUs that tend to co-occur, we

constructed an OTU co-occurrence network. Co-occurrence networks enable data-guided

identification of OTU assemblages and correlation of their abundance with continuous

environmental variables. In contrast to ANOVA-based permutation procedures, networks do not

carry a user bias of arbitrarily defining sample categories, which, even if chosen wisely, result in

information loss.

Our network captured the bulk of Baikal's OTUs. We constructed a co-occurrence network

for OTUs present in at least 80% of samples (105 nodes and 819 edges, Fig. 5; see

Supplementary Methods). Importantly, although 105 OTUs appeared to be a large reduction

from the total >38,000 detected in Baikal, the network OTUs amounted to over >81% of

cumulative relative abundance of every sample in >85% of samples. The remainder of the

samples had 63% ± 17% of relative abundance included in the networks. Environments that were

better sampled had more OTUs in the networks. Lower coverage environments included the

notable outliers collected at 500 m, 300 m in Southern Basin and at the surface of the eutrophic

Proval Bay and Selenga River plume. Thus, with network analyses we used non-categorized

continuous data to directly capture the vast majority of microbial community and environmental

covariate data, revealing the dominant groups of co-occurring OTUs and their association with

major environmental drivers.

Figure 5: (A) Co-occurrence network of OTUs across all samples. Edge grayscale hue reflects
pairwise correlation strength (darker edges show stronger correlations), and thickness indicates
edge betweenness score. Modules were defined using maximum modularity optimization. For
each module, the first principal component (eigenvector, PC1) of just that module's constituent
OTUs abundance matrix was used to summarize the dominant abundance trends across sampled
sites. (B) PC1 trends are summarized in a heatmap, where numbers are Spearman correlation
coefficients with p-values in parentheses below. Empty cells indicate non-significant results. (C)
Example PC1 trends are shown with respect to temperature.
326

The resulting network exhibited the "small world" properties (Fig. S10). Small world

networks are characterized by high connectivity between neighboring nodes and low

connectivity between distant nodes [40], creating clusters or modules of consistently co-

occurring OTUs. Within each module, OTUs with many connections (central nodes) are thought

to reflect the processes that bring together module members. Small world networks may be more

robust to perturbations but changes to the abundances of the central, well-connected OTUs may

disproportionately affect the whole modules [41–43].

Using the optimal modularity approach [24, 44], we identified four modules of OTUs that

tend to co-occur across sampled sites (Fig. 5A). Strong cohesion (see Supplementary Methods)

within Module 1 (M1; clustering coefficient, $CC_{M1}=0.68$; Table S1) and Module 3 (M3;

$CC_{M3}=0.82$) suggested common drivers for the consistently co-occurring OTUs. In contrast, low

cohesion within Module 2 (M2; $CC_{M2}=0.54$) and Module 4 (M4; $CC_{M4}=0.58$) pointed to a looser

internal structure.

Module eigenvectors [25] of M1 and M3 revealed opposing monotonic relationships with

respect to environmental covariates (detailed introduction in Supplementary Methods). M1

showed a negative monotonic relationship with depth (Spearman $\rho=-0.70$, $p=3\times10^{-6}$) and a

positive relationship with temperature ($\rho=0.71$, $p=3\times10^{-6}$). In contrast, M3 had a strong positive

relationship with depth ($\rho=0.82$, $p=5\times10^{-10}$) and a negative relationship with temperature ($\rho=-$

0.82, $p=5\times10^{-10}$). As expected, the direction of the opposing trends for the two clusters was

reversed for light, owing to its inverse relationship with depth (Fig. 5B). Thus, we found that a

large fraction of Baikal's planktonic prokaryotes can be classified into one of the two cohesive

clusters: mixed layer "warm" M1 and deeper waters "cold" M3 cluster.

Modules M2 and M4 (Fig. 5a) showed inverse opposing unimodal (non-monotonic) trends

with depth, temperature and light (Fig. 5B, C). Weaker clustering within these modules

suggested less cohesion, possibly due to the inclusion of taxa with weaker habitat preferences or

terrestrial or riverine dispersal. M2 showed highest cumulative relative abundance at an

intermediate temperature of about 10°C, with a strongly positive relationship with oxygen. In

contrast, M4 showed a negative relationship with oxygen, abundance peaks at low and high

temperature extremes and minimum abundance at approximately 11°C. Specifically, M4 was

high in abundance in the immediate surface (0 m) and deep water, but with a sharp drop in

abundance at 5 m depth.

Phylogenetic signal in modules. Membership in the modules had a weakly significant

phylogenetic signal (Fig. S11). This, combined with different habitat preferences among the

modules, suggests that different taxonomic groups displayed preferential environmental

associations, thus supporting the importance of niche differences and phylogenetic niche

conservatism.

M1 and M4 had a high percentage of *Bacteroidetes*, known as opportunistic degraders of

high molecular weight organic matter, such as proteins and carbohydrates, with genomes

containing numerous carbohydrate-active enzymes covering a large spectrum of substrates from

plant, algal and animal origin [45]. In aquatic systems, *Bacteroidetes* have been noted to follow

pulses of organic matter inputs and cyanobacterial blooms [46]. This further casts M1 as a warm

water ML module and M4 as the loose product of sediment and terrestrial input.

M1 is the only module to contain members (three OTUs) of the Chloroflexi phylum.

Although Chloroflexi have been found in diverse environments, including oxygenated (its type

genus) and anoxic [47] hot springs, the CL500-11 clade – a subclass of the deep-ocean SAR202

clade – has been suggested as characteristic of oxygenated hypolimnia in deep lakes, including

Crater Lake [48], Lake Biwa [49] and the Laurentian Great Lakes [50]. However, the three

Chloroflexi OTUs detected in our Lake Baikal network were from the Chloroflexi class (two

OTUs) and Roseiflexales order (one OTU), and, as part of M1, appear to prefer warmer and

shallower environments in the lake.

The cold deep-water cluster M3 had the highest relative abundance of OTUs in unclassified

phyla. While the module also contained characteristically terrestrial Gammaproteobacteria

Pseudomonas and *Acinetobacter*, their low connectivity suggested they are unlikely to play a

central role in M3 processes.

M2 is the only module with Cyanobacteria; the module's three most abundant OTUs –

indeed, second and fourth most abundant in the entire >38,000 OTU dataset – classified as

Synechococcus. Cumulative relative abundance of *Synechococcus* OTUs peaked at ~15 m,

reflecting the approximate location of deep chlorophyll maximum at stratified stations. To

further understand internal structure of the four modules, we focused on central OTUs that

played important roles in creating the module structure.

Central OTUs in each module reflect module ecology. The meaning of a node (in our case

OTU) position in the network is the subject of much discussion. A central OTU has a significant

positive correlation with a large number of that cluster's OTUs. We offer one abiotic and one

biotic interpretation, resulting from two explanations for the presence of network clusters in the

first place. First, clusters of OTUs that correlate with important environmental covariates, like

temperature, could reflect abiotic habitat filtering. In this case, OTUs in modules with higher

clustering coefficients would more consistently reflect environmental conditions associated with

that module. For example, every OTU in M3, which has the highest clustering coefficient (Table

S1), shows clear negative relationship with temperature (Fig. S12). A second explanation

requires an assumption that co-occurrences reflect biotic species interactions. Then, a central

OTU could be hypothesized to directly increase the abundance of other OTUs, e.g., by

synthesizing or otherwise making available a limiting resource. It is important to point out that

the two explanations are not mutually exclusive and could both be parts of ecological

mechanisms that give rise to observed clusters of co-occurring microbial species.

The most globally central (most connected) OTUs belong to the most populous module with

the greatest number of connections (M1). The most central OTU in the network is OTU137,

classified as an autotrophic methylotroph LD19, which was previously reported as a summer-fall

bloomer in Lakes Michigan and Muskegon, MI [51]. Although not very abundant in our dataset,

its high centrality suggests it is indicative of an ecological process that is at least partially

responsible for supporting other OTUs in the ML.

OTU001 (*Limnohabitans*) – also in M1 – has a high overall abundance and the network

position that straddles the balance between centrality in M1 and connectedness to M2. Its high

relative abundance and ubiquity across the lake suggest important roles in lake ecology.

Connectedness to two modules could reflect this ubiquity and further indicate an influence of

OTU001 on OTUs that occupy both niches. More broadly, *Limnohabitans* genus has many

ecotypes. Its fine-scale phylogeny and functioning, shown experimentally with isolates [52],

revealed diverse lifestyles. In fact, three OTUs out of 105 in our network were classified as

Limnohabitans, and one of the other OTUs (OTU104) was in the hypolimnetic M3. Separation of

different *Limnohabitans* OTUs into M1 and M3 with opposing spatial occurrences and

environmental preferences suggests they are indeed ecotypes [53] with distinct functions. Future

studies can experimentally assess *Limnohabitans* ecotype responses to temperature and other

environmental drivers.

The most connected OTU in the hypolimnetic M3 was Planctomycetes OTU022, classified

into the Phycisphaerales order. Little is known about the ecological role of Planctomycetes in

aquatic environments [46]. The order is known for its distinct visual appearance, and a recent

report of a 3D reconstruction of Phycisphaerales cellular membrane revealed characteristic deep

invaginations in the cellular envelope [54]. It is possible the increased surface area of

Phycisphaerales becomes useful in the oligotrophic hypolimnion of Baikal. In contrast to the

most central but not abundant OTU in M1, OTU022 is also the most abundant in M3, which is

the most tightly clustered module in the network (Table S1). These data suggest Phycisphaerales

may substantially contribute to nutrient cycling below the thermocline with direct impacts for the

rest of M3.

The most recent amplicon survey of Baikal included co-occurrence network analyses of

prokaryotes from the photic layer [15], calculated across samples collected at integrated depths

from 0 to 25 m. Authors reported a similar phylogenetic makeup in a network that was

characterized by predominantly positive interactions. Our sampling plan comprised separate

samples from discrete depths in both the photic and abyssal layers, making it possible to resolve

network clusters associated with the different conditions within the water column.

In summary, our survey of Lake Baikal's bacterioplankton revealed that temperature, along

with nutrients, are the major drivers of the microbial community composition and diversity,

consequently, the anthropogenic changes in these factors would likely significantly alter

planktonic microbial communities. The OTU co-occurrence network analysis identified two

major clusters, associated with the upper mixed layer and deep waters, with a detectable

phylogenetic signal in each cluster composition. The "small world" properties of the network

suggest that the communities may be resilient to perturbations but the changes in abundances of

the central OTUs may result in network rearrangement and substantial community shifts. Further

studies, including controlled field and laboratory experiments, are needed to experimentally test

the effects of different temperatures and other abiotic factors on the structure and diversity of

microbial communities in Lake Baikal.

Conflict of Interest Statement. The authors declare that the research was conducted in the

absence of any commercial or financial relationships that could be construed as a potential

conflict of interest.

ACKNOWLEDGMENTS. We thank Ted Ozersky for valuable discussions, Lev Yampolsky

for help with sampling plan design, and Pam Woodruff and Allyson Hutchens for assistance in

the laboratory with nutrient concentration measurements. We are grateful to the crew of the R.V.

Treskov of the Limnological Museum in Listvyanka and field technicians Elena Pislegina and

Alexander Pislegin at the Institute of Experimental Biology, Irkutsk State University for

facilitating fieldwork. We thank Ashley Shade and Jim Tiedje for comments on the manuscript.

This study was funded by National Science Foundation (NSF) Dimensions of Biodiversity grant

DEB-1136710.

REFERENCES

1. Lindeman RL. The Tophic-Dynamic Aspect of Ecology. *Ecology* 1942; **23**: 399–417.

2. Cotner JB, Biddanda BA. Small players, large role: Microbial influence on biogeochemical processes in pelagic aquatic ecosystems. *Ecosystems* 2002; **5**: 105–121.

3. Moore MVM, Hampton SES, Izmest'eva LLR, Silow E a., Peshkova E V., Pavlov BK. Climate Change and the World's "Sacred Sea"—Lake Baikal, Siberia. *Bioscience* 2009; **59**: 405–417.

4. Hampton SE, Izmest'Eva LR, Moore M V., Katz SL, Dennis B, Silow E a. Sixty years of environmental change in the world's largest freshwater lake - Lake Baikal, Siberia. *Glob Chang Biol* 2008; **14**: 1947–1958.

5. Team CW, Pachauri RK, Meyer LA. IPCC, 2014: climate change 2014: synthesis report. Contribution of Working Groups I. *II III to Fifth Assess Rep Intergov panel Clim Chang IPCC, Geneva, Switz* 2014; **151**.

6. Belykh OI, Gladkikh AS, Sorokovikova EG, Tikhonova I V., Potapov SA, Butina T V. Saxitoxin-Producing cyanobacteria in Lake Baikal. *Contemp Probl Ecol* 2015; **8**: 186–192.

7. Kadnikov V V., Mardanov A V., Beletsky A V., Shubenkova O V., Pogodaeva T V., Zemskaya TI, et al. Microbial community structure in methane hydrate-bearing sediments of freshwater Lake Baikal. *FEMS Microbiol Ecol* 2012; **79**: 348–358.

8. Lomakina A V, Pogodaeva T V, Morozov I V, Zemskaya TI. Microbial communities of the discharge zone of oil- and gas- gearing fluids in low-mineral Lake Baikal. *Microbiology* 2014; **83**: 278–287.

9. Gladkikh AS, Kalyuzhnaya O V, Belykh OI, Ahn TS, Parfenova V V. Analysis of

bacterial communities of two Lake Baikal endemic sponge species. *Microbiology* 2014; **83**: 787–797.

10. Zakharova YR, Adel'shin R V., Parfenova V V., Bedoshvili YD, Likhoshway Y V. Taxonomic characterization of the microorganisms associated with the cultivable diatom Synedra acus from Lake Baikal. *Microbiology* 2010; **79**: 679–687.

11. Bel'kova NL, Parfenova V V., Kostornova TY, Denisova LY, Zaichikov EF. Microbial biodiversity in the water of lake baikal. *Microbiology* 2003; **72**: 203–212.

12. Denisova LY, Bel'kova NL, Tulokhonov II, Zaĭchikov EF. Diversity of bacteria at various depths in the southern part of lake Baikal as detected by 16S rRNA sequencing. *Microbiology* 1999; **68**: 475–483.

13. Parfenova VV, Gladkikh AS, Belykh OI. Comparative analysis of biodiversity in the planktonic and biofilm bacterial communities in Lake Baikal. *Microbiology* 2013; **82**: 91–101.

14. Kurilkina MI, Zakharova YR, Galachyants YP, Petrova DP, Bukin YS, Domysheva VM, et al. Bacterial community composition in the water column of the deepest freshwater Lake Baikal as determined by next-generation sequencing. *FEMS Microbiol Ecol* 2016; **92**: 1–13.

15. Mikhailov IS, Zakharova YR, Bukin YS, Galachyants YP, Petrova DP, Sakirko M V., et al. Co-occurrence Networks Among Bacteria and Microbial Eukaryotes of Lake Baikal During a Spring Phytoplankton Bloom. *Microb Ecol* 2019; **77**: 96–109.

16. Kozhov M. Lake Baikal and Its Life. 1963. Springer Netherlands, Dordrecht.

17. Kozhova OM, Izmest'eva LR. Lake Baikal : evolution and biodiversity. *Biology of inland waters* . 1998. Backhuys Publishers.

18. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the miseq illumina sequencing platform. *Appl Environ Microbiol* 2013; **79**: 5112–5120.

19. Calcagno V, Mazancourt C De. glmulti : An R Package for Easy Automated Model Selection with (Generalized) Linear Models. *J Stat Softw* 2010; **34**: 1–29.

20. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan: Community Ecology Package. 2018.

21. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* 2013; **8**: e61217.

22. Friedman J, Alm EJ. Inferring correlation networks from genomic survey data. *PLoS Comput Biol* 2012; **8**: e1002687.

23. Berry D, Widder S. Deciphering microbial interactions and detecting keystone species with co-occurrence networks. *Front Microbiol* 2014; **5**: 219.

24. Csárdi G, Nepusz T. The igraph software package for complex network research. *InterJournal Complex Syst* 2006; **1695**: 1695.

25. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* 2008; **9**: 559.

26. Schmidt ML, White JD, Denef VJ. Phylogenetic conservation of freshwater lake habitat preference varies between abundant bacterioplankton phyla. *Environ Microbiol* 2016; **18**: 1212–1226.

27. Cáceres M De, Legendre P. Associations between species and groups of sites:\nindices and statistical inference. *Ecology* 2009; **90**: 3566–3574.

28. Sangwan P, Chen X, Hugenholtz P, Janssen PH. Chthoniobacter flavus gen.no., sp. nov.,

the First Pure-Culture Representative of Subdivision Two, Spartobacteria classis nov., of the Phylum Verrucomicrobia. *Appl Environ Microbiol* 2004; **70**: 5875–5881.

29. Walker CB, De La Torre JR, Klotz MG, Urakawa H, Pinel N, Arp DJ, et al. Nitrosopumilus maritimus genome reveals unique mechanisms for nitrification and autotrophy in globally distributed marine crenarchaea. *Proc Natl Acad Sci U S A* 2010; **107**: 8818–8823.

30. Hill M. Diversity and evenness: a unifying notation and its consequences. *Ecology* 1973; **54**: 427–432.

31. Jost L. Entropy and diversity. *Oikos* 2006; **113**: 363–375.

32. Leinster T, Cobbold CA. Measuring diveristy: the importance of species similarity. *Ecology* 2012; **93**: 477–489.

33. Klausmeier C, Litchman E, Daufresne T, Levin S. Optimal nitrogen-to-phosphorus stoichiometry of phytoplankton. *Nature* 2004; **429**: 171–174.

34. Logue JB, Langenheder S, Andersson AF, Bertilsson S, Drakare S, Lanzén A, et al. Freshwater bacterioplankton richness in oligotrophic lakes depends on nutrient availability rather than on species-area relationships. *ISME J* 2012; **6**: 1127–36.

35. Cushman SA, Landguth EL. Spurious correlations and inference in landscape genetics. *Mol Ecol* 2010; **19**: 3592–3602.

36. Cushman SA, Wasserman TN, Landguth EL, Shirk AJ. Re-evaluating causal modeling with mantel tests in landscape genetics. *Diversity* 2013; **5**: 51–72.

37. Logue JB, Lindström ES. Species sorting affects bacterioplankton community composition as determined by 16S rDNA and 16S rRNA fingerprints. *ISME J* 2010; **4**: 729–738.

38.   Lindström ES, Forslund M, Algesten G, Bergström A-K. External control of bacterial community structure in lakes. *Limnol Oceanogr* 2006; **51**: 339–342.

39.   Sjöstedt J, Martiny JBH, Munk P, Riemann L. Abundance of broad bacterial taxa in the sargasso sea explained by environmental conditions but not water mass. *Appl Environ Microbiol* 2014; **80**: 2786–2795.

40.   Watts DJ, Strogatz SH. Collectivedynamics of 'small-world' networks. *Nature* 1998; **393**: 440–442.

41.   Comte J, Lovejoy C, Crevecoeur S, Vincent WF. Co-occurrence patterns in aquatic bacterial communities. *BGD Biogeosciences Discuss* 2015; **12**: 10233–10269.

42.   Newman MEJ. The structure and function of complex networks. *Soc Ind Appl Math* 2003; **45**: 167–256.

43.   Newman M, Girvan M. Finding and evaluating community structure in networks. *Phys Rev E* 2004; **69**: 1–16.

44.   Brandes U, Delling D, Gaetler M. On Modularity Clustering. *IEEE Trans Knowl Data Eng* 2008; **20**: 172–188.

45.   Thomas F, Hehemann JH, Rebuffet E, Czjzek M, Michel G. Environmental and gut Bacteroidetes: The food connection. *Front Microbiol* 2011; **2**: 1–16.

46.   Newton RJ, Jones SE, Eiler A, McMahon KD, Bertilsson S, R. J. Newton, et al. A guide to the natural history of freshwater lake bacteria. *Microbiology and molecular biology reviews : MMBR* . 2011.

47.   Overmann J. Green Nonsulfur Bacteria. *Life Sci* 2008.

48.   Urbach E, Vergin KL, Larson GL, Giovannoni SJ. Bacterioplankton communities of Crater Lake, OR: Dynamic changes with euphotic zone food web structure and stable deep

water populations. *Hydrobiologia* 2007; **574**: 161–177.

49. Okazaki Y, Hodoki Y, Nakano SI. Seasonal dominance of CL500-11 bacterioplankton (phylum Chloroflexi) in the oxygenated hypolimnion of Lake Biwa, Japan. *FEMS Microbiol Ecol* 2013; **83**: 82–92.

50. Denef VJ, Mueller RS, Chiang E, Liebig JR, Vanderploeg HA. Chloroflexi CL500-11 populations that predominate keep lake hypolimnion bacterioplankton rely on nitrogen-rich DOM metabolism and C1 compound oxidation. *Appl Environ Microbiol* 2015.

51. Fujimoto M, Cavaletto J, Liebig JR, McCarthy A, Vanderploeg HA, Denef VJ. Spatiotemporal distribution of bacterioplankton functional groups along a freshwater estuary to pelagic gradient in Lake Michigan. *J Great Lakes Res* 2016; **42**: 1036–1048.

52. Salcher MM. Same same but different: Ecological niche partitioning of planktonic freshwater prokaryotes. *J Limnol* 2014; **73**: 74–87.

53. Fuhrman JA. Microbial community structure and its functional implications. *Nature* 2009; **459**: 193–199.

54. Santarella-Mellwig R, Pruggnaller S, Roos N, Mattaj IW, Devos DP. Three-Dimensional Reconstruction of Bacteria with a Complex Endomembrane System. *PLoS Biol* 2013; **11**.