A multi-omic analysis of optineurin proteinopathy in a yeast model suggests the

involvement of lipid metabolism in Amyotrophic Lateral Sclerosis

Daniel M. Bean[1, 2, 3], Silvia Hntova[2], Michael Mülleder[3], Sandra Magalhães[4, 5], Daniel J.

H. Nightingale[6], Kathryn S. Lilley[6], Markus Ralser[2,3], Alexandra Nunes[4,5], Brian J.

Goodfellow[5], Stephen G. Oliver[2, 3]

1. King's College London, Department of Biostatistics and Health Informatics, Institute

of Psychiatry, Psychology and Neuroscience, De Crespigny Park, London SE5 8AF,

United Kingdom

2. Cambridge Systems Biology Centre, University of Cambridge, Cambridge CB2 1GA,

United Kingdom

3. Department of Biochemistry, University of Cambridge, Cambridge CB2 1GA, United

Kingdom

4. iBiMED & Department of Medical Sciences, Campus Universitário de

Santiago, University of Aveiro, 3810-193 Aveiro, Portugal

5. CICECO & Department of Chemistry, Campus Universitário de Santiago, University

of Aveiro, 3810-193 Aveiro, Portugal

6. Cambridge Centre for Proteomics, Department of Biochemistry, University of

Cambridge, Cambridge CB2 1QR, United Kingdom

Corresponding author:
Brian J. Goodfellow
CICECO & Department of Chemistry, University of Aveiro, 3810-193 Aveiro, Portugal
Email: brian.goodfellow@ua.pt
Tel: +351 234 401 506

Abstract

Amyotrophic Lateral Sclerosis (ALS) is an incurable fatal neurodegenerative disease for which the precise mechanisms of toxicity remain unclear despite some significant advances in our understanding of the underlying genetic basis. A holistic, integrated view of cellular changes will be critical to understanding the processes of neurodegeneration and the development of effective treatments. Mutant forms of optineurin (a ubiquitin-binding protein involved in autophagy, membrane trafficking, and NF-κB activation) are found associated with cytoplasmic inclusions containing TDP43 or SOD1 in some ALS patients. We have taken a multi-omics approach to understand the cellular response to OPTN overexpression in a yeast model of ALS. We found that genetic interaction screens and metabolomics provided parallel, highly complementary data on OPTN toxicity. Genetic enhancers of OPTN toxicity in yeast relate directly to the native function of OPTN in vesicular trafficking and intracellular transport, suggesting the human OPTN protein is functional when expressed in yeast even though there is no yeast ortholog. Crucially, we find that the genetic modifiers and the metabolic response are distinct for different ALS-linked genes expressed in yeast. This lends strong support to the use of yeast as a model system and omics platform to study ALS.

Keywords

ALS; Systems Biology; Metabolomics

Introduction

Amyotrophic Lateral Sclerosis (ALS) is a fatal neurodegenerative disease for which there is no effective treatment available. ALS has been divided into familial (FALS) and sporadic (SALS) forms on the basis of family history, with FALS patients accounting for 5% of all ALS cases overall, but this dichotomy is now questioned[1]. Despite our incomplete picture of the genetic landscape of ALS, it is considered a genetic disease. Associated genetic variants, particularly the hexanucleotide repeat expansion of C9orf72[2,3] and mutations in SOD1[4], TDP43[5,6] and FUS[7,8], are the basis for experimental models of ALS in most model systems. Genetic variants and model organism studies implicate a wide range of cellular pathways in the neurodegenerative processes occurring in ALS, including oxidative stress, RNA metabolism, protein aggregation and degradation (autophagy, and the ubiquitin-proteasome system), and intracellular trafficking[9]. The hallmark histopathological feature of ALS is the presence of intracellular protein aggregates. In most cases, these aggregates contain the TDP43 protein, even though mutations in the TDP43 gene are only a rare cause of ALS. A notable exception is patients with SOD1 mutations, where intracellular aggregates contain the SOD1 protein, but not TDP43. With such a complex pathology underpinning ALS, it is vital to develop multi-omics approaches to understand how the interaction of multiple pathways is driving disease progression.

The OPTN gene encodes Optineurin, a ubiquitin-binding protein involved in autophagy[10–12] membrane trafficking[13] and NF-κB activation[14]. ALS-linked OPTN mutations were first detected in a Japanese cohort of ALS patients[15], where OPTN protein was also found colocalized in cytoplasmic inclusions containing TDP43 or SOD1, suggesting OPTN is broadly involved in ALS regardless of the underlying

mutation. Subsequent studies indicated that that OPTN mutations are relatively more

common in Asian populations[15–19], and more rare in Caucasian ALS patients[20–

23]. Mutations in OPTN are also linked to primary open-angle glaucoma[24] and

Paget's disease of bone[25–27], suggesting that OPTN itself is a key driver of toxicity.

Compared to other ALS-linked proteins, there is relatively little research focused on

OPTN-ALS. We therefore chose to focus our study on OPTN specifically. The

intracellular pathways in which OPTN is involved are strongly conserved between yeast

and human cells[28,29]; however, there is no yeast ortholog of optineurin.

Two previous studies have used a yeast model to study OPTN[30,31]. In the first such

study, Kryndushkin et al. found that both wild-type and mutant OPTN formed

intracellular aggregates and were toxic to yeast[31]. More recently, Jo et al. performed a

screen for yeast single-gene deletions that modify the toxicity of human OPTN in

yeast[30]. Their screen used a high-copy plasmid to express OPTN, resulting in a level

of toxicity that is too high to reliably detect enhancer phenotypes (as the authors also

note). Our study builds on this previous work in two main ways. First, we expressed the

OPTN coding sequence from a low copy-number CEN plasmid, to get lower toxicity

levels and enable the detection of both suppressors and enhancers. Secondly, we also

carried out a metabolomic screen to build up a more detailed multi-omic picture of

OPTN-ALS.

Results

Expression of wild-type human OPTN is toxic in yeast

In line with similar studies[30–35], we cloned human OPTN into yeast expression

vectors under the control of the GAL promoter to allow rapidly-inducible strong

expression of the transgene. We first confirmed previous results which showed OPTN

expression is toxic and that the protein forms aggregates when expressed in

yeast[30,31]. Spot tests and liquid culture growth assays both demonstrated the reduced

growth of cells expressing OPTN-YFP vs YFP controls (Figure 1). Microscopy

demonstrated that OPTN-YFP formed intracellular aggregates in yeast, whereas YFP

alone remained diffuse in the cytoplasm (Figure 1). As found previously[31], OPTN

tended to aggregate as a single point, whereas TDP-43 and FUS formed multiple diffuse

aggregates (not shown).

Although there appeared to be a growth phenotype for the E478G mutant in the spot

test, this was not confirmed in the liquid culture. Furthermore, the fluorescence of the

YFP tag was not detectable for OPTN-E478G-YFP under the same microscopy

conditions or in a Western blot. We therefore used only the wild-type OPTN construct

for all experiments in this study.

Genetic screening identifies protective genes

Genetic modifiers of OPTN toxicity were identified using high-throughput synthetic

genetic array (SGA) screens[36] to introduce the OPTN-YFP or control plasmids into

the BY4741 deletion library. Screens were carried out in biological triplicate, each of

which also contained 4 technical repeats. Hits from the screen were defined as strains

whose interaction score was greater than 2 standard deviations away from the mean

interaction score in all 3 biological repeats, with at least 3 of 4 technical repeats scored

and normal growth on SGlu.

We identified 30 suppressors and 64 enhancers of the OPTN growth phenotype

(Supplementary Table S1). The only enriched GO term for the suppressors of OPTN

toxicity was cytoplasmic translation, suggesting that these genes are not specific

suppressors of OPTN but instead affect the expression of OPTN from the plasmid. We

therefore focused on the enhancers of OPTN toxicity. Deletion of these enhancer genes

increases OPTN toxicity, which suggests their protein products exert a protective effect

when present.

The enriched GO terms for the enhancers are shown in Table 1. The most enriched

process was "mitochondrion-ER membrane tethering", which is known to be a key

regulator of cell death processes[37]. Several of the enriched terms relate to intracellular

trafficking, specifically ER-to-Golgi, which is striking as OPTN plays a major role in

Golgi transport and membrane trafficking processes in human cells. Of the 64

suppressors, 52 have at least one human ortholog. The human orthologs are also

enriched for vesicle-mediated transport pathways, including ER-to-Golgi transport and

Golgi vesicle transport, which suggests the screen results could be translatable to human

cells.

Jo et al. recently performed a related OPTN screen in which human OPTN was

expressed from a high copy plasmid and genetic interactions were measured using

pooled barcode sequencing[30]. Their detected 127 suppressors of toxicity they

identified, none is significant as a suppressor or enhancer in our screen (enhancers were

unreliable). However, several of the same pathways were implicated by the hits in both

studies, particularly lipid metabolism and vesicle-mediated transport.

Network analysis of protective genes

Given the coherent set of enriched GO terms for these genes, we expected that they, or their products, were likely to genetically or physically interact. Indeed 52/64 of the protective genes interact genetically, and 41/64 of their protein products interact physically (Figure S1). The genetic interaction network is a single component with 232 edges, whereas there are 4 separate components in the protein-protein interaction (PPI) network (containing 32, 4, 3 and 2 proteins, respectively). Together, the genetic and protein interactions connect 55/64 protective genes as a single component with 279 edges. The Syntaxin-like t-SNARE, TLG2 has high centrality (degree, betweeness, and closeness) in both the genetic and protein interaction networks. TLG2 is the yeast ortholog of mammalian Syntaxin-16, which functions in the ER-Golgi vesicle-mediated transport pathway. It is possible that deletion of TLG2 enhances the toxicity of OPTN in yeast by impairing trafficking to the vacuole, thus limiting protein degradation via this pathway. However, previous work in HEK cells found the ubiquitin-proteasome system to be the primary pathway responsible for degradation of OPTN[38].

Genetic modifiers of OPTN toxicity have little overlap with TDP43 or FUS modifiers in yeast

To determine whether the hits from our screen are specific to OPTN or simply represent a generic response to the expression of a toxic transgene, we compared our results to those of previous SGA screens for TDP43[35] (8 suppressors, 6 enhancers)and FUS[33] (36 suppressors, 24 enhancers) in yeast. These genetic modifiers have limited overlap, consistent with the relevance of our hits to the endogenous function of OPTN. All genes identified as modifiers in more than one screen are shown in Table 2.

The only overlapping genetic modifiers for OPTN were with FUS, with 3 common

enhancers and 2 common suppressors. Deletion of the MAP kinase gene SLT2

enhanced the toxic phenotype of both OPTN and FUS, potentially due to dysregulation

of peroxisome assembly or the unfolded protein response, both of which are regulated

by Slt2p. In a recent study, Jo et al. found that pharmacological inhibition of MAP2K5,

an upstream regulator of MAPK7, is a potential target for ALS therapy[30]. The 2 other

common enhancers, KGD2 and COX5A, encode mitochondrial proteins involved in the

TCA cycle (KGD2) and the inner mitochondrial membrane electron transport chain

(COX5A). Both deletions cause a severe reduction in growth rate and may be false

positives as the additional effect of OPTN is small.

Two suppressor deletions (rpl19b and rpp2b) were common to the OPTN and FUS

screens. Both genes encode ribosomal proteins and therefore affect cytoplasmic

translation. It is therefore possible that the protective phenotype is due to lowered

expression of the toxic transgene rather than a specific interaction. Finally, despite their

more similar cellular functions, only MRPL39, was identified (as an enhancer) in both

the TDP43 and FUS screens.

MS and NMR identification of altered metabolites and lipids

Given that the genetic modifiers of OPTN toxicity were distinct from those previously

identified for the ALS risk genes FUS and TDP43, we wondered whether strains

expressing these human proteins are also metabolically distinct. The stationary phase

(72hr) endometabolomes of yeast strains overexpressing OPTN, FUS and TDP43 were

therefore compared to controls using untargeted metabolic profiling by NMR and

targeted profiling by MS. A PCA analysis of OD- and TSP-normalized and pareto-

scaled NMR data matrices indicated that the profiles are, in fact, distinct

(Supplementary Figure S2A) and thus the metabolic differences are due to the specific

overexpressed protein. The targeted MS analysis of both the aqueous and lipid fractions

(Supplementary Figure S2B) from the same samples confirmed this.

The statistically significant metabolites (from MS and NMR data) that were altered in

OPTN samples compared to controls were identified and the effect size calculated

(Figure 2).

Metabolites whose levels increase when OPTN is overexpressed include those

associated with cellular and metabolic stress - such as γ-aminobutyric acid (GABA),

oxidized glutathione, glycerol, and trehalose. Many others are related to cellular energy

processes such as the TCA cycle, glycolysis and gluconeogenesis

(phosphoenolpyruvate, glutamine; Gln, glutamic acid, adenine nucleotides, and malate).

The most significantly increased metabolite was orotidylic acid followed by uridine-

diphosphate-N-acetylglucosamine (UDP-GlcNAc) and glycerol, while the metabolites

that decreased include the basic amino acids leucine and isoleucine; the TCA cycle

intermediates, succinate and citrate; and glycerophosphocholine (GPC).

Metaboanalyst 4.0[39] was used to identify the pathways most affected by the presence

of overexpressed OPTN compared to controls (Figure 3). The pathways affected that

have the highest impact include: alanine, aspartate and glutamate metabolism; glycine,

serine and threonine metabolism; arginine and proline metabolism and glutathione

metabolism. Pathways with high significance but lower impact include: butanoate

metabolism; pyrimidine/purine metabolism and glycerolipid metabolism.

To confirm the effect of OPTN on lipid metabolism indicated in the Metaboanalyst

results, we also profiled the endometabolome using LC-MS on the organic phase

extract. Distinct profiles were again observed (Figure S2B) for OPTN, FUS and TDP43

and 75 (37 negative ion + 38 positive ion) statistically significant m/z species were

identified as increased in cells overexpressing OPTN, while 78 (28 negative ion + 50

positive ion) species decreased. Of these, a reduced group was selected based on an

effect size above or below 4 and with VIPs larger than 1.56 from a UV-scaled PLS-DA

model using the positive ion mode data and ES above or below 4 and VIPS above 1.68

for the negative ion mode data. Identified lipid families are shown in Table 3 and the

associated m/z ions and tentative identification in Supplementary Table S2.

Discussion

In this study, we have generated genome-wide genetic interaction data and carried out a

metabolomic screen on the same OPTN yeast model. Significantly, we show that in

both profiles, yeast cells expressing OPTN are distinct from yeast cells expressing other

human genes associated with ALS (TDP-43 and FUS). We have also identified genetic

modifiers of OPTN toxicity that are directly related to the endogenous function of

OPTN in human cells.

The wild-type OPTN protein was toxic when expressed in yeast, consistent with related

studies of other ALS genes[30,31,33–35,40]. The ALS-linked OPTN mutant E478G

was not efficiently expressed in our system, and therefore we used the wild-type protein

for all screens. Although we are therefore unable to model the effect of the specific

mutations linked to ALS, a significant part of ALS pathology appears to be a

proteinopathy, which is recapitulated in this system. For example, Armakola et al. used

wild-type TDP-43 for a genome-wide yeast screen and identified dbr1 as a modifier

gene that was validated in human neuronal cell lines[35]. Therefore, whilst this system

models only part of the pathology of ALS, previous data suggest that results in this type

of system are translatable to human cells.

Crucially, the growth phenotype of this yeast proteinopathy model is not a generic

response to an overexpressed transgene, as might have been expected. This is shown by

the distinct modifiers identified for the ALS-linked genes FUS, TDP-43 and OPTN in

genetic interaction screens and in our metabolic profiling. One possible mechanism for

these distinct modifiers is that the expressed proteins retain at least part of their

endogenous function when expressed in yeast. Yeast is widely used for recombinant

protein production, both for research and commercial applications. Whilst the correct

folding of any particular recombinant protein is not guaranteed, and a substrate (and/or

cofactor) required to carry out that protein's molecular function may not be available in

yeast, it is possible that the OPTN protein retains its native molecular function when

expressed in yeast.

In a seminal work, Kachroo et al.[41] found that almost half of the deletion of 414

essential yeast genes could be complemented ("humanised") by the expression of their

human ortholog. Although yeast does not possess an OPTN ortholog, the function of

this human protein may be retained through interaction with conserved components of

the same pathways. For example, yeast also does not have any orthologs of the Bcl-2

apoptosis regulator proteins, yet expression of mammalian Bax protein induces cell

death[42,43], apparently though a mechanism conserved in human cells[44]. Taken

together with the direct relevance of the modifiers we identify in our genetic screen to

the endogenous function of OPTN (specifically intracellular trafficking and ER-Golgi

transport), and the differing metabolic profiles of FUS, TDP-43 and OPTN, we suggest

that the phenotype of yeast expressing transgenic OPTN is due to the native properties

or functions of the OPTN protein, and not simply due to its overexpression.

The metabolic profiling results suggest that OPTN is affecting the TCA cycle, which

may be related to general mitochondrial dysfunction. As overexpression of a non-native

protein in yeast would also be expected to induce the UPR (and ER stress), the

increased levels of some amino acids may be a result of this process causing increased

protein catabolism, a reduction in protein biosynthesis or a reduction in amino acid

utilization/biosynthesis. Oxidative stress was increased in OPTN yeast compared to

controls as the GSH_{red}/GSH_{ox} ratio is 1.4 times lower in these cells (Supplementary

Figure S3). Interestingly, an increase in the amino acid proline is seen in OPTN

expressing yeast. As the presence of this amino acid has been found to minimize protein

aggregation and the depletion of proline has been linked to the inhibition of the

UPR[45], our results are consistent with OPTN aggregation and triggering of the UPR

in our yeast system. Increased UDP-GlcNAc levels suggests that cell wall biosynthesis

may be decreased, and lipid metabolism also appears to be altered as indicated by a

decrease in GPC and an increase in Ser and in UDP-glucose, both involved in

sphingolipid biosynthesis. As UDP-GlcNAc is also intimately involved in the

production of N-glycans, with biosynthesis first taking place in the ER and subsequently

in the Golgi apparatus, any disruption in trafficking between these organelles could also

affect UDP-GlcNAc levels. This may corroborate our conclusion that the phenotype of

yeast cell expressing OPTN reflects the native function of the protein (ER-Golgi

transport) and is not a generic response to an exogenous protein. Our profiling data also

indicates that orotidylic acid had the largest increase of all the aqueous assigned

metabolites detected in our study. At this time, we do not have an explanation as to the

significance of this perturbation.

A number of metabolomic biomarkers have been proposed for neurological diseases

including ALS[46–49]. The results from Wuolikainen et al.[46] show some correlation

with our results where 3 of the top 5 positively correlated ALS metabolites (Pro, Trp,

AMP) in plasma are also seen as increased in our OPTN cells. A deficit in RNA

synthesis was also seen, suggesting a decrease in the PPP. The decrease in ribose-5-

phosphate seen in our yeast model is consistent with this. Basic amino acids were also

indicated as potential biomarkers in CSF and plasma. However, we see decreases in

yeast, while increases are seen in human fluids. A recent metabolomic study of a

neuronal cellular model of ALS[50] included analysis of metabolite variations seen for

cells overexpressing SOD1 and G93A SOD1 under serum deprivation. These results, in

general, compare well with those found in our OPTN stationary-phase yeast model.

Our lipid data from LC-MS profiling indicated that many lipid species were

significantly increased in OPTN-expressing cells (16 increasing above ES 7 compared

to 2 decreasing). In fact, only OPTN (and not FUS or TDP43) showed a larger number

of increased lipids compared to decreased lipids. Thus, OPTN appears to affect lipid

metabolism even as an exogenous protein in yeast. An LC-MS profiling study of the

lipidome for the CSF of ALS patients has identified a number of lipid biomarkers for

ALS[46]. Phosphatidylcholines, sphingomyelins, glucosylceramides and sterols were

found to be increased, while TAG was decreased in ALS patients. We also see increases

in lipids from these families in OPTN-overexpressing yeast cells. Although in a

different model, two ALS studies using different SOD1 mutated mice[51,52] also

demonstrated that lipids such as sphingolipids, ceramides and glucosylceramides are

increased in spinal cord fluid and skeletal muscle. Therefore, it appears that OPTN may

be producing a yeast phenotype that reflects lipidome effects seen in in vivo situations.

We have studied genetic interactions, integrating those results with protein interaction

data, and metabolites. However, additional 'omics approaches could, and should, be

added to build both a broad and deep intracellular understanding of ALS. This

panoramic view of ALS is necessary to predict the impact of perturbations to this

system either by mutation or, eventually, by treatment.

Methods

Yeast strains and media

The Synthetic Genetic Array (SGA)[36] starter strain Y7092 (MATα can1Δ::STE2pr-

Sp_his5 lyp1Δ his3Δ1 leu2Δ0 ura3Δ0 met15Δ0) was used for all experiments was

mated with the BY4741 deletion library in the SGA. Strains were manipulated, and

media prepared using standard microbiological techniques. Yeast were cultured in

synthetic minimal media without uracil, and with either 2% glucose (SGlu), 2%

raffinose (SRaf) or 2% galactose (SGal) as the carbon source.

To induce expression from the plasmid in liquid culture, single colonies were picked

into 50μl SGlu in a 96-well plate and incubated at 30˚C with shaking. After 24h, 10μl

of the SGlu cultures was inoculated into 200μl SRaf and incubated at 30˚C with

shaking. After 24h, 20μl of the SRaf cultures were inoculated into 100μl fresh SRaf and

incubated at 30˚C with shaking. SGal medium was inoculated with a 1:20 dilution of

this SRaf culture in either 96-well or 384-well plates.

For induction in the SGA, deletion mutants containing the plasmid were pinned onto

SGal plates at 384 colonies per plate. After incubation at 30˚C for 24h, these colonies

were pinned onto SRaf at 384 colonies per plate and incubated at 30˚C for 24h. Each

colony from the SRaf plates was pinned onto SGal 4 times (1536 colonies per plate),

and incubated at 30˚C. Plates were scanned at 300dpi after 48h of growth. All pinning

steps were performed using a Singer ROTOR HDA.

Plasmid construction

OPTN in pEGFP-C3 was kindly donated by Dr. Justin Yerbury. The OPTN ORF was

PCR amplified from pEGFP-C3 and cloned into the Gateway donor plasmid

pDONR221 (KanR) via the Gateway BP reaction according to the manufacturer's

instructions. The E478G mutation was introduced by site-directed mutagenesis using

the Aglient QuikChange II Site-Directed Mutagenesis Kit. Donor plasmids containing

FUS and TDP43 were kindly donated by the Gitler lab.

All pDONR221 Gateway donors were cloned into the Gateway destination vector

pAG416GAL-ccdb-EYFP[53] (CEN, URA3, AmpR, referred to as pAG416) obtained

from Addgene via the LR reaction, according to manufacturer instructions.

The empty pAG416 vector was modified to create the control (YFP only) plasmid by

removing the sequence between the GAL1 promoter and YFP. In the unmodified

pAG416, YFP is approximately 1750 bp from the GAL1 promoter, resulting in weak

expression. The YFP coding sequence was PCR-amplified from pAG416 and, in

parallel, pAG416 was digested with Kpn1 and Not1. The larger restriction fragment was

gel-purified and recombined with the PCR product in yeast. The resulting plasmid,

pAG416-short, was used as the control in all experiments.

All plasmid sequences were confirmed by restriction mapping and DNA sequencing.

YFP expression (either alone or as a tag) was confirmed by fluorescence microscopy

and Western blot using anti-GFP (AbCam antibody ab6556) and anti-histone H3

(AbCam ab1791) as a loading control. The presence of the expressed protein (OPTN,

TDP43 or FUS) was confirmed by in-gel digestion, followed by LC-MS/MS. Briefly,

yeast protein extraction for both western blotting and LC-MS/MS was performed

according to[54]. Total extract, corresponding to approximately 1.7×10^6 cells, was

resolved by SDS-PAGE gel electrophoresis. Following Coomassie staining, a band was

excised from the gel that corresponded to a ca. 20 kDa range around the predicted

molecular weight of the eYFP-tagged human protein. The band was dissected into cubes

of approximately 1 mm and destained (using ammonium bicarbonate), reduced (with

dithiothreitol) and alkylated (using iodoacetamide). The sample was digested for 16 h at

37C using a 1:50 (w/w) ratio of Sequencing Grade Modified Trypsin (Promega):protein

in the gel. The digest was analysed by LC-MS/MS on a nanoAcquity UPLC system

(Waters) coupled in-line to a LTQ Orbitrap Velos mass spectrometer (ThermoFisher

Scientific), essentially according to[55], but with the modification that MS2 scans were

performed on the twenty most intense ions per survey scan with a charge of 2+ or

above. In all cases, the human protein corresponding to the expressed transgene was

correctly identified in the LC-MS/MS data.

Mass spectrometry data processing

Raw mass spectrometry data files were converted to MGF format using MSConvert (version 3.0.9283, Proteowizard). MGF files were searched using an in-house Mascot server (version 2.6.0, Matrix Science) against three databases at the same time, which were a canonical S. cerevisiae database, downloaded from UniProt (March 2017; 6,749 sequences), a canonical isoformal version of the human SwissProt database (November 2016; 42,144 sequences) and the cRAPome database of common mass spectrometry contaminants[56] (January 2017; 115 sequences). Precursor tolerance was set to 20 ppm and fragment tolerance to 0.6 Da. Carbamidomethylation of cysteine was specified as a fixed modification and oxidation of methionine as a variable modification.

Genetic interaction screen

Genetic interactions were screened using Synthetic Genetic Array technology as described[36], the only modification being the use of URA3 as the selectable marker for the query strain (Y7092 transformed with OPTN-YFP or control plasmid) instead of NatMX4. Interactions were scored from images scanned at 300dpi after 48h of growth on SGal using Gitter[57] in SGAtools[58] (available at http://sgatools.ccbr.utoronto.ca/). Colonies that failed to grow on SGlu media were excluded from the analysis. Hits from the screen were defined as strains greater than 2 standard deviations away from the mean interaction score in all 3 biological repeats with at least 3 of 4 technical repeats scored and normal growth on SGlu.

Interaction networks

Yeast interaction data was retrieved from the 06/11/2017 update of YeastMine[59] (https://yeastmine.yeastgenome.org) and analysed in Cytoscape[60].

Quenching and extraction of the endometabolome

For all samples (TDP, FUS, OPTN) and controls two biological replicates were

prepared and two technical replicates used. After the final incubation in 25 ml galactose

medium (OD600 = 0.05) the samples were harvested at 72 hr and immediately placed

on ice. A protocol similar to Palomino-Schätzlein et al. (2013) was followed[61].

Pelleting and resuspension in cold phosphate buffer was carried out followed by

centrifugation and flash freezing in liquid nitrogen. For metabolite extraction, 500 μL of

methanol/chloroform (2:1) at 4°C was added to the frozen samples and the pellet was

resuspended by vortexing after 5 min. Five freeze/thaw cycles of 1 min in liquid

nitrogen and 2 min on ice were then carried out followed by addition of 250 μL of

chloroform and 250 μL of MilliQ water. Vortexing for 1 min was followed by a 30 min

centrifugation (20°C, 16000 x g). The upper aqueous layer was collected carefully with

a Gilson pipette. The lipophilic phase was collected into a glass vial. The samples were

dried using a stream of N_2 gas (NitroFlowLab) on a Techne Dri-Block(R) DB30

(aqueous 2-4 hr, lipid 10 min). The aqueous samples were further dried in a speedvac

(1-2 hr, Savant Speed Vac(R) SPD111V). All extracts were stored at 4°C.

NMR spectroscopy

For the NMR analysis, the aqueous extracts were re-suspended in 620 μl D_2O, 0.01%

TSP and 100 mM pH 7.0 phosphate buffer (Na_2HPO_4 100 mM, pH 7.0). All samples

were centrifuged (1 min, 20 °C, 16,000 g) before transferring to 5mm NMR tubes. The

^{1}H-NMR spectra were recorded at 298K on a Bruker Avance III 500 MHz spectrometer

using a TXI or TCI probe. 1D ^{1}H spectra were acquired using a NOESY pulse sequence

to suppress the water resonance, with a sweep width of 7002 Hz (14 ppm), 32k data

points, a recycle delay of 12 s, a mixing time of 100 ms and 128 scans per free

induction decay (FID).

Mass spectrometry

Aqueous fractions

Metabolite concentrations for the aqueous fractions were determined on a liquid

chromatography (Agilent 1290 Infinity) and tandem mass spectrometry (Agilent 6460)

system. All compounds were identified by comparing retention time and fragmentation

pattern with analytical standards. The instrument was operated in single reaction

monitoring mode. Ion transitions and analytical methods used for metabolite

identification and concentration determination are given in Supplementary Tables S3

and S4. Metabolite concentrations were determined by external calibration. Solvents

were of UPLC grade and chemicals of at least analytical grade. Specific conditions for

amino acids, other polar metabolites and UDP-N-acetylglucosamine are given in

supplementary methods.

Lipid fractions

Samples, re-suspended in 200μl of HPLC grade methanol, were analysed in both

positive and negative ion modes using a Waters Xevo G2 quadrupole time of flight (Q-

ToF) combined with an Acquity Ultra Performance Liquid Chromatogram (UPLC)

(Waters Corporation, Manchester, UK). Injection volumes and conditions along with

gradient parameters and data acquisition are given in supplementary methods.

Multivariate analysis

iNMR software (http://www.inmr.net) was used to process the NMR spectra with zero

filling to 64k data points and 0.3 Hz line broadening being applied before Fourier

transformation. The spectra were manually phased, baseline corrected, referenced to

TSP at 0.00ppm and exported (0.5-9.5 ppm for aqueous phase, 0.5-8 ppm for lipid

phase) as a matrix. The spectra were normalized to OD600 at the time of extraction and

then to TSP. The OD600 values at the time of extraction were (mean ± standard

deviation): control = 7.98 ± 0.14, TDP43 = 6.80 ± 0.09, FUS = 7.91 ± 0.08 and OPTN =

7.87 ± 0.15. The normalised spectra were then checked in iNMR and, if necessary,

alignment was carried out using using the 'speaq' package in R. The water region was

excluded from the alignment.

MVA were performed using the ropls package[62] in R. Initial Principal Components

Analysis identified outliers that were excluded from subsequent analyses. The

identification of metabolites for NMR was carried out by comparing the spectra with

those of standard compounds from the Biological Magnetic Resonance Data Bank, the

Yeast Metabolome Database and the Human Metabolome Database. The relative

amounts of the NMR metabolites and the effect size were determined by integrating the

area under the most well-separated metabolite peak in iNMR and then using in-house R

scripts. MS metabolite concentrations were used directly after normalisation to OD600.

Pairwise t-tests were carried out using the False Discovery Rate (FDR) to adjust for

multiple testing. Effect sizes were calculated and corrected for small sample sizes using

the formula:

$$Effect\ Size = (1 - (3/(4n1+n2-2)-1))((x1-x2)/pooledSD)$$

where pooled SD is the pooled standard deviation, x1 and x2 are the mean levels of

metabolite x and n1 and n2 are the number of replicates. Metaboanalyst 4.0[39] was

used to identify enriched metabolic pathways. The final list of metabolites used

included 51 from the targeted MS analysis and 7 (orotidylic acid,

glycerophosphocholine, trehalose, glycerol, betaine, uracil and acetate) non-duplicated

metabolites from NMR.

For the lipid LCMS data, positive and negative ion mode deisotoped results were

normalised to total area and analysed as for aqueous NMR and MS data. Tentative

identification of lipids with statistically significant effect size differences was carried

out using m/z data and the LIPID MAPS Online Tools[63] (Supplementary Table S2).

Data availability

The metabolomics data has been deposited at Metabolights with access code,

MTBLS796.

Acknowledgements

The work of D.M.B., G.F., S.H. and S.G.O. was supported by the Wellcome

Trust/MRC (grant code: 089703/Z/09/Z). D.J.H.N. was supported by a BBSRC

Strategic Longer and Larger grant, awarded to K.S.L. (award BB/L002817/1)

The authors are grateful to Dr. Justin Yerbury for the OPTN plasmid, Prof. Aaron Gitler

for the TDP-43 and FUS plasmids and Dr. Mike Deery for LC-MS/MS analysis of our

plasmid overexpression samples. We thank Steven Murfitt (University of Cambridge)

for the acquisition of the LCMS/MS data for the endometabolome lipid fraction

samples.

B.J.G. and A.N. were supported by grants UID/BIM/04501/2013 and

UID/CTM/50011/2013, co-funded by Fundação para a Ciência e Tecnologia I.P.

(PIDDAC) and by European Regional Development Fund (FEDER) and POCI-01-

0145-FEDER-007628 and POCI-01-0145-FEDER-007679, funded by the Operational

Programme Competitiveness and Internationalization COMPETE 2020. S.M. is

supported by Fundação para a Ciência e Tecnologia I.P. through the individual PhD

grant SFRH/BD/131820/2017. The NMR spectrometers used in this work are part of the

National NMR Network (PTNMR) and are partially supported by Infrastructure Project

Nº 022161 (co-financed by FEDER through COMPETE 2020, POCI and PORL and

FCT through PIDDAC).

This paper represents independent research part funded by the National Institute for

Health Research (NIHR) Biomedical Research Centre at South London and Maudsley

NHS Foundation Trust and King's College London. The views expressed are those of

the author(s) and not necessarily those of the NHS, the NIHR or the Department of

Health.

Author contributions

Conceptualisation – DMB, BJG, SGO

Formal Analysis – DMB, BJG

Funding Acquisition – SGO, BJG

Investigation – DMB, MM, SM, SH, DJHN

Supervision – KSL, MR, AN, BJG, SGO

Writing – Original Draft Preparation – DMB, BJG

Writing – Review and Editing – SGO, DMB, BJG, MM, SM, SH, DJHN, KSL, AN

Conflict of interest

The authors declare that no conflicts of interest exist.

References

1. Chiò A *et al.* 2014 Genetic counselling in ALS: facts, uncertainties and clinical

suggestions. *J. Neurol. Neurosurg. & Psychiatry* **85**, 478 LP-485.

2. Renton AE *et al.* 2011 A hexanucleotide repeat expansion in C9ORF72 is the

cause of chromosome 9p21-linked ALS-FTD. *Neuron*

(doi:10.1016/j.neuron.2011.09.010)

3. DeJesus-Hernandez M *et al.* 2011 Expanded GGGGCC Hexanucleotide Repeat

in Noncoding Region of C9ORF72 Causes Chromosome 9p-Linked FTD and

ALS. *Neuron* (doi:10.1016/j.neuron.2011.09.011)

4. Halper RDRSTPDFDASPHADDGJORJPDHXRDKAM-YDCAGSMBRTR. *et*

al. 1993 Mutations in Cu/Zn superoxide dismutase gene are associated with

familial amyotrophic lateral sclerosis. *Nature* **362**, 59–62.

(doi:10.1038/362059a0)

5. Kabashi E *et al.* 2008 TARDBP mutations in individuals with sporadic and

familial amyotrophic lateral sclerosis. *Nat. Genet.* (doi:10.1038/ng.132)

6. Sreedharan J *et al.* 2008 TDP-43 mutations in familial and sporadic amyotrophic

lateral sclerosis. *Science* (doi:10.1126/science.1154584)

7. Kwiatkowski TJ *et al.* 2009 Mutations in the FUS/TLS gene on chromosome 16

cause familial amyotrophic lateral sclerosis. *Science (80-.).*

(doi:10.1126/science.1166066)

8. Vance C *et al.* 2009 Mutations in FUS, an RNA Processing Protein, Cause

570  Familial Amyotrophic Lateral Sclerosis Type 6. *Science (80-.).*

571  (doi:10.1126/science.1165942)

572 9. Hardiman O, Al-Chalabi A, Chio A, Corr EM, Logroscino G, Robberecht W,

573  Shaw PJ, Simmons Z, Van Den Berg LH. 2017 Amyotrophic lateral sclerosis.

574  *Nat. Rev. Dis. Prim.* **3**. (doi:10.1038/nrdp.2017.71)

575 10. Dikic I *et al.* 2011 Phosphorylation of the Autophagy Receptor Optineurin

576  Restricts Salmonella Growth. *Science (80-.).* **333**, 228–233.

577  (doi:10.1126/science.1205405)

578 11. Wong YC, Holzbaur ELF. 2014 Optineurin is an autophagy receptor for damaged

579  mitochondria in parkin-mediated mitophagy that is disrupted by an ALS-linked

580  mutation. *Proc. Natl. Acad. Sci.* **111**, E4439–E4448.

581  (doi:10.1073/pnas.1405752111)

582 12. Sirohi K, Swarup G. 2016 Defects in autophagy caused by glaucoma-associated

583  mutations in optineurin. *Exp. Eye Res.* **144**, 54–63.

584  (doi:10.1016/j.exer.2015.08.020)

585 13. Nagabhushana A, Chalasani ML, Jain N, Radha V, Rangaraj N, Balasubramanian

586  D, Swarup G. 2010 Regulation of endocytic trafficking of transferrin receptor by

587  optineurin and its impairment by a glaucoma-associated mutant. *BMC Cell Biol.*

588  **11**. (doi:10.1186/1471-2121-11-4)

589 14. Zhu G, Wu CJ, Zhao Y, Ashwell JD. 2007 Optineurin Negatively Regulates

590  TNFα- Induced NF-κB Activation by Competing with NEMO for Ubiquitinated

591  RIP. *Curr. Biol.* **17**, 1438–1443. (doi:10.1016/j.cub.2007.07.041)

592 15. Maruyama H *et al.* 2010 Mutations of optineurin in amyotrophic lateral sclerosis.

593  *Nature* **465**, 223–226. (doi:10.1038/nature08971)

594 16. Li C, Ji Y, Tang L, Zhang N, He J, Ye S, Liu X, Fan D. 2015 Optineurin

mutations in patients with sporadic amyotrophic lateral sclerosis in China.

Amyotroph. Lateral Scler. Front. Degener. **16**, 485–489.

(doi:10.3109/21678421.2015.1089909)

17.   Iida A *et al.* 2012 Novel deletion mutations of OPTN in amyotrophic lateral

sclerosis in Japanese. *Neurobiol. Aging* **33**.

(doi:10.1016/j.neurobiolaging.2011.12.037)

18.   Iida A, Hosono N, Sano M, Kamei T, Oshima S, Tokuda T, Kubo M, Nakamura

Y, Ikegawa S. 2012 Optineurin mutations in Japanese amyotrophic lateral

sclerosis. *J. Neurol. Neurosurg. Psychiatry.* **83**, 233–235.

(doi:10.1136/jnnp.2010.234963)

19.   Naruse H *et al.* 2012 Mutational analysis of familial and sporadic amyotrophic

lateral sclerosis with OPTN mutations in Japanese population. *Amyotroph.*

Lateral Scler. **13**, 562–566. (doi:10.3109/17482968.2012.684213)

20.   Millecamps S *et al.* 2011 Screening of OPTN in French familial amyotrophic

lateral sclerosis. *Neurobiol. Aging* **32**.

(doi:10.1016/j.neurobiolaging.2010.11.005)

21.   Johnson L, Miller JW, Gkazi AS, Vance C, Topp SD, Newhouse SJ, Al-Chalabi

A, Smith BN, Shaw CE. 2012 Screening for OPTN mutations in a cohort of

British amyotrophic lateral sclerosis patients. *Neurobiol. Aging* **33**.

(doi:10.1016/j.neurobiolaging.2012.06.023)

22.   Del Bo R *et al.* 2011 Novel optineurin mutations in patients with familial and

sporadic amyotrophic lateral sclerosis. *J. Neurol. Neurosurg. Psychiatry* **82**,

1239–1243. (doi:10.1136/jnnp.2011.242313)

23.   Belzil V V., Daoud H, Desjarlais A, Bouchard JP, Dupré N, Camu W, Dion PA,

Rouleau GA. 2011 Analysis of OPTN as a causative gene for amyotrophic lateral

sclerosis. *Neurobiol. Aging* **32**. (doi:10.1016/j.neurobiolaging.2010.10.001)

24.   Rezaie T *et al.* 2002 Adult-Onset Primary Open-Angle Glaucoma Caused by

Mutations in Optineurin — Supplemental Data. *Science* **295**, 1077–1079.

(doi:10.1126/science.1066901)

25.   Albagha OME *et al.* 2010 Genome-wide association study identifies variants at

CSF1, OPTN and TNFRSF11A as genetic risk factors for Paget's disease of

bone. *Nat. Genet.* **42**, 520–524. (doi:10.1038/ng.562)

26.   Michou L, Conceição N, Morissette J, Gagnon E, Miltenberger-Miltenyi G, Siris

ES, Brown JP, Cancela ML. 2012 Genetic association study of UCMA/GRP and

OPTN genes (PDB6 locus) with Paget's disease of bone. *Bone* **51**, 720–728.

(doi:https://doi.org/10.1016/j.bone.2012.06.028)

27.   Silva IAL, Conceição N, Gagnon É, Caiado H, Brown JP, Gianfrancesco F,

Michou L, Cancela ML. 2018 Effect of genetic variants of OPTN in the

pathophysiology of Paget's disease of bone. *Biochim. Biophys. Acta - Mol. Basis*

Dis. **1864**, 143–151. (doi:https://doi.org/10.1016/j.bbadis.2017.10.008)

28.   Bonifacino JS, Glick BS. 2018 The Mechanisms of Vesicle Budding and Fusion.

Cell **116**, 153–166. (doi:10.1016/S0092-8674(03)01079-1)

29.   Finley D, Ulrich HD, Sommer T, Kaiser P. 2012 The ubiquitin-proteasome

system of Saccharomyces cerevisiae. *Genetics* **192**, 319–360.

(doi:10.1534/genetics.112.140467)

30.   Jo M *et al.* 2017 Yeast genetic interaction screen of human genes associated with

amyotrophic lateral sclerosis: Identification of MAP2K5 kinase as a potential

drug target. *Genome Res.* **27**, 1487–1500. (doi:10.1101/gr.211649.116)

31.   Kryndushkin D, Ihrke G, Piermartiri TC, Shewmaker F. 2012 A yeast model of

optineurin proteinopathy reveals a unique aggregation pattern associated with

645   cellular toxicity. *Mol. Microbiol.* **86**, 1531–1547. (doi:10.1111/mmi.12075)

646  32. Couthouis J *et al.* 2011 A yeast functional screen predicts new candidate ALS

647   disease genes. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 20881–90.

648   (doi:10.1073/pnas.1109434108)

649  33. Sun Z, Diaz Z, Fang X, Hart MP, Chesi A, Shorter J, Gitler AD. 2011 Molecular

650   determinants and genetic modifiers of aggregation and toxicity for the als disease

651   protein fus/tls. *PLoS Biol.* **9**. (doi:10.1371/journal.pbio.1000614)

652  34. Ju S *et al.* 2011 A Yeast Model of FUS/TLS-Dependent Cytotoxicity. *PLoS Biol.*

653   **9**, e1001052. (doi:10.1371/journal.pbio.1001052)

654  35. Armakola M *et al.* 2012 Inhibition of RNA lariat debranching enzyme suppresses

655   TDP-43 toxicity in ALS disease models. *Nat. Genet.* **44**, 1302–1309.

656   (doi:10.1038/ng.2434)

657  36. Baryshnikova A, Costanzo M, Dixon S, Vizeacoumar FJ, Myers CL, Andrews B,

658   Boone C. 2010 Synthetic genetic array (SGA) analysis in Saccharomyces

659   cerevisiae and schizosaccharomyces pombe. *Methods Enzymol.* **470**, 145–179.

660   (doi:10.1016/S0076-6879(10)70007-0)

661  37. Grimm S. 2012 The ER-mitochondria interface: The social network of cell death.

662   *Biochim. Biophys. Acta - Mol. Cell Res.* **1823**, 327–334.

663   (doi:10.1016/j.bbamcr.2011.11.018)

664  38. Mao J, Xia Q, Liu C, Ying Z, Wang H, Wang G. 2017 A critical role of Hrd1 in

665   the regulation of optineurin degradation and aggresome formation. *Hum. Mol.*

666   *Genet.* **26**, 1877–1889. (doi:10.1093/hmg/ddx096)

667  39. Xia J, Sinelnikov I V., Han B, Wishart DS. 2015 MetaboAnalyst 3.0-making

668   metabolomics more meaningful. *Nucleic Acids Res.* **43**, W251–W257.

669   (doi:10.1093/nar/gkv380)

40.   Johnson BS, McCaffery JM, Lindquist S, Gitler AD. 2008 A yeast TDP-43

proteinopathy model: Exploring the molecular determinants of TDP-43

aggregation and cellular toxicity. *Proc. Natl. Acad. Sci.* **105**, 6439–6444.

(doi:10.1073/pnas.0802082105)

41.   Kachroo AH, 1, Laurent JM, Yellman CM, Meyer AG, 1 2, Claus O. Wilke, 1, 2

3, Marcotte EM. 2015 Systematic humanization of yeast genes reveals conserved

functions and genetic modularity. *Science (80-.).* **348**, 921–925.

(doi:10.1126/science.aaa0769)

42.   William G, Christine S, Bhabatosh C. 1999 Role of mitochondria and C-terminal

membrane anchor of Bcl-2 in Bax induced growth arrest and mortality in

Saccharomyces cerevisiae . *FEBS Lett.* **380**, 169–175. (doi:10.1016/0014-

5793(96)00044-0)

43.   Xu Q, Reed JC. 2018 Bax Inhibitor-1, a Mammalian Apoptosis Suppressor

Identified by Functional Screening in Yeast. *Mol. Cell* **1**, 337–346.

(doi:10.1016/S1097-2765(00)80034-9)

44.   Manon S, Chaudhuri B, Guérin M. 1997 Release of cytochrome c and decrease

of cytochrome c oxidase in Bax-expressing yeast cells, and prevention of these

effects by coexpression of Bcl-xL. *FEBS Lett.* **415**, 29–32.

(doi:https://doi.org/10.1016/S0014-5793(97)01087-9)

45.   Liang X, Dickman MB, Becker DF. 2014 Proline biosynthesis is required for

endoplasmic reticulum stress tolerance in Saccharomyces cerevisiae. *J. Biol.*

Chem. **289**, 27794–27806. (doi:10.1074/jbc.M114.562827)

46.   Wuolikainen A, Jonsson P, Ahnlund M, Antti H, Marklund SL, Moritz T,

Forsgren L, Andersen PM, Trupp M. 2016 Multi-platform mass spectrometry

analysis of the CSF and plasma metabolomes of rigorously matched amyotrophic

lateral sclerosis, Parkinson's disease and control subjects. *Mol. BioSyst.* **12**,

1287–1298. (doi:10.1039/C5MB00711A)

47. Kori M, Aydın B, Unal S, Arga KY, Kazan D. 2016 Metabolic Biomarkers and

Neurodegeneration: A Pathway Enrichment Analysis of Alzheimer's Disease,

Parkinson's Disease, and Amyotrophic Lateral Sclerosis. *Omi. A J. Integr. Biol.*

20, 645–661. (doi:10.1089/omi.2016.0106)

48. Blasco H *et al.* 2017 Lipidomics Reveals Cerebrospinal-Fluid Signatures of ALS.

Sci. Rep. **7**. (doi:10.1038/s41598-017-17389-9)

49. H. B, F. P, B. MH, H. GP, P. V, R. AC, P. C. 2016 Metabolomics in amyotrophic

lateral sclerosis: how far can it take us? *Eur. J. Neurol.* **23**, 447–454.

(doi:10.1111/ene.12956)

50. Valbuena GN, Rizzardini M, Cimini S, Siskos AP, Bendotti C, Cantoni L, Keun

HC. 2016 Metabolomic Analysis Reveals Increased Aerobic Glycolysis and

Amino Acid Deficit in a Cellular Model of Amyotrophic Lateral Sclerosis. *Mol.*

Neurobiol. **53**, 2222–2240. (doi:10.1007/s12035-015-9165-7)

51. Dodge JC *et al.* 2015 Glycosphingolipids are modulators of disease pathogenesis

in amyotrophic lateral sclerosis. *Proc. Natl. Acad. Sci.*

(doi:10.1073/pnas.1508767112)

52. Henriques A *et al.* 2015 Amyotrophic lateral sclerosis and denervation alter

sphingolipids and up-regulate glucosylceramide synthase. *Hum. Mol. Genet.*

(doi:10.1093/hmg/ddv439)

53. Alberti S, Gitler AD, Lindquist S. 2007 A suite of Gateway® cloning vectors for

high-throughput genetic analysis in Saccharomyces cerevisiae. *Yeast* **24**, 913–

919. (doi:10.1002/yea.1502)

54. von der Haar T. 2007 Optimized Protein Extraction for Quantitative Proteomics

720   of Yeasts. *PLoS One* **2**, e1078. (doi:10.1371/journal.pone.0001078)

721 55. Chen C, Buhl E, Xu M, Croset V, Rees JS, Lilley KS, Benton R, Hodge JJL,

722   Stanewsky R. 2015 Drosophila Ionotropic Receptor 25a mediates circadian clock

723   resetting by temperature. *Nature* **527**, 516–520. (doi:10.1038/nature16148)

724 56. Mellacheruvu D *et al.* 2013 The CRAPome: a contaminant repository for affinity

725   purification-mass spectrometry data. *Nat. Methods* **10**, 730–6.

726   (doi:10.1038/nmeth.2557)

727 57. Wagih O, Parts L. 2014 gitter: A Robust and Accurate Method for Quantification

728   of Colony Sizes From Plate Images. *G3: Genes|Genomes|Genetics* **4**,

729   547–552. (doi:10.1534/g3.113.009431)

730 58. Wagih O *et al.* 2013 SGAtools: One-stop analysis and visualization of array-

731   based genetic interaction screens. *Nucleic Acids Res.* **41**.

732   (doi:10.1093/nar/gkt400)

733 59. Balakrishnan R, Park J, Karra K, Hitz BC, Binkley G, Hong EL, Sullivan J,

734   Micklem G, Cherry JM. 2012 YeastMine-An integrated data warehouse for

735   Saccharomyces cerevisiae data as a multipurpose tool-kit. *Database* **2012**.

736   (doi:10.1093/database/bar062)

737 60. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N,

738   Schwikowski B, Ideker T. 2003 Cytoscape: a software environment for

739   integrated models of biomolecular interaction networks. *Genome Res.* **13**, 2498–

740   504. (doi:10.1101/gr.1239303)

741 61. Palomino-Schätzlein M, Molina-Navarro MM, Tormos-Pérez M, Rodríguez-

742   Navarro S, Pineda-Lucena A. 2013 Optimised protocols for the metabolic

743   profiling of S. cerevisiae by1H-NMR and HRMAS spectroscopy. *Anal. Bioanal.*

744   *Chem.* **405**, 8431–8441. (doi:10.1007/s00216-013-7271-9)

62.   Thévenot EA, Roux A, Xu Y, Ezan E, Junot C. 2015 Analysis of the Human

Adult Urinary Metabolome Variations with Age, Body Mass Index, and Gender

by Implementing a Comprehensive Workflow for Univariate and OPLS

Statistical Analyses. *J. Proteome Res.* **14**, 3322–3335.

(doi:10.1021/acs.jproteome.5b00354)

63.   Fahy E, Sud M, Cotter D, Subramaniam S. 2007 LIPID MAPS online tools for

lipid research. *Nucleic Acids Res.* (doi:10.1093/nar/gkm324)

Figure Legends

Figure 1. Expression of OPTN is toxic to yeast. a) Growth in liquid culture showed a

reduced maximum exponential growth rate for cells expressing OPTN (red) compared

to YFP (black). There was no phenotype for cells expressing OPTN-E478G (blue). b)

Fluorescence microscopy showed diffuse cytoplasmic fluorescence for YFP alone,

whereas OPTN-YFP formed focal aggregates. There was no observable fluorescence for

OPTN-E478G with the same exposure (not shown). c) Spot-tests comparing growth on

Glucose (expression off) to Galactose (expression on) showed a growth phenotype for

OPTN that was similar to TDP43, and a weaker phenotype for OPTN-E478G.

Figure 2. Metabolic ES variations observed for yeast overexpressing OPTN versus

controls using data from targeted MS data and untargeted NMR profiles. Statistically

significant differences ($p < 0.05$) are shown as orange and green bars.

Figure 3. Enriched metabolic pathways observed for yeast overexpressing OPTN versus

controls using data from targeted MS data and untargeted NMR profiles in

Metaboanalyst 4.0.

Tables

774
775

	Reference	SGA screen enhancers (n=58)				
GO biological process term	#	#	expected	Fold Enrichment	+/-	P value
mitochondrion-endoplasmic reticulum membrane tethering	5	3	0.04	69.53	+	2.83E-02
organelle localization by membrane tethering	42	6	0.36	16.55	+	4.19E-03
membrane docking	42	6	0.36	16.55	+	4.19E-03
phospholipid transport	24	5	0.21	24.14	+	5.21E-03
ER to Golgi vesicle-mediated transport	88	7	0.76	9.22	+	2.54E-02
Golgi vesicle transport	195	14	1.68	8.32	+	2.09E-06
vesicle-mediated transport	402	17	3.47	4.9	+	6.77E-05
membrane organization	219	10	1.89	5.29	+	3.85E-02
Unclassified	702	4	6.06	0.66	-	0.00E+00

Table 1. GO biological process term enrichment for enhancer hits. Enrichment of terms

for the 58/64 enhancers that were mapped in the PANTHER database. The +/- column

indicates enrichment (+) or depletion (-) of the corresponding term in the enhancers

gene set. Enrichment calculated with PANTHER release 20170413 and GO release

2017-10-24. P values are shown after applying Bonferroni correction for multiple

testing.

Effect	ALS genes	Deletion systematic name	Deletion standard name	Human ortholog
Enhancer	OPTN, FUS	YHR030C	SLT2	MAPK7
Enhancer	OPTN, FUS	YDR148C	KGD2	DLST
Enhancer	OPTN, FUS	YNL052W	COX5A	COX4I1, COX4I2
Enhancer	TDP43, FUS	YML009C	MRPL39	MRPL33
Suppressor	OPTN, FUS	YBL027W	RPL19B	RPL19
Suppressor	OPTN, FUS	YDR382W	RPP2B	RPLP2

793 Table 2. Genetic modifiers identified for multiple ALS genes in yeast. OPTN modifiers

794 identified in this paper, TDP43 modifiers from Armakola et al.[35], FUS modifiers from

795 Sun et al.[33].

797
798

	Lipid family	Increased	Decreased
FA01	Fatty acids and conjugates	3	2
ST01	Sterols	3	
GL02	Diradylglycerols (DAG)	5	
GL03	Triradylglycerols (TAG)	2	
GP01	Glycerophosphocholines	1	
GP10	Glycerophosphates	1	
SP02	Ceramides	2	
SP05	Neutral glycosphingolipids	1	
Unknown	-	2	

815 Table 3. Identified lipid families[63] with differential levels in the LC-MS profiled

816 endometabolome of OPTN yeast vs control.

Figure 1. Expression of OPTN is toxic to yeast. a) Growth in liquid culture showed a reduced maximum exponential growth rate for cells expressing OPTN (red) compared to YFP (black). There was no phenotype for cells expressing OPTN-E478G (blue). b) Fluorescence microscopy showed diffuse cytoplasmic fluorescence for YFP alone, whereas OPTN-YFP formed focal aggregates. There was no observable fluorescence for OPTN-E478G with the same exposure (not shown). c) Spot-tests comparing growth on Glucose (expression off) to Galactose (expression on) showed a growth phenotype for OPTN that was similar to TDP43, and a weaker phenotype for OPTN-E478G.

Figure 2. Metabolic ES variations observed for yeast overexpressing OPTN versus controls using data from targeted MS data and untargeted NMR profiles. Statistically significant differences (p<0.05) are shown as orange and green bars.

Figure 3. Enriched metabolic pathways observed for yeast overexpressing OPTN versus controls using data from targeted MS data and untargeted NMR profiles in Metaboanalyst 4.0.