Inflammasome-independent role for NLRP3 in controlling innate anti-helminth immunity and tissue repair in the lung

AL Chenery[*][1,2], R Alhallaf[*][3], Z Agha[3], J Ajendra[1,2], J Parkinson[1,2], MM Cooper[3], BHK Chan[1,2], RM Eichenberger[3], LA Dent[4], AAB Robertson[5], A Kupz[3], D Brough[2], A Loukas[3], TE Sutherland[6], JE Allen[#][1,2] and PR Giacomin[#][3]

[*] Equal contribution
[#] Joint corresponding

Correspondence:

Paul R. Giacomin
paul.giacomin@jcu.edu.au
James Cook University, Cairns Campus, Building E4 McGregor Rd, Smithfield, QLD 4878, Australia.

Judith E. Allen
judi.allen@manchester.ac.uk
The University of Manchester, AV Hill Building, Oxford Road, Manchester, M13 9PT, United Kingdom.

[1] Wellcome Centre for Cell-Matrix Research, Faculty of Biology, Medicine & Health, Manchester Academic Health Science Centre, University of Manchester, Manchester, United Kingdom.

[2] Lydia Becker Institute for Immunology & Infection, Faculty of Biology, Medicine & Health, Manchester Academic Health Science Centre, University of Manchester, Manchester, UK.

[3] Centre for Molecular Therapeutics, Australian Institute of Tropical Health and Medicine, James Cook University, Smithfield, QLD 4878, Australia.

[4] Department of Molecular and Biomedical Science, School of Biological Sciences, University of Adelaide, Adelaide, SA, 5000, Australia.

[5] School of Chemistry and Molecular Biosciences, University of Queensland, St Lucia, QLD 4072, Australia.

The authors declare no conflicts of interest.

Abstract

45   Alternatively activated macrophages are essential effector cells during type 2

immunity and tissue repair following helminth infections. We previously showed that Ym1,

an alternative activation marker, can drive innate IL-1R-dependent neutrophil recruitment

during infection with the lung-migrating nematode, *Nippostrongylus brasiliensis* suggesting a

potential role for the inflammasome in the innate response to the infection. While NLRP3 has

50   important pro-inflammatory functions in macrophages, its role during type 2 responses and

repair is less defined. We therefore infected *Nlrp3*$^{-/-}$ mice with *N. brasiliensis*. Unexpectedly,

compared to WT mice, infected *Nlrp3*$^{-/-}$ mice had increased neutrophilia and eosinophilia,

correlating with enhanced worm killing but at the expense of increased tissue damage and

delayed lung repair. Transcriptional profiling showed that infected *Nlrp3*$^{-/-}$ mice exhibited

55   elevated type 2 gene expression compared to WT mice. Notably, there was a lack of

inflammasome activation found early post-infection with *N. brasiliensis* and in contrast to

Nlrp3$^{-/-}$ mice, anti-helminth responses were unaffected in caspase-1/11 deficient or WT mice

treated with the NLRP3-specific inhibitor MCC950. Together these data suggest that NLRP3

can constrain lung neutrophilia and helminth killing and negatively regulate type 2 immune

60   responses in an inflammasome-independent manner.

Introduction

The respiratory tract is a dynamic mucosal site that is constantly exposed to the external environment. As a vital barrier organ, the lungs must respond appropriately to both pathogens and innocuous antigens to maintain homeostasis. After tissue injury or an infectious insult, the lungs rapidly initiate immune resolution and repair mechanisms to maintain their essential physiological function. Alveolar macrophages (Mφs) help maintain homeostasis and can drive either pro-inflammatory or pro-resolution responses in the lung tissue microenvironment, depending on the stimulus. Classically activated Mφs upregulate anti-microbial factors such as NO, TNFα, IL-6, and IL-1β in response to pathogen or damage-associated molecular patterns (PAMPs or DAMPs respectively).[1] Conversely, during helminth infections, Mφs can become alternatively activated in response to IL-4Rα signalling, upregulating arginase, resistin-like molecule (RELM)-α, Ym1, and specific matrix metalloproteinases (MMPs), all important type 2 effector molecules with wound healing functions.[2,3] Furthermore, IL-4Rα signalling to Mφs is not only critical for tissue repair[4] but also controls acute airway inflammation and haemorrhage during infection with lung-migrating nematodes.[5] However, there is still little known about how tissue signals integrate to influence alveolar Mφ activation throughout the course of lung injury and repair.

Our previous work has shown that the murine chitinase like protein Ym1, a major product of IL-4Rα-activated macrophages, plays a prominent role in the early (day 2 post-infection) recruitment of neutrophils that swarm around *Nippostrongylus brasiliensis* larvae in the lungs and promote worm killing.[6] IL-17A-producing γδ T cells were key for this Ym1-mediated neutrophilia.[6] Ectopic overexpression of Ym1 drives *Il1b* expression whilst neutralising Ym1 prevents expression of *Il1b* following *N. brasiliensis* infection. Critically, blockade of the IL-1 receptor reduces the ability of Ym1 to induce IL-17-producing γδ T

cells and associated neutrophilia, leading us and others to hypothesize that Ym1 may activate an inflammasome.[7] The NLR family member NACHT, LRR and PYD domains-containing protein 3 (NLRP3) inflammasome is the most thoroughly characterised to date. TLR-NFκB activation induces expression of *Nlrp3* and *Il1b* (priming) which when coupled with the

90 presence of danger signals/DAMPs (e.g. reactive oxygen species, ATP, membrane rupture) results in the activation of NLRP3. NLRP3 oligomerises then associates with apoptosis-associated speck-like protein containing a CARD (ASC), thus creating the inflammasome platform.[8] The inflammasome recruits and activates the protease caspase-1,[9] which subsequently cleaves pro-IL-1β into its mature form.[8] Caspase-1 also cleaves pro-gasdermin

95 D and the resulting N-terminal fragments assemble to forms pores in the plasma membrane, facilitating secretion of IL-1β and triggering pyroptotic cell death.[10] Crystalline/particulate material can activate the NLRP3 inflammasome and Ym1 is well known to crystallize under conditions of chronic inflammation,[11] suggesting that Ym1 is a potential activator of the NLRP3 inflammasome.

100 While the NLRP3 inflammasome has been widely studied in the context of classically activated Mφs, comparatively little is known about whether NLRP3 plays a role during alternative activation and type 2 inflammatory settings in which Ym1 is highly induced. Our previous studies show that NLRP3 constrains type 2 responses during infection with the gut-dwelling helminth parasite *Trichuris muris*, with a major effect on the development of

105 adaptive Th2 cell immunity.[12] In the present study, we addressed whether the NLRP3 inflammasome contributed to acute lung injury, innate immune cell activation and lung repair during infection with the lung-migrating nematode *N. brasiliensis*. Counter to expectations, we discovered that NLRP3 constrained rather than promoted early lung neutrophilia, restricted helminth killing and other type 2 immune responses, but may have been important

110 for limiting infection-induced tissue damage.

Results

NLRP3 restricts early innate immune cell recruitment to the lung

To test our hypothesis that NLRP3 is required for the early recruitment of neutrophils

115   into the lung during infection with lung-migrating helminths, WT and *Nlrp3⁻/⁻* mice were

infected with *N. brasiliensis*. Lung tissue cells were isolated and analysed on day 2 post-

infection, where there was a peak of injury and neutrophilia. As expected, *N. brasiliensis*

infection of WT mice led to increased frequencies (**Figure 1a**) and total numbers (**Figure 1b**)

of eosinophils and neutrophils in the lung, compared to naïve mice. Unexpectedly, infected

120   *Nlrp3⁻/⁻* mice had increased numbers of recruited neutrophils and eosinophils compared to

infected WT mice. Total numbers of alveolar Mφs were not significantly different between

all groups (**Figure 1a-b** and **Figure S1** for gating strategy). Cellular content of

bronchoalveolar lavage (BAL) fluid was not analysed because all infected samples on day 2

were bloody due to lung damage. Together, these data suggest that rather than promote

125   granulocyte recruitment to the lung, NLRP3 may instead limit the recruitment of neutrophils

and eosinophils.

Figure 1 NLRP3 constrains lung innate cell recruitment during *N. brasiliensis* infection. WT and *Nlrp3*$^{-/-}$ mice were infected with *N. brasiliensis* (Nb) and day 2 post-infected lung alveolar (alv.) Mφ (CD11b^{lo}Siglec-F$^+$CD11c$^+$), neutrophil (CD11b$^+$Siglec-F$^-$Ly6G$^+$) and eosinophil (CD11b$^+$Siglec-F$^+$) (**a**) frequencies of live cells and (**b**) absolute numbers were measured by flow cytometry. Data are representative (mean ± s.e.m.) of 4 individual experiments with 3-5 mice per group (per experiment). ******P<0.01, ***P<0.001, (one-way ANOVA and Tukey-Kramer *post hoc* test).

NLRP3 limits innate anti-helminth immunity in the lung

We next addressed whether the increased granulocyte infiltration in infected *Nlrp3*$^{-/-}$ mice correlated with increased anti-helminth immunity. Rapid neutrophil recruitment to the lung in particular is known to be important for immunity to *N. brasiliensis*.[6,13] Thus, we

145 assessed numbers of larvae in WT and *Nlrp3*$^{-/-}$ mice during primary *N. brasiliensis* infection. Consistent with the enhanced early granulocyte response (**Figure 1b**), *Nlrp3*$^{-/-}$ mice displayed reduced numbers of lung-stage L4 larvae at both day 1 and day 2 post-infection compared to WT mice (**Figure 2a**). Consequently, *Nlrp3*$^{-/-}$ mice also displayed less parasite migration to the intestine at days 4 and day 6 post-infection (**Figure 2b**) and a decrease in

150 shed fecal eggs (**Figure 2c**). *Nlrp3*$^{-/-}$ mice also displayed increased presence of goblet cells in the small intestine (**Figure 2d**), consistent with an increased effector type 2 response. Supporting these *in vivo* data, we employed *in vitro* co-culture assays of *N. brasiliensis* L3 with sort-purified bone marrow neutrophils or Mφs from WT mice. We found that neutrophils were highly effective at immobilising and killing the larvae, although Mφs may

155 also contribute (**Figure 2e-f),** consistent with previous findings.[14] Together, these data

suggest that NLRP3 may constrain protective innate immune responses in the lung following *N. brasiliensis* infection, potentially by suppressing neutrophil and eosinophil accumulation.

Figure 2 Restriction of anti-helminth responses by NLRP3. WT and *Nlrp3⁻/⁻* mice were infected with 500 *N. brasiliensis* (Nb) L3s and (**a**) lung stage L4 larvae were quantified on days 1 and 2 post-infection. (**b**) Adult worms in the small intestine (days 4 and 6 post-infection) and (**c**) fecal parasite eggs shed in the stool (day 6) were quantified (per fecal pellet). (**d**) Representative images of proximal small intestine from day 6 infected mice, stained with Alcian Blue/Periodic Acid-Schiff. L3 larvae were co-incubated for 24 h with either no cells (Ctrl), Mφs or neutrophils *in vitro* and (**e**) imaged (scale bar = 100 μm). (**f**) Worm killing was measured based on reductions in larval motility. Data from **a-d** are representative (mean ± s.e.m.) of 3 individual experiments with 3-5 mice per group (per experiment). Data from **e-f** are representative (mean ± s.e.m.) of 2 individual experiments. *$P<0.05$, **$P<0.01$ (one-way ANOVA and Tukey-Kramer *post hoc* test or unpaired two-tailed student's *t* test).

NLRP3 regulates lung repair after helminth infection

The data thus far demonstrated that *Nlrp3*[−/−] mice displayed elevated neutrophilia and
eosinophilia in the lung and protective immunity in the early stages of primary infection,
accompanied by an enhanced type 2 effector response in the intestine. Both types of
granulocytes may contribute to lung tissue damage during *N. brasiliensis* infection and
allergic responses[5,6] but type 2 responses are typically tissue protective.[15] It was therefore
important to assess whether elevated granulocytic responses in *Nlrp3*[−/−] mice impacted the
resolution of inflammation and repair of lung tissue damage during the later stage of the
infection model.[16,17] To assess airway cell infiltration, BAL was performed on mice at day 7
post-*N. brasiliensis* infection, at a timepoint when haemorrhaging has normally ceased and
larvae have completely exited the lung tissue. As expected, eosinophils were the predominant
cell type elevated in the airways of infected WT mice compared to naïve controls along with
increased alveolar Mφs and neutrophils (**Figure 3a**). Critically, all three cell types were
significantly elevated in infected *Nlrp3*[−/−] mice compared to infected WT mice, suggesting a
persistence of cellular inflammation in the airways. This was consistent with evidence of
increased lung tissue pathology in *Nlrp3*[−/−] mice compared to WT mice after infection
(**Figure 3b-c**). Specifically, while lungs from infected WT mice displayed modest airway cell
infiltration and features of successful tissue repair, *Nlrp3*[−/−] lungs had more prominent and
dispersed immune cell foci and larger gaps in the alveoli indicating impaired repair (**Figure
3c**). To quantify lung damage, we performed fractal/lacunarity analysis of entire lung lobes
which has previously been shown to robustly measure lung pathology, correlating well with
traditional stereological measurements such as mean linear intercept.[18] Lacunarity (Λ), a
measure of heterogeneity/gaps in a structure that correlates with airway damage, was
increased for infected *Nlrp3*[−/−] lungs compared to infected WT lungs which showed no
significant differences over WT naïve controls (**Figure 3b**). We did not detect differences in

tissue pathology between WT and *Nlrp3*⁻/⁻ lungs as early as day 2 post-infection (**Figure S2**).

Together, these data reveal that NLRP3 is required for the resolution of inflammation and the

200 timely initiation of repair processes in the lung following injury by helminth infection.

Figure 3 NLRP3 regulates lung tissue repair following *N. brasiliensis* infection. WT and *Nlrp3*⁻/⁻ mice were infected with *N. brasiliensis* (Nb) and (**a**) day 7 post-infected BAL

205 alveolar (alv.) Mφ (CD11b⁻Siglec-F⁺CD11c⁺), neutrophil (CD11b⁺Siglec-F⁻Ly6G⁺) and eosinophil (CD11b⁺Siglec-F⁺) absolute numbers were measured by flow cytometry. Haematoxylin/eosin staining was performed on lung sections and imaged followed by (**b**) quantification of lacunarity (Λ) to assess lung damage on day 7. (**c**) Representative image insets are shown on top with low magnification scans of entire left lung lobes from individual

210 mice shown below (top panel scale bar = 1000 μm, bottom panel scale bar = 500 μm). Data were pooled (**a-b**; mean ± s.e.m.) from 3 individual experiments with 3-5 mice per group (per experiment). *$P<0.05$, **$P<0.01$, ***$P<0.001$ (one-way ANOVA and Tukey-Kramer *post hoc* test).

NLRP3 constrains type 2 cytokine responses and expression of Ym1

The findings above were not fully consistent with the enhanced eosinophilia and goblets cells in *Nlrp3*[−/−] mice, which would suggest that NLRP3 reduces type 2 immunity. Lung tissue repair following *N. brasiliensis* infection is dependent on type 2 responses including the expansion of IL-4Rα-activated Mφs.[5] Critically, while innate sources of Ym1 induce neutrophilia, during the later reparative stage, IL-4Rα-induced Ym1 has a direct role in repair.[2] We therefore examined whether these responses are dysregulated in *Nlrp3*[−/−] mice. Despite a failure to repair, compared to infected WT mice, infected *Nlrp3*[−/−] mice exhibited enhanced IL-4 protein levels in the lungs as early as day 2 post-infection (**Figure 4a**). By day 7 post-infection, using qRT-PCR we found that *Chil3* (Ym1) expression was significantly increased in the lungs of *Nlrp3*[−/−] mice compared to WT mice (**Figure 4b**). Similarly, there were significant increases in Ym1 protein levels in the BAL fluid of infected *Nlrp3*[−/−] mice (**Figure 4c**). To define the cellular source of Ym1, we performed intracellular cytokine staining and flow cytometry focussing on alveolar Mφs and neutrophils that are known sources of Ym1 during *N. brasiliensis* infection.[2] We found that infection caused a drop in the geometric mean fluorescence intensity (gMFI) of Ym1 staining in both alveolar Mφs and neutrophils (**Figure 4d)**, consistent with release of Ym1 into the airways. However, *Nlrp3*[−/−] alveolar Mφs had a greater drop in gMFI for Ym1 following infection compared to WT alveolar Mφs. Conversely, Ym1 release by neutrophils was equivalent in infected WT and *Nlrp3*[−/−] mice. Therefore, NLRP3 constrains expression of IL-4 and Ym1 in the lung during infection with *N. brasiliensis* and may control release of Ym1 by alveolar Mφs in the airways.

Figure 4 NLRP3 regulates Ym1 following infection with *N. brasiliensis*. After infection of WT and *Nlrp3*⁻/⁻ mice with *N. brasiliensis* (Nb), (**a**) IL-4 protein levels were measured in lung tissues by ELISA on day 2 post-infection (normalised to tissue weight). (**b**) *Chil3* (Ym1) gene expression was measured by qRT-PCR on day 7 post-infection (log$_2$ expression relative to *Rpl13a*). (**c**) BALF Ym1 was measured by ELISA and (**d**) intracellular expression by geometric mean fluorescence intensity (gMFI) (fold change over WT control) by alveolar (alv.) MΦs and neutrophils were quantified by flow cytometry on day 7 post-infection. Data are representative (**a**; mean ± s.e.m.) or pooled (**b-d**; mean ± s.e.m.) from 3 individual experiments with 3-5 mice per group (per experiment). *$P<0.05$, **$P<0.01$, (one-way ANOVA and Tukey-Kramer *post hoc* test).

Type 2 gene expression is regulated by NLRP3

To more broadly characterise the immune processes potentially dysregulated in *Nlrp3*⁻/⁻ mice, we performed differential gene expression analysis of whole lung RNA on day 7 post-infection with *N. brasiliensis* using the NanoString nCounter® Myeloid Innate Immunity panel. We observed differential expression of well-characterised genes associated with the later stages of *N. brasiliensis* infection such as *Rnase2a*, *Retnla* and *Chil3/4* that were, as expected, highly upregulated in infected WT mice relative to uninfected mice (**Figure S3** and **Table S1**). Under basal conditions, few genes were found to be downregulated in *Nlrp3*⁻/⁻ mice compared to naïve WT mice (**Figure S3**). However, in *N.*

brasiliensis-infected mice, 84 genes were differentially expressed between *Nlrp3*[−/−] mice

compared to WT ($P < 0.05$) (**Figure 5a-b**). Although no genes remained statistically

significant after Benjamini-Hochburg multiple test correction[19] (**Table S2**), principle

components analysis (PCA) and hierarchical clustering (**Figures 5c-d**) showed clear

separation of gene expression patterns between *N. brasiliensis*-infected WT and *Nlrp3*[−/−]

mice. Increased sample numbers were used to validate specific genes by qRT-PCR. Infected

Nlrp3[−/−] mice had significantly higher expression of the type 2 markers *Arg1* and *Ccl24* as

well as an increase in the neutrophil-associated *Cxcl3* compared to infected WT mice (**Figure

5e**). Despite not reaching significance in the NanoString analysis, qRT-PCR analysis showed

that *Il4* expression was found to be significantly increased in infected *Nlrp3*[−/−] mice

compared to infected WT mice, confirming the protein levels observed in **Figure 4a**.

Additional genes whose expression was increased in infected *Nlrp3*[−/−] mice included the

matrix metalloprotease-encoding *Mmp13* and the alveolar Mφ-expressed gene *Ear6*.[20] Thus,

NLRP3 appears to negatively regulate type 2 gene expression in the lung following tissue

injury caused by helminth infection and may control neutrophil chemotactic recruitment and

alveolar Mφ function.

Figure 5 NLRP3 regulates type 2 gene expression in the lung following *N. brasiliensis* infection. Whole lung RNA from WT and *Nlrp3$^{-/-}$* mice on day 7 post-infection with *N. brasiliensis* (Nb) was analysed by NanoString. (**a**) Volcano plot showing differentially expressed genes between Nb-infected WT and *Nlrp3$^{-/-}$* mice. (**b**) Top 20 differentially regulated genes between Nb-infected WT and *Nlrp3$^{-/-}$* mice (bar colour indicates statistical significance). (**c**) Unsupervised, hierarchically clustered heatmap of genes differentially expressed 7 days post Nb-infection in WT and *Nlrp3$^{-/-}$* mice. (**d**) Principle components analysis (PCA) of naïve and Nb-infected WT and *Nlrp3$^{-/-}$* mice. (**e**) Candidate differentially expressed genes validated by qRT-PCR (log$_2$ expression relative to *Rpl13a*). Data were from a single Nanostring run (**a-d**; n=2-4 mice per group) or pooled (**e**; mean ± s.e.m.) from 3 individual experiments with 3-5 mice per group (per experiment). *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$ (one-way ANOVA and Tukey-Kramer *post hoc* test).

Tissue resident Mφ alternative activation is regulated by NLRP3

Our finding that expression of *Chil3* (Ym1) was altered in *N. brasiliensis*-infected *Nlrp3$^{-/-}$* mice did not allow us to distinguish whether the change was due to the enhanced quantities of IL-4 observed or enhanced responsiveness of the Mφs to IL-4Rα signalling. To test this, we utilized IL-4-complex (IL-4c) delivery into the peritoneal cavity which robustly induces expression of Ym1 and RELM-α in peritoneal Mφs.[21] WT and *Nlrp3$^{-/-}$* mice were

injected with IL-4c i.p. following which we analysed the CD11b$^+$ myeloid compartment containing F4/80^{lo} monocyte-derived Mφs and F4/80^{hi}Tim4$^+$ tissue resident Mφs after 24 hours (**Figure 6a**). There would be no expectation that IL-4c delivery would induce the NLRP3 inflammasome and indeed we were unable to detect any evidence for inflammasome

300   activation in the peritoneal Mφs (data not shown). As expected, both Mφ populations showed increased intracellular expression of Ym1 and RELM-α in WT mice after IL-4c injection. IL-4c-stimulated *Nlrp3*$^{-/-}$ resident (F4/80^{hi}Tim4$^+$) Mφs had greater frequencies of both Ym1$^+$ and RELM-α$^+$ cells as well as an increased Ym1 gMFI compared to IL-4c-stimulated WT resident Mφs (**Figure 6b**). In contrast, we did not see enhanced expression of Ym1 or

305   RELM-α above WT levels in *Nlrp3*$^{-/-}$ monocyte-derived (F4/80^{lo}) Mφs after IL-4c stimulation (**Figure 6c**). Similarly, *in vitro* IL-4-stimulated BMDMs from *Nlrp3*$^{-/-}$ mice did not show increased alternative activation markers compared to WT BMDMs (**Figure S4**). Thus, it appears that *in vivo* NLRP3 can influence the Mφ response to IL-4, independent of the inflammasome, but these effects may be restricted to specific Mφs subpopulations, such

310   as tissue resident rather than monocyte-derived Mφs.

Figure 6 NLRP3 regulates resident Mφ alternative activation by IL-4. WT and *Nlrp3*^{–/–} mice were injected with either PBS or IL-4 complex (IL-4c) and (**a**) peritoneal CD11b⁺ cells containing F4/80^{lo} and F4/80^{hi}Tim4⁺ Mφs where analysed by flow cytometry after 24 hours. Frequency and geometric mean fluorescence intensity (gMFI) of intracellular Ym1 and RELM-α were determined in (**b**) resident F4/80^{hi}Tim4⁺ Mφs and (**c**) monocyte-derived F4/80^{lo} Mφs. Data are representative (**b** left panel; mean ± s.e.m.) or pooled (**b** right panel and **c**; mean ± s.e.m.) from 3 individual experiments with 3-5 mice per group (per experiment). *$P<0.05$, ****$P<0.0001$ (one-way ANOVA and Tukey-Kramer *post hoc* test).

Potential inflammasome-independent role of NLRP3 during lung anti-helminth responses

Thus far we have demonstrated that although NLRP3 suppressed type 2 immune responses, it also limited persistent lung inflammation after resolution of helminth infection. Our observation that NLRP3 constrained the early neutrophilic response was contrary to the expectation that infection with *N. brasiliensis* would result in inflammasome activation.

Although NLRP3 is most commonly associated with forming an inflammasome, there are some reports of inflammasome-independent roles for NLRP3.[22,23] Therefore, we investigated whether NLRP3 played inflammasome-dependent or -independent functions during *N.*

330   *brasiliensis* infection. First, we examined expression of infection-induced production of pro-IL-1β within alveolar Mφs, a precursor for inflammasome activation. *N. brasiliensis* infection caused increased levels of pro-IL-1β within alveolar Mφs, however this infection-induced response was unaffected by NLRP3 deficiency (**Figure 7a**). Further, levels of mature IL-1β released into the airways (an indicator of inflammasome activation), were nearly

335   undetectable in the BAL fluid in both WT and *Nlrp3*[−/−] mice (**Figure 7b**), suggesting that inflammasome activation in the lung is not a major feature of *N. brasiliensis* infection. To further delineate whether inflammasome activation occurred following *N. brasiliensis* infection, we quantified activated caspase-1 *ex vivo* by using a FAM-YVAD-FMK fluorochrome inhibitor of caspases (FLICA) probe on BAL cells on day 2 post-infection.

340   Critically, there was no evidence of infection-induced increases in caspase-1 activity in alveolar Mφs (**Figure 7c**). In contrast, we observed significant decreases in caspase-1 activity within neutrophils and eosinophils that were independent of NLRP3 deficiency (**Figure 7c**).

To more definitively establish whether inflammasome activation contributed to the observed phenotype of *Nlrp3*[−/−] mice, we compared the response to *N. brasiliensis* infection

345   in *Nlrp3*[−/−] mice to that of *Casp1/11*[−/−] mice which cannot mount either canonical or non-canonical NLRP3 inflammasome responses.[24] While *Casp1/11*[−/−] mice showed a trend towards increased immunity to lung-stage *N. brasiliensis* infection compared to WT mice, this was not statistically significant (**Figure 7d**) and adult worm burdens were equivalent to WT mice (**Figure 7e**). In contrast, *Nlrp3*[−/−] mice displayed increased anti-parasitic immunity

350   at the lung stage, characterised by significantly reduced lung larval numbers and adult worm burdens. In a final effort to establish a role for *N. brasiliensis*-induced inflammasome

activation we treated WT mice with MCC950, an NLRP3-specific inflammasome small

molecule inhibitor[25] throughout the course of infection (**Figure S5**). We failed to see changes

in myeloid cell numbers in the BAL throughout the infection following MCC950 treatment

355   (data not shown). By day 5 post-infection, we could see that mice treated with MCC950 had

a reduction in BAL neutrophil numbers but had no change in alveolar Mφ or eosinophil

numbers (**Figure S5**). However, we did observe increased expression of *Chil3* in the lungs of

infected mice treated with MCC950 (**Figure 7f**). Although consistent with our findings in

infected *Nlrp3*$^{-/-}$ mice, other type 2 markers including IL-4 itself were unchanged in

360   MCC950 treated or *Casp1/11*$^{-/-}$ mice (data not shown). Further, in contrast to NLRP3

deficiency, there was no difference in intestinal worm burden between untreated and

MCC950-treated mice (**Figure 7g**). Together, these data strongly suggest that NLRP3 plays

an inflammasome-independent role in constraining neutrophil and anti-helminth immunity in

the lung.

Figure 7 Inflammasome-independent role for NLRP3 during innate anti-helminth responses in the lung. WT and *Nlrp3*$^{-/-}$ mice were infected with *N. brasiliensis* (Nb) and on day 2 post-infection (**a**) intracellular pro-IL-1β levels were measured in alveolar (alv.) Mφs by flow cytometry, (**b**) released IL-1β in the BALF was quantified by ELISA and (**c**) frequency of active caspase-1 was measured *ex vivo* in BAL alveolar Mφs, neutrophils, and eosinophils by FAM-FLICA fluorescence. (**d**) On day 2 post-infection, L4 lung stage larvae and (**e**) day 6 adult intestinal worms were counted in WT, *Casp1/11*$^{-/-}$ and *Nlrp3*$^{-/-}$ mice. WT mice were treated with MCC950 during Nb infection and (**f**) *Chil3* (Ym1) expression in the lung was measured by qRT-PCR and (**g**) intestinal worm burdens were counted on day 6 post-infection. Data are representative (**a-e, g**; mean ± s.e.m.) or pooled (**f**; mean ± s.e.m.) from 2 or 3 individual experiments with 3-5 mice per group (per experiment). *$P<0.05$, **$P<0.01$, ***$P<0.001$ (one-way ANOVA and Tukey-Kramer *post hoc* test).

Discussion

380   Whilst the role of NLRP3 in classical inflammatory settings has been well-characterised, the function of NLRP3 during type 2 inflammation has been more contentious. An early controversy focussed on whether the pro-type 2 adjuvant activity of alum is dependent on its ability to activate the NLRP3 inflammasome.[26,27] Subsequently, a variety of studies have found that NLRP3 either promotes[28–30] or restricts[24,31] allergic responses and

385   Th2 cells. Thus, the role of NLRP3 in type 2 immunity remains largely unresolved.

        To help resolve this question, we have focussed on helminth infections, in which type 2 immunity is central for host protection. We previously found that the NLRP3 inflammasome played a major role in suppressing the adaptive type 2 immune response to intestinal helminth infection (in the caecum) with *T. muris*.[12] In that study, MCC950

390   treatment and Caspase-1/11 deficiency was able to phenocopy *Nlrp3*[−/−] mice during infection. In contrast, the present study showed that during the innate immune phase of *N. brasiliensis* infection, NLRP3 had an unexpected, and apparently inflammasome-independent, role in constraining the initial neutrophilia as well as the eosinophilia that results from the type 2 responses to the parasite and subsequent expulsion in the small intestine. These data suggest

395   that the NLRP3 inflammasome has differential roles during innate and adaptive immunity and may be tissue-specific, acting differently in two very distinct helminth infection models.

        We previously published that Ym1 was responsible for a more aggressive early neutrophilic response in the lung, which promotes *N. brasiliensis* larval killing but at the cost of increased host tissue damage.[6] Ym1 can form crystals in Mφs[11] which may be a trigger for

400   NLRP3 inflammasome assembly. Classically, neutrophilic influx, which can be induced by the presence of necrotic cells and DAMPs such as extracellular ATP during acute tissue injury, has been shown to be critically driven by NLRP3.[32] Additionally, NLRP3

inflammasome-dependent release of IL-1β is critical for neutrophil anti-bacterial responses to *Streptococcus pneumoniae* infection in the lungs.[33] We therefore hypothesized that Ym1

405 acted via NLRP3, and NLRP3 deficiency would reverse Ym1-mediated neutrophil recruitment. However, our data counterintuitively showed that during acute lung injury with *N. brasiliensis* infection, NLRP3 restricted neutrophilia. In addition, we saw no evidence for inflammasome activation during innate responses and neither MCC950-mediated inflammasome inhibition, nor Caspase 1/11 deficiency, replicated NLRP3-deficiency during

410 *N. brasiliensis* infection. It is therefore possible that NLRP3 has distinct roles regulating neutrophil recruitment that can either be dependent or independent of inflammasome activation. Our data are consistent with another report in which NLRP3, but not caspase-1/11 or ASC, limited early neutrophil influx during lung bacterial infection with *Francisella tularensis*.[23] CXCL3, a neutrophil chemotactic factor that can mobilise neutrophils from the

415 BM,[34] was increased in the lungs of *N. brasiliensis* infected NLRP3-deficient mice, suggesting a potential mechanism for NLRP3-dependent regulation of neutrophil recruitment. However, the increased *Cxcl3* expression we observed may just reflect increased neutrophil numbers as recruited neutrophils can express *Cxcl3*.[20]

In line with previous studies,[5,6] the enhanced neutrophilia we observed in the absence

420 of NLRP3 resulted in increased lung damage and impaired repair mechanisms. The initiation of repair responses in the lung following injury due to *N. brasiliensis* migration is dependent on IL-4Rα signalling.[5] Our transcriptional analysis revealed enhanced alternative Mφ activation and other type 2 markers such as *Il4* and *Arg1* in infected *Nlrp3$^{-/-}$* mice relative to WT controls. This correlated with enhanced *Ccl24* (Eotaxin-2) expression and increased

425 eosinophilia in infected *Nlrp3$^{-/-}$* mice. Further, *Mmp13* expression was also elevated in infected *Nlrp3$^{-/-}$* mice. As MMP13 has been shown to be regulated by IL-4Rα during *N. brasiliensis* infection,[5] the data further highlights possible type 2-dependent defects in tissue

repair processes in the absence of NLRP3. Thus, in our model NLRP3 limits the host-protective type 2 response and thus in its absence, one might expect enhanced repair.

430   However, paradoxically there was persistent damage in the gene deficient mice, suggesting NLRP3 is required for the timely resolution of the early inflammatory response. It is possible that type 2 responses are exaggerated in the absence of NLRP3 due to a compensatory need to repair the damaged tissues. Our data may also fit the hypothesis that early neutrophil responses are required to establish a type 2 response[35] and because we saw enhanced

435   neutrophilia in NLRP3-deficient mice, the magnitude of the resulting type 2 response could be exacerbated.

These findings highlight a complex relationship between NLRP3 and IL-4. It has been previously reported that IL-4 signalling negatively regulates NLRP3 by suppressing inflammasome assembly and caspase-1 activity as well as the subcellular localization of

440   NLRP3.[36] Additionally, in CD4[+] T cells it has been shown that NLRP3 can act as a transcription factor that supports differentiation to the Th2 cell lineage in an inflammasome-independent manner.[22] Our data suggest an additional mechanism by which NLRP3 can negatively regulate Mφ IL-4 responsiveness and controls expression of alternative activation markers during helminth infection in the lungs, independent of the inflammasome. Whether

445   NLRP3 is acting as a transcription factor in Mφs similar to its action in Th2 cells and how mechanistically, NLRP3 regulates IL-4 signalling, will be the subject of future work. It will also be important to clarify how inflammasome-dependent and -independent roles for NLRP3 relate to innate versus adaptive immune responses within type 2 disease contexts.

Materials and Methods

Mice and ethics statements

Nlrp3[tm1Vmd] (RRID:MGI:5468973), B6-*Nlrp3*[tm1Tsc]/Siec (RRID:IMSR_MUGEN:M153001) and *Casp1/11*$^{-/-}$ (RRID:IMSR_JAX:016621) mice were maintained on a C57BL/6J (RRID:IMSR_JAX:000664) background and bred in-house at the University of Manchester and James Cook University (JCU). All experiments were carried out in accordance with the United Kingdom Animals (Scientific Procedures) Act 1986 and under a Project License (70/8548) granted by the Home Office and approved by local Animal Ethics Review Group at the University of Manchester. Animal experimental protocols in Australia were approved by the JCU Animal Ethics Committee (A2213).

***N. brasiliensis* infection**

N. brasiliensis worms were propagated as previously described.[37] Infective L3 larvae were isolated and washed with sterile PBS and counted using a dissecting microscope. Mice were injected with 250 or 500 L3s subcutaneously. Upon culling the mice by pentobarbitone overdose i.p., BAL was performed with 10% FBS in PBS and lung lobes were collected. Lobes were either stored in RNAlater (Thermofisher), fixed in 10% formalin for histology or digested with Liberase TL (Roche). For lung-stage L4 counts, on days 1-2 post-infection, lungs were minced and incubated in PBS for 3 hours at 37 °C. Emergent larvae were counted using a dissecting microscope. For small intestinal worm burdens, worms were counted using a dissecting microscope following incubation at 37 °C. Fecal parasite eggs were enumerated from one fecal pellet/animal collected 6 days post-infection using a Whitlock paracytometer.

In vitro larval killing

CD11b$^+$ and Ly6G$^+$ neutrophils were sorted from BM of WT mice. For Mφ isolation, BM was isolated from WT mice and cells were cultured at 1 x 10^6 cells/ml in DMEM supplemented with 10% FBS, 5% horse serum (Invitrogen), 2 mM L-glutamine, 10 mM HEPES, 1 mM sodium pyruvate and 10 ng/ml M-CSF. On day 6, Mφs were plated overnight at a concentration of 2 x 10^5 cells/ml with RPMI 1640 supplemented with 10% heat-inactivated FBS and 100 U/ml penicillin plus 100 μg/ml streptomycin and incubated for a further 24 h. Normal mouse serum was added to plates (25%) as a source of complement. *In vitro* killing assays were performed by plating 100 *N. brasiliensis* L3s into 48-well plates with a 250 μl final volume containing Mφs or neutrophils (1 x 10^6 cells). Well contents were then assessed at 24 h for larval viability. Larvae were considered viable if they were visibly motile/curved. If larvae were straight and non-motile then they were classified as non-viable.

NLRP3 inhibition

One day prior to *N. brasiliensis* infection, mice were treated with either PBS vehicle or 10 mg/kg MCC950 (Sigma) by intraperitoneal injection in a 100 μl volume, a dose which has been previously shown to inhibit inflammasome activity.[38] Injections were repeated daily up until the experimental endpoint.

IL-4 complex injection and *in vitro* IL-4 stimulation

For *in vivo* IL-4 stimulation, mice were injected intraperitoneally with 1.5 μg recombinant IL-4 complexed with 7.5 μg anti-IL-4. After 24 hours, lavage was performed to isolate peritoneal exudate cells containing predominantly Mφs which were analysed by flow cytometry. For *in vitro* stimulation, day 6 cultured BM-derived Mφs (differentiated with

L929-conditioned media) were treated with 20 ng/ml IL-4 and analysed by flow cytometry

the following day.

Ex vivo **caspase-1 activation assay**

BAL cells were collected from mice at day 2 post-infection with *N. brasiliensis*. Cells were

then stained for inflammasome activation using a caspase-1 FAM-YVAD-FMK FLICA

probe kit (Bio-Rad). Cells were then processed for analysis by flow cytometry.

Flow cytometry

Single cell suspensions were washed in PBS and Live/Dead staining (Thermofisher) was

performed. Samples were Fc-blocked using α-CD16/32 (2.4G2, RRID:AB_394656) (BD

Biosciences) and mouse serum (BioRad). Blocking and subsequent surface staining was

performed using PBS containing 2 mM EDTA, 2% FBS, and 0.05% NaN$_3$. Antibodies used

for staining are listed in **Table S3**. Following surface staining, cells were incubated with IC

fixation buffer (ThermoFisher) prior to permeabilization for intracellular staining.

Intracellular antigens and antibodies used to detect them were. For secondary detection of

Ym1 and RELM-α, Zenon goat (RRID:AB_2753200) and rabbit (RRID:AB_2572214)

antibody labels (ThermoFisher) were used. For cell quantification, some samples were spiked

with 10 μm polystyrene beads (Sigma-Aldrich) prior to acquisition. Data were acquired on a

BD LSRFortessa flow cytometer and analysed using FlowJo v10 software.

ELISA

BAL supernatants were analysed for Ym1 using commercially available ELISA kits (R&D

systems and Peprotech, respectively). Lung homogenates were assayed for IL-4 using

standard sandwich ELISA protocols (eBioscience). Analytes were detected using horse radish

peroxidase-conjugated streptavidin and TMB substrate (BioLegend) and stopped with 1 M

525   HCl. Final absorbance at 450 nm was measured using a VersaMax microplate reader

(Molecular Devices).

RNA extraction and quantitative real-time PCR

Tissue samples stored in RNAlater (Thermo Fisher Scientific) were processed for RNA

530   extraction using a TissueLyser II and QIAzol reagent (Qiagen). Isolated RNA was quantified

using a Qubit fluorimeter and RNA BR kit (Qiagen). cDNA was synthesized using Tetro

reverse transcription kit (Bioline) and oligo dT 15-mers (Integrated DNA Technologies).

Quantitative real-time PCR was performed using SYBR green mix (Agilent Technologies)

and a LightCycler 480 II (Roche). A list of primer sequences used are shown in **Table S4**.

Transcriptional profiling

Quality control was performed on RNA samples with an Agilent 2200 TapeStation system

prior to downstream analyses. Samples were diluted and 100 ng of RNA was processed for

running on a Nanostring nCounter® FLEX system using the Myeloid Innate Immunity v2

540   panel. Raw counts were normalized to internal spike in controls and the expression of 13

stable housekeeping genes, as determined by geNorm algorithm within the nSolver Advanced

Analysis tool. Subsequent analyses were performed in R (version 3.5.3). After normalization,

transcripts with >15 counts were considered to be expressed and were $\log_2$ transformed.

Linear modelling using the limma R package[39] was used to calculate differential gene

545   expression. All expressed genes were used for principal components analysis (PCA).

Unsupervised hierarchical clustering was performed on significantly differentially expressed

genes between *N. brasiliensis*-infected WT and *Nlrp3*[−/−] mice the using the linkage method

complete and Euclidian distances.

Histology and fractal analysis

550 Proximal small intestine was fixed in 4% formaldehyde and embedded in paraffin, and 5 μM

sections were stained with PAS/Alcian Blue stains using the standard protocol of an

institutional histology service provider (QIMR Berghofer Medical Research Institute). Whole

left lung lobes were paraffin embedded and 5 μm sections were prepared for

haematoxylin/eosin staining. Slides were imaged using an Olympus slide scanner and high-

555 resolution image files were exported using Pannoramic Viewer software (3DHISTECH). The

images were then processed in a KNIME software workflow to obtain 50 random regions of

interests (ROIs) across the whole lung section. ROIs that contained lobe boundaries or

extensive artefacts were excluded from the analysis. The ROIs were then converted to binary

images and lacunarity (Λ) was quantified using the FracLac plugin for ImageJ (default

560 settings). The Λ values of all the ROIs were averaged to obtain estimates for the entire lobe.

Statistical analyses

Graphpad Prism 7 software was used for all statistical analyses. Data were assessed to be

normally distributed by the D'Agostino-Pearson omnibus normality test. Differences between

565 experimental groups were assessed by ANOVA (for normally distributed data) followed by

Tukey-Kramer *post hoc* multiple comparisons test or an unpaired two-tailed Student's *t* test.

In cases where data were not normally distributed, a Kruskal-Wallis test was used. For gene

expression data, values were $\log_2$ transformed to achieve normal distribution. Comparisons

with a *P* value of less than 0.05 were considered to be statistically significant.

Acknowledgements

This work was supported by the Wellcome Trust (106898/A/15/Z to JEA), the Medical Research Council UK (MR/K01207X/1 and MR/J001929/1 to JEA; MR/N003586/1 to DB), the Australian National Health and Medical Research Council (NHMRC grants 1117504 and 1132975 to AL), the Queensland Department of Science, Information Technology and Innovation (to PRG) and the Iraqi Cultural Attache in Australia (to RA). TES was supported by Medical Research Foundation UK joint funding with Asthma UK (MRFAUK-2015-302). We thank Vishva Dixit (Genentech) for the $Nlrp3^{tm1Vmd}$ mice. Thanks to Seth Masters (WEHI, Australia) for kindly providing B6-$Nlrp3^{tm1Tsc}$/Siec and Caspase1/11 double-deficient mice. We thank the Flow Cytometry, Bioimaging, Genomic Technologies, and Biological Services core facilities at the University of Manchester.

Author Contributions

ALC, RA, AK, AABR, TES, PRG and JEA designed the research plan. ALC, RA, ZA, JA, JP, MMC, BHKC and RME performed experiments, analysed data and revised the manuscript. LAD, AABR, AK, DB, AL and TES provided materials, intellectual input and revised the manuscript. ALC, PRG, and JEA wrote the manuscript. JEA, PRG and AL supervised the work and provided funding.

Disclosure

The authors declare no competing interests.

References

1. Martinez, F.O. Macrophage activation and polarization. *Front. Biosci.* **13**, 453 (2008).

2. Sutherland, T.E. *et al.* Ym1 induces RELMα and rescues IL-4Rα deficiency in lung repair during nematode infection. *PLOS Pathog.* **14**, e1007423 (2018).

3. Kreider, T., Anthony, R.M., Urban, J.F., Gause, W.C. & Gause, W.C. Alternatively activated macrophages in helminth infections. *Curr. Opin. Immunol.* **19**, 448–453 (2007).

4. Knipper, J.A. *et al.* Interleukin-4 receptor α signaling in myeloid cells controls collagen fibril assembly in skin repair. *Immunity* **43**, 803–816 (2015).

5. Chen, F. *et al.* An essential role for TH2-type responses in limiting acute tissue damage during experimental helminth infection. *Nat. Med.* **18**, 260–266 (2012).

6. Sutherland, T.E. *et al.* Chitinase-like proteins promote IL-17-mediated neutrophilia in a tradeoff between nematode killing and host damage. *Nat. Immunol.* **15**, 1116–1125 (2014).

7. Muallem, G. & Hunter, C.A. ParadYm shift: Ym1 and Ym2 as innate immunological regulators of IL-17. *Nat. Immunol.* **15**, 1099–1100 (2014).

8. Schroder, K. & Tschopp, J. The inflammasomes. *Cell* **140**, 821–832 (2010).

9. Boucher, D. *et al.* Caspase-1 self-cleavage is an intrinsic mechanism to terminate inflammasome activity. *J. Exp. Med.* **215**, 827–840 (2018).

10. Man, S.M., Karki, R. & Kanneganti, T.-D. Molecular mechanisms and functions of pyroptosis, inflammatory caspases and inflammasomes in infectious diseases. *Immunol. Rev.* **277**, 61–75 (2017).

11. Liu, Q. *et al.* p47phox deficiency induces macrophage dysfunction resulting in progressive crystalline macrophage pneumonia. *Am. J. Pathol.* **174**, 153–163 (2009).

12. Alhallaf, R. *et al.* The NLRP3 inflammasome suppresses protective immunity to gastrointestinal helminth infection. *Cell Rep.* **23**, 1085–1098 (2018).

13. Chen, F. *et al.* An essential role for TH2-type responses in limiting acute tissue damage during experimental helminth infection. *Nat. Med.* **18**, 260–266 (2012).

14. Chen, F. *et al.* Neutrophils prime a long-lived effector macrophage phenotype that mediates accelerated helminth expulsion. *Nat. Immunol.* **15**, 938–946 (2014).

15. Gause, W.C., Wynn, T.A. & Allen, J.E. Type 2 immunity and wound healing: evolutionary refinement of adaptive immunity by helminths. *Nat. Rev. Immunol.* **13**, 607–614 (2013).

625   16.   Reece, J.J., Siracusa, M.C. & Scott, A.L. Innate immune responses to lung-stage helminth infection induce alternatively activated alveolar macrophages. *Infect. Immun.* **74**, 4970–4981 (2006).

17.   Marsland, B.J., Kurrer, M., Reissmann, R., Harris, N.L. & Kopf, M. Nippostrongylus brasiliensis infection leads to the development of emphysema associated with the

630   induction of alternatively activated macrophages. *Eur. J. Immunol.* **38**, 479–488 (2008).

18.   Porzionato, A. *et al.* Fractal analysis of alveolarization in hyperoxia-induced rat models of bronchopulmonary dysplasia. *Am. J. Physiol. Cell. Mol. Physiol.* **310**, L680–L688 (2016).

635   19.   Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. R. Stat. Soc. Ser. B* **57**, 289–300 (1995).

20.   Shay, T. & Kang, J. Immunological Genome Project and systems immunology. *Trends Immunol.* **34**, 602–609 (2013).

21.   Gundra, U.M. *et al.* Alternatively activated macrophages derived from monocytes and

640   tissue macrophages are phenotypically and functionally distinct. *Blood* **123**, e110-22 (2014).

22.   Bruchard, M. *et al.* The receptor NLRP3 is a transcriptional regulator of TH2 differentiation. *Nat. Immunol.* **16**, 859–870 (2015).

23.   Periasamy, S., Le, H.T., Duffy, E.B., Chin, H. & Harton, J.A. Inflammasome-

645   independent NLRP3 restriction of a protective early neutrophil response to pulmonary tularemia. *PLOS Pathog.* **12**, e1006059 (2016).

24.   Madouri, F. *et al.* Caspase-1 activation by NLRP3 inflammasome dampens IL-33-dependent house dust mite-induced allergic lung inflammation. *J. Mol. Cell Biol.* **7**, 351–365 (2015).

650   25.   Coll, R.C. *et al.* A small-molecule inhibitor of the NLRP3 inflammasome for the treatment of inflammatory diseases. *Nat. Med.* **21**, 248–255 (2015).

26.   Kool, M. *et al.* Alum adjuvant stimulates inflammatory dendritic cells through activation of the NALP3 inflammasome. *J. Immunol.* **181**, 3755–3759 (2008).

27.   Franchi, L. & Núñez, G. The NLRP3 inflammasome is critical for aluminium

655   hydroxide-mediated IL-1beta secretion but dispensable for adjuvant activity. *Eur. J. Immunol.* **38**, 2085–2089 (2008).

28.   Besnard, A.-G. *et al.* NLRP3 inflammasome is required in murine asthma in the absence of aluminum adjuvant. *Allergy* **66**, 1047–1057 (2011).

29. Primiano, M.J. *et al.* Efficacy and pharmacology of the NLRP3 inflammasome inhibitor CP-456,773 (CRID3) in murine models of dermal and pulmonary inflammation. *J. Immunol.* **197**, 2421–2433 (2016).

30. Kim, R.Y. *et al.* Role for NLRP3 inflammasome-mediated, IL-1β-dependent responses in severe, steroid-resistant asthma. *Am. J. Respir. Crit. Care Med.* **196**, 283–297 (2017).

31. Huang, H. *et al.* Vitamin D receptor interacts with NLRP3 to restrict the allergic response. *Clin. Exp. Immunol.* **194**, 17–26 (2018).

32. Iyer, S.S. *et al.* Necrotic cells trigger a sterile inflammatory response through the NLRP3 inflammasome. *Proc. Natl. Acad. Sci.* **106**, 20388–20393 (2009).

33. Karmakar, M. *et al.* Neutrophil IL-1β processing induced by pneumolysin is mediated by the NLRP3/ASC inflammasome and caspase-1 activation and is dependent on K + efflux. *J. Immunol.* **194**, 1763–1775 (2015).

34. Burdon, P.C.E., Martin, C. & Rankin, S.M. The CXC chemokine MIP-2 stimulates neutrophil mobilization from the rat bone marrow in a CD49d-dependent manner. *Blood* **105**, 2543–2548 (2005).

35. Tacchini-Cottier, F. *et al.* An immunomodulatory function for neutrophils during the induction of a CD4+ Th2 response in BALB/c mice infected with Leishmania major. *J. Immunol.* **165**, 2628–2636 (2000).

36. Hwang, I. *et al.* Non-transcriptional regulation of NLRP3 inflammasome signaling by IL-4. *Immunol. Cell Biol.* **93**, 591–599 (2015).

37. Lawrence, R.A., Gray, C.A., Osborne, J. & Maizels, R.M. Nippostrongylus brasiliensis: cytokine responses and nematode expulsion in normal and IL-4-deficient mice. *Exp. Parasitol.* **84**, 65–73 (1996).

38. Krishnan, S.M. *et al.* Pharmacological inhibition of the NLRP3 inflammasome reduces blood pressure, renal damage, and dysfunction in salt-sensitive hypertension. *Cardiovasc. Res.* **115**, 776–787 (2019).

39. Ritchie, M.E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res.* **43**, e47–e47 (2015).