# Differential Metabolic Sensitivity of mTORC1- and mTORC2-Dependent Overgrowth

Maelle Devilliers[1], Damien Garrido[1,‡], Mickael Poidevin[1], Thomas Rubin[1,§], Arnaud Le Rouzic[2], and Jacques Montagne[1,*]

[1] Institute for Integrative Biology of the Cell (I2BC), CNRS, Université Paris-Sud, CEA, F-91190, Gif-sur-Yvette, France

[2] Laboratoire Evolution, Génomes, Comportement et Ecologie, CNRS, Université Paris-Sud, UMR 9191, F-91190, Gif-sur-Yvette, France

* Correspondence: Jacques.MONTAGNE@i2bc.paris-saclay.fr

‡ Present address: IRIC, Université de Montréal, Montréal, Québec H3T 1J4, Canada

§ Present address: Institut Curie, CNRS UMR 3215 / INSERM U-934, F-75248 Paris Cedex 5

Key words: Lipogenesis, Glycolysis, cell-autonomous effect, homeostasis

Short title: Metabolism and mTOR-induced overgrowth

**ABSTRACT**

The protein kinase mTOR is implicated in metabolic-related diseases and chiefly controls organismal growth and homeostasis in response to nutrients. Activation of mTOR promotes cell growth and enhances a glycolytic/lipogenic axis, suggesting that this metabolic axis is required to sustain mTOR-dependent growth. Here, we used *Drosophila* genetics to investigate this functional link both at the organismal and cell-autonomous levels. mTOR is present in two distinct complexes mTORC1 and mTORC2, which can be independently modulated in most *Drosophila* tissues. We confirm this independency in the fat body, the organ that fulfils hepatic and adipose functions. We show that ubiquitous mTOR over-activation affects carbohydrate and lipid metabolism, supporting the use of *Drosophila* as a powerful model to study the link between mTOR and metabolism. We show that targeted glycolytic or lipogenic restriction in fat body cells exclusively impedes mTORC2-induced overgrowth. Additionally, ubiquitous deficiency of lipogenesis (*FASN* mutants) results in a drop in mTORC1 but not mTORC2 signaling, whereas, at the cell-autonomous level, lipogenesis deficiency in fat body cells affects neither mTORC1 nor mTORC2 activity. These findings thus, reveal differential metabolic sensitivity of mTORC1- and mTORC2-dependent overgrowth. Furthermore, they suggest that local metabolic defects may elicit compensatory pathways between neighboring cells, whereas enzyme knockdown in the whole organism results in animal death. Importantly, our study weakens the use of single inhibitors to fight mTOR-related diseases and strengthens the use of drug combination and selective tissue-targeting.

**AUTHOR SUMMARY**

Cell growth is essential for animal development but is also deregulated in several human

diseases including cancers. The mTOR signaling network controls cell growth in response to

nutrients and growth factors. mTOR stimulation has been shown to promote basal metabolism,

including the glycolytic/lipogenic axis, which is essential to provide energetic cofactors and

building blocks to sustain cell growth. However, the requirement this metabolic axis for mTOR-

dependent growth has been poorly investigated yet. To address this issue, we used *Drosophila*

genetics and focused on the fat tissue that fulfils hepatic and adipose functions. Surprisingly, in

mosaic animals, where only a few cells were genetically stimulated for mTOR, we observed that

metabolic knockdown only moderates cell overgrowth. Additionally, this growth restriction

operates for only one of the two mTOR signaling branches, suggesting that mTOR-dependent

compensatory processes operate to circumvent metabolic defects. Given that ubiquitous

knockdown of this metabolic axis is essential for juvenile development and adult survival, our

study reveals that metabolic restriction is unlikely sufficient to counteract overgrowth disorders.

## INTRODUCTION

Growth of a multicellular organism is coordinated by signaling pathways that adjust intracellular processes to environmental changes. These signaling pathways include the mTOR (mechanistic Target Of Rapamycin) regulatory network that integrates the growth factor response as well as the nutritional and energetic status [1-7]. This network consists in two signaling branches in which the mTOR protein kinase is present in two distinct complexes, mTORC1 and mTORC2 that comprise raptor and rictor, respectively [8, 9]. Both mTORC1 and mTORC2 promote basal cellular functions, thereby providing building blocks to sustain cellular growth. However, despite a plethora of studies on mTORC1/2 signaling and functions, the requirement of basal metabolism—glycolytic/lipogenic axis— for mTOR-dependent cell growth has not been systematically investigated. The *Drosophila* model provides a powerful genetic system to address these issues [10], since both the intermediates of the mTOR signaling network and the basal metabolic pathways are conserved in the fruit fly [11-15].

Regulation of mTORC1 activity by ATP and amino acids depends on a multi-step process that results in the recruitment of an mTORC1 homodimer at the lysosomal membrane in the vicinity of the small GTPase Rheb (Ras homologue enriched in brain) [16-20]. Rheb stimulates mTORC1 activity [21], which in turn regulates several downstream targets. S6Kinase1 (S6K1) is one such kinase, sequentially activated through the phosphorylation of its T389 and T229 residues by mTORC1 and by PDK1 (Phosphoinositide-dependent protein kinase 1), respectively [22, 23]. Further, Rheb activation of mTORC1 is repressed by the tumor suppressor TSC (Tuberous sclerosis complex) that comprises subunits TSC1 and TSC2 [24-27].

The mTORC2 branch resides within the insulin signaling cascade [9]. Binding of insulin or related peptides to their cognate receptors results in recruitment of class I PI3K (Phosphoinositide 3-kinase) to the membrane. PI3K phosphorylates inositol lipids producing phosphatidylinositol-3,4,5-triphosphate (PIP3) [28, 29], while the tumor suppressor PTEN acts as a lipid phosphatase to counteract this process [30]. PIP3 constitutes a membrane docking

site for the protein kinase Akt whose subsequent activation depends on the phosphorylation of its S473 and T308 residues by mTORC2 and PDK1, respectively [31].

Constitutive activation of mTORC1 in MEFs (Mouse embryonic fibroblasts) has been shown to stimulate a metabolic network, including glycolysis, the pentose phosphate pathway and the biosynthesis of fatty acid (FA) and cholesterol [32]. Most of the genes encoding glycolytic enzymes are over-expressed in these cells as are those encoding LDH (lactate dehydrogenase) and Pdk1 (Pyruvate dehydrogenase kinase 1; an inhibitor of mitochondrial pyruvate processing). This suggests that mTORC1-activated MEFs potentiate anaerobic glycolysis and repress the tricarboxylic acid cycle (TCA). Conversely, adipose-specific knockout of raptor to impede mTORC1 formation, results in enhanced uncoupling of mitochondrial activity [33]. The increased lipogenesis observed in mTORC1 stimulated cells depends on a downstream transcriptional regulatory axis involving the cofactor Lipin 1 along with a SREBP (Sterol responsive element binding-protein) family member, which activates genes encoding lipogenic enzymes [32, 34]. Congruently, another study revealed that TSC2 mutant cells become addicted to glucose as a result of mTORC1 hyper-activity [35]. In addition, inhibition of mTORC1 activity revealed that these TSC2 mutant cells become also dependent on glutamine catabolism [36]; mTORC1 potentiates this catabolism to feed TCA anaplerosis, through 1) a S6K/eIF4B/Myc axis that increases glutaminase protein levels [37] and 2) the repression of SIRT4, a mitochondrial sirtuin that inhibits glutamine dehydrogenase [38].

Besides mTORC1 mediated regulation, mTORC2 also impinges on basal metabolism. Intracellular activation of Akt increases ATP levels [39, 40] through the stimulation of GLUT4-mediated glucose uptake [41] and the enhancement of the expression and activity of glycolytic enzymes [42, 43]. Akt also dampens glucose production by suppressing PEPCK (gluconeogenesis), glucose-6-phosphatase (glycogenolyse) and the glycogen synthesis repressor GSK3 [44, 45]. However, in contrast to mTORC1, Akt also promotes mitochondrial metabolism and oxidative phosphorylations [42, 46]. Conversely, hepatic knockout of the mTORC2 specific-subunit rictor results in constitutive gluconeogenesis and impaired glycolysis

and lipogenesis [47, 48]. Taken together, these studies strongly emphasize the role of mTOR in metabolic-related diseases and in adjusting metabolism to the nutritional and energetic status [7].

In the present study, we investigated the requirement of the glycolytic/lipogenic axis for the cellular growth induced by mTORC1 and mTORC2 hyper-activation in *Drosophila*. As previously demonstrated mTORC1 and mTORC2 reside on independent signaling branches in most *Drosophila* tissues [24, 49-52]. Here, we confirmed mTORC1 and 2 independency in the fat body (FB), the organ that fulfils hepatic and adipose functions to control body homeostasis in *Drosophila* [12, 13, 15]. We analyzed the effect of ubiquitous mTOR activation on glycolysis and lipogenesis. We show that in increase in mTOR activity provokes an apparent enhancement of metabolite consumption. Furthermore, our study reveals that metabolic restriction at the organismal level has dramatic consequences on animal survival, but no effect at the cell-autonomous level, suggesting that within an organism, alternative pathways may operate to compensate local metabolic defects. Nonetheless, at the cell-autonomous level, metabolic restriction can restrain mTORC2- but not mTORC1-induced overgrowth, indicating that the potential compensatory metabolic pathways do not fully operate in the context of mTORC2 over-activation.

**RESULTS**

**mTORC1 and mTORC2 independency in the fat body**

Previous studies showed that mTORC1 and mTORC2 reside on independent signaling branches in most *Drosophila* tissues [24, 49-52]. Activating either the mTORC1 or the mTORC2 signaling branch can be performed by overexpressing Rheb or depleting PTEN, respectively. To investigate mTORC1/2 independency in the FB, we generated somatic clones either over-expressing Rheb (*Rheb*+) [53] or homozygote for a *PTEN* mutation (*PTEN*-/-) [54]. The precursors of FB cells divide in the embryo; during larval life, the differentiated cells do not

divide but endoreplicate their DNA content to reach a giant size [55]. Therefore, to precisely

evaluate the effect on cell growth, somatic recombination events were induced during

embryogenesis at the stage of proliferation of the FB cell precursors and the resulting MARCM

clones were analyzed in the FB of late feeding L3 larvae, prior to the wandering stage that

precedes metamorphosis entry. Both $PTEN^{-/-}$ and $Rheb^+$ clonal cells were bigger than the

surrounding control cells and this cell size effect was dramatically increased in $PTEN^{-/-};Rheb^+$

combined clones (Figure 1A-D and 1M). We next analyzed this growth increase in the context of

the previously described $mTOR^{2L1}$ and $mTOR^{\Delta P}$ mutations. However, we could not find mutant

clones in the FB. Consistent with previous studies reporting that mTOR is critically required for

cell growth of endoreplicative tissues [54, 56], we reasoned that these clonal cells were likely

eliminated by cell competition [57]. Thus, we generated somatic clones in a *Minute* background

to slow down the growth of the surrounding control cells. In these conditions, *mTOR* mutant

clones could indeed be recovered. Both $mTOR^{2L1}$ and $mTOR^{\Delta P}$ mutant cells exhibited a

dramatic size reduction (Figure 1G and 1J) and this phenotype was dominant in $Rheb^+$

combined clonal cells (compare Figure 1E to 1H and 1K). In contrast, $mTOR^{\Delta P}$ but not $mTOR^{2L1}$

exhibited a clear dominant phenotype over the $PTEN^{-/-}$ mutation; the size of $mTOR^{\Delta P},PTEN^{-/-}$

clonal cells was dramatically reduced, whereas $mTOR^{2L1},PTEN^{-/-}$ clonal cells were giant

(compare Figure 1F to 1I and 1L). These findings indicate that the $mTOR^{2L1}$ mutation affects

mTORC1 but not mTORC2 signaling, whereas $mTOR^{\Delta P}$ affects both signaling branches.

Next, we used phospho-specific antibodies in immunostaining assays to analyze the

phosphorylation of Akt (P-Akt) and of the dS6K target, ribosomal protein rpS6 (P-S6). In $PTEN^{-/-}$

clonal cells, we observed an increase in the P-Akt intracellular signal (Figure 2A). Importantly

the P-Akt intracellular signal was absent in $mTOR^{\Delta P}$ cells (Figure 2B) but not affected in

$mTOR^{2L1}$ cells (Figure 2C). Staining with the rpS6 phospho-specific antibody revealed a patchy

signal, with only a subset of cells expressing the P-S6 signal in the FB (Figure 2E-J), a pattern

previously described in the wing imaginal disc [58]. Therefore, to evaluate mTORC1 activity, we

measured the ratio of P-S6 positive cells among the population of $GFP^+$ clonal cells. For control

clones, only labeled by GFP, about half of them were P-S6 positive (Figure 2E and 2K),

whereas most of the $mTOR^{2L1}$ and $mTOR^{\Delta P}$ clones were P-S6 negative (Figure 2F, 2G and 2K).

Importantly, almost all the $Rheb^+$ cells were P-S6 positive (Figure 2H and 2K), whereas the ratio

of P-S6 positive cells was slightly but not significantly increased in the $PTEN^{-/-}$ cell population

(Figure 2I and 2K). Taken together, these findings confirm that mTORC1 and mTORC2 operate

independently in FB cells and reveal that the $mTOR^{2L1}$ mutation affects only mTORC1, whereas

the $mTOR^{\Delta P}$ mutation affects both signaling branches.

**Activating mTORC1 or mTORC2 signaling impacts basal metabolism**

A number of studies support the notion that mTOR activity controls metabolism to sustain

cellular growth. To evaluate how mTORC1 and mTORC2 affect basal metabolism in *Drosophila*,

we analyzed various metabolites in whole animals that express the ubiquitous *da-gal4* driver to

direct Rheb overexpression ($Rheb^{++}$) or PTEN knockdown by RNA interference (*PTEN-RNAi*).

Larvae were fed either a standard or a 20%-sucrose supplemented diet (20%-SSD) and 0-5h

prepupae were collected, as this is a convenient phase to stage the animals after the feeding

period. When fed a standard diet, a high rate of lethality was observed for $Rheb^{++}$ and *PTEN-*

*RNAi* larvae, although a sufficient number of prepupae could be collected for metabolic analysis.

In contrast, none of the $Rheb^{++}$ and *PTEN-RNAi* larvae reached the prepupal stage when fed a

20%-SSD. Nonetheless, when $Rheb^{++}$ and *PTEN-RNAi* larvae were fed a standard diet during

early larval life and transferred onto a 20%-SSD at the L2/L3 molting transition, we could

recover a few prepupae for metabolic measurements. For both males and females fed a

standard diet, the body weight of $Rheb^{++}$ and *PTEN-RNAi* prepupae was roughly similar to that

of controls (Figure 3A and 3B). Conversely, providing a 20%-SSD resulted in a drop of the

prepupal weight of control animals that was significantly compensated in $Rheb^{++}$ and *PTEN-*

*RNAi* prepupae (Figure 3A and 3B).

Next, we measured the total amounts of protein, triacylglycerol (TAG), glycogen and trehalose—

the most abundant circulating sugar in *Drosophila*. Although variations in protein levels were

observed, none of them were statistically significant (Figure 3C). TAG levels in control prepupae were not affected by sucrose supplementation and did not vary in *PTEN-RNAi*, but were significantly decreased in *Rheb*[++] animals (Figure 3D). Feeding larvae a 20%-SSD since the L2/L3 molting transition resulted in a marked in increase in glycogen and trehalose levels in control prepupae (Figure 3E-F). In *Rheb*[++] and, in lower extent, in *PTEN-RNAi* prepupae, glycogen levels were significantly lower than those measured in controls (Figure 3E). Finally, trehalose levels were strongly decreased in both *Rheb*[++] and *PTEN-RNAi* prepupae fed either a standard or a 20%-SSD as compared to the control (Figure 3F). Taken together, these findings suggest that a ubiquitous increased activity of either mTORC1 or mTORC2 provokes an apparent increase in metabolite consumption. This metabolic rate is correlated with a relative increase in body weight for larvae fed a 20%-SSD, but not for those fed a standard diet. We previously observed that increasing dietary sucrose induced a reduction in food intake [59] that may account for the body weight reduction of control animals. Potentially, food intake could be less affected in *Rheb*[++] and *PTEN-RNAi* animals, thereby leading to a compensatory effect on body weight. Measuring food intake in *Rheb*[++] or *PTEN-RNAi* larvae was not applicable since most of them die during larval stage and thus, terminate feeding earlier. In sum, our data indicates that basal metabolism is altered in the few *Rheb*[++] or *PTEN-RNAi* larvae that survive and further suggests that in most cases stronger metabolic disruption happened, resulting in lethal homeostatic defects.

**Knocking-down glycolysis at the whole body level**

Since manipulating mTOR resulted in a decrease in the levels of TAG and glycogen stores and of circulating trehalose (Figure 3), we asked whether the basal energetic metabolism affected mTOR signaling. First, we ubiquitously expressed interfering RNA against phosphofructokinase1 (*PFK1-RNAi*), pyruvate kinase (*PK-RNAi*) pyruvate dehydrogenase (*PDH-RNAi*) and lactate dehydrogenase (*LDH-RNAi*). PFK1 catalyzes the third glycolytic reaction to form fructose 1,6-bisphophate; PK catalyzes the final glycolytic reaction to form

pyruvate; PDH directs the mitochondrial fate of pyruvate, whereas LDH directs its anaerobic fate (Figure 4A). When directed with the ubiquitous *da-gal4* driver, *PK-RNAi* provoked early larval lethality, *PFK1-RNAi* and *PDH-RNAi* provoked larval lethality at L2 or L3 stages, whereas *LDH-RNAi* induced a semi-lethal phenotype at larval or pupal stages (Figure 4B).

Second, to monitor mTOR activity following ubiquitous knockdown of glycolysis, the *da-gal4* driver was combined with a ubiquitous thermo-sensitive form of the Gal4 inhibitor, Gal80$^{ts}$ (*tub-gal80$^{ts}$*) that blocks Gal4 activity at 21°C but not at 29°C, thereby allowing RNAi expression after temperature shift. Each RNAi was ubiquitously induced at early L1 stage and protein extracts were prepared two days later using late L2 larvae. At this stage the larvae were still viable, although those expressing PK-RNAi did not undergo L2/L3 transition and eventually died (Figure 4B). Western-blotting using these L2 protein extracts revealed that RNAi-knockdown of PFK1, LDH or PDH did not affect Akt or dS6K phosphorylation (Figure 4C). In contrast, PK knockdown strongly decreased dS6K phosphorylation and to a lower extent Akt phosphorylation (Figure 4C). These results indicate that mTORC1 signaling may be affected when knocking down PK, but not when knocking down any other enzyme directly linked to glycolysis. Nonetheless, the lethal phenotype of PK-RNAi larvae occurring at the late L2 stage (Figure 4B) might weaken the larvae, inducing a subsequent effect on mTOR signaling.

To evaluate the requirement of glycolysis for adult survival, RNAi-knockdown was induced by temperature shift to 29°C in newly emerged flies and lethality was counted every second day. In both males and females, PK and PFK1 knockdown provoked lethality between 10 to 14 days after temperature shift (Figure 4D). Knockdown of PDH and LDH also induced adult lethality, although not as soon as PK and PFK1 knockdown (Figure 4D). As a comparison, we analyzed FASN knockdown in adults; about a quarter of *FASN-RNAi* flies died between 10 to 14 days, while the others survived nearly as well as control flies (Figure 4D). Taken together, these data indicate that glycolysis is essential for both larval development and adult survival. However, prior to the appearance of the deleterious phenotype, glycolysis knockdown is unlikely to impinge on mTOR signaling.

**Cell-autonomous requirement of glycolysis for mTORC2- but not mTORC1-dependent overgrowth**

To investigate the requirement of glycolysis to sustain mTOR-dependent growth at the cell-autonomous level, *PFK1-RNAi*, *PK-RNAi*, *PDH-RNAi* and *LDH-RNAi* were induced in *PTEN^-/-^* or *Rheb^+^* clones. Except a moderate effect of *PK-RNAi*, clones expressing interfering RNA against these metabolic enzymes did not significantly affect the growth of FB cells (Figure 5A-D and 5M). In combined clones, none of the RNAi affected the growth of *Rheb^+^* clones (Figure 5E-H and 5M). In contrast, the size of *PTEN^-/-^* clones was significantly decreased when co-expressing RNA against any of these metabolic enzymes (Figure 5I-M). These findings indicate that both aerobic and anaerobic glycolysis are required to sustain mTORC2-dependent overgrowth at the cell-autonomous level. In contrast, reducing glycolysis does not counteract mTORC1-dependent overgrowth, suggesting the existence of compensatory pathways.

**Effect of Lipogenesis on mTOR signaling**

Since glycolysis and FA synthesis are tightly connected metabolic pathways [59], we investigated whether lipogenesis affects mTOR signaling. FA synthesis is catalyzed by FASN (Figure 4A). The *Drosophila* genome encodes three *FASN* genes, *FASN1* is ubiquitously expressed but not *FASN2* or *FASN3* [60-62]. The deletion of the *FASN1* and *FASN2* tandem (*FASN^Δ24-23^* deletion, hereafter called *FASN^1-2^*) results in a lethal phenotype that can be rescued by feeding larvae a lipid-complemented diet (beySD) [59, 62]. We observed that beySD-rescued *FASN^1-2^* mutant larvae exhibited a delay in development, as measured by the duration of larval development to metamorphosis entry (Figure 6A). Further, when beySD-rescued *FASN^1-2^* mutant larvae were transferred at the L2/L3 larval transition onto a 10% sucrose-supplemented-beySD, only a few of them completed the third larval stage and, after an extreme developmental delay, entered metamorphosis (Figure 6A). Delay in development can be due to a default in

ecdysone production that results in giant pupae [63] or to impaired mTOR signaling that results in reduced body growth [54, 64]. Measurements of prepupal weight revealed that *FASN[1-2]* mutant prepupae exhibited a severe reduction in body weight, whether or not they were supplemented with sucrose (Figure 6B), suggesting a default in mTOR signaling. Therefore, we analyzed the phosphorylation of the *Drosophila* S6Kinase (dS6K) and Akt in protein extracts of late feeding L3 larvae. Western-blotting revealed that the dS6K protein resolved in several bands in *FASN[1-2]* extracts, whereas Akt protein was unchanged (Figure 6C). These results suggest that dS6K but not Akt might be degraded in the *FASN[1-2]* mutant background. In addition, dS6K phosphorylation decreased in *FASN[1-2]* extracts and became barely detectable when *FASN[1-2]* larvae were fed a sucrose-supplemented-beySD (Figure 6C). Conversely, the phosphorylation of Akt was unaffected in larvae fed a beySD, although it was slightly decreased in larvae fed a sucrose-supplemented-beySD (Figure 6C). This finding contrasts with our previous observation showing that FB explants of *FASN[1-2]* mutant larvae were hypersensitive to insulin [59]. However, *FASN[1-2]* mutants also exhibited a decrease in food intake [59], which might induce a systemic suppression of dS6K phosphorylation, while FB explant were cultured in nutrient media supplemented with insulin. Therefore, to determine whether *FASN* mutation affects mTOR signaling at the cell-autonomous level, we analyzed P-S6 and P-Akt in *FASN[1-2]* mutant clones in the FB. As for control clones, about half of the *FASN[1-2]* clonal cells were P-S6 positive (Figure 2J and 2K). Furthermore, no effect on P-Akt was observed in *FASN[1-2]* clonal cells (Figure 2D). In summary, these findings reveal that disrupting FA synthesis does not significantly affect mTORC1 and mTORC2 signaling at the cell-autonomous level, although it seems to impinge on mTORC1 signaling when inhibited in the whole animal whether directly or indirectly.

**Cell-autonomous requirement of FA synthesis for mTORC2- but not mTORC1-dependent overgrowth**

To determine, whether lipogenesis is required at the cell-autonomous level to sustain mTORC1

and/or mTORC2 dependent growth, we analyzed $FASN^{1-2}$ clones while enhancing either of the

mTOR signaling branch in FB cells. We previously reported [59] that $FASN^{1-2}$ clonal cells in the

FB were slightly reduced in size and that this effect was dramatically increased in larvae fed a

20%-SSD (Supplementary Figure S1 and Figure 7M). Therefore, we generated $PTEN^{-/-}$ and

$Rheb^+$ clones combined or not with the $FASN^{1-2}$ mutation and analyzed them in the FB of larvae

fed either a standard diet or a 20%-SSD. As compared to the standard diet, feeding larvae a

20%-SSD had no effect on the size of $Rheb^+$ clonal cells, but significantly reduced the size of

$PTEN^{-/-}$ and of $PTEN^{-/-}$;$Rheb^+$ clonal cells (Figure 7A-F and 7M). Further, when combined with

the $FASN^{1-2}$ mutation, $PTEN^{-/-}$ but not $Rheb^+$ clones were significantly reduced in size (Figure

7G-H and 7M). The $FASN^{1-2}$ mutation also provoked a severe size reduction of $PTEN^{-/-}$,$Rheb^+$

clones (Figure 7I and 7M). Moreover, as compared to the standard diet, feeding larvae a 20%-

SSD induced a significant size reduction of $FASN^{1-2}$;$Rheb^+$, $FASN^{1-2}$,$PTEN^{-/-}$ and $FASN^{1-2}$,$PTEN^{-}$

$^{/-}$;$Rheb^+$ clonal cells (Figure 7J-L and 7M). Of note, except for the $FASN^{1-2}$;$Rheb^+$ clonal cells in

larvae fed a 20%-SSD that exhibited a size roughly identical to that of the surrounding control

cells (Figure 7J), the cell size was always bigger than the controls (Figure 7M). These findings

indicate that, in larvae fed a standard diet, FA synthesis is at least in part required to sustain

over-growth induced by mTORC2, but not mTORC1. They also reveal that additional dietary

sucrose is rather detrimental for the growth of cells either deficient for FA synthesis or over-

active for mTORC2, suggesting that these cells have a restricted homeostatic ability to adjust to

an unbalanced diet, whereas mTORC1 activated cells at least in part maintain this ability.

**DISCUSSION**

In this study, we used the powerful *Drosophila* genetics to investigate the functional links

between mTOR-dependent growth and the glycolytic/lipogenic axis. On one hand, we

dampened this metabolic axis or enhanced mTORC1/2 activity in the whole organism to mimic

the effect that might be induced by drug treatment with a systemic inhibitor. On the other hand,

to monitor the cell growth process that spans the entire developmental program at the cell-

autonomous level, we analyzed clonal FB cells in mosaic animals. Intriguingly, our study reveals

apparent contradictory effects between perturbations at the whole body and cell-autonomous

levels.

At the organismal level, knockdown of glycolytic enzymes or deficiency of FASN result in animal

lethality. However, *FASN[1-2]* mutant animals supplemented with dietary lipids can survive but

exhibit a dramatic overall growth suppression. This growth defect might result from a decrease

in mTORC1 activity that is strongly reduced in *FASN[1-2]* mutant animals, suggesting that

mTORC1 but not mTORC2 signaling relies on lipogenesis. In contrast, at the cell autonomous

level, the mutation of *FASN[1-2]* restrains mTORC2 but not mTORC1 dependent overgrowth in FB

cells. These apparent contradictory findings, suggest that the growth defect and the reduction of

mTORC1 activity in *FASN[1-2]* mutants are not due to the addition of cell-autonomous effects but

rather to a systemic regulation. Potentially, FASN default might affect the activity of a specific

tissue, as for instance, the neurosecretory cells that synthesize and secrete insulin like peptides,

which promote systemic body growth [65]. Alternatively, considering that mTORC1 directly

responds to nutrients [18-20], the drop of mTORC1 activity may be a consequence of feeding,

since we previously reported a decrease in nutrient uptake in *FASN[1-2]* mutant animals [59].

Consistently, a previous study on the transcription factor mio —the *Drosophila* homologue of

mondoA and ChREBP that regulate the glycolytic/lipogenic axis in response to dietary sugar

[66, 67]— suggests the existence of a FASN-dependent effect in the FB on the control of food

intake [68]. FB-knockdown of *mio* results in the lack of sucrose-induced expression of *FASN1*

and in a decrease in food intake. This study suggests that the FASN default perturbs body

homeostasis and indirectly affects the neuronal control of feeding behavior. However, it does

not exclude that a lipogenic defect in neuronal cells may also directly impinge on feeding

behavior. Finally, the drop of mTORC1 activity observed in *FASN[1-2]* mutants may be a

consequence of malonyl-CoA accumulation, since mTOR malonylation has been reported to

inhibit mTORC1 but not mTORC2 activity [69]. Malonylation of mTOR may also account for the

size reduction of *FASN[1-2]* mutant cells over-expressing Rheb in animals fed a 20%-SSD, consistent with the increased expression of lipogenic enzymes induced by dietary sucrose [59]. Thus, mTOR malonylation and the subsequent decrease in mTORC1 activity might occur only when interfering with a context of high demand for lipogenesis.

Our study reveals that over-activation of mTORC1 and to a lesser extent of mTORC2, results in a decrease in glycogen and TAG stores and in circulating trehalose, suggesting that mTOR activation enhances metabolite consumption to sustain cell growth. It is therefore surprising that activation of neither mTORC1 nor mTORC2 induces an increase in body weight. Nonetheless, overall body growth depends on an intricate regulatory network that integrates cell-autonomous effects and humoral messages. For instance, previous studies reported that mTOR activation within the ring gland, results in a systemic decrease in body growth [70-72]. Therefore, mTOR ubiquitous activation is likely to promote the growth of most cells but might concurrently perturb endocrine signals dampening overall growth. Of note, we observed that larvae fed a 20%-SSD result in pupae with reduced body weight, an effect that is partially suppressed when either of the mTORC-signaling branch is over-activated. The fact that the overall body weight of mTOR-stimulated animals is maintained within a range likely compatible with organismal survival contrasts with the observed high rate of lethality. The decrease in stores and circulating sugars suggests that in these animals each cell tends to increase its basal metabolism evoking an egoist behavior that might perturb the equilibrium between cell-autonomous and systemic regulation. Thus, in a stressful situation, as when animals are fed a 20%-SSD, the need of a tight adjustment to an unbalanced diet may enhance the distortion between cell-autonomous effects and systemic regulation, resulting in an increased rate of lethality.

In agreement with other studies [24, 49-52], we show that the mTORC1 and mTORC2 branches work independently in the *Drosophila* FB and we provide evidence that the previously described *mTOR[2L1]* mutation [54] affects mTORC1 signaling only. A plethora of studies in mammalian cells indicate that mTOR activation directs metabolism towards glucose consumption, storage and anabolism [32, 34, 35, 39, 41-43, 73]. Our study rather suggests that in the *Drosophila*

larvae, mTOR promotes metabolite consumption through glycolysis but not storage. However,

at the cell-autonomous level, we observe that inhibition of lipogenesis or glycolysis restrains

neither larval FB cell growth nor overgrowth induced by mTORC1 stimulation in these cells.

These findings counteract the idea that mTORC1 potentiates a glycolytic/lipogenic axis [32] to

sustain cell growth. To overcome the lack of glycolytic products and of membrane lipids, these

cells may benefit of a transfer from neighboring cells and might favor alternative metabolic

pathways, including glutamine catabolism to feed TCA anaplerosis which has been shown to be

a crucial pathway in mTORC1-stimulated mammalian cells [36-38]. Nonetheless, such

compensatory processes do not fully operate to sustain mTORC2-dependent overgrowth. In

these cells, the mutation of PTEN results in mTORC2 hyper-activation and potentially impedes

the ability to modulate this signaling branch. Therefore, it is tempting to speculate that the

modulation of mTORC2 signaling at least in part contributes to the regulation of these

compensatory processes.

As a coordinator of growth and metabolism, mTOR plays a central role in tumor development [7,

74, 75]. PTEN, the tumor suppressor that counteracts PI3K activity upstream of mTORC2

signaling, is deficient in several human cancers [30]. Mutation of TSC1 or TSC2, the essential

subunits of the tuberous sclerosis complex, is associated with benign tumors but also with brain,

kidney and lung destructive diseases [76]. To investigate the role of mTOR regarding tumor

development, a recent study reported the generation of liver-specific double knockout mice for

TSC1 and PTEN [77]. These mice develop hepatic steatosis that eventually progresses to

hepatocellular carcinoma. Both processes are suppressed in mice fed the mTORC1/2 inhibitors

INK128, but not the mTORC1 inhibitor rapamycin, supporting an mTORC2 specific effect. The

combination of inhibitors against mTOR and metabolism is currently under clinical investigation

to fight cancers [7]. Importantly, our study reveals that ubiquitous inhibition of basal metabolism

produces dramatic effects during development, while it only moderates cell growth induced by

mTOR over-activation. Therefore, the use of drug therapy to fight cancer must be taken with

caution, in particular if development is not complete and most efforts should be made to selectively target sick tissues.

## MATERIAL & METHODS

### Genetics and fly handling

Fly strains: *P[w[+mC]=tubP-GAL80]LL10,P[ry[+t7.2]=neoFRT]40A*, *daughterless(da)-gal4*, *tub-gal80$^{ts}$*, *UAS-Dcr-2* (Bloomington Stock Center); *FASN$^{1-2}$* [59]; *mTOR$^{ΔP}$* [56]; *mTOR$^{2L1}$* and *PTEN* [54]; *EP(UAS)-Rheb* [53]; ); inducible RNA-interfering (*UAS-RNAi*) lines to *PTEN* (NIG 5671R-2), *FASN1* (VDRC 29349), *PFK1* (VDRC 3017), *PK* (VDRC 49533) *PDH* (VDRC 40410), *LDH* (VDRC 31192) [78]. The *Minute* stock used was previously referred to as *FRT40/P(arm-LacZ w$^+$)* [79] but exhibit both developmental delay and short and slender bristles, typically reported as *Minute* phenotype [57]. To generate MARCM clones in the *Minute* background, these flies were recombined with the *P[w[+mC]=tubP-GAL80]LL10,P[ry[+t7.2]=neoFRT]40A* chromosome.

The standard media used in this study contained agar (1g), polenta (6g) and yeast (4g) for 100ml. Lipid- (beySD) and sugar-complemented media were prepared as previously described [59].

To select *FASN$^{1-2}$* mutant larvae, we used a GFP-labelled CyO balancer chromosome. Flies were let to lay eggs on grape juice plates for less than 24 hrs. Then, some beySD media was put in the middle of the plates; larvae that do not express GFP were collected the next day and transferred to fresh tubes. Prepupae were collected once a day to evaluate developmental delay and to measure body weight.

### Molecular biology and Biochemistry

To test RNAi-knockdown efficacy to the glycolytic enzymes (Supplementary Figure S2), *UAS-Dcr-2;da-gal4,tub-gal80$^{ts}$* virgin females were mated with UAS-RNAi males. Flies were let to lay

eggs overnight and tubes were kept at 19°C for two days. Tubes were then transferred at 29°C

and two days later, larvae of roughly the same size were collected. Reverse transcription and

quantitative PCR were performed as previously described [60].

Protein extracts for western-blotting were prepared as previously described [51]. Antibody used

in for western-blotting have been previously described [51] or commercially provided for Akt

(Cell signaling 4054).

For metabolic measurements, parental flies were let to lay eggs in tubes for less than 24 hrs at

25°C. Tubes were then transferred at 29°C to strengthen the gal4/UAS effect, and using a *UAS-*

*Dcr-2* to strengthen the RNAi effect. Larvae were either maintained in the same tubes or

selected prior to L2/L3 transition and transferred on 20%-SSD. Collection of prepupae and

metabolic measurements were performed as previously described [59].

**Clonal analysis**

All the clones were generated using the MARCM strategy [80]. Parental flies were let to lay

eggs at 25°C for seven hrs. Tubes were then heat shocked for 65 minutes in a water bath at

38°C so that recombination happens while FB precursor cells are in dividing process. FB from

feeding larvae at the end of the L3 stage where dissected, fixed, membranes were labelled with

phalloidin and nuclei with DAPI, and FB were mounted as previously described [59]. Image

acquisitions were obtained using a Leica SP8 confocal laser-scanning microscope. For immuno

staining the phospho-S6 antibody has been previously described  [58] and the phospho-Akt

commercially provided (Cell signaling 4054). The cell size calculation have been performed as

previously described [59] and correspond to a set of experiments that spanned a two-year

period. It represent too many replicates, so that it was not possible to make them at the same

time. Therefore, for the graphs of cell size measurement (Figure 2M, 5M and 6G), values are

reused when they correspond to the same genotype and conditions. This allows a direct

comparison between the experiments.

## Statistical analysis

Statistical analyses were performed with R version 3.4.4, scripts are available on request. Significance for the statistical tests was coded in the following way based on the p-values: ***: $0 < p < 0.001$; **: $0.001 < p < 0.01$; *: $0.01 < p < 0.05$. P-values were corrected for multiple testing by a Holm-Bonferroni method [81]. Clone sizes were analyzed with a mixed-effect linear model on the logarithm of cell area, considering the treatment (Genotype and Sucrose conditions) as a fixed effect and Series/Larva as random effects (Figures 1, 5, and 7, Supplementary Table S1). The reported effects (and the corresponding P-values) were obtained from the difference between the (log) area of marked clonal cells and that of control surrounding cells from the same treatment, by setting the appropriate contrast with the "multcomp" package [82], according to the pattern: $E_{A,B} = \log(M_A) - \log(W_A) - [\log(M_B) - \log(W_B)]$, where $E_{A,B}$ is the difference between treatments (genotype and sucrose levels) A and B, $M_A$ and $M_B$ standing for the area of marked cells, and $W_A$, $W_B$ for the area of control cells in those treatments. This is equivalent to testing whether marked/control cell area ratios differ between treatments. PS6+ clone frequencies were treated as binomial measurements in a mixed-effect generalized linear model "lme4" package [83], featuring Genotype as a fixed effect, and Series/Larva as random effects. Both datasets of pupal weights were analyzed independently with linear models including Sex, Genotype, and Sucrose level effects and all their interaction terms (Figure 3A-B and Supplementary Table S3 for PTEN knockdown and Rheb overexpression; and Figure 6B and Supplementary Table S4 for *FASN$^{1-2}$* mutants). TAG, Protein, Glycogen, and Threalose concentrations were also analyzed with linear models involving Genotype, Sucrose level, and their interactions as fixed effects (Fig3 and Supplementary Table S3).

We wish to thank D Petit for preparing the fly media, H Stocker for fly stocks, A Teleman for the phospho-S6 antibody, M Gettings for editing the manuscript, and the NIG and VDRC stock centers for RNAi fly strains.

**REFERENCES**

1. Laplante M, Sabatini DM. mTOR signaling in growth control and disease. Cell. 2012;149(2):274-93. PubMed PMID: 22500797.

2. Howell JJ, Ricoult SJ, Ben-Sahra I, Manning BD. A growing role for mTOR in promoting anabolic metabolism. Biochem Soc Trans. 2013;41(4):906-12. PubMed PMID: 23863154.

3. Lamming DW, Sabatini DM. A Central role for mTOR in lipid homeostasis. Cell Metab. 2013;18(4):465-9. PubMed PMID: 23973332.

4. Caron A, Richard D, Laplante M. The Roles of mTOR Complexes in Lipid Metabolism. Annu Rev Nutr. 2015;35:321-48. PubMed PMID: 26185979.

5. Shimobayashi M, Hall MN. Making new contacts: the mTOR network in metabolism and signalling crosstalk. Nat Rev Mol Cell Biol. 2014;15(3):155-62. PubMed PMID: 24556838.

6. Saxton RA, Sabatini DM. mTOR Signaling in Growth, Metabolism, and Disease. Cell. 2017;168(6):960-76. doi: 10.1016/j.cell.2017.02.004. PubMed PMID: 28283069; PubMed Central PMCID: PMCPMC5394987.

7. Mossmann D, Park S, Hall MN. mTOR signalling and cellular metabolism are mutual determinants in cancer. Nat Rev Cancer. 2018;18(12):744-57. Epub 2018/11/15. doi: 10.1038/s41568-018-0074-8. PubMed PMID: 30425336.

8. Kim DH, Sarbassov DD, Ali SM, King JE, Latek RR, Erdjument-Bromage H, et al. mTOR interacts with raptor to form a nutrient-sensitive complex that signals to the cell growth machinery. Cell. 2002;110(2):163-75. PubMed PMID: 12150925.

9. Sarbassov DD, Guertin DA, Ali SM, Sabatini DM. Phosphorylation and regulation of Akt/PKB by the rictor-mTOR complex. Science. 2005;307(5712):1098-101. PubMed PMID: 15718470.

10. Ugur B, Chen K, Bellen HJ. Drosophila tools and assays for the study of human diseases.

Dis Model Mech. 2016;9(3):235-44. doi: 10.1242/dmm.023762. PubMed PMID: 26935102;

PubMed Central PMCID: PMCPMC4833332.

11. Montagne J, Radimerski T, Thomas G. Insulin signaling: lessons from the Drosophila

tuberous sclerosis complex, a tumor suppressor. Sci STKE. 2001;2001(105):PE36.

PubMed PMID: 11675514.

12. Padmanabha D, Baker KD. Drosophila gains traction as a repurposed tool to investigate

metabolism.    Trends    Endocrinol    Metab.    2014;25(10):518-27.    doi:

10.1016/j.tem.2014.03.011. PubMed PMID: 24768030.

13. Antikainen H, Driscoll M, Haspel G, Dobrowolski R. TOR-mediated regulation of

metabolism in aging. Aging Cell. 2017;16(6):1219-33. doi: 10.1111/acel.12689. PubMed

PMID: 28971552; PubMed Central PMCID: PMCPMC5676073.

14. Wangler MF, Hu Y, Shulman JM. Drosophila and genome-wide association studies: a

review and resource for the functional dissection of human complex traits. Dis Model Mech.

2017;10(2):77-88. doi: 10.1242/dmm.027680. PubMed PMID: 28151408; PubMed Central

PMCID: PMCPMC5312009.

15. Lehmann M. Endocrine and physiological regulation of neutral fat storage in Drosophila.

Mol Cell Endocrinol. 2018;461:165-77. doi: 10.1016/j.mce.2017.09.008. PubMed PMID:

28893568; PubMed Central PMCID: PMCPMC5756521.

16. Ma XM, Blenis J. Molecular mechanisms of mTOR-mediated translational control. Nat Rev

Mol Cell Biol. 2009;10(5):307-18. PubMed PMID: 19339977.

17. Goberdhan DC, Ogmundsdottir MH, Kazi S, Reynolds B, Visvalingam SM, Wilson C, et al.

Amino acid sensing and mTOR regulation: inside or out? Biochem Soc Trans. 2009;37(Pt

1):248-52. PubMed PMID: 19143641.

18. Groenewoud MJ, Zwartkruis FJ. Rheb and mammalian target of rapamycin in mitochondrial

homoeostasis. Open Biol. 2013;3(12):130185. PubMed PMID: 24352740.

19. Dibble CC, Manning BD. Signal integration by mTORC1 coordinates nutrient input with

biosynthetic output. Nat Cell Biol. 2013;15(6):555-64. PubMed PMID: 23728461.

20. Montagne J. A Wacky Bridge to mTORC1 Dimerization. Dev Cell. 2016;36(2):129-30. PubMed PMID: 26812011.

21. Yang H, Jiang X, Li B, Yang HJ, Miller M, Yang A, et al. Mechanisms of mTORC1 activation by RHEB and inhibition by PRAS40. Nature. 2017;552(7685):368-73. Epub 2017/12/14. doi: 10.1038/nature25023. PubMed PMID: 29236692; PubMed Central PMCID: PMCPMC5750076.

22. Montagne J, Thomas G. S6K integrates nutrient and mitogen signals to control cell growth. . In: Hall M, Raff, M., Thomas, G., editor. Cell growth: control of cell size.
 ed: Cold Spring Harbor Press; 2004. p. 265-98.

23. Magnuson B, Ekim B, Fingar DC. Regulation and function of ribosomal protein S6 kinase (S6K) within mTOR signalling networks. Biochem J. 2012;441(1):1-21. PubMed PMID: 22168436.

24. Radimerski T, Montagne J, Hemmings-Mieszczak M, Thomas G. Lethality of Drosophila lacking TSC tumor suppressor function rescued by reducing dS6K signaling. Genes Dev. 2002;16(20):2627-32. PubMed PMID: 12381661.

25. Garami A, Zwartkruis FJ, Nobukuni T, Joaquin M, Roccio M, Stocker H, et al. Insulin activation of Rheb, a mediator of mTOR/S6K/4E-BP signaling, is inhibited by TSC1 and 2. Mol Cell. 2003;11(6):1457-66. PubMed PMID: 12820960.

26. Inoki K, Li Y, Xu T, Guan KL. Rheb GTPase is a direct target of TSC2 GAP activity and regulates mTOR signaling. Genes Dev. 2003;17(15):1829-34. doi: 10.1101/gad.1110003. PubMed PMID: 12869586; PubMed Central PMCID: PMCPMC196227.

27. Dibble CC, Elis W, Menon S, Qin W, Klekota J, Asara JM, et al. TBC1D7 is a third subunit of the TSC1-TSC2 complex upstream of mTORC1. Mol Cell. 2012;47(4):535-46. doi: 10.1016/j.molcel.2012.06.009. PubMed PMID: 22795129; PubMed Central PMCID: PMCPMC3693578.

28. Engelman JA, Luo J, Cantley LC. The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism. Nat Rev Genet. 2006;7(8):606-19. doi: 10.1038/nrg1879. PubMed PMID: 16847462.

29. Haeusler RA, McGraw TE, Accili D. Biochemical and cellular properties of insulin receptor

signalling. Nat Rev Mol Cell Biol. 2018;19(1):31-44. doi: 10.1038/nrm.2017.89. PubMed

PMID: 28974775; PubMed Central PMCID: PMCPMC5894887.

30. Cully M, You H, Levine AJ, Mak TW. Beyond PTEN mutations: the PI3K pathway as an

integrator of multiple inputs during tumorigenesis. Nat Rev Cancer. 2006;6(3):184-92. doi:

10.1038/nrc1819. PubMed PMID: 16453012.

31. Lien EC, Dibble CC, Toker A. PI3K signaling in cancer: beyond AKT. Curr Opin Cell Biol.

2017;45:62-71. doi: 10.1016/j.ceb.2017.02.007. PubMed PMID: 28343126; PubMed Central

PMCID: PMCPMC5482768.

32. Duvel K, Yecies JL, Menon S, Raman P, Lipovsky AI, Souza AL, et al. Activation of a

metabolic gene regulatory network downstream of mTOR complex 1. Mol Cell.

2010;39(2):171-83. PubMed PMID: 20670887.

33. Polak P, Cybulski N, Feige JN, Auwerx J, Ruegg MA, Hall MN. Adipose-specific knockout

of raptor results in lean mice with enhanced mitochondrial respiration. Cell Metab.

2008;8(5):399-410. doi: 10.1016/j.cmet.2008.09.003. PubMed PMID: 19046571.

34. Peterson TR, Sengupta SS, Harris TE, Carmack AE, Kang SA, Balderas E, et al. mTOR

complex 1 regulates lipin 1 localization to control the SREBP pathway. Cell.

2011;146(3):408-20. doi: 10.1016/j.cell.2011.06.034. PubMed PMID: 21816276; PubMed

Central PMCID: PMCPMC3336367.

35. Inoki K, Zhu T, Guan KL. TSC2 mediates cellular energy response to control cell growth

and survival. Cell. 2003;115(5):577-90. PubMed PMID: 14651849.

36. Choo AY, Kim SG, Vander Heiden MG, Mahoney SJ, Vu H, Yoon SO, et al. Glucose

addiction of TSC null cells is caused by failed mTORC1-dependent balancing of metabolic

demand with supply. Mol Cell. 2010;38(4):487-99. doi: 10.1016/j.molcel.2010.05.007.

PubMed PMID: 20513425; PubMed Central PMCID: PMCPMC2896794.

37. Csibi A, Lee G, Yoon SO, Tong H, Ilter D, Elia I, et al. The mTORC1/S6K1 pathway

regulates glutamine metabolism through the eIF4B-dependent control of c-Myc translation.

Curr Biol. 2014;24(19):2274-80. doi: 10.1016/j.cub.2014.08.007. PubMed PMID: 25220053; PubMed Central PMCID: PMCPMC4190129.

38. Csibi A, Fendt SM, Li C, Poulogiannis G, Choo AY, Chapski DJ, et al. The mTORC1 pathway stimulates glutamine metabolism and cell proliferation by repressing SIRT4. Cell. 2013;153(4):840-54. doi: 10.1016/j.cell.2013.04.023. PubMed PMID: 23663782; PubMed Central PMCID: PMCPMC3684628.

39. Hahn-Windgassen A, Nogueira V, Chen CC, Skeen JE, Sonenberg N, Hay N. Akt activates the mammalian target of rapamycin by regulating cellular ATP level and AMPK activity. J Biol Chem. 2005;280(37):32081-9. doi: 10.1074/jbc.M502876200. PubMed PMID: 16027121.

40. Robey RB, Hay N. Is Akt the "Warburg kinase"?-Akt-energy metabolism interactions and oncogenesis. Semin Cancer Biol. 2009;19(1):25-31. doi: 10.1016/j.semcancer.2008.11.010. PubMed PMID: 19130886; PubMed Central PMCID: PMCPMC2814453.

41. Jaldin-Fincati JR, Pavarotti M, Frendo-Cumbo S, Bilan PJ, Klip A. Update on GLUT4 Vesicle Traffic: A Cornerstone of Insulin Action. Trends Endocrinol Metab. 2017;28(8):597-611. doi: 10.1016/j.tem.2017.05.002. PubMed PMID: 28602209.

42. Gottlob K, Majewski N, Kennedy S, Kandel E, Robey RB, Hay N. Inhibition of early apoptotic events by Akt/PKB is dependent on the first committed step of glycolysis and mitochondrial hexokinase. Genes Dev. 2001;15(11):1406-18. doi: 10.1101/gad.889901. PubMed PMID: 11390360; PubMed Central PMCID: PMCPMC312709.

43. Houddane A, Bultot L, Novellasdemunt L, Johanns M, Gueuning MA, Vertommen D, et al. Role of Akt/PKB and PFKFB isoenzymes in the control of glycolysis, cell proliferation and protein synthesis in mitogen-stimulated thymocytes. Cell Signal. 2017;34:23-37. doi: 10.1016/j.cellsig.2017.02.019. PubMed PMID: 28235572.

44. Nakae J, Kitamura T, Silver DL, Accili D. The forkhead transcription factor Foxo1 (Fkhr) confers insulin sensitivity onto glucose-6-phosphatase expression. J Clin Invest. 2001;108(9):1359-67. doi: 10.1172/JCI12876. PubMed PMID: 11696581; PubMed Central PMCID: PMCPMC209440.

45. McManus EJ, Sakamoto K, Armit LJ, Ronaldson L, Shpiro N, Marquez R, et al. Role that

phosphorylation of GSK3 plays in insulin and Wnt signalling defined by knockin analysis.

EMBO J. 2005;24(8):1571-83. doi: 10.1038/sj.emboj.7600633. PubMed PMID: 15791206;

PubMed Central PMCID: PMCPMC1142569.

46. Majewski N, Nogueira V, Bhaskar P, Coy PE, Skeen JE, Gottlob K, et al. Hexokinase-

mitochondria interaction mediated by Akt is required to inhibit apoptosis in the presence or

absence of Bax and Bak. Mol Cell. 2004;16(5):819-30. doi: 10.1016/j.molcel.2004.11.014.

PubMed PMID: 15574336.

47. Hagiwara A, Cornu M, Cybulski N, Polak P, Betz C, Trapani F, et al. Hepatic mTORC2

activates glycolysis and lipogenesis through Akt, glucokinase, and SREBP1c. Cell Metab.

2012;15(5):725-38. doi: 10.1016/j.cmet.2012.03.015. PubMed PMID: 22521878.

48. Yuan M, Pino E, Wu L, Kacergis M, Soukas AA. Identification of Akt-independent regulation

of hepatic lipogenesis by mammalian target of rapamycin (mTOR) complex 2. J Biol Chem.

2012;287(35):29579-88. doi: 10.1074/jbc.M112.386854. PubMed PMID: 22773877;

PubMed Central PMCID: PMCPMC3436168.

49. Radimerski T, Montagne J, Rintelen F, Stocker H, van der Kaay J, Downes CP, et al. dS6K-

regulated cell growth is dPKB/dPI(3)K-independent, but requires dPDK1. Nat Cell Biol.

2002;4(3):251-5. PubMed PMID: 11862217.

50. Dong J, Pan D. Tsc2 is not a critical target of Akt during normal Drosophila development.

Genes Dev. 2004;18(20):2479-84. PubMed PMID: 15466161.

51. Montagne J, Lecerf C, Parvy JP, Bennion JM, Radimerski T, Ruhf ML, et al. The nuclear

receptor DHR3 modulates dS6 kinase-dependent growth in Drosophila. PLoS Genet.

2010;6:e1000937. PubMed PMID: 20463884.

52. Pallares-Cartes C, Cakan-Akdogan G, Teleman AA. Tissue-specific coupling between

insulin/IGF and TORC1 signaling via PRAS40 in Drosophila. Dev Cell. 2012;22(1):172-82.

doi: 10.1016/j.devcel.2011.10.029. PubMed PMID: 22264732.

53. Stocker H, Radimerski T, Schindelholz B, Wittwer F, Belawat P, Daram P, et al. Rheb is an

essential regulator of S6K in controlling cell growth in Drosophila. Nat Cell Biol.

2003;5(6):559-65. PubMed PMID: 12766775.

54. Oldham S, Montagne J, Radimerski T, Thomas G, Hafen E. Genetic and biochemical

characterization of dTOR, the Drosophila homolog of the target of rapamycin. Genes Dev.

2000;14(21):2689-94. PubMed PMID: 11069885.

55. Edgar BA, Orr-Weaver TL. Endoreplication cell cycles: more for less. Cell.

2001;105(3):297-306. PubMed PMID: 11348589.

56. Zhang H, Stallock JP, Ng JC, Reinhard C, Neufeld TP. Regulation of cellular growth by the

Drosophila target of rapamycin dTOR. Genes Dev. 2000;14(21):2712-24. PubMed PMID:

11069888.

57. Morata G, Ripoll P. Minutes: mutants of drosophila autonomously affecting cell division rate.

Dev Biol. 1975;42(2):211-21. PubMed PMID: 1116643.

58. Romero-Pozuelo J, Demetriades C, Schroeder P, Teleman AA. CycD/Cdk4 and

Discontinuities in Dpp Signaling Activate TORC1 in the Drosophila Wing Disc. Dev Cell.

2017;42(4):376-87 e5. doi: 10.1016/j.devcel.2017.07.019. PubMed PMID: 28829945.

59. Garrido D, Rubin T, Poidevin M, Maroni B, Le Rouzic A, Parvy JP, et al. Fatty Acid

Synthase Cooperates with Glyoxalase 1 to Protect against Sugar Toxicity. PLoS Genet.

2015;11(2):e1004995. PubMed PMID: 25692475.

60. Parvy JP, Napal L, Rubin T, Poidevin M, Perrin L, Wicker-Thomas C, et al. Drosophila

melanogaster Acetyl-CoA-carboxylase sustains a fatty acid-dependent remote signal to

waterproof the respiratory system. PLoS Genet. 2012;8(8):e1002925. PubMed PMID:

22956916.

61. Chung H, Loehlin DW, Dufour HD, Vaccarro K, Millar JG, Carroll SB. A single gene affects

both ecological divergence and mate choice in Drosophila. Science. 2014;343(6175):1148-

51. PubMed PMID: 24526311.

62. Wicker-Thomas C, Garrido D, Bontonou G, Napal L, Mazuras N, Denis B, et al. Flexible

origin of hydrocarbon/pheromone precursors in Drosophila melanogaster. J Lipid Res.

2015;56(11):2094-101. PubMed PMID: 26353752.

63. Parvy JP, Wang P, Garrido D, Maria A, Blais C, Poidevin M, et al. Forward and feedback

regulation of cyclic steroid production in Drosophila melanogaster. Development.

2014;141(20):3955-65. PubMed PMID: 25252945.

64. Montagne J, Stewart MJ, Stocker H, Hafen E, Kozma SC, Thomas G. Drosophila S6

kinase: a regulator of cell size. Science. 1999;285(5436):2126-9. PubMed PMID:

10497130.

65. Rulifson EJ, Kim SK, Nusse R. Ablation of insulin-producing neurons in flies: growth and

diabetic phenotypes. Science. 2002;296(5570):1118-20. PubMed PMID: 12004130.

66. Mattila J, Havula E, Suominen E, Teesalu M, Surakka I, Hynynen R, et al. Mondo-Mlx

Mediates Organismal Sugar Sensing through the Gli-Similar Transcription Factor

Sugarbabe. Cell Rep. 2015;13(2):350-64. PubMed PMID: 26440885.

67. Richards P, Ourabah S, Montagne J, Burnol AF, Postic C, Guilmeau S. MondoA/ChREBP:

The usual suspects of transcriptional glucose sensing; Implication in pathophysiology.

Metabolism. 2017;70:133-51. doi: 10.1016/j.metabol.2017.01.033. PubMed PMID:

28403938.

68. Sassu ED, McDermott JE, Keys BJ, Esmaeili M, Keene AC, Birnbaum MJ, et al.

Mio/dChREBP coordinately increases fat mass by regulating lipid synthesis and feeding

behavior in Drosophila. Biochem Biophys Res Commun. 2012;426(1):43-8. doi:

10.1016/j.bbrc.2012.08.028. PubMed PMID: 22910416; PubMed Central PMCID:

PMCPMC3445662.

69. Bruning U, Morales-Rodriguez F, Kalucka J, Goveia J, Taverna F, Queiroz KCS, et al.

Impairment of Angiogenesis by Fatty Acid Synthase Inhibition Involves mTOR Malonylation.

Cell Metab. 2018;28(6):866-80 e15. Epub 2018/08/28. doi: 10.1016/j.cmet.2018.07.019.

PubMed PMID: 30146486.

70. Caldwell PE, Walkiewicz M, Stern M. Ras activity in the Drosophila prothoracic gland
regulates body size and developmental rate via ecdysone release. Curr Biol.
2005;15(20):1785-95. PubMed PMID: 16182526.

71. Colombani J, Bianchini L, Layalle S, Pondeville E, Dauphin-Villemant C, Antoniewski C, et
al. Antagonistic actions of ecdysone and insulins determine final size in Drosophila.
Science. 2005;310(5748):667-70. PubMed PMID: 16179433.

72. Mirth C. Ecdysteroid control of metamorphosis in the differentiating adult leg structures of
Drosophila melanogaster. Dev Biol. 2005;278(1):163-74. PubMed PMID: 15649469.

73. Wipperman MF, Montrose DC, Gotto AM, Jr., Hajjar DP. Mammalian Target of Rapamycin:
A Metabolic Rheostat for Regulating Adipose Tissue Function and Cardiovascular Health.
Am J Pathol. 2019;189(3):492-501. Epub 2019/02/26. doi: 10.1016/j.ajpath.2018.11.013.
PubMed PMID: 30803496.

74. Harachi M, Masui K, Okamura Y, Tsukui R, Mischel PS, Shibata N. mTOR Complexes as a
Nutrient Sensor for Driving Cancer Progression. Int J Mol Sci. 2018;19(10). Epub
2018/10/24. doi: 10.3390/ijms19103267. PubMed PMID: 30347859.

75. Tian T, Li X, Zhang J. mTOR Signaling in Cancer and mTOR Inhibitors in Solid Tumor
Targeting Therapy. Int J Mol Sci. 2019;20(3). Epub 2019/02/14. doi: 10.3390/ijms20030755.
PubMed PMID: 30754640; PubMed Central PMCID: PMCPMC6387042.

76. Henske EP, Jozwiak S, Kingswood JC, Sampson JR, Thiele EA. Tuberous sclerosis
complex. Nat Rev Dis Primers. 2016;2:16035. doi: 10.1038/nrdp.2016.35. PubMed PMID:
27226234.

77. Guri Y, Colombi M, Dazert E, Hindupur SK, Roszik J, Moes S, et al. mTORC2 Promotes
Tumorigenesis via Lipid Synthesis. Cancer Cell. 2017;32(6):807-23 e12. doi:
10.1016/j.ccell.2017.11.011. PubMed PMID: 29232555.

78. Dietzl G, Chen D, Schnorrer F, Su KC, Barinova Y, Fellner M, et al. A genome-wide
transgenic RNAi library for conditional gene inactivation in Drosophila. Nature.
2007;448(7150):151-6. PubMed PMID: 17625558.

79. Bohni R, Riesgo-Escovar J, Oldham S, Brogiolo W, Stocker H, Andruss BF, et al. Autonomous control of cell and organ size by CHICO, a Drosophila homolog of vertebrate IRS1-4. Cell. 1999;97(7):865-75. PubMed PMID: 10399915.

80. Lee T, Luo L. Mosaic analysis with a repressible cell marker (MARCM) for Drosophila neural development. Trends Neurosci. 2001;24(5):251-4. PubMed PMID: 11311363.

81. Holm S. A simple sequentially rejective multiple test procedure. Scand J Statist. 1079;6(65-70).

82. Hothorn T, Bretz F, Westfall P. Simultaneous inference in general parametric models. Biom J. 2008;50(3):346-63. Epub 2008/05/16. doi: 10.1002/bimj.200810425. PubMed PMID: 18481363.

83. Bates D, Mächler M, Bolker BM, Walker S. Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software. 2015;67(1):1-48. doi: 10.18637/jss.v067.i01.

**FIGURE LEGENDS**

**Figure 1: mTORC1 and mTORC2 dependent growth in FB cells.** (**A-L**) MARCM clones labeled by GFP (green) in the FB of L3 larvae. Nuclei were labeled with DAPI (silver) and membranes with phalloidin (red). Control (A), $Rheb^+$ (B), $PTEN^{-/-}$ (C) and $PTEN^{-/-};Rheb^+$ (D) clones were generated in a wild type background. $Rheb^+$ (E), $PTEN^{-/-}$ (F), $mTOR^{2L1}$ (G) $mTOR^{2L1}, Rheb^+$ (H) $mTOR^{2L1},PTEN^{-/-}$ (I), $mTOR^{\Delta P}$ (J), $mTOR^{\Delta P},Rheb^+$ (K) and $mTOR^{\Delta P},PTEN^{-/-}$ (L) clones were generated in a *Minute* (*M*) background. Scale bars: 50μm. (**M**) Relative size of control (Co), $Rheb^+$, $PTEN^{-/-}$, and $PTEN^{-/-};Rheb^+$ clonal cells generated in a wild type background.

**Figure 2: mTORC1 and mTORC2 activity in FB cells.** (**A-J**) MARCM clones labeled by GFP (green) in the FB of L3 larvae. Clones were generated in a wild type (A,D,E,H,I,J) or a *Minute* (B,C,F,G) background and nuclei were labeled with DAPI (silver). FB tissues with $PTEN^{-/-}$ (A),

*mTOR^ΔP* (B), *mTOR^2L1* (C) and *FASN^1-2* (J) clones were stained with a phospho-AKT antibody.

FB tissues with control (D), *mTOR^ΔP* (E), *mTOR^2L1* (F) *Rheb^+* (G), *PTEN^-/-* (H) and *FASN^1-2* (I)

clones were stained with a phospho-S6 antibody. Scale bars: 50µm. (**K**) Percentage of P-S6

positive clones with respect to the total number of MARCM clones for control, *FASN^1-2*, *PTEN^-/-*,

*Rheb^+*, *mTOR^2L1* and *mTOR^ΔP* genotypes.

**Figure 3: Enhanced mTORC1 or mTORC2 signaling affects larval metabolism.** (**A-B**) Body

weight of female (A) and male (B) prepupae formed from larvae fed either a standard (0%) or a

20%-SSD (20%) as from the L2/L3 transition. (**C-F**) Measurement of total protein (C), TAG (D),

glycogen (E) and trehalose (F) levels in prepupae fed either a standard or a 20%-SSD.

Prepupae used in these measurements were the F1 progeny from *da-gal4* virgin females mated

to either control (Co), *EP(UAS)-Rheb* (*Rheb^++*) or *UAS-PTEN-RNAi* (*PTEN-Ri*) males.

**Figure 4: Glycolysis knockdown in whole organisms.** (**A**) Scheme of basal metabolism.

Glucose and trehalose enter glycolysis as glucose-6P, whereas fructose follows a distinct

pathway to triose-P. Enzymes investigated in the present study are marked in red. (**B**)

Phenotype of ubiquitous RNAi knockdown of PFK1, PK, LDH and PDH. Flies were left to lay

eggs overnight either at 29°C (column 0h) or at 19°C and transferred to 29°C the day after

(column 24h); then development proceeded at 29°C (i.e. the temperature that inactivates

Gal80). (**C**) Western-blot analysis of total (top) or phosphorylated (mid) dS6K (left) or Akt (right)

proteins; tubulin (bottom) was used as a loading control. Protein extracts were prepared with

late L2 control larvae (Co) or L2 larvae expressing RNAi against the indicated metabolic

enzymes. (**E-F**) Survival at 29°C of male (top) and female (bottom) control flies or flies

expressing RNAi against the indicated metabolic enzymes as from adult eclosion.

**Figure 5: Cell-autonomous requirement of glycolytic enzymes for mTOR-dependent overgrowth.** (**A-G**) MARCM clones labeled by GFP (green) in the FB of L3 larvae. Nuclei were labeled with DAPI (silver) and membranes with phalloidin (red). Genotypes of MARCM clones are: *PFK1-RNAi* (A), *PK-RNAi* (B), *LDH-RNAi* (C), *PDH-RNAi* (D), *Rheb$^+$,PFK1-RNAi* (E), *Rheb$^+$,PK-RNAi* (F), *Rheb$^+$,LDH-RNAi* (G), *Rheb$^+$,PDH-RNAi* (H), *PTEN$^{-/-}$;PFK1-RNAi* (I), *PTEN$^{-/-}$;PK-RNAi* (J), *PTEN$^{-/-}$;PDH-RNAi* (K) and *PTEN$^{-/-}$;PDH-RNAi* (L). Scale bars: 50µm. (**M**) Relative size of clonal cells corresponding to the clones shown in A-L, and in Figure 2A for control (Co).

**Figure 6: *FASN$^{1-2}$* mutation affects developmental growth and mTOR signaling.** (**A**) Developmental duration from egg laying to metamorphosis onset of *w$^{1118}$* control (Co) and *FASN$^{1-2}$* (FASN) larvae fed either a beySD (0%) or a 10% sucrose-supplemented-beySD as from the L2/L3 transition (10%); n: total number of larvae collected for each condition. (**B**) Prepupal weight of females (left) and males (right) as listed in 4A; the numbers of weighted prepupae are indicated above each sample. (**C**) Western-blot analysis of (from top to bottom) total dS6K, phosphorylated dS6K, total Akt, phosphorylated Akt and total tubulin as a loading control. Protein extracts were prepared from feeding L3 larvae prior to the wandering stage as listed in 4A. For each condition, at least 30 larvae were used to prepare protein extracts.

**Figure 7: Cell-autonomous requirement of FASN activity for mTOR-dependent overgrowth.** (**A-L**) MARCM clones labeled by GFP (green) in the FB of L3 larvae fed either a standard (A-C, G-I) or a 20%-SSD (D-E, J-L). Nuclei were labeled with DAPI (silver) and membranes with phalloidin (red). Genotypes of MARCM clones are: *Rheb$^+$* (A,D), *PTEN$^{-/-}$* (B,E) *PTEN$^{-/-}$,Rheb$^+$* (C,F), *FASN$^{1-2}$;Rheb$^+$* (G,J) *FASN$^{1-2}$,PTEN$^{-/-}$* (H,K) and the *FASN$^{1-2}$,PTEN$^{-/-}$;Rheb$^+$* (I,L). Scale bars: 50µm. (**M**) Relative size of clonal cells corresponding to the clones shown in A-L and in supplementary Fig S1 for *FASN$^{1-2}$* and Figure 1A for control (Co).

**SUPPORTING INFORMATIONS**

**S1 Table. Statistical tests corresponding to figures 1M (A), 5M (B), and 7M (C).** The model tests for the difference is the log ratio of *GFP⁺/GFP⁻surrounding* cells between pairs of genotypes (Co = control), accounting for larvae and series random effects (see the Methods section). P-values were corrected for multiple testing (Holm-Bonferroni method) and quantify the risk of rejecting the true null hypothesis at the table level.

**S2 Table. Statistical test of the difference in frequency between MARCM clones to control.** The model is a mixed effect, generalized linear model considering frequencies as binomial data and genotype as a fixed effect, while larvae / series were random effects. P-values were adjusted for multiple testing by a Holm-Bonferroni correction.

**S3 Table. Statistical tests corresponding to Figure 3.** Differences in male (A) or female (B) weight, in protein (C), in TAG (D), in glycogen (E) and in trehalose (F) between control (Co) and prepupae that ubiquitously express either Rheb (*Rheb⁺⁺*) or an RNAi to PTEN (*PTENi*). The column at the right indicates P-values corrected for multiple testing (Holm-Bonferoni correction).

**S4 Table. Statistical tests corresponding to Figure 6B.** Prepupal weight differences in males and females (Estimate and standard error, in mg) between control (*w⁻*) and *FASN[1-2]* animals fed either a standard (0%) or a sucrose enriched diet (10%). The **Pr** column (right) indicates P-values corrected for multiple testing (Holm-Bonferoni correction).

**S1 Fig. Effect of dietary sugar on *FASN^1-2* mutant cells.** *FASN^1-2* MARCM clones labeled by

GFP (green) in the FB of L3 larvae fed either a standard (A) or a 20%-SSD (B). Nuclei are

labeled with DAPI (silver) and membranes by phalloidin (red). Scale bars: 50μm.

**S2 Fig. Knockdown efficacy of the RNAi to the glycolytic enzymes.** (**A**) mRNA expression

in control larvae (blue bars, +/- standard error) was adjusted to 100. mRNA levels after RNAi

knockdown (red bars, +/- standard error) was calculated as a percentage of corresponding

expression in control larvae. (**B**) Oligonucleotides used in RT-Q-PCR for each glycolytic

enzyme.

Figure 1

Figure 2

Figure 3

Figure 5

A

B

C

Figure 6

Figure 7

Figure 4