Diverse phenotypic responses and phosphate content in foxtail millet genotypes under greenhouse and field conditions

S. Antony Ceasar[1, 2,*#], M. Ramakrishnan[1, 3*], K. K. Vinod[4], G. Victor Roch[1], Hari D. Upadhyaya[5], Alison Baker[2, #], S. Ignacimuthu[1,]

Affiliations

1. Division of Plant Biotechnology, Entomology Research Institute, Loyola College, Chennai, India

2. Centre for Plant Sciences, School of Molecular and Cellular Biology, Faculty of Biological Sciences, University of Leeds, Leeds, UK

3. The State Key Laboratory of Subtropical Silviculture, Zhejiang A&F University, Hangzhou, Zhejiang, China

4. ICAR-Indian Agricultural Research Institute, Rice Breeding and Genetics Research Centre, Aduthurai, Tamil Nadu, India

5. International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru, Telangana, India and Center of Excellence for Advanced Materials Research, King Abdulaziz University Jeddah 21589, Saudi Arabia

***These authors contributed equally to this work**

Author Emails

MR: ramakrishnan11deri004@loyolacollege.edu,

KKV: kkvinod@iari.res.in,

GVR: rochyesj@gmail.com,

HDU: harideo.upadhyaya@gmail.com

SI: eriloyola@hotmail.com

[#]**Correspondence**

Alison Baker	**S. Antony Ceasar**
Centre for Plant Sciences,	Division of Plant Biotechnology,
School of Molecular and Cellular Biology,	Entomology Research Institute,
Faculty of Biological Sciences,	Loyola College,
University of Leeds, Leeds, UK	Chennai, India
Phone: +44 (0)113 3433045	Phone: +91-44-2817 4644
Email: a.baker@leeds.ac.uk	**Email**: antony_sm2003@yahoo.co.in

Running title: Phenotype responses of foxtail millet under low phosphate

Date of submission: April 12, 2019

Number of tables: 4

Number of figures: 7

Word count (start of the introduction to the end of the acknowledgements): 6631

Color figures online and in print: All (7) (Both online and print)

Supplementary data: Figures = 4; Tables = 3

Highlight

Enormous phenotypic and phosphate content variation of foxtail millet under low-phosphate supply in greenhouse and natural field conditions identifies genotypic plasticity for future breeding for improved P use efficiency.

Abstract

Phosphorous (P) is an important macronutrient for the growth of all agricultural crops. This study reports phenotype analysis for P responses in field (two different seasons, monsoon and summer) and greenhouse, using 54 genotypes of foxtail millet (*Setaria italica*) under P-fertilized (P+) and unfertilized (P-) conditions. Variation was seen for plant height, leaf number and length, tillering ability and seed yield traits. Genotypes ISe 1234 and ISe 1541 were P+ responders, and the genotypes ISe 1181, ISe 1655, ISe 783 and ISe 1892 tend more towards low P tolerance for total seed yield. Genotypes that performed well under P-conditions were almost as productive as genotypes that performed well under P+ conditions suggesting some genotypes are well adapted to nutrient-poor soils. In the greenhouse, significant variation was seen for root hair density and root hair number and for fresh and dry weights of shoot and root under P- stress. However, there was not much difference in the shoot and root total P and inorganic phosphate (Pi) levels of five selected high and low responding genotypes. In the root and leaf tissues, total P and Pi contents of five high responding genotypes were higher than the five low responding genotypes.

Keywords: foxtail millet; field phenotyping; low phosphorous stress; phenotyping; root hair density; root hair length

Abbreviations:

P - Phosphorous

Pi - Inorganic phosphate

PUE - P use efficiency

PAE - Pi acquisition efficiency

QTL - Quantitative trait loci

PUtE - Pi utilization efficiency

RCBD - Randomized complete block design

SDW - Shoot dry weight

RDW - Root dry weight

SL - Shoot length

RL - Root length

RHD - Root hair density

RHL - Root hair length

PH - Plant height

NT - Number of tillers

NPT – Number of productive tillers

NL - Leaf number

LL - Leaf length

LF - Flower length

NC - Cluster number

SPC - Seeds per cluster

25SW - 25 seed weight

TSE - Total seed yield

REML - Restricted maximum likelihood

Introduction

Phosphorous (P) plays essential functions in living systems and it is an irreplaceable element for crop production. Plants uptake P from the soil as inorganic phosphate (Pi) and the process is affected by external factors such as soil pH, microbial activity, cationic abundance and presence of mycorrhizal fungi (*Schachtman et al.,* 1998). Although soils are rich in P, paradoxically it remains highly limiting for plant growth due to chemical fixation in organic or in inorganic form. Globally, 5.7 billion hectares of land are affected by Pi-deficiency that significantly hampers crop production (Niu *et al.,* 2013). In continuous cropping systems, replacement of depleted nutrients is carried out through external application for fertilizers to sustain crop production. However, projections suggest reserves of rock phosphate, the prime source of inorganic phosphate fertilizers, may only last for the next 200 years (Cordell *et al.,* 2009; Sattari *et al.,* 2012). The low Pi use efficiency (PUE) of the modern cultivars (around 20%), is a major problem in intensive cropping systems, where the input requirement of phosphatic fertilizers is high, leading to fixation in the soil and/or environmental leaching of the unutilised Pi in water bodies (Vinod & Heuer, 2012). Under circumstances of continuous cropping and Pi fixation, input reduction will exacerbate Pi starvation in crop plants (Vinod, 2015). To tackle this problem, there is a concurrent need for Pi input reduction and crop PUE improvement.

PUE in plants comprises Pi acquisition efficiency (PAE); the extent to which plants acquire Pi from the soil, and Pi utilization efficiency (PUtE); the efficient use of internal P resources. Since both the components are complementary, simultaneous improvement of PAE and PUtE is an important objective for sustainable crop production in order to achieve reduced reliance on external Pi supplementation (Veneklaas *et al.,* 2012; Wang *et al.,* 2010). Several approaches have been adopted with the goal of improving both the PAE and PUtE of crop plants. These include both transgenic and breeding based approaches (reviewed in Baker *et al.,* 2015; Lopez-Arredondo *et al.,* 2014). Extensive phenotyping studies have been performed to identify the genotypes with greater PUtE and to map the associated quantitative trait loci (QTL) for use in breeding programs (reviewed in Wiel *et al.,* 2015). Most of these studies focus on P starvation tolerance because improvement of P acquisition from soil and utilisation is the prime objective of breeding for PUE in crops. Under P starvation, P homeostasis is sustained through adaptive mechanisms such as improved nutrient foraging in the rhizosphere through root architectural manifestations that include root proliferation and deeper penetration (Lynch & Brown, 2001).

In *Arabidopsis* as well as several crops such as rice, maize, common bean, white lupin, tomato, black mustard and finger millet, root modifications such as primary root length reduction, improved branching, increased lateral root length, enhancement in number of lateral roots and root hair proliferation were reported as Pi starvation responses (Dinkelaker *et al.,* 1995; Carswell *et al.,* 1996; Borch *et al.,* 1999; Kim *et al.,* 2008; Lambers *et al.,* 2011; Jin *et al.,* 2012; Ramakrishnan *et al.*, 2017; Niu *et al.,* 2013). Additionally, roots play a major role in boosting rhizosphere biology by improving microfloral symbiosis and rhizochemical reactions through root exudates which aids P solubilization and thereby Pi uptake. Root level adaptation augments the innate ability of plants to uptake more Pi from the soil, thereby making it a potential trait targeting improvement in PAE (Lynch & Brown, 2008; Rouached *et al.,* 2010; Péret *et al.,* 2011). Aerial parts of cereal crops also show key phenotypic modifications in response to low P starvation. These include stunted shoot growth, dark green leaf and reduced yield in oat (Andersson *et al.,* 2003), increased primary root length and reduced photosynthesis in rice (He *et al.,* 2003; Wissuwa *et al.,* 2005), reduction in leaf and primary root growth and photosynthesis in maize (Mollier & Pellerin, 1999) and suppressed shoot growth in sorghum (Yoneyama *et al.,* 2007). Reduced shoot growth and lower seed yield were reported in foxtail millet genotype Maxima grown under low P in the greenhouse (Ceasar *et al.,* 2014).

Millets are small-seeded cereals in the family Poaceae, mainly cultivated and consumed by people in the tropical world. The millet grains are rich in essential nutrients like calcium, magnesium, zinc and iron and form a major source of nutrition and food security for millions in less developed nations (Ceasar & Ignacimuthu, 2009). Foxtail millet (*Setaria italica*) is widely cultivated in the semi-arid regions of Asia (India, China and Japan) and also in Southern Europe and has the longest history of cultivation among all the millets (Diao *&* Jia, 2017). Unlike the major cereals, breeding interventions in millets are limited especially using biotechnological tools. Foxtail millet is a genetically amenable model crop since it is a diploid and possesses a relatively small genome of ~515 Mb. Currently, the genome sequence information of foxtail millet varieties is available (Li & Brutnell, 2011; Bennetzen *et al.,* 2012; Zhang *et al.,* 2012). Foxtail millet is also closely related to bio-fuel grasses such as switchgrass and Napier grass (Muthamilarasan & Prasad, 2015). Together with its wild relative, green foxtail (*S. viridis*), both *Setaria* species are considered good models for nutrient management in other millets (Huang *et al.,* 2016; Ceasar 2018). There are no reports to date exploring the responses to P starvation in foxtail millet, except for a regional report

(Kalaghatagi *et al.,* 2000). We hypothesise that given similar cultural conditions, with
contrasting P levels, those foxtail millet genotypes that fare well under limited P nutrition
would be P starvation tolerant and those do well under high P nutrition would be fertilisation
responsive. This information would be of value for future breeding programmes. In this
study, we report the response of fifty-four genotypes of foxtail millet to P sufficiency (P+)
and starvation (P-) under both greenhouse and natural field conditions.

Materials and methods

Plant material

Seeds of fifty genotypes used in this study were obtained from the International Crops
Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru, Hyderabad, India. The
genotype 'Maxima1' (Acc.No: Bs 3875; Welsh Plant Breeding Station, Aberystwyth
University, UK) previously used in genetic studies on P response (Ceasar *et al.,* 2014; Ceasar
et al., 2017) and three local genotypes of Southern India (CO5, CO6 and CO7) procured from
Tamil Nadu Agricultural University (TNAU), Coimbatore, India) were also included in the
trial, and two of them, CO5 and CO7 were used as local checks for comparison. The details
of all the fifty-four genotypes are given in **Supplementary Table S1**. These genotypes were
evaluated under greenhouse and natural field conditions in two separate experiments
conducted in two different seasons. Both the experiments were conducted using a randomized
complete block design (RCBD) with two replications.

Field experimentation

The growth and yield studies were conducted under natural field conditions at Paiyur,
Tamil Nadu, India (12°25' N, 78°13' E) located at an elevation of 460 meters above sea level
and repeated for two years, in the summer season during 2015 (April to July) and in the
monsoon season in 2017 (August to November). The experimental station received an
average rainfall of ~105.17 mm in April to July 2015 and ~188.72 mm in August to
November 2017. This location has a history of traditional small millet cultivation for over
several hundred years. An unfertilized field that was left barren for several previous seasons
was chosen for the study. Prior to the experiment, soil samples were collected from around
the field, and the available P was found to be 5.5 mg/kg, which is normally considered as Pi-
deficient under natural field conditions (Gemenet *et al.,* 2016). Two treatments were
maintained, (i) in which no P fertilizer was applied (P-) and (ii) wherein P application was
done at double the recommended doses (P+). The crop was raised by sowing the seeds in

shallow furrows and plants were grown with a spacing of 30 cm between rows and 10 cm
between plants within a row. A block size of 11 x 1 meter was maintained with a spacing of
60 cm between the blocks. Each block consisted of 27 genotypes (25 tests and two checks
CO5 and CO7), with one check alternating every three test genotypes. The borders rows were
planted with check varieties. There were two such blocks per each P regime (P- and P+). Two
replications were maintained. Pi fertilization was done by applying 100g of diammonium
phosphate (DAP) only into each of the P+ plots. To compensate the extra N supplied through
DAP, 40 g of urea was applied in the P- plots such that the amount of N was the same for
both Pi treatments. K was also the same between the Pi treatments. The field was irrigated
once a week and the weeding done in every alternate week. The plants were allowed to grow
to maturity under normal recommended agronomic practices. Agro-morphological data were
collected on per plant basis at maturity on plant height (PH), tiller number (NT), productive
tiller number (NPT), leaf number (NL), leaf length (LL), panicle length (LF), cluster number
(NC)/panicle, 25 seed weight (25SW), seeds per cluster (SPC) and total seed yield
(TSE)/panicle.

Greenhouse experimentation

Greenhouse experiments were conducted at the Entomology Research Institute (ERI),
Loyola College, Chennai, Tamil Nadu, India. Seeds were sown in the 4 L pots (10 seeds/pot)
and allowed to germinate and grow in perlite medium (Astrra Chemicals, Chennai, Tamil
Nadu, India) supplied with a basal nutrient solution (Ceasar et al., 2014) containing 300 μM
of Pi (P+) or 10 μM of Pi (P-). For supplying different Pi regimens in nutrient solutions
different ratios of KH_2PO_4 and K_2SO_4 were used to maintain a constant concentration of
potassium. Three replicates were maintained for each of the Pi treatments. The whole
experiment was maintained in a greenhouse at 26ºC, 16 h light with 85% relative humidity
under well-lit and aerated conditions. The shoot dry weight (SDW) and root dry weight
(RDW) of 15-day old seedlings for each treatment were observed and combined into a single
trait, biomass to avoid reduced variance due to fractionation. Shoot length (SL), root length
(RL), root hair density (RHD) and root hair length (RHL) were determined at 28 days. For
the analysis of RHD and RHL, the seedlings were carefully removed from perlite, washed
twice with distilled water and blotted dry. Root hair measurements were performed according
to (Slabaugh *et al.,* 2011) with some modifications. About 5 cm from the primary root cap
was photographed using a Canon Coolpix digital camera under a Leica Stereo Microscope

(Wetzlar, Germany) and measured using ImageJ image processing and analysis software
v1.80_112 (Schneider *et al.,* 2012).

Analysis of phosphate content

The total P and Pi contents of the root and shoot tissues of the five most extreme
genotypes in each experiment were analysed according to Chiou *et al.,* (2006). Selection of
extreme genotypes was done independently on both P- and P+ treatments following the same
procedure described below.

Data analyses

Analysis of variance was conducted on the field observations for two seasons using
restricted maximum likelihood (REML) approach implemented in 'lme4' package in the R
statistical environment. The genotypes and season were treated as random factors to estimate
the unbiased predictors from the model. Those traits having non-significant genotypic effects
were dropped, and the predicted genotype effects of different traits were used for further
analyses. To identify the extreme 10% genotypes for the overall phenotypic performance, the
trait predictors were ranked in descending order individually for all the traits, so that the
highest value gets the rank of one. The cumulative ranking (rank sum) of each of the
genotype was computed by adding the ranks of that genotype for all the traits. For instance,
under P+, the genotype ISe1387 was ranked 1st for PH, NL, LF, NC and TSE, 2nd for LL, 3rd
for SPC and 37th for NPT making its rank sum as 47. Similarly, the rank sums were worked
out for the rest of the genotypes. A similar approach was implemented for greenhouse data.
From the rank sums, extreme five genotypes were picked having low and high-rank sums,
under each category of P- and P+ treatments. Four classes of extreme genotypes were
selected, namely, lowest rank sum (high performers) under P+, highest rank sum (low
performers) under P+, high and low performers under P-.

Genotype comparison for P response under field and greenhouse conditions

Field and greenhouse data was used to identify genotypic similarity in P response
both under P+ and P- conditions. Before clustering, the data were centred by subtracting the
trait means and scaled by dividing by the respective standard deviation. A popular
unsupervised centroid-based clustering known as k-means clustering (Hartigan & Wong,
1979) was performed by R using *kmeans* function. The clusters were graphically determined
by the package 'gplots'.

Cluster analysis based on plasticity

Response to P availability in plants was assumed to affect all the phenotypic traits observed. Therefore, the genotypic P response among different traits (plasticity) was worked out as the percentage difference under both P conditions over the mean P response. Hierarchical clustering was done on the plasticity matrix, using correlations as the distance measure and by performing multiscale bootstrap resampling (Shimodaira, 2004). The *p*-values computed for bootstrap events for each cluster was used for identifying significant clusters. There are two *p*-values computed, BP and AU, BP being the bootstrap probability and AU being the approximately unbiased *p*-value, a better measure of approximation (Shimodaira, 2002, 2004). The computations were performed using the 'pvclust' package (Suzuki & Shimodaira, 2006) in R using correlations and average agglomeration with 3000 bootstrap resampling. All the analyses were done using R Studio 1.1.463 running R version 3.5.1.

Results

Foxtail millet genotypes show differential response for Pi nutrition under natural field and greenhouse conditions

Analysis of variance revealed that significant genotypic variation existed for all the traits under both P- and P+ conditions except for 25 seed weight, which was dropped from further analyses (**Table 1**). The season and genotype x season component showed non-significant variance. The heritability calculated from the estimated genotypic variance obtained from the vector of genetic effects and the error variance obtained from the vector of residuals showed profound influence of genotypic variance to the total variance for all the traits in the population. Although these estimates are not accurate, they indicate the general trend of influence of different sources of variation to the heritable component of the trait (Gilmour *et al.,* 1995).

Based on the cumulative ranking, genotypes ISe 1387 and ISe 1687 were in the top 5 in both Pi fertilized and unfertilised field plots. Other genotypes that performed well under both P- and P+ conditions were CO7 (4[th], 7[th]), ISe132 (10[th], 2[nd]), ISe1851 (7[th], 3[rd]) and ISe 869 (5[th], 14[th]). Other top genotypes were ISe 663 (3[rd] under P-) and ISe 132 and ISe 907 (2[nd] and 4[th] under P+), (**Figure 1A**). Maxima1 was the poorest among all the genotypes under P-,

followed by ISe1335, ISe1234, ISe1037 and ISe1302. Maxima1, ISe1335, and ISe1234 were

also in the bottom 5 under P+ treatment.

The means for the field experimentation clearly illustrated the response of genotypes

for each trait analysed (**Table 2 and Supplementary Table S2**). Statistically significant

differences were seen for all the traits when the good and poor performing genotypes based

on the rank sums for the P+ plot were compared. In the case of P- conditions also,

conspicuous differences were recorded for all the traits. However, when the selection was

done for extreme five genotypes in both the directions in both the treatments, differences in

phenotypic performance was found reduced. The inflorescence images depicting the variation

in some of the genotypes with varied responses to P fertilization is given in **Figure 2**.

Greenhouse grown plants also showed a high degree of variation with respect to

growth on differing levels of Pi (**Supplementary Table S3**). Genotype ISe 1851 had the

lowest rank sum (high performer) under P-, this was followed by CO6, ISe 1338, ISe 1305

and ISe 1254. (**Figure 1B**). Genotypes ISe 1474 and ISe 748 had the highest (low performer)

rank sum under P-. Under high phosphate, the cumulative rank order was very different,

though CO6 was 2nd. Genotype ISe 748 performed poorly under P+ as well as P- (**Figure**

1B). Collectively, the high performing genotypes under P- conditions had significantly higher

biomass, SL, RL, RHD and RHL than low performers (**Table 3**.) The low performer ISe

1687, registered significantly higher RHD and RHL than other low performers under P-

condition. ISe 1687 was interesting because despite RHL and RHD that were comparable to

the high performers all other responses were poor, probably because the roots themselves

were very small. Under P+ conditions, the five low performers were comparable to the five

low performers under P- conditions for all parameters. Genotypes ISe748, ISe1254 and

ISe1687 were common to both treatments as low performers. ISe748, ISe1254 and ISe1736

grown under P+ treatment as well as under P- condition lacked root hairs (**Table 3** and

Figure 3). In contrast, the root hairs in the other genotypes (ISe 1851, ISe 1305, ISe 1209

and ISe 1387) shown in **Figure 3** were sparse or undiscernible under P+ conditions but were

increased in length and number in P-conditions. The high responding genotypes under P+

were different from the high responding genotypes under P- with the exception of the local

genotype CO6 which was ranked 2nd under both P- and P+ (**Figure 1B**). It attained 90 % of

the SL and 76 % of the biomass under the P- conditions as compared to P+. For the root traits

under P-, CO6 attains 69 % of the RL, 209 % of the RHD and 113 % of the RHL when

compared to P+ conditions.

ISe 1387 which was the highest ranking genotype under field conditions in both Pi regimes, ranked low when grown under greenhouse conditions (**Figure 1A, B**). Maxima1, which showed a poor field performance under both fertilised and unfertilised conditions, was the fourth best performer under P+ treatments, and mid-ranking under P- in the greenhouse screen. Similarly, CO6 which performed very well under both P+ and P- in the greenhouse experiments indicated a moderate response under field conditions (37^{th} under P- conditions and 24^{th} under P+ conditions). ISe 1851 performed well under both field as well as greenhouse assays and was the most consistent high-responding genotypes in the study. Other high performers were CO7 and ISe 869. Further, the genotype CO5 which was placed at the 13^{th} position in both P- and P+ soil, was placed at the first position under P+ and 6^{th} position under P- situations under greenhouse evaluation.

P contents of high-responding genotypes were higher than low performers under field and greenhouse conditions

The total P and Pi contents were assayed in the leaf and root tissues of each of the five high and low responding genotypes under P- and P+ conditions, grown in the greenhouse (**Figure 4**) as well as in the field (**Figure 5**). Good responders under the P+ condition in the glasshouse generally had much higher (1.5-3 fold higher) total leaf P and leaf Pi contents than low performers. Root total P and Pi was also higher although there was more variation and some overlap with some of the poor responders such as ISe748 and ISe 1387 (**Figure 4A and B**). The low responding genotype ISe 748 had the same level of Pi in root tissues as some of the high-responding genotypes (CO5 and ISe 869) but much less in the shoot tissues. This may suggest that ISe 748 had low ability in exporting Pi from the root to shoot tissues (compare **Figure 4A and B**). Under P-, the high responding genotypes maintained higher levels of both root and shoot P compared to poor responders (**Figure 4C and D**). As expected, in both high and low performers, more plant Pi is present under P+ conditions than under P- conditions (compare **Figure 4A and C and 4B and D**), but the good responders on P- maintained similar levels of tissue P to poor responders on P+ suggesting higher PAE (compare **Figure 4B and C**).

Under field conditions, the best performers had higher (>50 %) total and Pi content than the poor performers (**Figure 5A and B**). The levels of shoot total P and Pi were similar between the good responders under glasshouse and field conditions but root total P and Pi was higher in the best performers under field conditions compared to glasshouse (**Figure 4A and Figure 5A**). Most of the good responding genotypes grown under P- condition in the

field maintained similar levels of total P and Pi in shoots compared to poor responders (compare **Figure 5C and 5D**). The good responders on P- maintained similar levels of leaf Pi to poor responders on P+ but root total P and Pi was much lower, again pointing to effective export from root to shoot (**Figure 5B and C**). For example, the high responding genotype ISe 1687 had more than 100 µmoles/gm total P which was significantly greater than other high performers and all low performers under P- treatment.

Interrelations between P content among the extreme genotypes showed that shoot and root Pi contents were significantly correlated to total P in the respective tissues both under field and greenhouse screening (**Table 4**). Under greenhouse situations, Pi content of shoot tissues was also related to root Pi content but not to total root P content. Overall, under field situations, both the root Pi and shoot Pi were correlated as well as to the total P content in both the plant tissues. Conspicuously, there was good agreement between P content of the same genotypes grown under field and greenhouse conditions in both the root and shoot tissues. Analysis of the correlations between tissue-specific P content assayed under field and greenhouse, within different categories of genotypes falling within the four response classes indicated discernible deviation from the overall pattern. Shoot Pi of the good and poor responders under P- condition showed no apparent relation between field and greenhouse conditions, whereas the same under P+ conditions showed significant but negative relations. Negative correlations were also recorded between field and greenhouse-based assays of total shoot P as well as root Pi, under P+ regime. In poor performers, however, total root and shoot P showed a significant positive correlation between field and greenhouse data.

Genotypic plasticity for seed yield identifies low P tolerant genotypes based on P response

Genotypic plasticity, the ability of the plants to withstand the extreme variations in P availability, could identify three sets of genotypes such as low P tolerant, high P tolerant (P+ responders) and intermediates under both P+ and P- conditions. The genotypes ISe 1181, ISe 1655, ISe 783, ISe 1892 were low P tolerant based on seed yield (on the left side of **Figure 6**). Genotypes ISe 1234, ISe 1541, ISe 1563, ISe 1820, and ISe1888 were P+ responders and hence were identified as high P tolerant types (on the right side of **Figure 6**). It is of particular interest to understand which genotypes were most responsive to P fertilisation. Although plasticity over the mean performance was low (19.3%), the genotype ISe 1387 showed a particularly strong response to P application in both greenhouse and field

conditions, in terms of absolute phenotypic performance (**Figure 4 and 5**). But, this genotype was high P tolerant, not low P tolerant based on plasticity analysis (**Figure 6**).

Genotype clusters based on P level response in field and greenhouse experiments

The standardised phenotypic data-based analysis of 54 foxtail millet genotypes in both greenhouse and field conditions under P- and P+, revealed two to four clusters of genotypes by *k*-means clustering. The cluster pattern was displayed as a heatmap in **Supplementary Figures S1, S2, S3 and S4,** where yellow equates to a high value and red to a low one. Comparing performance in the field, some of these varieties showed that variabilities in agronomical traits patterned the grouping of foxtail millet genotypes in response to Pi conditions. There were four clusters of genotypes under field conditions in P- level (**Supplementary Figure S1**). The genotypes ISe 1181, ISe 375 and ISe 376 were low P responders for total seed yield, and these three genotypes were found grouped together into one cluster (**Supplementary Figure S1**). But, these three genotypes were grouped in separate clusters based on the bootstrap probability (BP) and the approximately unbiased (AU) *p*-value (**Figure 7**). The P+ responders ISe 1387, ISe 1687, ISe 663, ISe 869 and CO7, were clustered with genotypes which showed an intermediate response to the unfertilized soil (**Supplementary Figure S1**). The yield-related traits such as NPT, TSE and SPC showed higher variabilities while the remaining traits such as PH, LF, LL, NC and NL exhibited lesser variabilities in the analysis (**Supplementary Figure S1 and S2**). We observed similar variations in the analysis for genotypes grown in P-fertilized soil. The most notable example was the genotype ISe 1387 on the top line of the supplementary figure S2, and it was placed in a separate node. This genotype performed relatively well under both greenhouse and field conditions in both phosphate conditions (**Figure 1A, 5** and **Supplementary Figure S2**). The low performers Maxima1, ISe 1037 and ISe 1335 were grouped at the bottom of figure 7 with genotypes ISe 1541 and ISe 1234 which were P+ responders for total seed yield (**Figure 7**).

The traits measured under greenhouse conditions under P+ treatment (**Supplementary Figure S3**) showed that higher variability in biomass influenced the genotypic clustering pattern more than the remaining traits SL, RL, RHD and RHL that showed lesser variability. Under P- conditions (**Supplementary Figure S4**), however, the biomass and SL traits were the major traits that diversified genotypes than RHD and RHL. Three groups in the foxtail millet genotypes were identified under P+ condition, with the genotypes CO5, CO6, ISe 2 and Maxima forming a single cluster and are recognised as high

performers under P+ (**Supplementary Figure S3).** Most of the low performers were grouped

with genotypes that were intermediate in response to the Pi conditions. Within this cluster, we

could see all of the high performers (CO5, CO6, ISe 2 and Maxima1) under P+ which gave

us confidence in the analysis. Whereas, the traits measured under P- (**Supplementary Figure**

S4) created only two groups in the foxtail millet genotypes, and SL showed higher variability.

The RHD and RHL showed the same variability in both P- and P+ with good responders,

such as ISe 1851, CO6, ISe 1338, ISe 1305 and ISe 254, which were found grouped with

genotypes which showed above medium response under P- (**Figure 1B** and **Supplementary**

Figure S4). The genotype Maxima1 was clustered in a separate node and these genotypes

showed an intermediate response in P-.

Discussion

Evaluation of the P response of foxtail millet genotypes under field conditions is

important because little information on intrinsic P response of foxtail millet is available and

there is potential for translation in practical agriculture. Field situations are complicated by

environmental and edaphic factors and biotic influences of the co-existing flora and fauna,

however root system responses and rhizosphere environment are difficult to observe and

measure. Conversely, greenhouse studies allow evaluation under relatively controlled

conditions, and with closer observations, but what plants experience under natural

environments such as light, extreme ranges of temperature, nutrient availability and biotic

interactions are not recreated under greenhouse environment. Greenhouse based

measurements of root system parameters, plant biomass by dry weight and shoot length were

used to complement field data to have a comprehensive evaluation of the genotype

performance.

The agromorphologic data obtained showed high levels of variation for P response as

expected in the field trials. Although many genotypes gave consistent response under both the

seasons, a few showed variations due to environmental variations. This can well be attributed

to genotypic homeostasis that varies between genotype to genotype. Moreover, this is not

unexpected with the current panel of genotypes that are obtained from different countries of

origin (Upadhyaya *et al.,* 2008). This effect was more apparent with the line Maxima1 which

belongs to the race Maxima which prevails in eastern China, Georgia (Eurasia), Japan and

Korea. Since the variation imposed was largely due to P input, the remaining effects either

through genotypic or through genotype x environment interaction were assumed to be

uniform in both the experimental systems employed in this study. Therefore, the genotypes

were ranked for growth in terms of higher values of different traits measured. The rank sums were then used as the criteria for selection of extreme genotypes to divulge the contrast in response to P nutrition in foxtail millet genotypes. The ranking is a simple but efficient non-parametrical method in genotype selection (Kang, 1988; Huehn, 1996) but lacks statistical properties. Therefore, to ensure statistical significance between selected individuals, only those traits which showed statistical significance at 95% confidence level were used for ranking, a rank sum calculated and extreme 5% genotypes were selected based on the rank sum. Since the traits of measurement differed under both field and greenhouse screens, the rank sums were computed separately. This identified extreme genotypes under two systems and under two levels of P nutrition. Some of the genotypes such as ISe 1687, Maxima1 and ISe 1387 showed contrasting ranking patterns under field and greenhouse evaluation, which are to be considered as more unstable genotypes, wherein the pattern of P response in one system cannot be construed for the response under the other. Using this context, however, no stable genotype could be identified that was ranked at extreme under both field and greenhouse conditions. The exception to this was ISe 1851, which performed well under both P+ and P- conditions in the field and P- condition in the greenhouse.

P within the plant is translocated easily between the source and sink. The genotypes that possess an active sink of P are P use efficient. There was good agreement between plant P content under field and greenhouse conditions indicating that the P response pattern of the genotypes was consistent irrespective of the growing systems. However, the relationship did not hold for extreme genotypic groups, because of the high magnitude of specific P responses in these sets of genotypes. P uptake efficient genotypes can efficiently harness P under P+ conditions. Furthermore, good responders under P+ and P- conditions accumulated more P in the plant system than the poor responders. Those with high P in the plant system also have better utilisation efficiency if they can effectively trade P between different utilization points of growth and metabolism. Hence, in this experiment good responders under P- can be construed as low P tolerant genotypes, while good responders under P+ can be identified as P loving genotypes. Low P tolerant genotypes with good seed yield such as ISe 663 and ISe 869 are highly preferred for future breeding programmes in foxtail millet targeting P starvation tolerance. Further, the extensive genetic variation for P responses observed in this study, signal the possibility of genetic improvement of foxtail millet for P use efficiency as seen in other cereal crops such as rice (Wissuwa & Ae, 2001), wheat (Batten, 1986), barley (Gahoonia & Nielsen, 2004) and maize (Zhu *et al.,* 2005).

Under Pi insufficiency conditions plants reallocate resource to roots increasing the root-shoot ratio and increasing RHD and RHL to promote top soil foraging for P in the growth substrates because more P is available in this layer due to the presence moisture and organic matter, increased microbial activity and oxygen availability (Lynch & Brown, 2001; Péret *et al.*, 2014). In soil, root proliferation in response to low P (Péret *et al.*, 2014) has been reported in rice conditioned by the QTL, *Pup1* (Wissuwa *et al.,* 2005; Heuer *et al.,* 2017). Root traits are considered important targets for breeding for better nutrient acquisition (Gahoonia & Nielsen, 2004; Wissuwa, 2003). The formation of root hairs has been considered one of the key strategies to overcome Pi starvation response with minimal carbon cost (Lynch & Ho, 2005). In *Arabidopsis*, increase in number and length of root hairs improved the Pi uptake under Pi-limiting conditions (Bates and Lynch, 2000). All the high responding foxtail millet genotypes (ISe 1851, CO6, ISe 1338, ISe 1305 and ISe 254) produced abundant long root hairs under P- in the greenhouse experiment. RHL and RHD were associated with higher Pi content under P- conditions and Pi content is a good indicator to gain the better information on plant response under low P conditions.

Higher biomass in the genotypes which produced more and longer root hairs is an indication of better P utilization. However, having a high number of root hairs per root was not sufficient. Genotype ISe 1387 had good root hairs under P- conditions but produced extremely small biomass in the glasshouse experiment. However it was the best performer in the field experiment. Other factors such as efficient symbioses or P acquisition ability from organic sources may influence its potential biomass and effect efficient uptake of Pi from the soil. Some foxtail genotypes had very little or minute root hairs produced under P+ conditions, which suggested that proliferation of root hairs was a low P response. In barley, a root hairless mutant reduced the phosphate uptake under the P- condition and was also associated with decreased biomass production (Gahoonia & Nielsen, 2003; Gahoonia *et al.,* 2001). The good responders under P- condition in the study had both high RHD and RHL. However, in some crops such as soybean, RHD and RHL under low P conditions showed a negative association (Wang *et al.,* 2004). This could be considered as an adaptive response in foxtail millet, a cereal with the adventitious root system, as against the tap root system of soybean wherein a trade-off in terms of carbon use efficiency has been maintained, because combining both RHL and RHD will be too costly in terms of carbon usage. However, in common bean, both higher RHL and RHD were found in P-efficient genotypes (Yan *et al.,* 2004).

In the field-based study, the high responding genotypes under unfertilized conditions
did almost as well as the top performers under P-fertilized conditions, suggesting high
phosphate acquisition and/or use efficiency of these genotypes. Studies of pearl millet and
sorghum showed enormous genetic variation for PUtE in natural field conditions based on
their responses under P starvation (Leiser *et al.,* 2014; Gemenet *et al.,* 2014). The changes in
the growth, biomass and yield have been recognized as important indicators of Pi deficiency
tolerance as reported in other cereals including oat (Andersson *et al.,* 2003), rice (He *et al.,*
2003; Wissuwa *et al.,* 2005), maize (Mollier & Pellerin, 1999) and sorghum (Yoneyama *et*
al., 2007).

Since the seed yield is the ultimate target of any crop based study, plasticity analysis
of P response was limited to this trait. Plasticity is the difference between P+ and P- response
of any genotype under a given cultural condition expressed as the percentage over the
average of that genotype in both the conditions. Since the response under low P is subtracted
from high P, positive plasticity indicates better performance under fertilised conditions and
vice-versa. The genotypes ISe 1387 and ISe 1687 performed well in the natural field in P-
fertilized and unfertilized conditions. However, ISe 1387 and ISe 1687 are P+ responders for
TSE based on plasticity analysis, because they were more productive under fertilised
condition. However, the plasticity of these genotypes is low and closer to the axis (19.3% and
6.6% respectively), which indicated that they could perform equally well under low as well as
high P conditions, irrespective of the magnitude of yield, with a slight edge towards high P
response. Such genotypes are highly preferable, as they can be grown both under varying P
situations.

Clustering analysis and representation of the data as heat maps allowed an overview
of the performance of all genotypes under both greenhouse and field conditions.
Performances in the greenhouse and in the field were not well correlated in the clusters, and
no convincing correlation between any greenhouse measured parameter and seed yield in
unfertilized conditions could be ascertained. However, seed yield under P-fertilized and
unfertilized conditions was correlated, which suggested that genetic control of seed yield was
largely independent of P supply. The total and Pi contents of high-responding genotypes were
much higher than that of low responding genotypes, under both the P-fertilized and
unfertilized field. . Although all the genotypes grown in the natural field were able to grow in
the under unfertilized soil, their uptake efficiency varied considerably resulting in growth
differences. This was reflected in contrast between good and poor responding genotypes.

This data is a pertinent reminder that conditions in the field are very different from those in
more controlled greenhouse conditions and therefore such data should be used as
complementary to the field data to draw conclusions that are practically useful. In an earlier
study on rice genotypes grown under natural soil and solution trials, none of the parameters
was found to be significantly correlated between soil and solution experiments for relative P-
use efficiency (Ni *et al.,* 1998). Further, the importance of field-based testing in two different
seasons is emphasised to obtain an unbiased performance of genotypes. Several factors like
pH of the soil, microorganisms, presence of cations like calcium and aluminium, organic
matter substances and mycorrhizal colonisation can influence the availability of Pi to the
plants (Gemenet *et al.,* 2016). Genotypes that produce fewer root hairs might be compensated
for by mycorrhizal colonisation in the field. A recent meta-analysis of mycorrhizal
colonisation and responsiveness across a range of crop plants and wild relatives concluded
that both colonisation and responsiveness vary markedly (Lehmann *et al.,* 2012). While the
investigation of interaction with mycorrhizal fungi was beyond the scope of this study it is an
important consideration for future study, as is the investigation of root traits such as root
branching and angle which are important traits for nutrient efficient root systems (York &
Lynch, 2015; Zhu *et al.,* 2005).

Details of supplementary data

Supplementary Table S1. Details of foxtail millet genotypes used in the present study

Supplementary Table S2. The raw data, the difference in mean values and plasticity
analysis between Pi fertilized (P+) and unfertilized (P-) treatments of seedlings grown in
natural field condition.

Supplementary Table S3. The difference in mean values between P+ (300 μM Pi) and P-
(10 μM Pi) treatments of seedlings grown in greenhouse condition

Supplementary Figure S1. A clustered heatmap for 54 genotypes of foxtail millet response
under unfertilized (P-) soil. Genotypes are scaled and hierarchically clustered by Euclidean
distance. Yellow equates to a high value and red to a low value. PH, plant height; NPT,
productive tiller number; NL, leaf number; LL, leaf length; LF, flower length; NC, cluster
number; SPC, seeds per cluster; TSE total seed yield.

Supplementary Figure S2. A clustered heatmap for 54 genotypes of foxtail millet response
under Pi fertilized (P+) soil. Genotypes are scaled and hierarchically clustered by Euclidean
distance. Yellow equates to a high value and red to a low value. PH, plant height; NPT,

productive tiller number; NL, leaf number; LL, leaf length; LF, flower length; NC, cluster

number; SPC, seeds per cluster; TSE total seed yield.

Supplementary Figure S3. A clustered heatmap for 54 genotypes of foxtail millet response

for the greenhouse under P+ (300 μM Pi). Genotypes are scaled and hierarchically clustered

by Euclidean distance. Yellow equates to a high value and red to a low value. SL, shoot

length; RL, root length; RHD, root hair density; RHL, root hair length.

Supplementary Figure S4. A clustered heatmap for 54 genotypes of foxtail millet response

for the greenhouse under P- (10 μM Pi). Genotypes are scaled and hierarchically clustered by

Euclidean distance. Yellow equates to a high value and red to a low value. SL, shoot length;

RL, root length; RHD, root hair density; RHL, root hair length.

Acknowledgements

We thank the Tamil Nadu Agricultural University (TNAU), Coimbatore, Tamil Nadu, India,

for supplying the seeds of three local genotypes of foxtail millet. We thank TNAU at

Regional Research Station (RRS), Paiyur, Tamil Nadu, India, and the field workers Mr Palani

and his family members, Ms Vadamalli, Ms Kalaivani, Mr, Murugan, and others at Paiyur,

Tamil Nadu, India for the helps with the field study.

Conflict of interest statement

The authors declare that there is no conflict of interest.

Authors and Contributors

Conceived and designed the experiments: SAC, MR, SI, AB; Performed the

experiments: SAC, MR, VR; Analyzed the data: SAC, MR, KKV; Contributed

reagents/materials/analysis tools: HDU, SI, AB; Wrote the paper: SAC, MR, KKV, AB, SI,

HDU.

References

Andersson MX, Stridh MH, Larsson KE, Liljenberg C, Sandelius AS. 2003. Phosphate-deficient oat replaces a major portion of the plasma membrane phospholipids with the galactolipid digalactosyl diacyl glycerol. Federation of European Biochemical Societies Letters **537,** 128-132.

Baker A, Ceasar SA, Palmer AJ, Paterson JB, Qi W, Muench SP, Baldwin SA. 2015. Replace, reuse, recycle: improving the sustainable use of phosphorus by plants. Journal of Experimental Botany **66,** 3523-3540.

Bates TR, Lynch JP. 2000. The efficiency of *Arabidopsis thaliana* (Brassicaceae) root hairs in phosphorus acquisition. American Journal of Botany **87,** 964-970.

Batten GD. 1986. The uptake and utilization of phosphorus and nitrogen by diploid, tetraploid and hexaploid wheats (*Triticum* spp.). Annals of Botany **58,** 49-59.

Bennetzen JL, Schmutz J, Wang H, *et al.* 2012. Reference genome sequence of the model plant Setaria. Nature Biotechnology **30,** 555-561.

Borch K, Bouma TJ, Lynch JP, Brown KM. 1999. Ethylene: a regulator of root architectural responses to soil phosphorus availability. Plant, Cell & Environment **22,** 425-431.

Carswell C, Grant BR, Theodorou ME, Harris J, Niere JO, Plaxton WC. 1996. The fungicide phosphonate disrupts the phosphate-starvation response in *Brassica nigra* seedlings. Plant Physiology **110,** 105-110.

Ceasar SA, Hodge A, Baker A, Baldwin SA. 2014. Phosphate concentration and arbuscular mycorrhizal colonisation influence the growth, yield and expression of twelve PHT1 family phosphate transporters in foxtail millet (*Setaria italica*). PLoS ONE **9,** e108459.

Ceasar SA. 2018. Genome-wide Identification and *in silico* analysis of PHT1 family genes and proteins in *Setaria viridis*: The best model to study nutrient transport in millets. Plant Genome, doi:10.3835/plantgenome2018-04-0019

Ceasar SA, Ignacimuthu S. 2009. Genetic engineering of millets: current status and future prospects. Biotechnology Letters **31,** 779-788.

Ceasar SA, Baker A, Ignacimuthu S. 2017. Functional characterization of the PHT1 family transporters of foxtail millet with development of a novel *Agrobacterium*-mediated transformation procedure. Scientific Reports **7,** 14064.

Chiou TJ, Aung K, Lin SI, Wu CC, Chiang SF, Su CL. 2006. Regulation of phosphate homeostasis by microRNA in *Arabidopsis*. Plant Cell **18,** 412-421.

Cordell D, Drangert JO, White S. 2009. The story of phosphorus: Global food security and food for thought. Global Environmental Change **19,** 292-305.

Diao X, Jia G. 2017. Origin and domestication of foxtail Millet. In: Doust A, Diao X. (eds.), Genetics and Genomics of Setaria, Plant Genetics and Genomics: Crops and Models Springer **19,** doi: 10.1007/978-3-319-45105-3_4

Dinkelaker B, Hengeler C, Marschner H. 1995. Distribution and function of proteoid roots and other root clusters. Plant biology **108,** 183-200.

Gahoonia TS, Nielsen NE. 2003. Phosphorus (P) uptake and growth of a root hairless barley mutant (bald root barley, brb) and wild-type in low- and high-P soils. Plant, Cell & Environment **26,** 1759-1766.

Gahoonia TS, Nielsen NE. 2004. Barley genotypes with long root hairs sustain high grain yields in low-P field. Plant and Soil **262,** 55-62.

Gahoonia TS, Nielsen NE, Joshi PA, Jahoor A. 2001. A root hairless barley mutant for elucidating genetic of root hairs and phosphorus uptake. Plant and Soil **235,** 211-219.

Gemenet DC, Hash CT, Sy O, Zangre RG, Sanogo MD, Leiser WL. 2014. Pearl millet inbred and testcross performance under low phosphorus in West Africa. Crop Science **54,** 2574-2585.

Gemenet DC, Leiser WL, Beggi F, Herrmann LH, Vadez V, Rattunde HFW, Weltzien E, Hash CT, Buerkert A, Haussmann BIG. 2016. Overcoming phosphorus deficiency in West African pearl millet and sorghum production systems: promising options for crop improvement. Frontiers in Plant Science **7,** 1389.

Gilmour AR, Thompson R, Cullis BR. 1995. Average information REML: An efficient algorithm for variance parameter estimation in linear mixed models. Biometrics **51,** 1440-1450.

Hartigan JA, Wong MA. 1979. Algorithm AS 136: A K-means clustering algorithm. Applied Statistics **28,** 100–108.

He Y, Liao H, Yan X. 2003. Localized supply of phosphorus induces root morphological and architectural changes of rice in split and stratified soil cultures. Plant and Soil **248,** 247-256.

Heuer S, Gaxiola R, Schilling R, Herrera-Estrella L, López-Arredondo D, Wissuwa M, Delhaize E, Rouached H. 2017. Improving phosphorus use efficiency: a complex trait with emerging opportunities. The Plant Journal **90,** 868-885.

Huang P, Shyu C, Coelho CP, Cao Y, Brutnell TP. 2016. *Setaria viridis* as a model system to advance millet genetics and genomics. Frontiers in Plant Science **7,** 1781.

Huehn M. 1996. Non-parametric analysis of genotype x environment interactions by ranks. In: Kang MS, Gauch HG. (eds.), Genotype by Environment Interaction. CRC Press, Boca Raton, 213-228.

Kang MS. 1988. A rank-sum method for selecting high-yielding, stable corn genotypes. Cereal Research Communications **16,** 113-115.

Jin J, Tang C, Armstrong R, Sale P. 2012. Phosphorus supply enhances the response of legumes to elevated CO_2 (FACE) in a phosphorus-deficient vertisol. Plant and Soil **358,** 91-104.

Kalaghatagi SB, Jirali DI, Wali SY, Nagod MS. 2000. Response of foxtail millet (*Setaria italica*) to nitrogen and phosphorus under rainfed conditions of northern dry zone of Karnataka. Annals of Arid Zone **39,** 169-171.

Kim HJ, Lynch JP, Brown KM. 2008. Ethylene insensitivity impedes a subset of responses to phosphorus deficiency in tomato and petunia. Plant, Cell & Environment **31,** 1744-1755.

Lambers H, Finnegan PM, Laliberté EMH, Shane MW, Veneklaas EJ. 2011. Phosphorus nutrition of proteaceae in severely phosphorus-impoverished soils: are there lessons to be learned for future crops? Plant Physiology **156,** 1058-1066.

Lehmann A, Barto EK, Powell JR, Rillig MC. 2012. Mycorrhizal responsiveness trends in annual crop plants and their wild relatives-a meta-analysis on studies from 1981 to 2010. Plant and Soil **355,** 231-250.

Leiser WL, Rattunde HFW, Weltzien E, Haussmann BIG. 2014. Phosphorus uptake and use efficiency of diverse West and Central African sorghum genotypes under field conditions in Mali. Plant and Soil **377,** 383-394.

Li P, Brutnell TP. 2011. *Setaria viridis* and *Setaria italica*, model genetic systems for the Panicoid grasses. Journal of Experimental Botany **62,** 3031-3037.

Lopez-Arredondo DL, Leyva-Gonzalez MA, Gonzalez-Morale SI, Lopez-Bucio J, Herrera-Estrella L. 2014. Phosphate nutrition: improving low-phosphate tolerance in crops. Annual Review of Plant Biology **65,** 95-123.

Lynch JP, Brown KM. 2001. Topsoil foraging-an architectural adaptation of plants to low phosphorus availability. Plant and Soil **237,** 225-237.

Lynch JP, Brown KM. 2008. Root strategies for phosphorus acquisition. In: White PJ, Hammond JP. (eds.), The Ecophysiology of Plant-Phosphorus Interactions. Springer, 83-116.

Lynch JP, Ho MD. 2005. Rhizoeconomics: Carbon costs of phosphorus acquisition. Plant and Soil **269,** 45-56.

Mollier A, Pellerin S. 1999. Maize root system growth and development as influenced by phosphorus deficiency. Journal of Experimental Botany **50,** 487-497.

Muthamilarasan M, Prasad M. 2015. Advances in *Setaria* genomics for genetic improvement of cereals and bioenergy grasses. Theoretical and Applied Genetics **128,** 1-14.

Ni JJ, Wu P, Lou AC, Tao QN. 1998. Rice seedling tolerance to phosphorus stress in solution culture and soil. Nutrient Cycling in Agroecosystems **51,** 95-99.

Ni X, Xia Q, Zhang H, *et al.* 2017. Updated foxtail millet genome assembly and gene mapping of nine key agronomic traits by resequencing a RIL population. GigaScience **6,** 1-8.

Niu YF, Chai RS, Jin GL, Wang H, Tang CX, Zhang YS. 2013. Responses of root architecture development to low phosphorus availability: a review. Annals of Botany **112,** 391-408.

Péret B, Clément M, Nussaume L, Desnos T. 2011. Root developmental adaptation to phosphate starvation: better safe than sorry. Trends in Plant Science **16,** 442-450.

Péret B, Desnos T, Jost R, Kanno S, Berkowitz O, Nussaume L. 2014. Root Architecture Responses: In Search of Phosphate. Plant Physiology **166,** 1713-1723.

Ramakrishnan M, Ceasar SA, Vinod KK, Duraipandiyan V, Ajeesh Krishna TP, Upadhyaya HD, NA Al-Dhabi, Ignacimuthu S. 2017. Identification of putative QTLs for seedling stage phosphorus starvation response in finger millet (*Eleusine coracana* L. Gaertn.) by association mapping and cross species synteny analysis. PLoS ONE **128,** e0183261.

Rouached H, Arpat AB, Poirier Y. 2010. Regulation of phosphate starvation responses in plants: Signaling Players and Cross-Talks. Molecular Plant **3,** 288-299.

Sattari SZ, Bouwman AF, Giller KE, Van-Ittersum MK. 2012. Residual soil phosphorus as the missing piece in the global phosphorus crisis puzzle. Proceedings of the National Academy of Sciences of the United States of America **109,** 6348-6353.

Schachtman DP, Reid RJ, Ayling SM. 1998. Phosphorus uptake by plants: from soil to cell. Plant Physiology **116,** 447-453.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. Nature methods **9,** 671-675.

Slabaugh E, Held M, Brandizzi F. (2011) Control of root hair development in *Arabidopsis thaliana* by an endoplasmic reticulum anchored member of the R2R3-MYB transcription factor family. The Plant Journal **67,** 395-405.

Shimodaira H. 2002. An approximately unbiased test of phylogenetic tree selection. Systematic Biology **51,** 492-508.

Shimodaira H. 2004. Approximately unbiased tests of regions using multistep-multiscale bootstrap resampling. The Annals of Statistics **32,** 2616-2641.

Suzuki R, Shimodaira H. 2006. Pvclust: an R package for assessing the uncertainty in hierarchical clustering. Bioinformatics **22,** 1540-1542.

Upadhyaya HD, Pundir RPS, Gowda CLL, Reddy VG, Singh S. 2008. Establishing a core collection of foxtail millet to enhance the utilization of germplasm of an underutilized crop. Plant Genetic Resources **7,** 177–184.

Veneklaas EJ, Lambers H, Bragg J, Finnegan PM, Lovelock CE, Plaxton WC, Price CA, Scheible WR, Shane MW, White PJ, Raven JA. 2012. Opportunities for improving phosphorus-use efficiency in crop plants. New Phytologist **195,** 306-320.

Vinod KK, Heuer S. 2012. Approaches towards nitrogen and phosphorus-efficient rice. AoB Plant **2012,** pls028.

Vinod KK. 2015. Enhancing nutrient starvation tolerance in rice. In: Jaiwal PK, Singh RP, Dhanker OP. (eds.), Genetic manipulation in plants for mitigation of climate change, Springer, doi: 10.1007/978-81-322-2662-8_6

Wang L, Liao H, Yan X, Zhuang B, Dong Y. 2004. Genetic variability for root hair traits as related to phosphorus status in soybean. Plant and Soil **261**, 77-84.

Wang X, Shen J, Liao H. 2010. Acquisition or utilization, which is more critical for enhancing phosphorus efficiency in modern crops? Plant Science **179,** 302-306.

Wiel CCM, Linden CG, Scholten OE. 2015. Improving phosphorus use efficiency in agriculture: opportunities for breeding. Euphytica **207,** 1-22.

Wissuwa M. 2003. How do plants achieve tolerance to phosphorus deficiency? small causes with big effects. Plant Physiology **133,** 1947-1958.

Wissuwa M, Gamat G, Ismail AM. 2005. Is root growth under phosphorus deficiency affected by source or sink limitations? Journal of Experimental Botany **56,** 1943-1950.

Wissuwa M, Ae N. 2001. Genotypic variation for tolerance to phosphorus deficiency in rice and the potential for its exploitation in rice improvement. Plant Breeding **120,** 43-48.

Yan X, Liao H, Beebe SE, Blair MW, Lynch JP. 2004. QTL mapping of root hair and acid exudation traits and their relationship to phosphorus uptake in common bean. Plant and Soil **265,** 17-29.

Yoneyama K, Xie X, Kusumoto D, Sekimoto H, Sugimoto Y, Takeuchi Y, Yoneyama K. 2007. Nitrogen deficiency, as well as phosphorus deficiency in sorghum, promotes the production and exudation of 5-deoxystrigol, the host recognition signal for arbuscular mycorrhizal fungi and root parasites. Planta **227,** 125-132.

York LM, Lynch JP. 2015. Intensive field phenotyping of maize (*Zea mays* L.) root crowns identifies phenes and phene integration associated with plant growth and nitrogen acquisition. Journal of Experimental Botany **66,** 5493-5505.

Zhang G, Liu X, Quan Z, *et al.* 2012. Genome sequence of foxtail millet (*Setaria italica*) provides insights into grass evolution and biofuel potential. Nature Biotechnology **30,** 549-554.

Zhu J, Kaeppler SM, Lynch JP. 2005. Topsoil foraging and phosphorus acquisition efficiency in maize (*Zea mays*). Functional Plant Biology **32,** 749-762.

Tables

Table 1. Testing for the significance of variance component effects by restricted maximum likelihood (REML) method.

Traits	Variance			AIC	Heritability
	Genotype	Season	Genotype x Season		
H-PH	269.5*	7.50	4.05	2701.53	0.95
H-NPT	5.68*	0.11	0.54	1638.07	0.91
H-NL	1.21*	0.05	0.00	1251.44	0.87
H-LL	37.8*	0.34	0.64	2085.77	0.95
H-LF	16.23*	0.48	1.05	1702.24	0.97
H-NC	270.73*	4.21	5.98	2446.55	0.97
H-SPC	379.2*	4.52	34.4	2310.76	0.95
H-TSE	41.12*	0.32	0.18	880.99	0.99
L-PH	511.32*	5.46	2.05	2307.75	0.98
L-NPT	6.54*	0.32	0.02	1503.11	0.92
L-NL	1.99*	0.03	0.00	1120.4	0.93
L-LL	52.24*	0.26	0.04	1918.04	0.95
L-LF	15.6*	0.34	0.47	1531.03	0.96
L-NC	344.5*	14.04	9.74	2104.48	0.97
L-SPC	285.97*	1.62	6.61	1905.93	0.98
L-TSE	45.76*	0.42	0.21	656.35	0.99

*Significant at $p<0.05$ by chi-square test. AIC, Akaike information criterion. Trait names are prefixed H- for P+ response and L- for low P response. PH, plant height in cm; NT, number of tillers; NPT, number of productive tillers; NL number of leaves; LL, length of leaf in cm; LF, length of panicle in cm; NC, number of clusters/panicle; SPC, seeds per cluster; TSE total seed yield in g.

Table 2. Mean values of the five high and low responding genotypes of foxtail millet in growth assays under unfertilised (P-) and P fertilised (P+) natural field conditions

Genotypes	PH	NPT	NL	LL	LF	NC	SPC	TSE
Good responders under unfertilised plot (P-)								
ISe 1387*	146.33±17.41[a]	4.5±1.76[a]	10.5±1.05[a]	47.71±3.28[a]	23.93±2.83[a]	142.33±15.86[a]	49.87±8.96[a]	25.15±1.31[a]
ISe 1687*	133±10.77[b]	8.33±1.75[b]	10±1.26[b]	39.21±9.82[b]	17.75±2.31[b]	74.67±8.64[b]	60.33±6.8[b]	14.79±0.9[b]
ISe 663*	131.17±7.83[b]	13.33±1.37[c]	10.67±0.82[b]	32.56±3.08[c]	20.45±1.55[c]	79±8.41[c]	28.33±3.72[c]	17.93±1.01[c]
CO7*	132.86±4.6[b]	10.97±1.47[d]	10±0.63[c]	35.45±1.65[d]	20.96±0.92[c]	85±2.34[d]	30.33±2.94[c]	14.13±1.09[b]
ISe 869*	140.83±14.96[c]	8.67±3.44[e]	9.17±1.6[d]	29.21±6.46[e]	18.46±3.78[d]	73.67±13.69[b]	41.5±5.01[d]	12.66±0.48[d]
Poor responders under unfertilised plot (P-)								
Maxima1#	72.83±10.4[a]	4.33±1.21[a]	6.33±1.51[a]	17.3±3.75[a]	8.83±5.4[a]	28.33±12.24[a]	20.04±1.89[a]	1.15±0.38[a]
ISe 1335#	66.83±16.49[b]	5.0±1.79[b]	5.17 ± 1.6[b]	11.51±4.05[b]	5.96±2.4[b]	33.83±3.66[b]	32.5±5.05[b]	2.28±0.61[a]
ISe 1037#	68.33±17.11[b]	4.83±2.14[b]	7.33±1.75[c]	19.23±6.68[c]	8.9±1.78[c]	45.33±5.2[c]	10.83±2.32[c]	4.28±0.79[b]
ISe 1234#	72.67±17.49[a]	5.17±1.94[c]	6.33±1.63[a]	18.63±7.21[d]	8.89±3.73[c]	48.33±5.92[c]	11.33±1.21[c]	3.84±0.62[b]
ISe 1302#	80±15.57[c]	5.17±1.6[c]	6.5±0.84[a]	18.15±2.58[d]	10.01±2.12[d]	40.5±3.21[d]	14.67±3.14[c]	3.04±0.18[b]
Good responders under P fertilised plot (P+)								
ISe 1387*	173.17±16.67[a]	6.67±1.37[a]	11.67±1.75[a]	48.26±8.14[a]	27.63±3.2[a]	152.67±16.28[a]	90±11.76[a]	30.52±1.15[a]
ISe 132*	142.83±8.04[b]	10.67±5.5[b]	8.17±1.47[b]	35.86±3.74[b]	23.25±1.78[b]	82±6.16[b]	29.5±5.96[b]	28.48±1.43[b]
ISe 1851*	151.5±15.64[c]	7±2.37[c]	9.5±0.55[c]	33.89±3.87[c]	21.46±2.68[c]	69.67±7.99[c]	43.83±4.4[c]	22.39±1.08[c]
ISe 907*	131±10.04[d]	10.67±3.33[b]	8.33 ± 0.52[b]	39.87±13.54[d]	18.77±3.21[d]	80.5±1.52[b]	40.13±11.14[c]	15.91±0.47[d]
ISe 1687*	131.33 ± 5.68[d]	8.83±3.19[d]	9 ± 1.26[d]	33.1±3.79[c]	19.61±2.61[d]	68.17±9.06[d]	74±6.23[d]	15.8±0.76[d]
Poor responders under P fertilised plot (P+)								
Maxima1#	87.17±16.8[a]	4±1.67[a]	5±1.41[a]	16.48±6.01[a]	9.05±3.38[a]	34.33±8.85[a]	25.89±2.4[a]	0.83±0.34[a]
ISe 1335#	84.33±12.55[a]	5±1.67[b]	5.17±2.32[a]	12.52±5.68[b]	6.41±1.14[b]	29.67±9.73[b]	32±4.6[b]	2.94±0.4[b]
ISe 1299#	83.17±8.4[b]	7.5±0.84[c]	6.83±0.41[b]	22.54±3.04[c]	10.03±3.17[c]	47.83±7.39[c]	12.83±1.94[c]	2.97±0.41[b]
ISe 1234#	93.67±5.39[c]	4.67±2.5[d]	6.17 ±1.33[c]	22.86±4.22[c]	9.03±3.43[a]	41.5±8.17[d]	22.33±3.27[d]	28.26±0.87[c]
ISe 1161#	107.67±23.31[d]	9.17±0.75[e]	6.67±2.25[b]	21.34±7.39[c]	13.65±4.33[d]	53.33±15.81[e]	29.5±6.66[a]	5.98±0.13[d]

Values are Mean ± SD; *, high ranking genotypes; #, low ranking genotypes. The genotypes showing variation between total and productive tillers are in bold.

In each column, values followed by different letters are significantly different (*p<0.05*) based on students t-test. PH, plant height in cm; NT, number of tillers; NPT, number of productive tillers; NL number of leaves; LL, length of leaf in cm; LF, length of panicle in cm; NC, number of clusters/panicle; SPC, seeds per cluster; TSE, total seed yield in   g.

Table 3. Mean values for the five high and low responding genotypes of foxtail millet in growth assays under low (P-) and high (P+) in greenhouse conditions

Genotypes	BIO	SL	RL	RHD	RHL
Good responders under P+					
CO-5*	8.00 ± 1.00^a	21.73 ± 7.98^a	13.13 ± 0.87^a	41.67 ± 4.04^a	3.16 ± 0.01^a
CO-6*	10.47 ± 0.15^b	15.87 ± 5.24^b	19.60 ± 0.82^b	21.67 ± 3.06^b	3.26 ± 0.00^a
ISe 2*	7.10 ± 1.00^c	17.30 ± 0.61^c	15.00 ± 1.00^c	13.67 ± 1.53^c	3.04 ± 0.04^a
ISe 869*	6.70 ± 0.62^c	18.07 ± 5.97^c	8.80 ± 1.25^d	23.33 ± 2.52^d	3.56 ± 0.07^b
Maxima1*	15.30 ± 1.00^d	10.20 ± 2.35^d	10.60 ± 3.08^e	24.67 ± 3.51^d	2.57 ± 0.04^c
Poor responders under P+					
ISe 748#	2.13 ± 0.60^a	2.60 ± 0.92^a	5.17 ± 0.40^a	0.00 ± 0.00	0.00 ± 0.00
ISe 1687#	2.80 ± 2.27^b	5.17 ± 1.67^b	5.40 ± 0.26^{ab}	1.00 ± 1.73^a	0.49 ± 0.00^a
ISe 1387#	3.00 ± 0.56^b	4.53 ± 0.95^b	6.70 ± 0.62^c	0.00 ± 0.00	0.00 ± 0.00
ISe 376#	3.50 ± 0.30^c	8.43 ± 2.40^c	5.93 ± 1.55^b	0.00 ± 0.00	0.00 ± 0.00
ISe 1254#	6.10 ± 1.15^d	2.93 ± 0.51^a	4.87 ± 0.67^{ab}	1.67 ± 2.89^a	0.40 ± 0.00^a
Good responders under P-					
ISe 1851*	7.03 ± 3.67^a	9.13 ± 2.81^a	12.43 ± 3.77^a	51.33 ± 3.21^a	4.85 ± 0.10^a
CO-6*	8.00 ± 0.46^c	14.40 ± 3.27^b	13.70 ± 1.62^b	45.33 ± 3.79^b	3.70 ± 0.11^b
ISe 1338*	6.66 ± 0.32^a	15.90 ± 2.25^c	16.10 ± 2.21^c	49.33 ± 1.53^a	3.75 ± 0.12^b
ISe 1305*	6.07 ± 2.15^b	13.00 ± 0.50^d	9.50 ± 4.74^d	50.67 ± 4.51^a	6.24 ± 0.05^c
ISe 254*	5.87 ± 0.81^b	10.43 ± 4.56^e	9.13 ± 1.21^d	44.00 ± 3.61^b	8.01 ± 0.10^d
Poor responders under P-					
ISe 1474#	2.07 ± 0.59^a	4.50 ± 2.18^b	2.47 ± 0.45^a	1.20 ± 0.00^a	1.06 ± 0.18^a
ISe 748#	2.90 ± 0.36^b	5.90 ± 1.91^b	1.67 ± 0.76^b	0.00 ± 0.00	0.00 ± 0.00
ISe 1687#	2.40 ± 1.01^a	5.17 ± 2.02^b	3.50 ± 1.37^c	31.00 ± 3.00^b	2.37 ± 0.03^b
ISe 1254#	4.43 ± 0.78^c	6.67 ± 0.38^c	4.73 ± 0.25^d	0.00 ± 0.00	0.00 ± 0.00
ISe 1736#	3.90 ± 0.96^d	5.40 ± 1.89^b	7.10 ± 1.21^e	0.00 ± 0.00	0.00 ± 0.00

Values are Mean ± SD; *= High ranking genotypes; # = Poor ranking genotypes. Within each column, values followed by different letters are significantly different ($p<0.05$) based on students t-test.
BIO, seedling biomass in mg; SL, shoot length in cm; RL, root length in cm; RHD, root hair density per 10µm length, RHL, root hair length in µm.

Table 4. Interrelations between P content in shoot and roots among the extreme genotypes showing P response under greenhouse (upper diagonal) and field (lower diagonal) conditions. The diagonal values (in bold) are correlations between screen house and field parameters. Values shown against genotype response groups are the correlations between field and greenhouse parameters under each category.

	Pi shoot	*Total P shoot*	*Pi root*	*Total P root*
Pi shoot	**0.705***	0.761*	0.820*	0.294
Total P shoot	0.949*	**0.842***	0.790*	0.730*
Pi root	0.940*	0.953*	**0.808***	0.526*
Total P root	0.919*	0.971*	0.978*	**0.501***
Good P-	-0.157	0.747*	0.407	0.418
Poor P-	-0.250	0.215	-0.190	-0.692*
Good P+	-0.668*	-0.746*	-0.878*	0.237
Poor P+	-0.507*	0.870*	0.066	0.698*

*significant as $p<0.05$

Legends for Figures

Figure 1. Rank sums of genotypes based on (A) field and (B) greenhouse evaluation. The 10 extreme genotypes are selected under each P regime, P+ and P-. P+ ranks are ordered from poor (having high-rank sum) to good (having low-rank sum). P ranks are ordered from good (low-rank sum) to poor (high-rank sum). Five genotypes with the lowest rank sum in each group are selected as best genotypes (green), and five genotypes each with highest rank sum are selected as poor genotypes (red).

Figure 2. Flowers with mature seeds of selected genotypes (good, intermediates and poor responding) of foxtail millet grown in a natural field Pi fertilized (P+) and unfertilized (P-) soil conditions (P-, low phosphate; P+, high phosphate). The genotypes in the top row are ISe 663, CO7, ISe 1338, ISe 869 and ISe 1736 and bottom row are ISe 1037, ISe 1335, ISe 1234, ISe 1563 and Maxima1.

Figure 3. Root hair images of selected genotypes of foxtail millet showing a response to P- and P+ in the greenhouse**.** The image was taken after 15 days of growth under P- and P+ in the greenhouse. The genotypes ISe 1851 and ISe 1305 are the top in the cumulative ranking and high responding genotypes for root hair formation under P- condition. The genotypes ISe 748, ISe 1687 and ISe 1736 are low performers under P-. The genotype ISe 1387 is a low performer under P+. The genotype ISe 1820 is an intermediate responder in both P- and P+.

Figure 4. Inorganic (green) and total P assay (brown) for the five each of high performers and low performers of foxtail millet genotypes under P- (10 μM Pi) and P+ (300 μM Pi) conditions in greenhouse conditions. A, high performers under P+; B, low performers under P+; C, high performers under P-; D, low performers under P-. The letters over the bars indicate statistically significant difference if the letters are different as p<0.05. The vertical lines indicate error bars.

Figure 5. Inorganic (green) and total P assay (brown) for the five each of high performers and low performers of foxtail millet genotypes under unfertilized (P-) and P fertilized (P+) conditions in the natural field. A, high performers under P+; B, low performers under P+; C, high performers under P-; D, low performers under P-. The letters over the bars indicate statistically significant difference if the letters are different as p<0.05. The vertical lines indicate error bars.

Figure 6. Differential response of genotypes for total seed yield. The genotypes on the left side of the figure (Scale 0 to -150 and red colour bars) tend more towards low P tolerance.

Genotypes with long bars on right side of the figure (Scale 0 to 200 and green colour bars) are P+ responders, those closer to the axis are more plastic to P level variations and show stable performance under both P+ and P- conditions.

Figure 7. Hierarchical cluster showing the approximately unbiased probability (AU) for the P response genotypes. The value greater than 95% indicates that clusters crated by the variables of P response and node is strongly supported by data. Red boxes indicate the clustering of the genotypes.

A Rank sums of genotypes based on P- and P+ in the field

B Rank sums of genotypes based on P- and P+ in the greenhouse

P -   P+   P -   P+

ISe 1851   ISe 1305

ISe 1820   ISe 1209

ISe 1387   ISe 1736

ISe 1687   ISe 748

Cluster dendrogram with AU/BP values (%)

AU - Approximately Unbiased *p*-vlaue

Distance: uncentered
Cluster method: average