A Pandas complex adapted for piRNA-guided transposon silencing

Kang Zhao[#,1], Sha Cheng[#,2], Na Miao[#,1], Ping Xu[#,1,3], Xiaohua Lu[#,1], Ming Wang[1], Yuhan Zhang[2], Xun Yuan[2], Weiwei Liu[1], Xin Lu[1], Xuan Ouyang[1], Peng Zhou[1], Jiaqi Gu[1,4], Yiqun Zhang[1], Ding Qiu[1], Shan Wang[1], Zhaohui Jin[2], Youzhong Wan[3], Jinbiao Ma[4], Hong Cheng[2], Ying Huang[2,5], Yang Yu[1,5]

[1]Institute of Biophysics, Key Laboratory of RNA Biology, Chinese Academy of Sciences, University of Chinese Academy of Sciences, Beijing, China, 100101
[2]State Key Laboratory of Molecular Biology, Shanghai Key Laboratory of Molecular Andrology, CAS Center for Excellence in Molecular Cell Science, Shanghai Institute of Biochemistry and Cell Biology, Chinese Academy of Sciences, University of Chinese Academy of Sciences, Shanghai, China, 200031
[3]School of Life Sciences, National Engineering Laboratory of AIDS Vaccine, Key Laboratory for Molecular Enzymology and Engineering of Ministry of Education, Jilin University, Changchun, Jilin, China, 130012
[4]State Key Laboratory of Genetic Engineering, Department of Biochemistry, School of Life Sciences, Fudan University, Shanghai, China, 200438

[#]contributed equally

[5]correspondence: yuyang@ibp.ac.cn; huangy@sibcb.ac.cn

Keywords: piRNA, transposon silencing, dNxf2, Panoramix, crystal structure, *Drosophila*, dNxf1/TAP, dNxt1/p15

Abstract

The repression of transposons by the Piwi-interacting RNA (piRNA) pathway is essential to protect animal germ cells. In *Drosophila* ovaries, Panoramix (Panx) enforces transcriptional silencing by binding to the target-engaged Piwi-piRNA complex, although the precise mechanisms by which this occur remain elusive. Here, we show that Panx functions together with a germline specific paralogue of a nuclear export factor, dNxf2, and its cofactor dNxt1 (p15) as a ternary complex to suppress transposon expression. Structural and functional analysis demonstrate that dNxf2 binds Panx via its UBA domain and play an important role in transposon silencing through binding to transposon transcripts directly. Unexpectedly, dNxf2 interacts directly with dNxf1 (TAP), which is also essential for transposon silencing. Transient tethering of dNxf2 to nascent transcripts leads to their nuclear retention. Therefore, we propose that dNxf2 may function as a Pandas (Panoramix-dNxf2 dependent TAP/p15 silencing) complex, which counteracts the canonical RNA exporting machinery (TAP/p15) and restricts transposons to nuclear peripheries. Our findings may have broader implications for understanding how RNA metabolism modulates epigenetic gene silencing and heterochromatin formation.

Transposons are highly abundant in eukaryotes, which make up nearly half of human genome. To maintain eukaryotic genome integrity, nascent transcripts of transposons are often targeted by nuclear Argonaute proteins for transcriptional gene silencing (TGS)[1-5]. In animal gonads, the PIWI-clade Argonautes guided by piRNAs (PIWI-interacting RNA) are thought to recognize nascent transposon transcripts and direct sequence-specific heterochromatin formation[3]. As a critical cofactor of *Drosophila* nuclear Piwi, Panoramix (Panx, also known as Silencio) links the target-engaged Piwi-piRNA complex to the general silencing machinery[6,7]. Enforced tethering of Panx to nascent transcripts leads to cotranscriptional silencing and correlates with deposition of histone H3 lysine 9 trimethylation (H3K9me3) marks[6,7]. However, the mechanism by which Panx mediates the repression remains unknown. An equally important question is why H3K9me3 marks are not always sufficient for transposon silencing[8].

To understand this fundamental question, we cross-examined proteins that co-immunoprecipitated with Panx (Extended Data Fig. 1a) with piRNA pathway candidate genes from RNA interference (RNAi) screens[9-12]. Unexpectedly, dNxf2 was identified as a potential cofactor of Panx (Extended Data Fig.1a-c). dNxf2 belongs to an evolutionarily conserved NXF (underline{n}uclear underline{e}xport underline{f}actor) family of proteins, yet depletion of dNxf2 had no effect on polyadenylated mRNA export[13,14]. Instead, dNxf2 and its cofactor dNxt1 (also known as p15) were both identified in two published RNAi screens as being essential for transposon silencing[9,10]. Similar to Panx, dNxf2 is specifically expressed in female gonads (Extended Data Fig.1d).

To validate the interaction between Panx and dNxf2, GFP immunoprecipitation from ovaries expressing GFP-Panx fusion under its native promoter was performed. Results of mass spectrometry analysis (Extended Data Fig. 1b) and western blotting demonstrated that endogenous dNxf2 is associated with Panx (Fig. 1a). Likewise, Halo-tagged dNxf2 was able to precipitate

endogenous Panx from Ovarian Somatic Cell (OSC) lysates with a Halo-tag integrated into the C-

terminus of Nxf2 (Fig. 1b and Extended Data Fig. 6e). We next tested if dNxf2 is functionally

required for Panx-mediated silencing. The luciferase transcripts with BoxB sites in their 3′

untranslated regions are repressed if λN-Panx is tethered[6,7]. The expression level of luciferase was

measured upon germline-specific knockdowns of either dNxf2 or dNxt1 (Fig. 1c). Despite

constitutive tethering of λN-Panx, loss of either dNxf2 or dNxt1 significantly weakened the ability of

Panx to repress the reporter, as compared to the controls (Zuc or attp2, Fig. 1c). Consistent with

the reporter de-repression, transposon transcripts are elevated upon dNxf2 RNAi (Extended Data

Fig. 1e). Taken together, our data suggested that dNxf2 and dNxt1 may function as a heterodimer

either with or downstream of Panx to suppress transposon expression.

Next, we used RNA sequencing (RNA-seq) to examine global effects on transposon

expression with germline-specific knockdowns of dNxf2, compared with Panx RNAi (Fig. 1d-f). As

expected, dNxf2 knockdown triggered a dramatic increase of transposon transcripts (Fig. 1d),

similar to that of Panx (Fig. 1e-f), suggesting that dNxf2 is specifically required for silencing of

transposons repressed by Panx. To rule out off-target effects of RNAi, mutants of dNxf2 were

generated using CRISPR/Cas9 technique (Extended Data Fig. 2a)[15]. The dNxf2 mutant female

flies were completely sterile with a deletion of 20 amino acids at the N-terminus of dNxf2 (Extended

Data Fig. 2b), similar to other core piRNA pathway mutants[6]. Loss of dNxf2 showed little effect on

Piwi nuclear localization or stability (Extended Data Fig.2c-d), indicating that dNxf2 functions as an

effector protein. Furthermore, dNxf2 mutants showed global up-regulation of transposons

(Extended Data Fig. 2e-g) and de-repression of the luciferase reporter despite λN-Panx tethering

(Extended Data Fig. 3a). The protein level of endogenous Panx is noticeably reduced in the

absence of dNxf2 (Extended Data Fig. 2d). To rule out the possibility that dNxf2 might affect

transposons indirectly via Panx stability, transposon expression levels were measured upon

overexpression of λN-Flag-Panx under the dNxf2 mutant background (Extended Data Fig. 3a-e).

Still, dNxf2 mutant female flies lost transposon control (Extended Data Fig. 3c-d) and were

completely sterile as if Panx did not exist (Extended Data Fig. 3e).

The striking phenotypic similarities between dNxf2 and Panx prompted us to test whether these

two proteins interact directly. We used yeast two-hybrid (Y2H) assays to determine the interacting

regions. The domain architecture of dNxf2 is very similar to that of the canonical RNA export factor,

dNxf1/TAP (also known as sbr, Fig. 2a). Both proteins contain leucine-rich repeats (LRR), an RNA

recognition motif (RRM), a nuclear transport factor 2 (NTF2)-like domain, and a ubiquitin-

associated domain (UBA). Y2H assays were performed to examine the interactions. Interestingly,

Panx only interacts with the UBA domain of dNxf2 but not that of dNxf1 (Fig. 2b), indicating that

this interaction between Panx and dNxf2 is specific (Fig. 2b). Surprisingly, neither full length nor

the NTF2+UBA fragment of dNxf2 could bind Panx (Fig. 2b), suggesting that the UBA domain of

dNxf2 might be in a "closed" conformation in the presence of the NTF2 domain. In this regard, the

interactions between dNxt1 and the NTF2 domains of either dNxf2 or dNxf1 are also weakened in

the presence of its UBA domain (Fig. 2c). Since *Drosophila* Nxt1 itself is absent in the Y2H system,

we tested whether dNxt1 might release the UBA domain from the NTF2 domain to permit Panx

binding. Indeed, ectopic expression of dNxt1 is sufficient to allow full length dNxf2 to interact with

Panx in a Y2H assay (Fig. 2d). Next, we mapped the minimum region of Panx down to residues

315-343 (NIR, d<u>N</u>xf2 <u>i</u>nteracting <u>r</u>egion) as sufficient for UBA binding (Fig. 2e and Extended Data

Fig. 3f). Consistent with the fact that dNxt1 forms a heterodimer with dNxf2[13], we found that dNxt1

co-migrates with a fusion protein of dNxf2$^{NTF2+UBA}$-(Gly-Ser)$_4$-Panx^{NIR} by size-exclusion

chromatography (Fig. 2f), suggesting that Panx, dNxf2, and dNxt1 may exist as a ternary complex.

We were not able to crystalize the dNxf2^{NTF2} domain, instead, we crystallized dNxf1^{NTF2} in complex

with dNxt1 and determined the structure (Extended Data Fig. 4a). Residues that may involve in the

binding of dNxf2^{NTF2} to dNxt1 were modeled according to the sequence alignment result and the

structure of dNxf1^{NTF2} (Extended Data Fig. 4a). dNxf2^{NTF2} maintained most, if not all, residues that

interact with dNxt1 as validated by the Y2H and co-immunoprecipitation assays (Extended Data

Fig. 4a-e), indicating that the interaction mode of dNxf2^{NTF2} and dNxt1 complex is similar to that of

dNxf1^{NTF2} and dNxt1.

Purified Panx^{NIR} forms a stable complex with the UBA domain of dNxf2 with a dissociation

constant of ~3.2 μM as measured by isothermal titration calorimetry (ITC) (Fig. 2g). To further

explore the molecular basis of interactions between dNxf2 and Panx, the crystal structure of the

dNxf2-Panx complex was determined (Fig. 3a-d). Despite many efforts, only the fusion protein of

dNxf2^{UBA}-(Gly-Ser)$_4$-Panx^{NIR} could be successfully crystallized. The structure was solved to a 1.5

Å resolution (Table S1). dNxf2^{UBA} forms a compact three-helix bundle (α1–α3) with a 3_{10}-helix (η1)

at the C-terminus (Fig. 3b). The Panx^{NIR} is folded into a long α-helix and lays on the hydrophobic

surface formed by α2 and α3 (Fig. 3b-c and Extended Data Fig. 4f). A324, A328, V331, L332, and

I335 on Panx interact with V800, F819, F826, F840, L823 and I827 on dNxf2 via hydrophobic

interactions (Fig. 3d). Moreover, R321 and R327 on Panx form salt bridges with D837 and E799

on Nxf2^{UBA}, respectively (Fig. 3d). To validate the intermolecular interactions between dNxf2 and

Panx, key residues on the interacting interface were mutated (Fig. 3e). While either L823A or

D837A single point mutation affected the binding between Panx and dNxf2^{UBA}, the double point

mutation of dNxf2^{UBA} (F826A/I827A) nearly abolished its interaction with Panx in both Y2H and co-

immunoprecipitation assays (Fig. 3e-f), highlighting the significant contribution of these residues in

Panx binding.

In contrast to the highly charged surface of the Nxf1-type UBA (i.e. hsNxf1 or scMex67),

dNxf2^{UBA} favors hydrophobic binding with Panx (Fig. 3c and Extended Data Fig. 5a-b). Key residues

on the interacting interface are highly conserved among different *Drosophila* species but altered in

the Nxf1-type UBA (Extended Data Fig. 5a). On the opposite surface of the Nxf1-type UBA, a

hydrophobic pocket is formed to accommodate the FxFG peptide of the nuclear pore complex (NPC)

(Extended Data Fig. 5c-d). However, this pocket is missing in dNxf2^{UBA} due to a salt bridge formed

between K829 and E814 (Extended Data Fig. 5c). Additionally, the bulky side chain of L825 on

dNxf2^{UBA} may hinder FG binding (Extended Data Fig. 5c). In contrast, the corresponding amino

acids in hsNxf1 (A602) or scMex67 (G583) contain much smaller side chains (Extended Data Fig.

5c-d), therefore, give space for FxFG interactions. Consistent with the structural prediction,

dNxf2^{UBA} failed to interact with the FG-repeats of Nup214, a NPC component known to interact with

dNxf1^{UBA} (Extended Data Fig. 5e)[14]. Since two copies of FG binding domains (NTF2 and UBA) are

minimally required for proper RNA export[16], dNxf2 lacks at least one copy of the FG binding domain

(UBA) and thus cannot export RNAs.

To validate the importance of the direct interactions between dNxf2 and Panx *in vivo*, a

previously described λN/BoxB luciferase reporter system was used to check if artificial tethering of

dNxf2 could lead to repression. As expected, significant repression upon tethering of a λN-dNxf2

fusion protein (Fig. 3g) was observed, unlike that of the negative controls (λN-YFP or GFP-dNxf2

lacking a λN-tag). Similar to Panx, the level of λN-dNxf2 mediated repression was in a dosage-

dependent manner, which is correlated with the number of BoxB binding sites (Fig. 3h). Most

importantly, the repression is dependent on the presence of the dNxf2 UBA domain (Fig. 3g, dNxf2-

ΔUBA).

The domain architecture of dNxf2 is similar to Nxf1 proteins which also contains RNA binding

domains at the N-terminus, implying that dNxf2 might directly bind to transposon transcripts *in vivo*

(Fig. 3a)[13]. To validate this hypothesis, we performed GoldCLIP/RT-qPCR experiments, which

reply on a covalent attachment of the Halo-tag to its ligand on beads, affording a denaturing

purification of crosslinked protein-RNA complexes[17]. A Halo-tag was inserted to the C-terminus of

dNxf2 (dNxf2-Halo) using CRISPR/Cas9 (Extended Data Fig. 6a-f)[18]. Strikingly, following UV

crosslinking, transposon transcripts (mdg1), but not rp49, remained attached to the dNxf2-Halo

fusion protein despite denaturing washes (Extended Data Fig. 6b-c). The association is dependent

on UV crosslinking, demonstrating a direct binding between mdg1 and dNxf2 (Extended Data Fig.

6b-c). Interestingly, the interaction was only observed when both histone H1 and Heterochromatin

Protein 1a (HP1a) were depleted by RNAi, but not in the control knockdown (Extended Data Fig.

6b-c), supporting that the majority of transposon transcripts remain suppressed in a wildtype

background. Low steady-state levels of transposon nascent RNAs make it extremely difficult to

obtain reasonable signals in an already inefficient UV crosslinking experiment. Nevertheless, upon

removal of the downstream silencing factors (H1 and HP1a), transposon transcripts accumulate

and bind to dNxf2 (Extended Data Fig. 6b). In contrast, Frogger didn't show any detectable CLIP

signal despite of the dramatic up-regulation of its transcripts upon H1 and HP1a double RNAi

(Extended Data Fig. 6d). This result suggests that the binding of dNxf2 to transposon is likely

piRNA-guided, since Frogger is not targeted by Piwi-piRNAs in OSCs[19]. In order to obtain enough

material, we performed GoldCLIP-seq experiments using dNxf2-Halo knock-in OSCs depleted of

Maelstrom, a piRNA pathway effector component either parallel or downstream of H3K9me3

establishment on transposons (personal communications with Mikiko Siomi)[8]. Consistent with the

RT-qPCR results, CLIP-seq demonstrated that dNxf2 preferentially binds to Piwi-targeted

transposons (Fig. 4a and Extended Data Fig. 7). Moreover, the RNA binding domains (RBD) of

dNxf2 are essential for silencing in the tethering assays (Extended Data Fig. 8). Therefore,

dNxf2^{RBD} are likely to be involved in the effector step of silencing rather than the Piwi-dependent

recruitment of dNxf2. Collectively, our data strongly argues that Panx and dNxf2/dNxt1 function

together as a stable complex to directly suppress transposons that are targeted by Piwi-piRNAs.

Loss of Panx leads to significantly decreased H3K9me3 marks over transposons[6,7]. We next tested whether removing dNxf2 could result in a similar phenomenon. Since Panx is unstable in the absence of dNxf2 (Extended Data Fig. 2d), we performed H3K9me3 ChIP-qPCR assays over several transposons as well as the Firefly-10xBoxB reporter tethered with λN-Flag-Panx. H3K9me3 showed rather mild changes upon removal of dNxf2. This is in direct contrast to the dramatic upregulation of transposon transcripts in the absence of dNxf2 (Extended Data Fig. 3c-d and 9). This result suggested that transposon silencing and H3K9me3 deposition are uncoupled in *dNxf2* mutants. Interestingly, we found that Het-A chromatin left nuclear peripheries upon loss of either Panx or dNxf2 (Extended Data Fig. 10)[20]. Transposon transcripts (I element) which are targeted by piRNAs have been previously shown to accumulate within nucleus[21]. Thus, we proposed that certain RNA export machineries might be modulated by the piRNA pathway to prevent transposon transcripts from getting translated. Human Nxf1/TAP has been reported to dimerize with most NXF family members to regulate RNA export[22,23]. We wondered whether dNxf2 might interact with dNxf1 *in vivo*. GFP-tagged dNxf2 can co-immunoprecipitate Halo-tagged dNxf1 from OSC lysates (Fig. 4b). Either NTF2 or UBA domain of dNxf1 is sufficient to interact with dNxf2^{NTF2} (Fig. 4c and Extended Data Fig. 11). Most importantly, GST pull-down assays demonstrated a direct binding between dNxf1 and dNxf2 (Fig. 4d and Extended Data Fig. 12). Consistent with the involvement of dNxf1 in transposon silencing, dNxf1 was found in a genome-wide RNAi screen for piRNA pathway components[11]. Female flies with mutant alleles of dNxf1 are sterile and have defects in meiotic spindles[24], a phenotype reminiscent of core piRNA pathway mutants[25,26]. Therefore, we tested if dNxf1 is required for Panx-mediated silencing. Indeed, germline specific depletion of dNxf1 significantly impaired the repression of the reporter mediated by tethering λN-Panx, thereby genetically placing dNxf1 at the effector step of transcriptional silencing (Fig. 4e). To rule out any indirect or off-target effects of dNxf1 knockdowns, we introduced either wild-type dNxf1 or a

truncated version missing the RNA binding domains (ΔRBD=RRM+LRR) as transgenic flies. As

expected, wild-type dNxf1 suppressed the transposon overexpression caused by the shRNA

targeting the 5' untranslated region of endogenous dNxf1 (Fig. 4f). Importantly, the truncated dNxf1

lacking the RNA cargo binding domains (ΔRBD) maintained the ability to rescue transposon

silencing (Fig. 4f). Since the RBD of dNxf1 is required for RNA export[16], our results strongly suggest

a specialized involvement of dNxf1 in transposon silencing. Next, we sought to directly visualize

potential changes in RNA localization by using a rapamycin-inducible tethering system (Fig. 4g and

Extended Data Fig. 13). GFP mRNAs containing 10x copies of BoxB binding sites tethered with

λN-FKBP are mostly localized in the cytoplasm (Fig. 4g and Movie S5-12). Upon Rapamycin

treatment, λN-FKBP dimerizes with FRB-dNxf2 fusion protein, allowing tethering of dNxf2 to the

GFP mRNAs. Intriguingly, GFP mRNAs start to accumulate at nuclear peripheries upon binding of

FRB-dNxf2 (Fig. 4g). The effect is specific to dNxf2 since FRB alone failed to cause any change.

Given this, we named this multi-protein complex as Pandas (Panx-dNxf2 dependent TAP/p15

silencing). Our data raise the possibility that deterring the function of dNxf1 in transposon RNA

export may be a key intermediate event in piRNA-guided silencing (Extended Data Fig. 14). In the

absence of dNxf2, Panx failed to efficiently suppress either transposons or the tethered reporters,

despite the overexpression of λN-Flag-Panx (Extended Data Fig. 3a-e). In this regard, λN-Flag-

Panx was unable to stay bound to the de-repressed transposon chromatin as measured by Flag

ChIP-qPCR (Extended Data Fig. 13c). Similarly, the Flag-ChIP signals over the tethered reporter

were also diminished even when λN-Flag-Panx was constitutively tethered to the nascent RNAs.

This is consistent with the observation that direct tethering of HP1a to nascent RNAs via λN/BoxB

system failed to install silencing[7]. This is in stark contrast to HP1a's ability to induce

heterochromatin formation if tethered via DNA[27-30]. Our data provided a possible explanation why

RNA tethering of HP1a could not induce silencing. Therefore, sequestering nascent transposon

transcripts within nucleus might be important to fully establish heterochromatin and enforce

silencing. We believed that, by removing dNxf2, we had uncovered an intermediate state of

silencing during heterochromatin formation.

Like any coding mRNA, if not restrained by Piwi-piRNAs, transposon transcripts would likewise

be transported into the cytoplasm by the dNxf1/dNxt1 exporting machinery[14,21]. In piRNA-guided

TGS, dNxf2/dNxt1 may function together with Panx as a stable complex to counteract this process

(Extended Data Fig. 13). Our structure provided mechanistic insights into why dNxf2[UBA] prefers to

bind the silencing factor Panx rather than the FG repeats of NPCs (Extended Data Fig. 5).

Accordingly, part of dNxf2's silencing function may be hijacking the RNA export machinery and

repurposing dNxf1 into a "dead-end" complex, hence trapping transposon transcripts within the

nucleus (Fig. 4g and Extended Data Fig. 12). Dam-ID showed that both Piwi and NPCs contacts

chromatin at the same regions[31]. In this regard, dNxf1 can localize to nuclear peripheries in which

most constitutive heterochromatin resides[13,32-34]. The mechanism of how dNxf2 prevents dNxf1

from exporting transposon transcripts requires further investigation. The fact that dimerization of

hsNxf1/TAP is important for RNA export raised a possibility that dNxf2 might block dNxf1-mediated

RNA export by preventing its dimerization[22,23]. Nevertheless, our data indicated that sequestering

transposons to nuclear peripheries via the Pandas complex may help to establish/maintain their

heterochromatic state (Extended Data Fig. 10 and Movie 1-4)[33,34]. Interestingly, recent evidence

demonstrated that hsNxf1 is required for efficient elongation of RNA polymerase II[35]. It is tempting

to speculate that the Pandas complex might also inhibit transcriptional elongation of transposons

via neutralizing dNxf1. In summary, we uncovered an unexpected role of transposon RNA export

blockage prior to transcriptional gene silencing. Our results will have broader implications for how

RNA metabolism modulates epigenetic gene silencing and heterochromatin formation (Extended

Data Fig. 14).

REFERENCES

1. Ge, D. T. & Zamore, P. D. Small RNA-directed silencing: the fly finds its inner fission yeast? *Curr. Biol.* **23,** R318–20 (2013).
2. Martienssen, R. & Moazed, D. RNAi and heterochromatin assembly. *Cold Spring Harb Perspect Biol* **7,** a019323 (2015).
3. Czech, B. & Hannon, G. J. One Loop to Rule Them All: The Ping-Pong Cycle and piRNA-Guided Silencing. *Trends Biochem. Sci.* **0,** (2016).
4. Ozata, D. M., Gainetdinov, I., Zoch, A., Carroll, D. X. N. O. X. & Zamore, P. D. PIWI-interacting RNAs: small RNAs with big functions. *Nat. Rev. Genet.* 1–20 (2018). doi:10.1038/s41576-018-0073-3
5. Gainetdinov, I., Colpan, C., Arif, A., Cecchini, K. & Zamore, P. D. A Single Mechanism of Biogenesis, Initiated and Directed by PIWI Proteins, Explains piRNA Production in Most Animals. *Mol. Cell* **71,** 775–790.e5 (2018).
6. Yu, Y. *et al.* Panoramix enforces piRNA-dependent cotranscriptional silencing. *Science* **350,** 339–342 (2015).
7. Sienski, G. *et al.* Silencio/CG9754 connects the Piwi-piRNA complex to the cellular heterochromatin machinery. *Genes Dev.* **29,** 2258–2271 (2015).
8. Sienski, G., Dönertas, D. & Brennecke, J. Transcriptional silencing of transposons by Piwi and maelstrom and its impact on chromatin state and gene expression. *Cell* **151,** 964–980 (2012).
9. Czech, B., Preall, J. B., McGinn, J. & Hannon, G. J. A transcriptome-wide RNAi screen in the Drosophila ovary reveals factors of the germline piRNA pathway. *Mol. Cell* **50,** 749–761 (2013).
10. Muerdter, F. *et al.* A genome-wide RNAi screen draws a genetic framework for transposon control and primary piRNA biogenesis in Drosophila. *Mol. Cell* **50,** 736–748 (2013).
11. Handler, D. *et al.* The genetic makeup of the Drosophila piRNA pathway. *Mol. Cell* **50,** 762–777 (2013).
12. Guruharsha, K. G. *et al.* A protein complex network of Drosophila melanogaster. *Cell* **147,** 690–703 (2011).
13. Herold, A., Klymenko, T. & Izaurralde, E. NXF1/p15 heterodimers are essential for mRNA nuclear export in Drosophila. *RNA* **7,** 1768–1780 (2001).
14. Katahira, J. Nuclear export of messenger RNA. *Genes (Basel)* **6,** 163–184 (2015).
15. Port, F., Chen, H.-M., Lee, T. & Bullock, S. L. Optimized CRISPR/Cas tools for efficient germline and somatic genome engineering in Drosophila. *Proc. Natl. Acad. Sci. U.S.A.* **111,** E2967–76 (2014).
16. Braun, I. C., Herold, A., Rode, M. & Izaurralde, E. Nuclear export of mRNA by TAP/NXF1 requires two nucleoporin-binding sites but not p15. *Mol. Cell. Biol.* **22,** 5405–5418 (2002).
17. Gu, J. *et al.* GoldCLIP: Gel-omitted Ligation-dependent CLIP. *Genomics Proteomics Bioinformatics* **16,** 136–143 (2018).
18. Savic, D. *et al.* CETCh-seq: CRISPR epitope tagging ChIP-seq of DNA-binding proteins. *Genome Res.* **25,** 1581–1589 (2015).
19. Iwasaki, Y. W. *et al.* Piwi Modulates Chromatin Accessibility by Regulating Multiple Factors Including Histone H1 to Repress Transposons. *Mol. Cell* **63,** 408–419 (2016).
20. Radion, E. *et al.* Key role of piRNAs in telomeric chromatin maintenance and telomere nuclear positioning in Drosophila germline. *Epigenetics Chromatin* **11,** 40 (2018).
21. Chambeyron, S. *et al.* piRNA-mediated nuclear accumulation of retrotransposon transcripts in the Drosophila female germline. *Proc. Natl. Acad. Sci. U.S.A.* **105,** 14964–14969 (2008).
22. Matzat, L. H., Berberoglu, S. & Lévesque, L. Formation of a Tap/NXF1 homotypic complex is mediated through the amino-terminal domain of Tap and enhances interaction with

nucleoporins. *Mol. Biol. Cell* **19,** 327–338 (2008).

23. Aibara, S., Katahira, J., Valkov, E. & Stewart, M. The principal mRNA nuclear export factor NXF1:NXT1 forms a symmetric binding platform that facilitates export of retroviral CTE-RNA. *Nucleic Acids Res.* **43,** 1883–1893 (2015).

24. Golubkova, E. V. *et al.* Dm nxf1/sbr gene affects the formation of meiotic spindle in female Drosophila melanogaster. *Chromosome Res.* **17,** 833–845 (2009).

25. Czech, B. & Hannon, G. J. A Happy 3' Ending to the piRNA Maturation Story. *Cell* **164,** 838–840 (2016).

26. Khurana, J. S. & Theurkauf, W. piRNAs, transposon silencing, and Drosophila germline development. *J. Cell Biol.* **191,** 905–913 (2010).

27. Danzer, J. R. & Wallrath, L. L. Mechanisms of HP1-mediated gene silencing in Drosophila. *Development* **131,** 3571–3580 (2004).

28. Hines, K. A. *et al.* Domains of heterochromatin protein 1 required for Drosophila melanogaster heterochromatin spreading. *Genetics* **182,** 967–977 (2009).

29. Li, Y., Danzer, J. R., Alvarez, P., Belmont, A. S. & Wallrath, L. L. Effects of tethering HP1 to euchromatic regions of the Drosophila genome. *Development* **130,** 1817–1824 (2003).

30. Azzaz, A. M. *et al.* Human heterochromatin protein 1α promotes nucleosome associations that drive chromatin condensation. *J. Biol. Chem.* **289,** 6850–6861 (2014).

31. Ilyin, A. A. *et al.* Piwi interacts with chromatin at nuclear pores and promiscuously binds nuclear transcripts in Drosophila ovarian somatic cells. *Nucleic Acids Res.* **45,** 7666–7680 (2017).

32. Kerkow, D. E. *et al.* The structure of the NXF2/NXT1 heterodimeric complex reveals the combined specificity and versatility of the NTF2-like fold. *J. Mol. Biol.* **415,** 649–665 (2012).

33. van Steensel, B. & Belmont, A. S. Lamina-Associated Domains: Links with Chromosome Architecture, Heterochromatin, and Gene Repression. *Cell* **169,** 780–791 (2017).

34. Towbin, B. D., Meister, P. & Gasser, S. M. The nuclear envelope--a scaffold for silencing? *Curr. Opin. Genet. Dev.* **19,** 180–186 (2009).

35. Chen, S. *et al.* The mRNA Export Receptor NXF1 Coordinates Transcriptional Dynamics, Alternative Polyadenylation, and mRNA Export. *Mol. Cell* **74,** 118–131.e7 (2019).

ACKNOWLEDGEMENTS

We thank Shanghai Science Research Center for their instrumental support and technical assistance. We thank the staff from BL19U1 beamline at Shanghai Synchrotron Radiation Facility (SSRF) for assistance during data collection. We thank Ms. Shuoguo Li from Center for Biological Imaging (CBI), Institute of Biophysics, Chinese Academy of Science for her help of taking and analyzing SIM images. This work was supported in part by grants from the Ministry of Science and Technology of China (2017YFA0504200 to Y.Y.) and by grants from the National Natural Science Foundation of China (91640102 and 31870741 to Y.H.; 91640105 and 31770875 to Y.Y.), and the Chinese Academy of Sciences Facility-based Open Research Program and the State Key Laboratory of Molecular Biology.

AUTHOR CONTRIBUTIONS

Y.Y. and Y.H. conceived the project and wrote the manuscript. K.Z. constructed the dNxf2-Halo knock-in OSCs and established the dNxf2 mutant. K.Z., P.X., W.W.L. and X.H.L. performed co-immunoprecipitations, tethering assays, transgenic fly constructions, RNA-seq, RT-qPCR experiments. S.C. and X.Y. performed structural studies, beta-gal activity assays, and ITC experiments; K.Z., Z.J., F.Q., P.Z., S.W., X.O., J.Q.G. and P.X. performed cloning. S.C. and X.L. performed Y2H assays. K.Z. and X.O. performed FACS analysis. N.M. preformed FISH and RNAi experiments. M.W. and Y.Z. performed bioinformatics analysis. Y.Z.W., J.M. and H.C. provided critical reagents and advice; All authors discussed the results and commented on the manuscript.

COMPETING INTERESTS

The authors declare that there is no conflict of interests.

MATERIALS & CORRESPONDENCE

All materials and correspondence requesta should be addressed to yuyang@ibp.ac.cn.

Figure 1

Figure 1. dNxf2 functions as a cofactor of Panx in the piRNA pathway.

a, Western blots showing co-immunoprecipitation of endogenous dNxf2 with GFP-Panx from ovary lysates. Two different rabbit polyclonal monospecific dNxf2 (dNxf2-N and dNxf2-C) antibodies were used to detect endogenous dNxf2. **b**, Halo-ligand staining and western blots showing co-immunoprecipitation of endogenous Panx with dNxf2-Halo driven by the endogenous dNxf2 promoter from OSC lysates. A rabbit polyclonal monospecific Panx antibody was used to detect endogenous Panx. Left panel shows depletion of dNxf2-Halo proteins in the unbound sample by Halo beads, and the anti-Tubulin blots serve as loading controls; Right panel shows endogenous Panx proteins. **c**, The effects of germline (*nanos*-GAL4) knockdown of the indicated genes on Renilla-normalized Firefly luciferase activity of the reporter while tethering λN-Panx. For comparison, the relative value of the attp2 control was used for normalization. Data show mean ± s.d. (n = 15; *P = 1.41387E-07). **d**, Comparison of steady-state RNA levels are shown as reads per million (rpm) mapping to the sense strand of each transposon consensus from the *nanos*-GAL4 driven knockdown for dNxf2 (*Y* axis) versus control (*X* axis). Dashed lines indicate two-fold changes. The average of two replicates is shown. KD = knockdown. Red dots indicate transposon elements with significant changes. **e**, Comparison of steady-state RNA levels (RNA-seq; shown as RPM) mapping to the sense strands of each transposon consensus from the *nanos*-GAL4 driven knockdowns of the indicated genes. Red dots indicate transposon elements with significant changes from **d**. **f**, Heat map displaying steady-state RNA levels (RNA-seq) as reads per million (rpm) for top 70 detected transposons from the *nanos*-GAL4 driven knockdowns of the indicated genes in a blue-white scale.

Figure 2

Figure 2. The UBA domain of dNxf2 interacts with Panx directly.

a, Domain architectures of Panx, dNxf2, dNxf1, and dNxt1 are shown. Numbers above the diagrams correspond to amino acid residues of each protein. Domain names are abbreviated within respective colored region. **b-e**, Y2H assays mapping the interacting regions between *Drosophila* Nxf1/2 and Panx or dNxt1. Interactions were determined by either measuring the beta-galactosidase activity produced by the reporter gene or growth on YSD media lacking the indicated essential amino acid or Uracil. Data are averages of three independent experiments (n=3). Proteins or fragments shown above the dashed line are used as preys in the assays. **b**, Y2H assays mapping regions of *Drosophila* Nxf1/2 that interact with Panx. **c**, Y2H assays mapping regions of *Drosophila* Nxf1/2 that interact with dNxt. **d**, Yeast three hybrid determining the requirement of dNxt1 for Nxf2:Panx interaction. **e**, Y2H assays mapping minimum regions of Panx that interact with dNxf2^{UBA}. **f**, Left panel shows the size exclusive chromatography profile of the NTF2 and UBA domains of dNxf2 forming heterodimers with dNxt1 and Panx^{NIR} in solution, respectively. A dNxf2 fragment spanning the NTF2 and UBA domains that is covalently linked to Panx^{NIR} forms a stable complex with dNxt1. Right panel, SDS-PAGE shows the components of the peak in the elution profile. Color schemes used for the three complexes are indicated in the key. **g**, Quantification of the dissociation constant for the interaction between dNxf2^{UBA} and Panx^{NIR} as measured by an isothermal titration calorimetry assay.

Figure 3

Figure 3. Structure of dNxf2^{UBA} in complex with Panx^{NIR}.

a, Schematic of interacting region between dNxf2 and Panx. Numbers above or below the diagrams correspond to amino acid residues of dNxf2 or Panx, respectively. Domain names are abbreviated within respective colored region. NIR, d<u>N</u>xf2 <u>i</u>nteracting <u>r</u>egion. **b**, Left, cartoon representation of dNxf2^{UBA} in complex with Panx^{NIR}. The dNxf2^{UBA} and Panx^{NIR} fragment are colored in wheat and teal, respectively. Right, a view rotated 90° around the vertical axis. **c**, Electrostatic potential analysis showing the Panx-binding surface of dNxf2^{UBA}. Panx^{NIR} is shown in cartoon mode. **d**, A detailed view showing the interactions between dNxf2^{UBA} and Panx^{NIR}. Key residues involved in binding are shown in sticks. Close-up views of hydrophobic interactions between dNxf2^{UBA} and Panx^{NIR} are shown on the right. **e**, Y2H assays measuring the binding of wild-type or mutant dNxf2^{UBA} with Panx^{NIR}. Mutations of key residues are indicated along the bars. **f**, Western blots and Halo-ligand staining showing co-immunoprecipitation of GFP-tagged Panx or its NIR deletion mutant (ΔNIR) with Halo-tagged dNxf2 or its F826A/I827A double mutant from OSC cells. GFP serves as a negative control. **g**, Effects of the indicated λN fusion proteins or a non-tethering control (GFP-dNxf2) on luciferase activity of the reporters integrated into the attP2 landing site. Data show mean ± s.d. (n = 15; *P = 1.41387E-07). **h**, The effects of λN-dNxf2 tethering on luciferase activity of reporters with increasing number of BoxB sites. All reporters are integrated into the same genomic locus (attP2 landing site). Fold repression is calculated as total protein-normalized Firefly luciferase luminescent values of the control (no tethering) divided by that of the indicated experiments. Data show mean ± s.d. (n = 15; *P = 1.41387E-07).

Figure 4

Figure 4. The Pandas complex is required for piRNA-guided transposon silencing.

a, Comparison of GoldCLIP-seq are shown as reads per million (rpm) mapping to the sense strand of each transposon consensus for Maelstrom (*X* axis) versus a LacZ control knockdown (*Y* axis). A mammalian spike-in was used to normalize different samples. Dashed lines indicate two-fold changes. The pooled reads from two replicates are shown. Red dots indicate Piwi-targeted transposons, blue dots indicate H1-dependent transposons and green dots indicate other transposons. **b**, Western blots and Halo-ligand staining showing co-immunoprecipitation of GFP-tagged dNxf2 with Halo-tagged dNxf1 from OSC cells. ΔNTF2, Halo-dNxf1 lacking the NTF2 domain, and GFP serves as a negative control. **c**, Western blots and Halo-ligand staining showing co-immunoprecipitation of Halo-tagged dNxf2-NTF2 domain with different domain truncations of GFP-tagged dNxf1 from OSC cells. ΔRBD, dNxf1 lacking the N-terminus RRM and LRR domains; GFP serves as a negative control. **d**, SDS-PAGE showing pulldown results of the $Nxf2^{NTF2+UBA}$-$(GS)4-Panx^{NIR}$/dNxt1 complex by either GST-tagged dNxf1-NTF2 or dNxf1-(NTF2+UBA) respectively, compared to a GST control. **e**, The effects of germline specific knockdown (nanos-GAL4) of the indicated genes on Renilla-normalized Firefly luciferase activity of the reporter while tethering λN-Panx. For comparison, the results from Fig. 1C (Egg, attp2 and Zuc) were shown here again. Data show mean ± s.d. (*n* = 15; *P = 1.41387E-07). **f**, RT-qPCR results showing the effects of germline (*nanos*-GAL4) knockdown of the indicated genes in the presence or absence of different transgenes. Fold changes are calculated as rp49-normalized RNA levels divided by that of the corresponding controls. Mean values ± s.d. from 3 independent experiments are shown. **g**, SIM super-resolution microscopy of RNA FISH. Data showing the comparison of the localizations of the reporter mRNAs tethered with λN-FKBP, upon transient recruitment of either FRB-dNxf2 or FRB alone when treated with rapamycin for 2 hours to induce FKBP:FRB dimerization. DMSO serves as a control. Top panel, RNA signal (red) with DAPI staining; middle, DAPI staining (blue); bottom, RanGAP-GFP fusion protein (green) were shown. The scale bars represent 5 μM in length.

a Physical interactions (DPiM)

b

c GFP Pull down

d Gene expression across different tissues (FlyAtlas)

e

Extended Data Figure 2. dNxf2 mutant resembles Panx mutant. a, A schematic of CRISPR/Cas9-induced mutation
in the open reading frame of dNxf2 is shown. Arrowheads indicate the cleavage sites targeted by the gRNA with the
corresponding PAM site highlighted in light blue. The resulting mutant is shown below the wild-type sequence with the
mutated/deleted region shown in red. **b,** Vials showing pupae from equal number of either heterozygous or mutant
dNxf2 female flies crossed with OregonR males. **c,** Representative immunostaining of endogenous Piwi protein (green)
from stage 5 ovaries of either heterozygous or mutant dNxf2. DAPI staining (blue) was used to indicate nuclei. **d,**
Western blots showing protein expression level of endogenous dNxf2, Panx and Piwi from either heterozygous or
mutant dNxf2 ovary lysates. Beta-tubulin serves as a loading control. **e,** RT-qPCR results showing relative steady-state
RNA levels of the indicated transposons for the mutant or heterozygous dNxf2 ovaries, compared to the wildtype control
(w1118). Fold changes are calculated as *rp49*-normalized RNA levels divided by that of the corresponding controls.
Mean values ± s.d. from 3 independent experiments are shown. **f,** Comparison of steady-state RNA levels are shown
as reads per million (rpm) mapping to the sense strands of each transposon consensus from dNxf2 mutant versus
w1118 wild-type fly ovaries. Dashed lines indicate two-fold changes. The average of two replicates is shown. Red dots
indicate transposon elements with significant changes. **g,** Comparison of steady-state RNA levels (RNA-seq; shown
as RPM) mapping to the sense strands of each transposon consensus from dNxf2 mutant versus Panx mutant fly
ovaries. Red dots indicate transposon elements with significant changes from **f**.

363
Extended Data Figure 3. dNxf2 is required for Panx-mediated transposon silencing. a, The expression of the
total protein-normalized Firefly luciferase activity of the reporter under the indicated background (heterozygous control
or transheterozygous mutant) while overexpressing λN-Flag-Panx driven by a Ubiquitin promoter. Data show
mean ± s.d. (n = 15; *P = 1.41387E-07). **b,** Western blots showing protein expression level of endogenous dNxf2, Panx
from either heterozygous or mutant dNxf2 ovary lysates while overexpressing λN-Flag-Panx driven by a Ubiquitin
promoter. Beta-tubulin serves as a loading control. **c,** RT-qPCR results showing relative steady-state RNA levels of
the indicated transposons for the mutant or heterozygous dNxf2 ovaries, compared to the wildtype control while
overexpressing λN-Flag-Panx driven a Ubiquitin promoter. Fold changes are calculated as *rp49*-normalized RNA levels
divided by that of the corresponding controls. Mean values ± s.d. from 3 independent experiments are shown. **d,**
Comparison of steady-state RNA levels are shown as reads per million (rpm) mapping to the sense strands of each
transposon consensus from dNxf2 mutant versus wild-type fly ovaries when overexpressing ubiquitin driven Panx.
Dashed lines indicate two-fold changes. The average of two replicates is shown. Red dots indicate transposon
elements with significant changes. **e,** Vials showing pupae from equal number of either heterozygous or mutant dNxf2
female flies when overexpressing ubiquitin driven Panx, mated with OregonR males. **f,** Schematic of full-length Panx
and its truncated fragments used in the Y2H assays in Fig. 2e. Numbers corresponding to amino acid residues at either
the N- or C-terminus of each protein fragment.

380
381

382
Extended Data Figure 4. dNxf2 Structural analysis of dNxf1^{NTF2} in complex with dNxt1. a, Cartoon representation
of dNxf1 NTF2 domain (orange) in complex with dNxt1 (green). **b,** Electrostatic surface potential representation of
dNxt1, highlighting hydrophobic and electrostatic surfaces continuously from acidic (red) to non-polar (grey), to basic
(blue). Key residues of dNxf1 are shown in stick mode (orange). The corresponding residues in dNxf2 (yellow) are
modelled in Coot by mutation operation based on the dNxf1 structure in this study. **c,** dNxf1 NTF2 (orange) and dNxf2
NTF2 (yellow, model based on dNxf1 structure) are superimposed with human TAP NTF2 domain (cyan, PDB
accession code 1JN5). Key residues that interact with the Phenylalanine-Glycine (FG) peptide (magenta) are labelled
and shown in sticks. **d,** Sequence alignment of the NTF2 domains of dNxf1, dNxf2, human TAP (Q9UBU9), and ceNxf2
(Q9XVS8). The secondary structure annotation of dNxf1 is shown on the top. Residue positions of the reference
sequence (dNxf1) are also indicated above the sequences. Arrows and spirals indicate beta-strands and alpha-helices,
respectively. Red letters indicate similar residue; white letters in red shading indicate identical residues; blue boxes
indicate conserved positions. **e,** Western blots and Halo-ligand staining showing co-immunoprecipitation of GFP-
tagged dNxt1 with Halo-tagged dNxf1 or dNxf2 from OSC cells. GFP serves as a negative control. **f,** Sequence
alignment of the region of Panx that interacted with dNxf2. Residues involved in the interactions are marked by teal
diamond.

Extended Data Figure 5. Structural comparison of the UBA domains from different NXF family proteins. a,
Sequence alignment of the UBA domains of Nxf family proteins. Residues within the dNxf2-Panx interaction interface
are indicated in brown diamond. Residues that blocks the entry of the FG peptides are indicated in green circle.
Residues that are identical are shown in white with red shading. Residues with high similarities are shown in red.
Secondary structural elements are indicated above. **b,** Superimposition of the UBA domains of dNxf2 (wheat), human
Nxf1 (green, PDB ID: 1OAI), and yeast Mex67 (purple, PDB ID: 2KHH). Electrostatic potential analysis showing that
the corresponding surfaces in the UBA domains of dNxf2 (left), human Nxf1 (middle) and yeast Mex67 (right) are

positively charged and negatively charged, respectively. **Fig. 3c** was shown here again for comparison. **c,** Detailed
view, left to right, of the UBA domains of dNxf2, human TAP, yeast Mex67, and dNxf1 (model). The dNxf1 model was
generated by SWISS-MODEL server. In human Nxf1 and yeast Mex67, the FG peptides bind in the concave passage
formed by α2 and α3. **d,** Electrostatic potential analysis showing a closed surface on dNxf2 UBA that blocks the entry
of FG-nup. In the corresponding surface of dNxf2 UBA, the passage for the peptide is blocked (left). Human Nxf1 UBA
(PDB ID: 1OAI) and yeast Mex67 UBA (PDB ID: 2KHH) show a hydrophobic pocket to accommodate the two
phenylalanines of the FxFG peptides (middle). A model of dNxf1 UBA (right) indicates that a peptide binding pocket
may exist on the dNxf1 surface. **e,** Y2H assays show that dNxf1 UBA domain but not dNxf2 UBA interacts with Nup214
FG-repeat region. Interactions were determined by either measuring the beta-galactosidase activity produced by the
reporter gene. Data are averages of three independent experiments (n=3). Proteins or fragments of Nup214 are shown
above the dashed line.
417
418

419
Extended Data Figure 6. dNxf2 specifically binds to transposons *in vivo*. a, Schematic representation of the
GoldCLIP experiments. OSC, Ovarian Somatic Cell. TE, transposon. H, Halo-tag. **b,** RT-qPCR results showing relative
levels of the mdg1 transposon or rp49 for dNxf2-Halo GoldCLIP sample, compared to either no UV or a Halo control.
Fold enrichments are calculated as a mammalian spike-in normalized RNA levels divided by that of the input samples.
For comparison, the value of Halo control was used for normalization. Mean values ± s.d. from 3 independent
experiments are shown. **c,** Data from **b** was redrawn in $\log_2$ scale showing relative enrichment or depletion of the
indicated transcripts, normalized to the Halo alone control. **d,** RT-qPCR results showing relative steady-state RNA
levels of the indicated transposons for OSC from the H1+HP1a double siRNA knockdown, compared to a control siRNA
knockdown. Fold changes are calculated as *rp49*-normalized RNA levels divided by that of the corresponding controls.
Mean values ± s.d. from 3 independent experiments are shown. **e,** A Schematic of the knock-in strategy is displayed.
Agarose gel showing PCR results confirming a correct integration of Halo-tag to the C-terminus of the dNxf2 locus.
PCR primers amplifying the indicated regions are shown on the top and two sgRNAs targeting near the stop codon of
dNxf2 are indicated at the bottom of the schematic. **f,** Western blots and Halo-ligand staining showing depletion of
dNxf2-Halo or Halo alone by Halo pull down. Beta-tubulin blots serve as a loading control. The asterix indicates non-
specific bands stained by Halo-ligand.

435
436 **Extended Data Figure 7. dNxf2 binds to Piwi/piRNA targeted transposons**. **a,** Density plots for dNxf2-Halo
437 GoldCLIP reads from Mael knockdown OSCs (top panel), dNxf2-Halo GoldCLIP reads from LacZ knockdown OSCs
438 (middle panel), and Panx co-immunoprecipitated smRNA reads (bottom panel, from previously published data[20]) over
439 the transposon consensus of mdg1. **b** and **c** are the same as in **a** except for gypsy (**b**) and 297 (**c**).
440

441
442
Extended Data Figure 8. The Pandas complex is required for piRNA-guided transposon silencing. a, Effects of
the indicated λN fusion proteins or a non-tethering control (GFP-dNxf2) on luciferase activity of the reporters integrated
into the attP2 landing site. Data from **Fig. 3g** was shown here again for comparison. Data show mean ± s.d. (n = 15).
b, FACS sorting showing counts and fluorescence of GFP positive cells gated by high RFP expressing OSCs electro-
transfected with the indicated plasmids expressing different λN-fusion proteins. OSC (grey), autofluorescence of OSCs
without any GFP reporter; Tethered (red), GFP reporter OSCs transfected with the indicated expression plasmids;
GFP-10xBoxB (green), GFP reporter cells with no transfection.
450
451

452
Extended Data Figure 9. ChIP-qPCR assays measuring H3K9me3 changes upon loss of dNxf2. a, H3K9me3
mark enrichments at transposons comparing heterozygous and mutant for dNxf2. **b,** H3K9me3 mark enrichments at
the indicated transposons comparing heterozygous and mutant for dNxf2 when overexpressing Panx driven by a
Ubiquitin promoter. Green and brown bar, polyclonal antibody specific to H3K9me3; Black and grey bar, pre-immune
rabbit IgG. Mean values ± s.d. from 3 independent experiments are shown. **c,** H3K9me3 mark enrichments at the
Firefly-10xBoxB reporter tethered with λN-flag-Panx in the same genetic background as in **b**. Schematic of qPCR
primer pair positions is shown on the right.

a

HeT-A / DAPI / Lamin   3D reconstruction

b

HeT-A / DAPI / Lamin

461
462
Extended Data Figure 10. Het-A left nuclear peripheries upon loss of either Panx or dNxf2. a, Left three columns
showing 3D SIM super-resolution microscopy of DNA FISH (HeT-A, green) and Lamin A (red) double staining from
either Panx or dNxf2 heterozygote versus mutant ovaries. Right three columns, estimation of the positioning of
clustered HeT-A signals relative to the nuclear membrane of nurse cells. 10-20 nuclei were quantified for each condition.
The distance was displayed in a blue-red scale with red indicating longer distance between the green dots and nuclear
membrane. **b,** Projections of the 3D videos on the *XY* surface either in combination (bottom right) or in split channels
for DAPI (blue, upper right), Lamin (red, bottom left) and HeT-A (green, upper left) on the selected ovaries from **a**. See
also Supplementary Movies 1-4.

471
472

473
Extended Data Figure 11. dNxf2 interacts with dNxf1 in *Drosophila* cell lines. a, Western blots and Halo-ligand
staining showing co-immunoprecipitation of GFP-tagged various truncated dNxf2 with Halo-tagged full length dNxf1
from OSC cells. ΔRBD, dNxf2 lacking N-terminus RNA binding domains; NTF2, dNxf2^{NTF2} domain; UBA, dNxf2^{UBA}
domain. **b,** Western blots and Halo-ligand staining showing co-immunoprecipitation of GFP-tagged dNxf1^{NTF2} domain
with Halo-tagged various truncated dNxf1 from S2 cells. ΔRBD, dNxf1 lacking all N-terminus RNA binding domains;
NTF2, dNxf1^{NTF2} domain; UBA, dNxf1^{UBA} domain. GFP serves as a negative control. **c,** The same as in B except the
experiments were done using OSC cells.
481
482

483
484
Extended Data Figure 12. dNxf1 directly interacts with the NTF2 domain of dNxf2 via either NTF2 or UBA
domains. a, dNxf2$^{NTF2+UBA}$, dNxt1, and Panx^{NIR} form a ternary complex in solution as shown on the size exclusion
chromatography profile. Left panel, OD280 trace; Right panel, SDS-PAGE shows the components of the peak in the
elution profile. **b-d,** GST pull-down assays showed the direct interactions between dNxf1 and dNxf2. Each of 60 μg of
purified GST-dNxf1^{NTF2}/dNxt1 (**b**), GST-dNxf1^{UBA} (**c**), GST-dNxf1$^{NTF2+UBA}$/TS-dNxt1 complex (**d**) and GST protein was
immobilized on GSH beads, respectively. Increasing amounts of dNxf2^{NTF2}/dNxt1 complex (0, 15, 30, 60 μg) were
added and incubated. The beads were washed and aliquots of the bound fractions (8%) were analyzed by SDS-PAGE.
Each input protein of 2 μg was loaded on SDS-PAGE as well. Positions of molecular weight markers are indicated on
the left in kDa.

494
Extended Data Figure 13. Rapamycin inducible tethering system. a, Schematic representation of a rapamycin-
inducible system for λN-FKBP:FRB tethering. **b,** FACS sorting showing counts and fluorescence of GFP positive cells
gated by high RFP expressing OSCs (as in **Extended Data Fig. 7b**) transfected with the plasmids expressing both λN-
FKBP and the indicated FRB-fusion proteins. 24 hours after transfection, cells were treated with rapamycin for
additional 24 hours. Two independent biological replicates were shown. **c,** Flag-Panx enrichments at the indicated
transposons comparing heterozygous and mutant for dNxf2 when overexpressing Panx driven by a Ubiquitin promoter.
Green and brown bar, monoclone antibody specific to Flag (M2); Black and grey bar, pre-immune mouse IgG. Mean
values ± s.d. from 3 independent experiments are shown. **d,** Flag-Panx enrichments at the Firefly-10xBoxB reporter
tethered with λN-flag-Panx in the same genetic background as in (**c**). Schematic of qPCR primer pair positions is shown
in **Extended Data Fig. 9c**.
505
506

507
508
509 **Extended Data Figure 14. A model for the Pandas (Panoramix-dNxf2 dependent TAP/p15 silencing) complex**
510 **in piRNA-guided silencing, preventing transposon RNA export when targeted by Piwi:piRNAs.**
511
512
513
514