1 **Long title:**
2 Neogenin-1 distinguishes between myeloid-biased and balanced *Hoxb5*[+] mouse long-term
3 hematopoietic stem cells
4
5 **Short title:**
6 NEO1 marks lineage-biased hematopoietic stem cells
7
8 **Authors**
9 Gunsagar S. Gulati[1,*], Monika Zukowska[1,2,*], Joseph Noh[1], Allison Zhang[1], Rahul Sinha[1],
10 Benson M. George[1], Daniel J. Wesche[1], Irving L. Weissman[1,3,#], Krzysztof Szade[1,2,#]
11
Affiliations
[1] Institute for Stem Cell Biology and Regenerative Medicine, Stanford University, Stanford, CA,
94305, USA.
[2] Department of Medical Biotechnology, Faculty of Biochemistry, Biophysics and
Biotechnology, Jagiellonian University, Krakow, Poland.
[3] Department of Developmental Biology, Stanford University School of Medicine, Stanford, CA,
94305, USA.
19
20
21 *These authors contributed equally.
22
23 **[#]Co-senior authors to whom correspondence should be addressed:**
24 Krzysztof Szade, Ph.D.
25 krzysztof.szade@uj.edu.pl
26
27 Irving L. Weissman, M.D.
28 irv@stanford.edu
29 Tel: 650-723-6520
30

31 **Keywords:** hematopoietic stem cell, transplantation, myeloid-bias, aging, Neogenin-1, blood

ABSTRACT

Hematopoietic stem cells (HSCs) self-renew and generate all blood cells. Recent studies with single-cell transplants (1-3) and lineage tracing (4, 5) suggest that adult HSCs are diverse in their reconstitution and lineage potentials. However, prospective isolation of these subpopulations has remained challenging. Here, we identify Neogenin-1 (NEO1) as a unique surface marker on a fraction of mouse HSCs labeled with *Hoxb5*, a specific reporter of long-term HSCs (LT-HSCs) (6). We show that NEO1+*Hoxb5*+ LT-HSCs expand with age and respond to myeloablative stress, while NEO1−*Hoxb5*+ LT-HSCs exhibit no significant change in number. NEO1+*Hoxb5*+ LT-HSCs are more often in the G_2/S cell cycle phase compared to NEO1−*Hoxb5*+ LT-HSCs in both young and old bone marrow. Upon serial transplantation, NEO1+*Hoxb5*+ LT-HSCs exhibit myeloid-biased differentiation and reduced reconstitution, while NEO1−*Hoxb5*+ LT-HSCs are lineage-balanced and stably reconstitute recipients. Gene expression comparison reveals increased expression of cell cycle genes and evidence of lineage-priming in the NEO1+ fraction. Finally, transplanted NEO1+*Hoxb5*+ LT-HSCs rarely generate NEO1−*Hoxb5*+ LT-HSCs, while NEO1−*Hoxb5*+ LT-HSCs repopulate both LT-HSC fractions. This supports a model in which dormant, balanced, NEO1−*Hoxb5*+ LT-HSCs can hierarchically precede active, myeloid-biased NEO1+*Hoxb5*+ LT-HSCs.

SIGNIFICANCE STATEMENT

Hematopoietic stem cells (HSCs) are rare cells that have the unique ability to regenerate themselves and produce all blood cells throughout life. However, HSCs are functionally heterogeneous and several studies have shown that HSCs can differ in their contribution to major blood lineages. In this study, we discovered that the surface marker, Neogenin-1, can divide mouse HSCs into two subpopulations—one that is more active but biased towards producing myeloid cells and another that is more dormant and capable of equally producing all blood lineages. Neogenin-1 reveals the diversity and hierarchical relationship of HSCs in the mouse bone marrow, enables the prospective isolation of myeloid-biased and balanced HSCs, and opens opportunities to do the same in humans.

INTRODUCTION

The hematopoietic system is hierarchically organized into distinct cell types and cellular states with unique functions and regenerative potentials ([7](#)). Residing at the apex of this hierarchy is the hematopoietic stem cell – the master orchestrator of all blood and immune development, maintenance, and regeneration. HSCs have the unique ability to self-renew and give rise to all major lineages of blood and immune cells throughout life. Over the years, combinations of surface markers ([8-12](#)) and reporter genes ([6](#), [13-15](#)) have refined the definition of mouse HSCs and enabled the purification of long-term hematopoietic stem cells (LT-HSCs), a refined subset of HSCs capable of serially reconstituting irradiated recipients in a transplantation model. Recently, we identified *Hoxb5* as a specific marker of long-term repopulating cells and generated a *Hoxb5*-mCherry reporter mouse strain for the prospective isolation of these cells ([6](#)). We demonstrated that the *Hoxb5*-mCherry reporter significantly improves the precise isolation of serially transplantable LT-HSCs compared to immunophenotypically defined HSCs (phenotypic HSCs, or pHSCs). As previously shown, only 7-35% of pHSCs are *Hoxb5*-mCherry[+] and the potential for long-term multilineage reconstitution is restricted to this fraction ([6](#)).

Although *Hoxb5*[+] LT-HSCs are multilineage contributing, self-renewing cells ([6](#)), the functional heterogeneity within this compartment has not yet been characterized. Understanding the composition of LT-HSCs may offer valuable insights into the mechanism of HSC expansion with age, as well the competition of diverse HSCs for bone marrow niches ([16](#), [17](#)). On a per cell basis, HSCs from older mice exhibit biased differentiation towards myeloid lineages and reduced stem cell activity, presumably due to weaker responses to SDF-1 ([17](#), [18](#)) and poorer engraftability of HSCs in G_1 ([19](#)) and $S/G_2/M$ ([20](#)). These aged HSCs may arise from either the cell-intrinsic transition of balanced to myeloid-biased LT-HSCs or the clonal expansion of pre-existing fractions of myeloid-biased LT-HSCs ([16](#), [17](#), [21-23](#)). Several studies support the presence of pre-existing myeloid-biased LT-HSCs by demonstrating that myeloid-biased subpopulations of LT-HSCs in young, healthy mice respond to environmental challenges, such as inflammation and infection ([24](#), [25](#)). Results from lineage tracing with genetic barcodes ([4](#), [5](#)) and single cell transplants of LT-HSCs also support the notion of inherent functional diversity among long-term repopulating HSCs ([1-3](#)). However, these studies did not identify markers to prospectively isolate distinct sub-populations of HSCs. Other groups have identified markers, such as CD150 ([21](#)), CD41 ([26](#)), vWF ([27](#)), and CD61 ([24](#)) that enrich for self-renewing lineage-biased subpopulations of HSCs. However, these markers were shown to segregate fractions of pHSCs, which contain both short-term, *Hoxb5*[−] and long-term, *Hoxb5*[+] HSCs. Our previous study showed that *Hoxb5*[−] pHSCs are homogenously lymphoid-biased ([6](#)), but the diversity of *Hoxb5*[+] LT-HSCs has not yet been fully explored. Therefore, we sought to interrogate the heterogeneity among purified *Hoxb5*[+] LT-HSCs and identify a strategy to prospectively isolate these cells with phenotypic markers.

Here, we find that Neogenin-1 (NEO1), a transmembrane receptor of the immunoglobulin family ([28](#)), is expressed on a fraction of *Hoxb5*[+] LT-HSCs and decreases with differentiation. Although NEO1 has been extensively investigated as a receptor for axon guidance ([29](#), [30](#)), neuronal survival ([31](#)), skeletal myofiber differentiation ([32](#)), intracellular iron homeostasis ([33](#)), mammary epithelial development ([34](#)), and endothelial migration ([35](#)), as of yet, little is known about the cells that express this marker in the bone marrow. We find that NEO1[+]*Hoxb5*[+] LT-HSCs represent a myeloid-biased subset of LT-HSCs that responds to myeloablative stress and expands with age. Contrastingly, NEO1[−]*Hoxb5*[+] LT-HSCs exhibit greater reconstitution potential, balanced lineage output, and a more quiescent cell cycle status compared to NEO1[+]*Hoxb5*[+] LT-HSCs. After transplant, NEO1[−]*Hoxb5*[+] LT-HSCs give rise to NEO1[+]*Hoxb5*[+] LT-HSCs, but the reverse transition is rarely observed. We, therefore, propose a model of early

112 long-term hematopoiesis in which balanced, quiescent LT-HSCs self-renew and generate long-
113 term myeloid-biased LT-HSCs in response to stress and during the course of aging.
114
115 **RESULTS**
116
Neogenin-1 (NEO1) marks a subpopulation of mouse *Hoxb5*⁺ LT-HSCs and human HSCs
Functional heterogeneity within *Hoxb5*⁺ LT-HSCs is poorly understood. To identify surface
candidates that fractionate *Hoxb5*⁺ LT-HSCs, we first pattern-searched 64 microarray
expression profiles of 23 distinct mouse hematopoietic cell types ([36]) for (1) genes annotated to
code for cell surface proteins (GO Biological Process: 0009986) and (2) genes specifically
expressed in HSCs compared to downstream cell types (**Fig. 1A**; *SI Appendix*, Table S1). We
found several known HSC-specific markers, including *Robo4* ([37]), *Slamf1* ([11]), *Ly6a* ([8])*, Vwf*
([27])*, TEK* ([38])*,* and a member of the *Gpcr5* family ([39]), validating the utility of our approach. We
also identified several novel markers of HSCs that have not been previously reported (**Fig. 1A**).
Among the top 3 most enriched surface markers on HSCs, Neogenin-1 (*Neo1*) was more highly
expressed on HSCs compared to the other two candidates (**Fig. 1 *A* and *B***). Single-cell RNA-
sequencing data of hematopoietic stem and progenitor cells validated the enriched expression
of *Neo1* in LT-HSCs compared to downstream short-term HSCs and progenitors (*SI Appendix,*
Fig. S1). *Neo1* was also expressed on subsets of bone marrow stromal and endothelial cells
and demarcated the contours of trabecular bone (*SI Appendix,* Fig. S2), suggesting its
expression is not restricted to hematopoietic cells in the bone marrow.
133
We next used flow cytometry to measure the relative protein levels of NEO1 on the surface of 2-
to-3-month-old early hematopoietic progenitors, multipotent progenitor subset A (MPPa),
multipotent progenitor subset B (MPPb), phenotypic HSCs defined as Lin⁻c-KIT⁺SCA1⁺CD48⁻
FLK2⁻CD150⁺CD34⁻ (hereafter referred to as pHSCs), and two populations among pHSCs,
including *Hoxb5*⁺ LT-HSCs and *Hoxb5*⁻ short-term HSCs (ST-HSCs) (**Fig. 1C**; *SI Appendix,* Fig.
S3). Consistent with its gene expression, the relative protein levels of NEO1 and the frequency
of NEO1⁺ cells progressively decreased with differentiation. NEO1⁺ cells comprised a
significantly higher fraction of *Hoxb5*⁺ LT-HSCs compared to downstream cells (**Fig. 1 *C* and
*D***). NEO1 was also expressed on a fraction of long-term reconstituting Lin⁻CD34⁺CD38⁻
CD45RA⁻CD90⁺ HSCs from human bone marrow ([40]), although NEO1 enrichment in human
HSCs was diminished compared to that observed in mouse HSCs (*SI Appendix*, Fig. S4).
145
NEO1⁺*Hoxb5*⁺ LT-HSCs selectively expand with age and respond to myeloablative stress
Previous studies have shown that a subpopulation of pHSCs expands with age ([2], [21]) and
responds to environmental challenge ([24], [25]). However, the effect of aging and stress on LT-
HSCs and their subpopulations have not yet been evaluated. To that end, we first measured the
number and frequency of *Hoxb5*⁺ LT-HSCs (**Fig. 2 *A* to *C***; *SI Appendix,* Fig. S5) and NEO1⁺
and NEO1⁻ fractions (**Fig. 2 *D* to *F***; *SI Appendix,* Fig. S5) in 2-month, 5-month, 13-month, and
22-month-old bone marrow. Consistent with the overall expansion of pHSCs (*SI Appendix,* Fig.
S5 *A* and *B*), we observed that the total number of *Hoxb5*⁺ LT-HSCs and *Hoxb5*⁻ ST-HSCs was
significantly increased (**Fig. 2 *B* and *C***; *SI Appendix,* Fig. S5 *C* to *E*). The frequency of *Hoxb5*⁺
LT-HSCs among pHSCs, although on average higher in bone marrow from older (13-month-old
and 22-month-old) than younger (2-month-old and 5-month-old) mice, was highly variable in
aged mice (**Fig. 2*A***; *SI Appendix,* Fig. S5*C*).
158
Despite the variable expansion of *Hoxb5*⁺ LT-HSCs, the frequency of NEO1⁺ cells among
Hoxb5⁺ LT-HSCs progressively increased with age in a consistent manner (**Fig. 2*D***; *SI*
Appendix, Fig. S5*F*). Fewer than 20% of *Hoxb5*⁺ LT-HSCs in 2-month-old mice expressed
surface NEO1, while more than 80% of 22-month-old *Hoxb5*⁺ LT-HSCs were NEO1⁺ (**Fig. 2*D*;**

SI Appendix, Fig. S5*F*). However, while the number of NEO1$^+$ cells per million whole bone
marrow cells increased with age (**Fig. 2*E***; *SI Appendix,* Fig. S5*G*), the number of NEO1$^-$*Hoxb5*$^+$
LT-HSCs did not significantly change (**Fig. 2*F***; *SI Appendix,* Fig. S5*H*). This suggests that the
NEO1$^+$ fraction selectively expands among *Hoxb5*$^+$ LT-HSCs in the bone marrow, while the
number of NEO1$^-$ *Hoxb5*$^+$ LT-HSCs remains stable with age.

We next evaluated the response of *Hoxb5*$^+$ LT-HSCs and the NEO1$^+$ and NEO1$^-$
subpopulations to myeloablative stress. 4-month-old adult mice were treated with 150 mg/kg 5-
fluorouracil (5-FU) and their bone marrow was analyzed 5 days post-treatment when HSC
proliferation is maximum. ([41]) (**Fig. 2*G***). While the frequency of *Hoxb5*$^+$ and *Hoxb5*$^-$ cells
among pHSCs did not change (**Fig. 2*H***), a significantly higher percentage of *Hoxb5*$^+$ LT-HSCs
were NEO1$^+$ than NEO1$^-$ after treatment (**Fig. 2*I***). This response may be due to the proliferation
of NEO1$^+$*Hoxb5*$^+$ LT-HSCs, differentiation of NEO1$^-$ *Hoxb5*$^+$ to NEO1$^+$*Hoxb5*$^+$, or a combination
of the two.

NEO1$^+$*Hoxb5*$^+$ LT-HSCs are more proliferative than NEO1$^-$*Hoxb5*$^+$ LT-HSCs in young and
old mice
We next asked whether the difference in expansion of NEO1$^+$ versus NEO1$^-$*Hoxb5*$^+$ LT-HSCs
during aging and in response to myeloablative stress can be partially explained by differences in
proliferation ([42]). To address this, we measured the percent of each cell population in G_0, G_1,
and G_2/S by KI-67 and DAPI staining at both 2-to-3 months and 12-to-14 months of age (**Fig. 3**;
SI Appendix, Fig. S6). Consistent with previous reports ([43]), 12-to-14-month-old HSCs were
more often in G_0 compared to 2-to-3-month-old HSCs, suggesting increased exhaustion and
senescence with age in the absence of leukemia (**Fig. 3**; *SI Appendix,* Fig. S6). In support of
the limited expansion of *Hoxb5*$^+$ LT-HSCs among pHSCs with age, we did not observe
significant differences in G_0, G_1, or G_2/S status between *Hoxb5*$^+$ LT-HSCs and *Hoxb5*$^-$ ST-HSCs
(**Fig. 3 *A to C***; *SI Appendix,* Fig. S6). NEO1$^+$ and NEO1$^-$*Hoxb5*$^+$ LT-HSCs also did not differ in
the proportion of G_0 cells in both 2-to-3-month-old and 12-to-14-month-old bone marrow (**Fig.**
3*D*; *SI Appendix,* Fig. S6)

However, NEO1$^+$*Hoxb5*$^+$ LT-HSCs were significantly more often in G_2/S compared to NEO1$^-$
Hoxb5$^+$ LT-HSCs in both young and old bone marrow (**Fig. 3*F***; *SI Appendix,* Fig. S6). Moreover,
in the young bone marrow, there was a significantly smaller percentage of NEO1$^+$*Hoxb5*$^+$ LT-
HSCs in G_1 compared to NEO1$^-$*Hoxb5*$^+$ LT-HSCs (**Fig. 3*E***; *SI Appendix,* Fig. S6). Taken
together, this suggests that NEO1$^+$*Hoxb5*$^+$ LT-HSCs are more proliferative, which may partially
contribute to their selective expansion during aging and in response to myeloablative stress.

Neogenin-1 marks a less regenerative, myeloid-biased fraction of *Hoxb5*$^+$ LT-HSCs
Given the aging and cell cycle differences between NEO1$^+$ and NEO1$^-$*Hoxb5*$^+$ LT-HSCs, we
next evaluated their reconstitution potential and lineage output by 10-cell transplants into
congenic irradiated primary recipients (**Fig. 4*A***). Over the course of 16 weeks, the percent of
total chimerism among peripheral blood that was donor-derived, was similar between NEO1$^+$
and NEO1$^-$*Hoxb5*$^+$ LT-HSC transplants (**Fig. 4*B***). However, among all donor-derived peripheral
blood, NEO1$^+$*Hoxb5*$^+$ LT-HSCs gave rise to a higher percentage of granulocytes and monocytes
(myeloid) and lower percentage of B and T cells (lymphoid) compared to NEO1$^-$*Hoxb5*$^+$ LT-
HSCs (**Fig. 4 *C* and *D***).

To evaluate the long-term reconstitution potential and the stability of lineage bias, we serially
transplanted 1000 donor-derived Lin$^-$c-KIT$^+$SCA1$^+$ (KLS) cells from primary recipients into
congenic irradiated secondary hosts (**Fig. 4*A***). Although KLS cells from both donors
repopulated all major lineages in secondary hosts, NEO1$^-$*Hoxb5*$^+$-derived cells exhibited

significantly higher reconstitution compared to NEO1⁺*Hoxb5*⁺-derived cells (**Fig. 4*E***). Moreover,
as during primary transplant, NEO1⁺*Hoxb5*⁺-derived cells maintained significant bias towards
granulocytes and monocytes and away from B and T cells compared to NEO1⁻*Hoxb5*⁺-derived
cells. (**Fig. 4 *F* and *G***). This suggests that both myeloid-biased and balanced phenotypes are
long-term maintained on secondary transplantation.
219
Transcriptional programs recapitulate functional differences between NEO1⁺ and NEO1⁻
***Hoxb5*⁺ LT-HSCs**
We next sought to understand the transcriptional programs that drive the observed functional
differences between NEO1⁺*Hoxb5*⁺ and NEO1⁻*Hoxb5*⁺ LT-HSCs. We isolated 250-500
NEO1⁺*Hoxb5*⁺ and NEO1⁻*Hoxb5*⁺ cells from female, 8-to-12-week old *Hoxb5*-mCherry mice and
performed low-input full-length RNA-sequencing using the Smart-Seq2 protocol ([44](#)) (**Fig. 5*A***).
Paired gene expression comparison of the two populations identified 1,036 differentially
expressed genes (false discovery rate *P-adjusted* < 0.1; **Fig. 5*B*;** *SI Appendix,* Fig. S7A and
Table S2) ([45](#)). Genes implicated in activation, cell cycle, and differentiation, such as *Fanca,*
Fancb, and *Mycn* ([46](#), [47](#)) were enriched in NEO1⁺*Hoxb5*⁺ LT-HSCs, while genes involved in
anti-redox, e.g. *Sod1* ([48](#)), and regulation of stem cell potency, e.g. *Klf4* and *Malat1* ([49](#), [50](#))
were enriched in NEO1⁻*Hoxb5*⁺ LT-HSCs (**Fig. 5*B***). Gene set enrichment analysis (GSEA) ([51](#))
and hypergeometric test with gene ontology (GO): biological processes ([52](#), [53](#)) revealed that
the driving differences (FDR < 0.05, P value < 0.05) between NEO1⁺ and NEO1⁻ cells were cell
cycle and ribosomal RNA expression (**Fig. 5*C***; *SI Appendix,* Fig. S7).
235
We also searched for the expression of lineage-specific transcripts that may indicate signs of
early myeloid and lymphoid priming in LT-HSCs. Among the genes significantly enriched in
NEO1⁺ compared to NEO1⁻*Hoxb5*⁺ LT-HSCs, we found several myeloid genes, including *Lrg1*
and lineage-related transcription factors, such as *Meis2, Hoxb6,* and *Cebpe* (**Fig. 5*E***), and
platelet genes (**Fig. 5*F***), such as *Vwf, Clu,* and *Selp*. We also find that NEO1⁺ LT-HSCs are
significantly enriched (*Q* < 0.05) for previously reported gene signatures of megakaryocyte
progenitors (MkP) and pre-erythrocyte colony-forming units (preCFU-E) (*SI Appendix,* Fig. S7*C*)
([27](#)). Moreover, the gene expression signature of NEO1⁺*Hoxb5*⁺ LT-HSCs significantly aligned
with expression profiles of previously reported myeloid-biased LT-HSCs (**Fig. 5*D, top***), while
NEO1⁻ LT-HSCs were enriched for the balanced LT-HSC signature (**Fig. 5*D, bottom***).
Altogether, these data suggest that LT-HSCs may sample regions of the transcriptome
associated with their lineage fate decisions.
248
Finally, we also compared NEO1⁺ and NEO1⁻*Hoxb5*⁺ LT-HSC gene expression with respect to
known stemness-associated genes. Overall, NEO1⁺*Hoxb5*⁺ LT-HSCs had higher expression of
known stem-related genes, including *Ctnnal1* ([14](#)), *Fgd5* ([13](#)), *Bmi1* ([54](#)), *Gprc5c* ([39](#)), and
Slamf1 ([11](#)) (**Fig. 5*G***), the latter of which was confirmed by flow cytometry (*SI Appendix,* Fig.
S8). This suggests that although previously identified stemness genes enrich for a self-renewing
phenotype, their expression may also be associated with myeloid-bias.
255
Lineage-balanced NEO1⁻*Hoxb5*⁺ LT-HSCs outcompete NEO1⁺*Hoxb5*⁺ LT-HSCs in
reconstitution and reside at the apex of hematopoiesis
To directly compare the relative fitness of NEO1⁺*Hoxb5*⁺ LT-HSCs and NEO1⁻*Hoxb5*⁺ LT-HSCs,
we co-transplanted 200 cells from each fraction with host supporter cells into irradiated 2-to-3-
month-old congenic recipients (**Fig. 6*A***). Donor origin was distinguished by CD45.2 and EGFP
expression using Hoxb5-mCherry and *CAG*-EGFP;*Hoxb5*-mCherry mice (**Fig. 6*A***). Unlike the
primary transplants above, NEO1⁻*Hoxb5*⁺ LT-HSCs exhibited significantly higher reconstitution
potential compared to NEO1⁺*Hoxb5*⁺ LT-HSCs in the co-transplantation setting (**Fig. 6*B***). This

suggests that NEO1⁻*Hoxb5*⁺ LT-HSCs are more fit to reconstitute young recipients compared to
NEO1⁺*Hoxb5*⁺ LT-HSCs.
266
Co-transplantation also confirmed that NEO1⁺*Hoxb5*⁺ LT-HSCs contribute significantly more to
granulocytes and monocytes (myeloid) and less to B and T cells (lymphoid) compared to NEO1⁻
Hoxb5⁺ LT-HSCs (**Fig. 6 *C* and *D***). We also quantified platelet fractions among EGFP⁺ donors.
Relative platelet contribution from NEO1⁺*Hoxb5*⁺ was not significantly different from NEO1⁻
Hoxb5⁺ donors during the first 16 weeks post-transplant, but significantly increased among
NEO1⁺*Hoxb5*⁺-derived PB at 20 weeks post-transplant (**Fig. 6*E***).
273
Finally, we also measured the composition of LT-HSCs between NEO1⁺ and NEO1⁻ derived
bone marrow in the co-transplantation setting. Both NEO1⁺ and NEO1⁻*Hoxb5*⁺ cells produced
on average equal numbers of *Hoxb5*⁺ LT-HSCs per million bone marrow cells (*SI Appendix,* Fig.
S9), and among *Hoxb5*⁺ LT-HSCs, more NEO1⁺ than NEO1⁻*Hoxb5*⁺ cells (**Fig. 6*F***). However,
NEO1⁺*Hoxb5*⁺ cells produced significantly fewer NEO1⁻*Hoxb5*⁺ cells compared to NEO1⁻
Hoxb5⁺ cells ($P = 0.006$). This suggests limited transition from NEO1⁺ to NEO1⁻, while NEO1⁻
Hoxb5⁺ cells are capable of giving rise to high percentages of both populations (**Fig. 6*F***).
Therefore, NEO1⁻*Hoxb5*⁺ LT-HSCs likely precede NEO1⁺*Hoxb5*⁺ LT-HSCs in the differentiation
hierarchy (**Fig. 6*G***).
283
284 **DISCUSSION**
285
Previously, we showed that phenotypic HSCs (pHSCs) are variable in their reconstitution
potential and that *Hoxb5* expression distinguishes long-term from short-term repopulating HSCs
([6]). While *Hoxb5*- pHSCs were unable to repopulate secondary recipients and were
homogeneously lymphoid-biased, Hoxb5+ pHSCs serially reconstituted recipients and exhibited
variable contribution to hematopoietic lineages. Therefore, we sought to understand the
diversity of self-renewing HSCs in the mouse bone marrow using *Hoxb5* as a reporter to mark
long-term HSCs (LT-HSCs).
293
To accomplish this, we screened gene expression profiles for candidate surface markers that are
strictly enriched in HSCs and stratify *Hoxb5*⁺ LT-HSCs into subpopulations. We identified
Neogenin-1 (*Neo1*; NEO1) as a transmembrane receptor specifically expressed on a sub-fraction
of *Hoxb5*⁺ LT-HSCs. Although NEO1 is well known for its role in the brain ([29-31]), skeletal muscle
([32]), breast epithelia ([34]), endothelia ([35]), and other tissues ([33]), its expression in the bone
marrow and association with LT-HSCs has not yet been extensively studied. We find that NEO1⁺
cells comprise a minor fraction of *Hoxb5*⁺ LT-HSCs in young mice that progressively expands with
age and represents >80% of 22-month-old *Hoxb5*⁺ LT-HSCs. This expansion can be partially
explained by the higher frequency of NEO1⁺*Hoxb5*⁺ in the G_2/S cell cycle phase compared to
NEO1⁻*Hoxb5*⁺ LT-HSCs. Both NEO1⁺ and NEO1⁻*Hoxb5*⁺ LT-HSCs are long-term self-renewing
and contribute to all major blood lineages during primary and secondary transplantations into
irradiated mice. However, NEO1⁺*Hoxb5*⁺ LT-HSCs exhibit a stable bias towards myeloid lineages
and are less productive in secondary transplants compared to NEO1⁻*Hoxb5*⁺ LT-HSCs. Gene
expression comparison reveals higher cell cycle and lower ribosomal gene expression in NEO1⁺
than NEO1⁻ LT-HSCs. This is consistent with the higher incidence of NEO1⁺*Hoxb5*⁺ LT-HSCs in
the G_2/S than G_1 cell cycle phase ([55]). NEO1⁺*Hoxb5*⁺ LT-HSCs also exhibited early sampling of
myeloid- and platelet-related genes, including lineage-related transcription factors, at the
transcript-level.
312
313 Several previous studies have used single cell transplants to describe HSC heterogeneity and
314 the existence of different lineage-primed states. Single cell HSC transplants by Dykstra and

colleagues described two fractions of long-term self-renewing HSCs, 'α cells' and 'β cells', which were myeloid-biased and lineage-balanced, respectively ([1]). Yamamoto and colleagues also have demonstrated the presence of myeloid-restricted progenitors with long-term repopulating activity (MyRPs) that are derived directly from balanced HSCs and expand with age ([2], [3]). While single HSC transplants evidenced the presence of balanced and myeloid-biased LT-HSCs, these studies did not identify surface or transcriptional markers to distinguish these populations. We have demonstrated a strategy for the prospective isolation of balanced and myeloid-biased LT-HSCs using Neogenin-1 and *Hoxb5*. The functional potential of NEO1⁺ and NEO1⁻*Hoxb5*⁺ LT-HSCs matches the characteristics of the myeloid-biased and balanced LT-HSC fractions previously predicted by single cell transplants ([1-3]).

Other markers have also been proposed to enrich for myeloid-biased HSCs. For example, our group previously showed that high CD150 surface expression enriches for myeloid-biased HSCs ([21]). However, the association of CD150 with myeloid-bias was not evaluated in long-term repopulating cells and additional marker combinations can improve the purification of balanced from myeloid-biased HSCs. Nevertheless, *Hoxb5*⁺NEO1⁺ LT-HSCs indeed express higher CD150 (*SI Appendix,* Fig. S7), validating our initial attempts to prospectively isolate myeloid-biased HSCs ([21]). CD41 has also been suggested to mark myelo-erythroid HSCs ([26]), although it likely separates different fractions. CD41⁻ HSCs are lymphoid biased and proliferative, while NEO1⁻*Hoxb5*⁺HSCs are balanced and quiescent. The vWF reporter mouse is another system used to isolate platelet-biased and myeloid-biased HSCs ([27]). However, like the CD41⁻ HSCs, vWF⁻ HSCs are lymphoid-biased HSCs that are phenotypically distinct from the balanced, quiescent NEO1⁻*Hoxb5*⁺ cells we describe in this study. Finally, CD61 was recently described as a surface marker on long-term repopulating myeloid-biased LT-HSCs that respond to inflammatory stress and expand with age ([24]). The CD61⁺ and CD61⁻ LT-HSCs are transcriptionally similar to NEO1⁺ and NEO1⁻*Hoxb5*⁺ LT-HSCs, suggesting that these markers may capture similar cell types (**Fig 5 *Ea* and *F***). Leveraging combinations of both surface markers will likely improve the purification of balanced LT-HSCs from lineage-biased LT-HSCs.

Our results also bring to question the hierarchical relationship between lineage-primed and balanced LT-HSCs. Previous single cell transplants studies suggest some degree of plasticity between LT-HSC fractions with different differentiation potentials ([1]). Our co-transplantation experiment indicates that NEO1⁻*Hoxb5*⁺ LT-HSCs are likely upstream of NEO1⁺*Hoxb5*⁺ LT-HSCs, as NEO1⁻*Hoxb5*⁺ LT-HSCs produced NEO1⁺ *Hoxb5*⁺ LT-HSCs, while conversion of NEO1⁺ *Hoxb5*⁺ LT-HSCs to NEO1⁻*Hoxb5*⁺ HSCs was rare. The few instances of NEO1⁺ donors producing NEO1⁻*Hoxb5*⁺ LT-HSCs may be attributable to impurity among the 200 cells that were injected. This is also consistent with our gene expression and cell cycle analysis demonstrating that NEO1⁺*Hoxb5*⁺ LT-HSCs are more often cycling compared to NEO1⁻*Hoxb5*⁺ cells ([56]). Moreover, NEO1⁻*Hoxb5*⁺ LT-HSCs contributed more to total hematopoiesis in secondary transplants and co-transplants compared to NEO1⁺*Hoxb5*⁺ LT-HSCs. Therefore, our data suggest that balanced quiescent LT-HSCs reside at the apex of the hematopoietic hierarchy, corroborating a recent study showing that myeloid-biased MyRPs are derived from balanced HSCs ([2], [3]). However, this contrasts with the view that vWF⁺ and CD41⁺ platelet/myeloid-biased cells reside at the apex of the hematopoietic hierarchy ([26], [27]).

Additional work will also be required to delineate the differentiation path that distinct LT-HSC fractions follow to generate various blood cells. Our study suggests that balanced NEO1⁻*Hoxb5*⁺ LT-HSCs contribute to myeloid lineage through a NEO1⁺*Hoxb5*⁺ LT-HSC intermediate. However, it is unclear whether NEO1⁻*Hoxb5*⁺ LT-HSCs require the NEO1⁺*Hoxb5*⁺ intermediate state or can independently produce lineage progenitors through alternative routes. Moreover, it remains to be answered whether all NEO1⁺*Hoxb5*⁺ LT-HSCs are derived from NEO1⁻*Hoxb5*⁺ LT-HSCs. To fully

reveal the hierarchical order and sequence of differentiation events, single cell tracking
experiments will be required. Critically, our experimental results are based on the behavior of cells
upon transplantation into young, irradiated mice. A recent study using individually barcoded HSCs
showed that lineage biases are more pronounced after transplantation into lethally irradiated mice
compared to unirradiated or anti-c-KIT-depleted syngeneic mice ([4]). This suggests that post-
transplant lineage bias may be either due to plasticity in lineage output or the selective
engraftment of pre-existing HSC subsets. Therefore, it will be important to evaluate the potential
and hierarchical relationship between NEO1$^+$ and NEO1$^-$*Hoxb5*$^+$ LT-HSCs during *in situ,*
unperturbed hematopoiesis with *in vivo* lineage tracing.
375
Furthermore, we note that comparing NEO1$^+$ and NEO1$^-$ fractions within pHSCs without gating
Hoxb5$^+$ LT-HSCs may mislead the significance of NEO1 to separate myeloid-biased from
balanced cells. The vast majority of NEO1$^-$ cells among pHSCs are short-term, lymphoid-biased
Hoxb5$^-$ ST-HSCs ([6]) that far outnumber the long-term, balanced NEO1$^-$*Hoxb5*$^+$ LT-HSCs we find
in this study. NEO1 is also rarely expressed in *Hoxb5*$^-$ downstream cells and bone marrow niche
cells and its functional role in these populations has not been evaluated in this study.
382
383 The functional differences between NEO1$^+$ and NEO1$^-$ Hoxb5+ LT-HSCs may be influenced by
384 intrinsic programs, external cues, or both. As NEO1 is a receptor to many known ligands, ongoing
385 studies are evaluating the role of the bone marrow niche and particular ligands to NEO1 in
386 influencing lineage bias and stem cell maintenance.
387
388 Finally, our antibody against NEO1 also identified higher proportions of cells in human HSCs
389 compared to MPPs, LMPPs, and downstream progenitors, suggesting a possibly conserved role
390 of NEO1 in human HSC biology. Therefore, further evaluation of NEO1$^+$*Hoxb5*$^+$ LT-HSCs and the
391 receptor-ligand interactions may offer insights into evolutionarily conserved mechanisms of
392 lineage bias during long-term hematopoiesis.
393
394 Taken together, we have identified a novel marker on the surface of *Hoxb5*$^+$ LT-HSCs, Neogenin-
395 1, that enables the separation of myeloid-biased LT-HSCs from quiescent, balanced LT-HSCs
396 with the highest long-term repopulation potential. Our findings reveal a previously undescribed
397 layer of functional heterogeneity among strictly defined functional LT-HSCs and enable the
398 precise and prospective study of LT-HSCs and their fractions.

MATERIALS AND METHODS

Mice

2-to-3-month-old female *Hoxb5*-mCherry mice (MGI:5911679; available through RIKEN BioResource Research Center: RBRC09733) were used as donors (CD45.2) for transplant experiments and bulk RNA-sequencing. Additionally, 2-to-3-month-old *CAG-EGFP*;*Hoxb5*-mCherry mice (in-house colony) were used for co-transplant assays. 2-to-3-month-old female B6.SJL-*Ptprc*ᵃ *Pepc*ᵇ/BoyJ mice (Jackson Laboratory) were used as recipients (CD45.1) for transplant experiments and for supporter bone marrow. 4-month-old female *Hoxb5*-mCherry mice were used for experiments with 5-fluorouracil. 5-month, 13-month, and 22-month-old female *Hoxb5*-mCherry mice were used for aging analysis. 2-to-3-month-old and 12-to-14-month-old female *Hoxb5*-mCherry mice were used for cell cycle analysis. 2-to-3-month-old C57BL/6J female mice (Jackson Laboratory) were used for fluorescent-minus-one (FMO) controls for *Hoxb5*-mCherry expression.

Gene expression profiles of mouse hematopoietic cells and the bone marrow niche

All microarray data used in this study are accessible through the Gene Expression Commons platform (http://gexc.stanford.edu) and the Gene Expression Omnibus (GEO) accession, GSE34723. We analyzed 64 microarray gene expression profiles (GEPs) of 23 distinct mouse hematopoietic cell types (*SI Appendix,* Table S1) for surface markers enriched in HSCs compared to downstream progeny. GEPs were normalized against a large common reference of >11,939 Affymetrix Mouse Genome 430 2.0 microarrays as described before(36). For each gene, the probeset with the largest dynamic range was selected and transformed to percentile ranks (range: -100% to +100%) based on its relative expression to the reference.

Genes were further subset based on two main criteria (1) positive percentile expression in HSCs and (2) annotation as a cell surface protein based on GO:0009986, leaving 186 gene candidates. Fold-change enrichment in HSCs was calculated as the average percentile rank for each gene among HSCs divided by the average percentile rank for that gene across all other cells.

Single cell transcriptomes of hematopoietic stem and progenitors (HSPCs) were acquired from GEO accessions, GSE90742, and from the Single-Cell Gene Expression Atlas for hematopoietic cells (http://blood.stemcells.cam.ac.uk/single_cell_atlas.html). In the dataset from Rodriguez-Fraticelli et al. (5), cells with greater than 0 unique molecular identifiers (UMIs) were considered *Neo1*⁺. In the dataset from Nestorowa et al. (57), cells with greater than 4 $\log_2$ counts were considered *Neo1*⁺. A threshold of 4 was chosen based on the color gradient thresholds for the diffusion maps in the Single-Cell Gene Expression Atlas.

tSNE plots of single cell RNA-sequencing data from the mouse bone marrow niche were generated using the online tool nichExplorer (https:// https://compbio.nyumc.org/niche/)(58).

Bone marrow imaging

Femurs were collected and fixed in 4% PFA with 10% EDTA solution at 4°C overnight. After decalcification in 20% EDTA solution at 4°C for two days, bones were saturated in increasing concentrations of sucrose solution (10% for 1h, 20% for 1h, and 30% overnight) at 4°C. Prepared bones were frozen in OCT compound using an isopentane bath cooled on dry ice. For staining, 50-70 µm thick sections were cut on microscope slides and blocked in 3% BSA with 10% FBS solution (blocking buffer) for 1h at room temperature. Sections were stained with primary antibodies against NEO1 (R&D cat. no. AF1079; dilution: 1:100) and Endomucin (ThermoFisher

Scientific; clone: V.7C7; dilution: 1:100) in 10% blocking buffer at 4°C overnight and washed with
PBS. For secondary staining, sections were incubated with donkey anti-rat IgG (H+L) AF488-
conjugated and donkey anti-goat IgG (H+L) Cy3-conjugated (Jackson ImmunoResearch; dilution:
1:400) in 10% blocking buffer at room temperature for 1h. After another wash with PBS, samples
were counterstained with DAPI and sealed for imaging. Images were acquired using a LSM880
confocal microscope (Zeiss) with LD LCI Plan-Apochromat 25x/0.8 objective. Images were
analyzed using the Fiji software.
457
Transplantation assays
2-to-3-month-old female B6.SJL-*Ptprc^a Pepc^b* /BoyJ (CD45.1) recipient mice were lethally
irradiated at a single dose of 9 Gy. For reconstitution assays, 10 NEO1$^+$*Hoxb5$^+$* or NEO1$^-$
Hoxb5$^+$ LT-HSCs were isolated from donor CD45.2$^+$ *Hoxb5*-mCherry mice (MGI:5911679) as
described in sections *'Bone marrow isolation'* and *'Flow cytometry'* and co-injected with 2x10^5
recipient whole bone marrow cells in 200 µl of PBS with 2% FBS into the retro-orbital venous
plexus. For secondary transplants, 1000 CD45.2$^+$ Lin$^-$cKIT$^+$SCA1$^+$ (KLS) cells were isolated by
flow cytometry and transplanted together with 2 x 10^5 recipient (CD45.1) whole bone marrow
cells into lethally irradiated recipient CD45.1$^+$ mice as described above. For co-transplantation
assays, 200 NEO1$^+$ and NEO1$^-$*Hoxb5$^+$* LT-HSCs were isolated from either CD45.2$^+$ *Hoxb5*-
mCherry mice (MGI:5911679) or an in-house strain of EGFP$^+$CD45.2$^+$ *Hoxb5*-mCherry mice and
transplanted into lethally irradiated recipient CD45.1$^+$ mice at a split dose of 9 Gy with a 4-hour
interval, controlling for donor strain biases by transplanting the same condition from both strains.
In all cases, recipients with lower than 1% total chimerism were considered failed
transplantations and excluded from analysis.
473
Peripheral blood analysis for chimerism
Peripheral blood collections for assessment of donor chimerism were performed at 4, 8, 12, and
16 weeks after primary and secondary transplantations and 8, 12, 16, and 20 weeks after co-
transplantations. At each time point, 50-100 µl of blood was collected from the retro-orbital
venous plexus using heparinized capillary tubes (Fisher Scientific) and added to K$_2$/EDTA-
coated MiniCollect tubes (Greiner Bio-One). Red blood cells were depleted with two rounds of
ACK lysis buffer by incubating at RT for 5 min each. Cells were then washed with cold PBS.
Cells were Fc-blocked with of rat IgG (LifeSpan BioSciences) and stained with 5 µg/ml of rat
anti-mouse antibodies (catalog no., concentrations, and clone provided in Table S3 of the *SI*
Appendix) to CD45.1, CD45.2, CD11B, GR1, B220, CD3, and only in the co-transplantation
assay, CD41. 7-Aminoactinomycin D (7-AAD; BD Bioscience) was added for live and dead cell
discrimination.
486
For reconstitution assays, total donor chimerism was defined as the percentage of CD45.1$^-$
CD45.2$^+$ cells among total CD45.1$^-$CD45.2$^+$ and CD45.1$^+$CD45.2$^-$ cells. For co-transplantation
assays, total donor chimerism was defined as the percentage of either CD45.1$^-$CD45.2$^+$EGFP$^-$
cells or CD45.1$^-$CD45.2$^+$EGFP$^+$ cells among total CD45.1$^-$CD45.2$^+$EGFP$^-$, CD45.1$^-$
CD45.2$^+$EGFP$^+$, and CD45.1$^+$CD45.2$^-$EGFP$^-$ cells. For all cases, lineage chimerism was
evaluated as the percentage of lymphoid (B220$^+$ B cells and CD3$^+$ T cells) or myeloid
(GR1$^+$CD11B$^+$ granulocytes and monocytes among donor-derived cells. Only EGFP$^+$ cells were
evaluated for platelet chimerism. In these cases, platelet chimerism was calculated as the
percent of CD41$^+$ platelets among all donor-derived EGFP$^+$ cells and platelets.
496
Mouse hematopoietic stem cell isolation by flow cytometry
HSCs were isolated from 2-to-3-month-old, 5-month-old, 13-month-old, and 22-month-old bone
marrow of *Hoxb5*-mCherry mice (MGI:5911679). For bone marrow isolation, tibia, femur, and
pelvis were dissected, crushed with mortar and pestle in FACS buffer (2% fetal bovine serum

(FBS) in PBS with 100 U/ml DNase), and the supernatant was collected. For WBM isolation, red
blood cells were depleted with ACK lysis buffer by incubating at RT for 10 min and Fc-blocked
by incubating with rat IgG (LifeSpan BioSciences) for 10 min. For c-KIT$^+$ cell isolation, samples
were Fc-blocked with rat IgG for 10 min, incubated in c-KIT magnetic beads (Miltenyi) with 100
U/ml DNase, and MACS-isolated using LS magnetic columns (Miltenyi) as per manufacturer's
protocol. For the cell cycle analysis, samples were blocked with TruStain FcX (CD16/32) instead
of rat IgG, and enriched for c-KIT$^+$ cells. Each sample was normalized to an equal number of
cells (4.5 million c-KIT$^+$ cells) and processed following a previously published protocol for cell
cycle analysis of HSCs ([59]). Sample collections with fewer than 10 total *Hoxb5$^+$* LT-HSCs after
flow cytometry analysis were considered underpowered for determining the distribution of cells
in G_0, G_1, and G_2/S and therefore excluded.
512
Samples for mouse HSC isolation were stained with a cocktail of antibodies against lineage
markers, i.e. CD3, Gr-1, CD11B, B220, and TER119 (AF700), c-KIT (APC-Cy7), SCA-1 (PE-
Cy7), CD48 (BV711), FLK2 (PerCP-Cy5.5), CD150 (BV421), CD34 (primary: biotin; secondary:
Strep-BUV737), and NEO1 (primary: goat anti-mouse/human (R&D cat. no. AF1079);
secondary: donkey anti-goat IgG (H+L) cross-absorbed AF488-conjugated; negative control:
normal goat IgG). Although the antibody to NEO1 is polyclonal, results were consistent across
multiple reagent lots and experiments. Antibody-labeled cells were shown to express higher
NEO1 mRNA compared to unlabeled cells (*SI Appendix,* Fig. S7*A*).
521
522 Primary and secondary antibody incubations were 20-30 min each with 5 min wash step in
523 between. Catalog number, concentrations, and clone information are provided in Table S3 of *SI*
524 *Appendix.*
525
526 Flow cytometry and cell sorting were performed on the BD FACSAria and BD LSRFortessa.
527 Gating strategy for the different populations is shown in *SI Appendix,* Fig. S3. 7-AAD or DAPI
528 were used as a viability dye for dead cell exclusion, depending on the assay. All cells were
529 suspended in FACS buffer (2% FBS in PBS) on ice unless otherwise indicated.
530
Myeloablative stress with 5-fluorouracil (5-FU)
4-month-old female *Hoxb5*-mCherry mice were injected with 150 mg of 5-FU (Sigma-Aldrich)
per kg body weight from a stock solution of 10 mg/ml in PBS ([41]). Bone marrow populations
were isolated and analyzed 5 days after treatment as described above. Notably, given the
upregulation of CD11B in HSCs post-treatment with 5-FU ([41]), the antibody to CD11B was
omitted from the lineage staining panel.
537
Human hematopoietic stem cell analysis by flow cytometry
Human CD34$^+$ bone marrow cells were purchased from AllCells, Inc and their use were
approved by the Stanford University Institutional Review Boards. To analyze human
hematopoietic stem cell and progenitor populations, CD34$^+$ bone marrow cells were stained
against lineage markers, i.e. CD3, CD4, CD8, CD11B, CD14, CD19, CD20, CD56, and GPA
(PE-Cy5), stem and progenitor markers, i.e. CD34 (APC-Cy7), CD38 (APC), CD45RA (BV785),
and CD90 (FITC), and NEO-1 (primary: goat anti-mouse/human (R&D cat. No. AF1079);
secondary: donkey anti-goat IgG (H+L) Cy3-conjugated; negative control: normal goat IgG).
Propidium iodide (PI; Sigma-Aldrich) was added for live and dead cell discrimination. Catalog
number, concentrations, and clone information are provided in Table S3 of *SI Appendix*. Gating
strategy for the different populations is shown in **Figure 1***E*. Flow cytometry was performed as
described above.
550
RNA extraction and library preparation for bulk RNA sequencing

For RNA-sequencing experiments, 250-500 cells from two pooled mice per sample were sorted
directly into 100 µL of lysis buffer (Buffer RL) and RNA was isolated with the Single Cell RNA
Purification Kit (Norgen Biotek Corp.) according to the manufacturer's protocol. RNA quality was
measured by capillary electrophoresis using the Agilent 2100 Bioanalyzer with Nano mRNA
assay at the Stanford Protein and Nucleic Acid (PAN) Facility.
557
Libraries were prepared using the Smart-seq2 protocol by Picelli et al., 2014 with minor
modifications. Briefly, cDNA was generated by oligo-dT primed reverse transcription with MMLV
reverse transcriptase (SMARTScribe, Clontech) and a locked template-switching
oligonucleotide (TSO). This was followed by 18 cycles of PCR amplification using KAPA HiFi
hotStart ReadyMix and ISPCR primers. Amplified cDNA was then purified using 0.7x volume
Agencourt AMPure XP beads to remove smaller fragments. The resulting cDNA concentration
and size distribution for each well was determined on a capillary electrophoresis-based Agilent
2100 Bioanalyzer with High Sensitivity DNA chip at the Stanford PAN facility. 40 ng of cDNA
was then tagmented, uniquely barcoded, and PCR enriched using the Nextera DNA Library
Prep Kit (Illumina, San Diego, CA). Libraries were then pooled in equimolar amounts and
purified of smaller fragments using 0.7x Agencourt AMPure XP beads. Pooled libraries were
checked for quality using the Agilent Bioanalyzer with High Sensitivity DNA chip at the Stanford
PAN facility. 10 samples were sequenced with 151 bp paired-end reads on a single lane of
NextSeq 500 (Illumina, San Diego, CA) at the Stanford Functional Genomics Facility.
572
573 After sequencing, bcl2fastq2 v2.18 (Illumina) was used to extract the data and generate FASTQ
574 files for each sample by using unique barcode combinations from the Nextera preparation. Raw
575 reads were trimmed for base call quality (PHRED score >=21) and for adapter sequences using
576 Skewer v0.2.2 (60). Trimmed reads were then aligned to the mouse genome assembly (mm10)
577 from UCSC (http://genome.ucsc.edu) using STAR v2.4 with default setting (61).
578
RNA sequencing analysis
Count normalization and differential gene expression analysis was performed using the DESeq2
v1.22.2 package in R (45). Raw counts from STAR were inputted into a DESeqDataSet object
indicating NEO1 status ('status') and mouse subject ('subject') as factors ('design = ~subject +
status'). Counts were size-factor normalized using the 'DESeq' function and $\log_2$-transformed.
Pairwise differential gene expression analysis was performed using the lfcShrink function and
indicating 'type = apeglm', which applies the adaptive t prior shrinkage estimator. As
recommended (45), a threshold of P-adjusted < 0.1 was used to define significance for
differentially expressed genes (*SI Appendix,* Table S2).
588
Gene set enrichment analysis (GSEA) was performed using the GSEA software provided by the
Broad Institute (51) and the clusterProfiler v3.10.0 (52) and HTSanalyzer v2.34.0 (62) packages
in R. Hypergeometric test with GO: Biological Processes and dot plots in Figure S7 were
generated using the clusterProfiler package in R. Gene expression signatures of myeloid and
non-myeloid LT-HSCs were acquired from the original study by Mann et al., 2018 (24). Gene
expression signatures of lineage-restricted progenitors, including megakaryocyte progenitors
(MkP), pre-erythrocyte colony-forming units (preCFU-E), pre-granulocyte/macrophage
progenitors (preGM), pre-megakaryocyte/erythrocyte progenitors (preMegE), and common
lymphoid progenitors (CLP), were acquired from the original study by Sanjuan-Pla et al., 2013
(27).
599
Statistics
Statistical significance between two groups was determined using a paired or unpaired
Student's *t* test, as appropriate. For comparison of two groups across multiple time points,

statistical significance was determined using a two-way ANOVA using the groups and time
points as factors. Multiple hypothesis correction was applied to gene expression comparisons
using the Benjamini-Horchberg procedure. Results with *P-adjusted* < 0.1 were considered
significant. Data analyses were performed with R 3.5.1, Prism v7 (GraphPad Software, Inc.),
and FlowJo v10 (FlowJo, LLC). The investigators were not blinded to allocation during
experiments and outcome assessment. No sample-size estimates were performed to ensure
adequate power to detect a pre-specified effect size.

Acknowledgements
We thank R. Yamamoto, C.K.F. Chan, J. Xiang, V. Mascetti, V.G. Alvarado, A. Chandra, and A.
Manjunath for technical assistance and discussion. We are grateful to T. Naik for lab
management, A. McCarty and C. Wang for mouse breeding and management, P. Lovelace and
S. Weber for their support and assistance with FACS, and Stanford Functional Genomic Facility
(SFGF) for assistance with sequencing scRNA-seq libraries. We thank K. Kao, T. Sakamaki,
and M. Miyanishi for assistance with breeding the *Hoxb5*-mCherry and *CAG*-EGFP;*Hoxb5*-
mCherry mouse strains.

Funding
This study was supported by the California Institute for Regenerative Medicine (RT3-07683) to
I.L.W.; Stanford-UC Berkeley Stem Cell Institute, anonymous donors; National Cancer Institute,
DHHS (PHS Grant Number CA09302), the National Heart, Lung, and Blood Institute Ruth L.
Kirschstein National Research Service Award (F30HL147460), and the Stanford Medical
Science Training Program to G.S.G. Foundation For Polish Science (HOMING POIR.04.04.00-
00-5F16/18-00) and MOBILITY PLUS Fellowship from the Polish Ministry of Science and Higher
Education to K.S. HARMONIA UMO-2015/18/M/NZ3/00387 to Alicja Jozkowicz and Leading
National Research Center (KNOW) to the Faculty of Biochemistry, Biophysics, and
Biotechnology of Jagiellonian University, supporting M.Z. Stanford Vice President for
Undergraduate Research office (VPUE) grant and Bio-X summer funding to J.N.

Author contributions
K.S. identified Neogenin-1. K.S., G.S.G., M.Z., and I.L.W. designed the experiments; K.S.,
G.S.G., M.Z., J.N., A.Z., R.S., B.G., and D.W. performed the experiments. K.S., G.S.G., M.Z.,
and J.N. analyzed the data. G.S.G., K.S., M.Z., and I.L.W. wrote the manuscript. I.L.W.
supervised the study. All authors commented on the manuscript at all stages.

Competing interests:
Authors declare no competing interests.

Data and materials availability
All expression datasets from the public domain analyzed in this work are described in **Materials
and Methods**. The bulk RNA-sequencing data generated in this study has been deposited in
GEO under accession code GSE130504. All data generated or analyzed in this study have also
been deposited in the Dryad Digital Repository under doi:10.5061/dryad.376gn3k.

References:

1. Dykstra B, *et al.* (2007) Long-term propagation of distinct hematopoietic differentiation programs in vivo. *Cell Stem Cell* 1(2):218-229.
2. Yamamoto R, *et al.* (2018) Large-Scale Clonal Analysis Resolves Aging of the Mouse Hematopoietic Stem Cell Compartment. *Cell Stem Cell* 22(4):600-607 e604.
3. Yamamoto R, *et al.* (2013) Clonal analysis unveils self-renewing lineage-restricted progenitors generated directly from hematopoietic stem cells. *Cell* 154(5):1112-1126.
4. Lu R, Czechowicz A, Seita J, Jiang D, & Weissman IL (2019) Clonal-level lineage commitment pathways of hematopoietic stem cells in vivo. *Proc Natl Acad Sci U S A* 116(4):1447-1456.
5. Rodriguez-Fraticelli AE, *et al.* (2018) Clonal analysis of lineage fate in native haematopoiesis. *Nature* 553(7687):212-216.
6. Chen JY, *et al.* (2016) Hoxb5 marks long-term haematopoietic stem cells and reveals a homogenous perivascular niche. *Nature* 530(7589):223-227.
7. Seita J & Weissman IL (2010) Hematopoietic stem cell: self-renewal versus differentiation. *Wiley Interdiscip Rev Syst Biol Med* 2(6):640-653.
8. Spangrude GJ, Heimfeld S, & Weissman IL (1988) Purification and characterization of mouse hematopoietic stem cells. *Science* 241(4861):58-62.
9. Ikuta K & Weissman IL (1992) Evidence that hematopoietic stem cells express mouse c-kit but do not depend on steel factor for their generation. *Proc Natl Acad Sci U S A* 89(4):1502-1506.
10. Osawa M, Hanada K, Hamada H, & Nakauchi H (1996) Long-term lymphohematopoietic reconstitution by a single CD34-low/negative hematopoietic stem cell. *Science* 273(5272):242-245.
11. Kiel MJ, *et al.* (2005) SLAM family receptors distinguish hematopoietic stem and progenitor cells and reveal endothelial niches for stem cells. *Cell* 121(7):1109-1121.
12. Christensen JL & Weissman IL (2001) Flk-2 is a marker in hematopoietic stem cell differentiation: a simple method to isolate long-term stem cells. *Proc Natl Acad Sci U S A* 98(25):14541-14546.
13. Gazit R, *et al.* (2014) Fgd5 identifies hematopoietic stem cells in the murine bone marrow. *J Exp Med* 211(7):1315-1331.
14. Acar M, *et al.* (2015) Deep imaging of bone marrow shows non-dividing stem cells are mainly perisinusoidal. *Nature* 526(7571):126-130.
15. Sawai CM, *et al.* (2016) Hematopoietic Stem Cells Are the Major Source of Multilineage Hematopoiesis in Adult Animals. *Immunity* 45(3):597-609.
16. de Haan G & Lazare SS (2018) Aging of hematopoietic stem cells. *Blood* 131(5):479-487.
17. Morrison SJ, Wandycz AM, Akashi K, Globerson A, & Weissman IL (1996) The aging of hematopoietic stem cells. *Nat Med* 2(9):1011-1016.
18. Wright DE, Bowman EP, Wagers AJ, Butcher EC, & Weissman IL (2002) Hematopoietic stem cells are uniquely selective in their migratory response to chemokines. *J Exp Med* 195(9):1145-1154.
19. Passegue E, Wagers AJ, Giuriato S, Anderson WC, & Weissman IL (2005) Global analysis of proliferation and cell cycle gene expression in the regulation of hematopoietic stem and progenitor cell fates. *J Exp Med* 202(11):1599-1611.
20. Fleming WH, *et al.* (1993) Functional heterogeneity is associated with the cell cycle status of murine hematopoietic stem cells. *J Cell Biol* 122(4):897-902.
21. Beerman I, *et al.* (2010) Functionally distinct hematopoietic stem cells modulate hematopoietic lineage potential during aging by a mechanism of clonal expansion. *Proc Natl Acad Sci U S A* 107(12):5465-5470.

22. Rossi DJ*, et al.* (2007) Deficiencies in DNA damage repair limit the function of
haematopoietic stem cells with age. *Nature* 447(7145):725-729.
23. Rossi DJ*, et al.* (2005) Cell intrinsic alterations underlie hematopoietic stem cell aging.
Proc Natl Acad Sci U S A 102(26):9194-9199.
24. Mann M*, et al.* (2018) Heterogeneous Responses of Hematopoietic Stem Cells to
Inflammatory Stimuli Are Altered with Age. *Cell Rep* 25(11):2992-3005 e2995.
25. MacLean AL*, et al.* (2017) Single Cell Phenotyping Reveals Heterogeneity Among
Hematopoietic Stem Cells Following Infection. *Stem Cells* 35(11):2292-2304.
26. Gekas C & Graf T (2013) CD41 expression marks myeloid-biased adult hematopoietic
stem cells and increases with age. *Blood* 121(22):4463-4472.
27. Sanjuan-Pla A*, et al.* (2013) Platelet-biased stem cells reside at the apex of the
haematopoietic stem-cell hierarchy. *Nature* 502(7470):232-236.
28. Wilson NH & Key B (2007) Neogenin: one receptor, many functions. *Int J Biochem Cell
Biol* 39(5):874-878.
29. Wilson NH & Key B (2006) Neogenin interacts with RGMa and netrin-1 to guide axons
within the embryonic vertebrate forebrain. *Dev Biol* 296(2):485-498.
30. Xu K*, et al.* (2014) Neural migration. Structures of netrin-1 bound to two receptors
provide insight into its axon guidance mechanism. *Science* 344(6189):1275-1279.
31. Matsunaga E*, et al.* (2004) RGM and its receptor neogenin regulate neuronal survival.
Nat Cell Biol 6(8):749-755.
32. Bae GU*, et al.* (2009) Neogenin regulates skeletal myofiber size and focal adhesion
kinase and extracellular signal-regulated kinase activities in vivo and in vitro. *Mol Biol
Cell* 20(23):4920-4931.
33. Zhang AS, West AP, Jr., Wyman AE, Bjorkman PJ, & Enns CA (2005) Interaction of
hemojuvelin with neogenin results in iron accumulation in human embryonic kidney 293
cells. *J Biol Chem* 280(40):33885-33894.
34. Srinivasan K, Strickland P, Valdes A, Shin GC, & Hinck L (2003) Netrin-1/neogenin
interaction stabilizes multipotent progenitor cap cells during mammary gland
morphogenesis. *Dev Cell* 4(3):371-382.
35. Park KW*, et al.* (2004) The axonal attractant Netrin-1 is an angiogenic factor. *Proc Natl
Acad Sci U S A* 101(46):16210-16215.
36. Seita J*, et al.* (2012) Gene Expression Commons: an open platform for absolute gene
expression profiling. *PLoS One* 7(7):e40321.
37. Smith-Berdan S*, et al.* (2011) Robo4 cooperates with CXCR4 to specify hematopoietic
stem cell localization to bone marrow niches. *Cell Stem Cell* 8(1):72-83.
38. Yano M*, et al.* (1997) Expression and function of murine receptor tyrosine kinases, TIE
and TEK, in hematopoietic stem cells. *Blood* 89(12):4317-4326.
39. Cabezas-Wallscheid N*, et al.* (2017) Vitamin A-Retinoic Acid Signaling Regulates
Hematopoietic Stem Cell Dormancy. *Cell* 169(5):807-823 e819.
40. Majeti R, Park CY, & Weissman IL (2007) Identification of a hierarchy of multipotent
hematopoietic progenitors in human cord blood. *Cell Stem Cell* 1(6):635-645.
41. Randall TD & Weissman IL (1997) Phenotypic and functional changes induced at the
clonal level in hematopoietic stem cells after 5-fluorouracil treatment. *Blood* 89(10):3596-
3606.
42. Bernitz JM, Kim HS, MacArthur B, Sieburg H, & Moore K (2016) Hematopoietic Stem
Cells Count and Remember Self-Renewal Divisions. *Cell* 167(5):1296-1309 e1210.
43. Pietras EM, Warr MR, & Passegue E (2011) Cell cycle regulation in hematopoietic stem
cells. *J Cell Biol* 195(5):709-720.
44. Picelli S*, et al.* (2014) Full-length RNA-seq from single cells using Smart-seq2. *Nat
Protoc* 9(1):171-181.

45. Love MI, Huber W, & Anders S (2014) Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* 15(12):550.
46. Du W*, et al.* (2013) Inflammation-mediated notch signaling skews fanconi anemia
hematopoietic stem cell differentiation. *J Immunol* 191(5):2806-2817.
47. Laurenti E*, et al.* (2008) Hematopoietic stem cell function and survival depend on c-Myc
and N-Myc activity. *Cell Stem Cell* 3(6):611-624.
48. Pervaiz S, Taneja R, & Ghaffari S (2009) Oxidative stress regulation of stem and
progenitor cells. *Antioxid Redox Signal* 11(11):2777-2789.
49. Park CS, Shen Y, Lewis A, & Lacorazza HD (2016) Role of the reprogramming factor
KLF4 in blood formation. *J Leukoc Biol* 99(5):673-685.
50. Ma XY*, et al.* (2015) Malat1 as an evolutionarily conserved lncRNA, plays a positive role
in regulating proliferation and maintaining undifferentiated status of early-stage
hematopoietic cells. *BMC Genomics* 16:676.
51. Subramanian A*, et al.* (2005) Gene set enrichment analysis: a knowledge-based
approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*
102(43):15545-15550.
52. Yu G, Wang LG, Han Y, & He QY (2012) clusterProfiler: an R package for comparing
biological themes among gene clusters. *OMICS* 16(5):284-287.
53. Ashburner M*, et al.* (2000) Gene ontology: tool for the unification of biology. The Gene
Ontology Consortium. *Nat Genet* 25(1):25-29.
54. Park IK*, et al.* (2003) Bmi-1 is required for maintenance of adult self-renewing
haematopoietic stem cells. *Nature* 423(6937):302-305.
55. Derenzini M*, et al.* (2005) Key role of the achievement of an appropriate ribosomal RNA
complement for G1-S phase transition in H4-II-E-C3 rat hepatoma cells. *J Cell Physiol*
202(2):483-491.
56. Morrison SJ & Weissman IL (1994) The long-term repopulating subset of hematopoietic
stem cells is deterministic and isolatable by phenotype. *Immunity* 1(8):661-673.
57. Nestorowa S*, et al.* (2016) A single-cell resolution map of mouse hematopoietic stem
and progenitor cell differentiation. *Blood* 128(8):e20-31.
58. Tikhonova AN*, et al.* (2019) The bone marrow microenvironment at single-cell resolution.
Nature.
59. Szade K, Bukowska-Strakova K, Zukowska M, Jozkowicz A, & Dulak J (2016) Analysis
of Cell Cycle Status of Murine Hematopoietic Stem Cells. *Methods Mol Biol* 1516:91-99.
60. Jiang H, Lei R, Ding SW, & Zhu S (2014) Skewer: a fast and accurate adapter trimmer
for next-generation sequencing paired-end reads. *BMC Bioinformatics* 15:182.
61. Dobin A*, et al.* (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*
29(1):15-21.
62. Wang X, Terfve C, Rose JC, & Markowetz F (2011) HTSanalyzeR: an R/Bioconductor
package for integrated network analysis of high-throughput screens. *Bioinformatics*
27(6):879-880.
786
787

Overview of Figures and Tables

Main Figures:

Supporting Information (SI) Appendix:

Figure 1 | Identification of Neogenin-1 as a unique surface marker on mouse hematopoietic
stem cells (HSCs). (**A and B**) *In silico* screen to identify unique surface markers on HSCs from
64 microarray expression profiles of 23 distinct mouse hematopoietic cell types. (**A**) Heatmap
showing normalized gene expression of gene-ontology-annotated (GO: 0009986) surface
markers across different hematopoietic compartments. 186 surface markers expressed on HSCs
and 23 phenotypes categorized as 'HSCs', 'MPPs', 'Early progenitors', 'Late progenitors',
'Myeloid', and 'Lymphoid' are displayed (*SI Appendix*, Table S1). Genes are ordered from top to
bottom by $\log_2$ fold enrichment in HSCs compared to downstream cells and the top most enriched
genes (>2 $\log_2$ fold enrichment) are highlighted in a box. For further details, see methods. (**B**)
Barplots showing normalized gene expression of *Neo1* across the cell type categories shown in
A. Statistical significance was calculated by unpaired, two-tailed Student's *t*-test between 'HSC'
and each cell type. ****P < 0.0001. (**C and D**) Flow cytometry analysis of NEO1 surface
expression in the mouse bone marrow (*n* = 5 mice). (**C**) Contour plots with outliers showing the
gating scheme for early progenitors, MPPb, MPPa, pHSCs, *Hoxb5*⁻ ST-HSCs, and *Hoxb5*⁺ LT-
HSCs (top) and the corresponding surface expression of NEO1 for select populations (bottom).
Colors correspond to populations shown. Goat IgG isotype control for fluorescence staining with

goat anti-mouse/human NEO1 antibody is highlighted in orange (bottom). (**D**) Barplots showing
the percent of NEO1$^+$ cells for each cell type gated in **c**. Statistical significance was calculated by
a paired, two-tailed Student's t-test between '*Hoxb5*$^+$ LT-HSC' and each cell type. ** P < 0.01, *** P
< 0.001. Barplots in this figure indicate mean ± SD.

Figure 2 | NEO1⁺*Hoxb5*⁺ LT-HSCs selectively expand during aging and respond to myeloablative stress. (**A to C**) Frequency and number of *Hoxb5*⁺ LT-HSCs and *Hoxb5*⁻ ST-HSCs at 2 (*n* = 7 mice), 5 (*n* = 7 mice), 13 (*n* = 6 mice), and 22 (*n* = 9 mice) months of age. (**A**) Percent of cells among pHSCs that are *Hoxb5*⁺ (orange) and Hoxb5⁻ (green). Statistical significance was calculated using an unpaired, two-tailed Student's *t*-test between 2 months and each time point. ns = non-significant, *P* ≥ 0.05. (**B**) Number of *Hoxb5*⁺ LT-HSCs per million whole bone marrow (WBM) cells. Statistical significance was calculated using an unpaired, two-tailed Student's *t*-test between 2 months and 22 months of age. **P < 0.05. (**C**) Number of *Hoxb5*⁻ ST-

HSCs per million whole bone marrow (WBM) cells. Statistical significance was calculated using
an unpaired, two-tailed Student's *t*-test between 2 months and 22 months of age. ***$P < 0.001$.
(**D to F**) Frequency and number of NEO1$^+$ and NEO1$^-$*Hoxb5*$^+$ LT-HSCs at 2 ($n = 7$ mice), 5 ($n =$
7 mice), 13 ($n = 6$ mice), and 22 ($n = 9$ mice) months of age. (**D**) Percent of cells among *Hoxb5*$^+$
LT-HSCs that are NEO1$^+$ (red) and NEO1$^-$ (blue). Statistical significance was calculated using an
unpaired, two-tailed Student's *t*-test between 2 months and each time point. ***$P < 0.001$, ****$P <$
0.0001 (**E**) Number of NEO1$^+$ *Hoxb5*$^+$ LT-HSCs per million whole bone marrow (WBM) cells.
Statistical significance was calculated using an unpaired, two-tailed Student's *t*-test between 2
months and 22 months of age. *$P < 0.05$. (**F**) Number of NEO1$^-$ *Hoxb5*$^+$ LT-HSCs per million
whole bone marrow (WBM) cells. Statistical significance was calculated using an unpaired, two-
tailed Student's *t*-test between 2 months and 22 months of age. ns = non-significant, $P \geq 0.05$. (**G**
to I) Response of HSC subpopulations from 4-month-old *Hoxb5*-mCherry mice 5 days after
treatment with 5-fluoruracil (5-FU). (**G**) Experimental design of myeloablative stress with 5-FU (n
$= 6$ mice) with PBS control ($n = 5$ mice). (**H**) Frequency of *Hoxb5*$^+$ LT-HSCs and *Hoxb5*$^-$ ST-HSCs
among all pHSCs 5 days after treatment with 5-FU or PBS. Statistical significance was calculated
using an unpaired, two-tailed Student's *t*-test. ns = non-significant, $P \geq 0.05$. (**I**) Frequency of
NEO1$^+$ and NEO1$^-$ *Hoxb5*$^+$ LT-HSCs among all *Hoxb5*$^+$ LT-HSCs 5 days after treatment with 5-
FU or PBS. Statistical significance was calculated using an unpaired, two-tailed Student's *t*-test.
**$P < 0.01$. All barplots in this figure indicate mean ± SD.

873
Figure 3 | Neogenin-1 marks a more proliferative fraction of LT-HSCs. (**A to F**) Cell cycle
analysis of 2-to-3-month-old ($n = 10$ mice) and 12-to-14-month-old ($n = 10$ mice) LT-HSC fractions
with Ki-67 and DAPI staining. (**A to C**) Percent of *Hoxb5*+ LT-HSCs or *Hoxb5*− ST-HSCs in (**A**)
G_0, (**B**) G_1, and (**C**) G_2/S in 2-to-3-month-old and 12-to-14-month-old mouse bone marrow.
Statistical significance was calculated using an unpaired, two-tailed Student's *t*-test. ns = non-
significant, $P \geq 0.05$. (**D to F**) Percent of NEO1+ or NEO1− *Hoxb5*+ LT-HSCs in (**D**) G_0, (**E**) G_1, and
(**F**) G_2/S in 2-to-3-month-old and 12-to-14-month-old mouse bone marrow. Statistical significance
was calculated using an unpaired, two-tailed Student's *t*-test. ns = non-significant, $P \geq 0.05$, *$P <$
0.05, **$P < 0.01$. All barplots in this figure indicate mean ± SD.

Figure 4 | NEO1⁺ *Hoxb5*⁺ LT-HSCs exhibit myeloid bias and reduced reconstitution potential upon serial transplantation. (**A**) Experimental design for primary and secondary transplantations of NEO1⁺ and NEO1⁻*Hoxb5*⁺ LT-HSCs. (**B to D**) Measurements of reconstitution potential and lineage priming after primary transplantation ('NEO1⁺', n = 14 mice; 'NEO1⁻', n = 11 mice). (**B**) Percent of donor-derived (CD45.2⁺) cells among all peripheral blood cells at 4, 8, 12, and 16 weeks post-transplant. (**C**) Percent of myeloid cells (GR1⁺CD11B⁺ granulocytes and monocytes) among donor-derived (CD45.2⁺) cells at 4, 8, 12, and 16 weeks post-transplant. (**D**) Percent of lymphoid cells (B220⁺ B cells and CD3⁺ T cells) among donor-derived (CD45.2⁺) cells at 4, 8, 12, and 16 weeks post-transplant. (**E to G**) Same as in **B to D** but analyzing peripheral blood in secondary recipients transplanted with 1000 donor-derived Lin⁻c-KIT⁺SCA1⁺ (KLS) cells from primary hosts ('NEO1⁺', n = 8; 'NEO1⁻', n = 9). Statistical significance for **B to G** was calculated using two-way ANOVA with time post-transplant and NEO1 status as factors. **P < 0.01, ns = non-significant. All line plots in this figure indicate mean ± SEM.

Figure 5 | Distinct transcriptional signatures of NEO1⁺ and NEO1⁻ *Hoxb5*⁺LT-HSCs. (**A**) Experimental design for bulk RNA-sequencing of NEO1⁺ and NEO1⁻ *Hoxb5*⁺ LT-HSCs. (**B**) Heatmap of differentially expressed genes (n = 1,036 genes; FDR < 0.1) after pairwise comparison of NEO1⁺ (n = 5 samples) and NEO1⁻ (n = 5 samples) *Hoxb5*⁺ LT-HSC transcriptomes using DESeq2. Select genes are highlighted. Genes are ordered from left to right by log$_2$ fold enrichment in NEO1⁺ over NEO1⁻ *Hoxb5*⁺ LT-HSCs. (**C and D**) Gene set enrichment analysis (GSEA) plots of molecular signatures significantly enriched (Q value < 0.05) over a gene list ordered by log$_2$ fold change, including (**C**) 'G2_M_HALLMARK' (*top*), 'RIBOSOME_KEGG' (*bottom*), (**D**) Myeloid LT-HSC signature (*top*), and non-myeloid LT-HSC signature (*bottom*) from

Mann et al., 2018 ([24](#)). NES, normalized enrichment score. (**E to G**) Barplots showing $\log_2$ and
DESeq2-normalized gene expression for select genes associated with (**E**) granulocyte or
monocyte, (**F**) platelet, or (**G**) stem programs. Statistical significance was calculated using a
paired, two-tailed Student's *t*-test adjusted for multiple hypothesis testing with Benjamini-
Hochberg procedure. *P-adjusted < 0.05, **P-adjusted < 0.01, ***P-adjusted < 0.001, *****P-*
adjusted < 0.0001. All barplots in this figure indicate mean ± SEM.

Figure 6 | NEO1⁻*Hoxb5*⁺ LT-HSCs outcompete NEO1⁺*Hoxb5*⁺ LT-HSCs in reconstitution potential and reside at the apex of the hematopoietic hierarchy. (**A**) Experimental design for co-transplantation of NEO1⁺ and NEO1⁻*Hoxb5*⁺ LT-HSCs (*n* = 12 mice). (**B to D**) Measurements of reconstitution potential and lineage priming after co-transplantation. (**B**) Percent of donor-derived cells among all peripheral blood cells at 8, 12, 16, and 20 weeks post-transplant. (**C**) Percent of myeloid cells (GR1⁺CD11B⁺ granulocytes and monocytes) among donor-derived cells at 8, 12, 16, and 20 weeks post-transplant. (**D**) Percent of lymphoid cells (B220⁺ B cells and CD3⁺ T cells) among donor-derived cells at 8, 12, 16, and 20 weeks post-transplant. Statistical significance for **B to D** was calculated using two-way ANOVA with time post-transplant and NEO1 status as factors. **$P < 0.01$. (**E**) Percent of platelets (EGFP⁺CD41⁺) among donor-derived cells at 8, 12, 16, and 20 weeks post-transplant. Statistical significance at 20 weeks post-transplant

was calculated using an unpaired, two-tailed Student's t-test. $*P < 0.05$. (**F**) Percent of NEO1$^+$
and NEO1$^-$ $Hoxb5^+$ LT-HSCs derived from donor NEO1$^+$ and NEO1$^-Hoxb5^+$ LT-HSCs in the
mouse bone marrow 20 weeks post-transplant. Only samples for which $Hoxb5^+$ LT-HSCs were
present are shown ('NEO1$^+$ derived', $n = 9$; 'NEO1$^-$ derived', $n = 8$). Statistical significance was
calculated using a paired, two-tailed Student's t-test between the percent of NEO1$^+$ and NEO1$^-$
$Hoxb5^+$ LT-HSCs derived from the same donor and an unpaired, two-tailed Student's t-test
between the percent of NEO1$^-Hoxb5^+$ LT-HSCs between NEO1$^+$ and NEO1$^-$ donors. P values
are indicated on the graph. (**G**). Schema depicting a revised model of long-term hematopoiesis
with a lineage-balanced, quiescent NEO1$^-Hoxb5^+$ LT-HSC residing above a downstream
myeloid-biased NEO1$^+Hoxb5^+$ LT-HSC. The thickness of the lines indicates the degree of
contribution and dashed lines mark putative differentiation paths. All line plots in this figure
indicate mean ± SEM. Scatter dot plot in **F** indicates mean ± SD.

Figure S1 | Single cell RNA sequencing shows selective expression of Neogenin-1 in a subset of LT-HSCs. Percent of *Neo1*[+] cells among each phenotype defined by the authors of two independent scRNA-seq studies of mouse hematopoietic stem and progenitor cells (HSPCs) (5, 57).

Figure S2 | NEO1 expression in the bone marrow niche. (A) tSNE plot of *Neo1* expression (*top*) among different bone marrow niche populations (*bottom*) from a published scRNA-seq dataset of mouse osteoblasts and bone marrow stromal cells (58). (B) Representative images of bone marrow sections showing NEO1 (red) signal on the surface of trabecular bone (*) and stromal cells. Sinusoids are highlighted with an antibody to Endomucin (green) and nuclei were counterstained with DAPI (blue). "#" marks the bone surface. Scale bar is shown on the graph and boxes indicate corresponding magnified images below.

Figure S3 | Gating scheme for the prospective isolation of NEO1⁺ and NEO1⁻ *Hoxb5*⁺ LT-HSCs by flow cytometry. Arrows indicate the gating sequence. All gates were drawn with respective to fluorescence-minus-one (FMO) controls. SSC, side scatter; FSC, forward scatter; 7-AAD, 7-Aminoactinomycin D.

Figure S4 | Flow cytometry analysis of NEO1 surface expression in the human bone marrow. (**A**) Contour plots with outliers showing the gating scheme for human HSCs, MPPs, and LMPPs (*n* = 3). (**B**) Barplots showing the percent of NEO1[+] cells for each cell type gated in **A**. MPP, multipotent progenitor, LMPP, lymphoid-primed multipotent progenitor. Barplots indicate mean ± SD.

Figure S5 | Fold-change in pHSCs, *Hoxb5*⁻ ST-HSCs, *Hoxb5*⁺ LT-HSCs, and NEO1⁺ and NEO1⁻ *Hoxb5*⁺ LT-HSCs with age. (**A and B**) Number and fold-change in pHSCs from mouse bone marrow at 2 ($n = 7$ mice), 5 ($n = 7$ mice), 13 ($n = 6$ mice), and 22 ($n = 9$ mice) months of age. Barplots showing (**A**) the number of pHSCs per million whole bone marrow (WBM) cells and (**B**) fold-change in pHSCs from 2-month-old mice. (**C**) Flow cytometry diagrams of *Hoxb5*-mCherry (*y*-axis) expression in the mouse bone marrow at 2 ($n = 7$ mice), 5 ($n = 7$ mice), 13 ($n = 6$ mice), and 22 ($n = 9$ mice) months of age. (**D and E**) Barplots showing the fold-change in (**D**) *Hoxb5*⁺ LT-HSCs and (**e**) *Hoxb5*⁻ ST-HSCs from 2-month-old mice. (**f**) Flow cytometry diagrams of NEO1 (*x*-axis) expression in the mouse bone marrow. (**G and H**) Barplots showing the fold-change in (**G**) NEO1⁺ *Hoxb5*⁺ LT-HSCs and (**H**) NEO1⁻ *Hoxb5*⁺ LT-HSCs from 2-month-old mice. For all barplots, statistical significance was calculated using an unpaired, two-tailed Student's *t*-test between 2 months and 22 months of age. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ns = non-significant, $P \geq 0.05$. Bars and error bars indicate mean ± SD.

A

B

Figure S6 | Cell cycle analysis of *Hoxb5⁻* ST-HSCs, *Hoxb5⁺* LT-HSCs, and NEO1⁺ and NEO1⁻
***Hoxb5⁺* LT-HSCs.** (**A**) Pie charts (*top*) showing the average percent of each cell type from 2-to-
3-month-old mouse bone marrow in G_0 (green), G_1 (yellow), and G_2/S (orange) as measured by
flow cytometry analysis of KI-67 and DAPI staining (*bottom*). Same as in (**B**) but with 12-to-14-
month-old mouse bone marrow. KI-67 gates were drawn with respective to a fluorescence-minus-
one (FMO) control.

Figure S7 | Enrichment of GO biological processes and lineage-restricted progenitor signatures in NEO1⁺ and NEO1⁻ *Hoxb5*⁺ LT-HSC transcriptomes. (**A**) Barplots showing log$_2$ and DESeq2-normalized gene expression of *Neo1* in NEO1⁺ and NEO1⁻ cells isolated by flow cytometry. Statistical significance was calculated using a paired, two-tailed Student's *t*-test adjusted for multiple hypothesis testing with Benjamini-Hochberg procedure. **P < 0.01. Bars and error bars indicate mean ± SEM. (**B**) Dot plots showing the top 5 'GO: Biological Process' pathways enriched in NEO1⁺ and NEO1⁻ *Hoxb5*⁺ LT-HSCs by hypergeometric test. *P*-adjusted values are indicated by color gradient and gene ratios by dot size. Number of genes significantly enriched in NEO1⁺ or NEO1⁻ are indicated in parenthesis below each column. (**C**) GSEA plots showing enrichment of gene set signatures of lineage restricted progenitors from a previous study (27) over a gene list ordered by log$_2$ fold change in NEO1⁺ versus NEO1⁻ *Hoxb5*⁺ LT-HSCs. NES, normalized enrichment score and *Q* values are indicated on the graph. MkP, megakaryocyte progenitor; preCFU-E, pre-erythrocyte colony-forming units; preGM, pre-granulocyte/macrophage progenitors; preMegE, pre-megakaryocyte/erythrocyte progenitors; CLP, common lymphoid progenitor.

A

B

Figure S8 | Association between NEO1 and CD150 in *Hoxb5*+ LT-HSCs by flow cytometry. (**A**) Histogram (*left*) and bar plots (*right*) showing the percent of NEO1+ cells in three bins of CD150 expression, including 'High', 'Mid', and 'Low'. Statistical significance was calculated by a paired, two-tailed Student's *t*-test between 'High' and 'Low'. ***P < 0.001. Bars and error bars (*right*) indicate mean ± SEM. (**B**) Representative flow cytometry diagram of NEO1 fluorescent intensity (*x*-axis) and CD150 fluorescent intensity (*y*-axis) with a linear regression line and Pearson correlation coefficient shown.

Post-transplant contribution to
Hoxb5⁺ LT-HSCs

Figure S9 | Post-transplant contribution of NEO1⁺ and NEO1⁻ *Hoxb5*⁺ LT-HSCs to *Hoxb5+ LT-HSCs*. Bar plot showing the number of *Hoxb5*⁺ LT-HSCs per million whole bone marrow (WBM) cells from each donor population 20 weeks after co-transplant of NEO1⁺ and NEO1⁻ *Hoxb5*⁺ LT-HSCs ($n = 10$). Bars and error bars indicate mean ± SD. Statistical significance was calculated by a paired, two-tailed Student's *t*-test. ns = not significant, $P \geq 0.05$.