1 Untangling the Indonesian tangle net fishery: describing a data-poor

2 fishery targeting large threatened rays (Order Batoidea)

Brooke M. D'Alberto[1, 2*], William T. White[2, 3], Andrew Chin[1], Dharmadi[4], Colin A.

Simpfendorfer[1]

[1] Centre for Sustainable Tropical Fisheries and Aquaculture & College of Science and Engineering,

James Cook University, Townsville, 4810 Australia

[2] CSIRO Oceans and Atmosphere, Hobart, 7004 Australia.

[3] Australian National Fish Collection, CSIRO National Research Collections Australia, Hobart, 7004

Australia.

[4] Research Centre for Fisheries Management and Conservation, Ministry of Marine Affairs and

Fisheries, Government of Indonesia, Jakarta, 10110 Indonesia

[*]Corresponding author: brooke.dalberto@my.jcu.edu.au (B.M.D.)

18 Abstract

1. Shark-like rays (Order Rhinopristiformes) are among the most threatened families of marine

fish. Yet little is known about their populations, as these rays are normally taken as

opportunistic catch in fisheries targeting other species and are thus poorly reported. One

exception is the Indonesian tangle net fishery, which targets shark-like rays.

2. Market surveys of Muara Angke landing site in Jakarta, north-western Java (including one

frozen shipment from Benoa Harbour, Bali), were conducted between 2001 and 2005, and

recorded landed catch for this fishery. Recent catch data from Indonesian Capture Fisheries

(2017 – 2018) were also examined to provide contemporary information about landed catch.

3. 1,559 elasmobranchs (sharks and rays) were recorded, comprised of 24 species of rays and

nine species of sharks. The most abundant species landed were the pink whipray *Pateobatis*

fai and the bottlenose wedgefish *Rhynchobatus australiae*, the latter being the main target

species.

4. Catch composition varied based on differences in species catchability and may also be

indicative of localized declines. The fishery was highly selective for larger sized individuals,

however smaller size classes of target species were also caught in other Indonesian fisheries

resulting in fishing pressure across all age classes.

5. Evidence of substantial declines in global landings of wedgefish species, and the observed shift in catch composition in the Indonesian tangle net fishery, increases concerns about the status of shark-like rays and stingrays in Indonesia.

Keywords: Conservation, giant guitarfish, liong bun net, South-East Asia, stingrays, threatened species, wedgefish.

1. Introduction

Rays (Superorder Batoidae) are among the most threatened groups of chondrichthyans (sharks, rays, and chimaeras) (Dulvy et al., 2014a). Substantial declines in populations, catch rates and landings, as well as localised extinctions have been reported for several ray species (Moore, 2015). Declines in elasmobranchs populations are mainly the result of the rapid expansion of chondrichthyan catch in target and non-target fisheries (Dulvy et al., 2014a, Clarke et al., 2006), and the globalisation of trade (Lack and Sant, 2009, Clarke et al., 2007). Recently, the global chondrichthyan catch has been increasingly dominated by rays (Dulvy et al., 2014a). This increase is likely the result of (a) improved reporting, and (b) declines in shark catches due to stronger national and international regulations (Dulvy et al., 2014a). Rays, like most chondrichthyans, have intrinsically low biological productivity, due to their slow growth, late maturity, long generation times, and low fecundity, and are therefore slow to recovery from population declines (Fowler et al., 2002). Most ray species (excluding deep sea skates) are exposed to high levels of intense and expanding fishing pressure due to their strong association with soft bottom habitats in shallow (<100 m depth) tropical and temperate coastal waters (Last et al., 2016). Many species of rays play an important trophic role in soft sediment ecosystems as bioturbators (Kyne and Bennett, 2002, White et al., 2013, Flowers et al., 2020), and their coastal habitats are under threat from additional anthropogenic influences (Dulvy et al., 2016, Compagno and Cook, 1995).

Wedgefish (Family Rhinidae, 10 species), and giant guitarfish (Family Glaucostegidae, 6 species), are large (maximum size 300 cm total length, TL) benthic rays collectively referred to as shark-like rays (Last et al., 2016). They are found throughout the Indo-Pacific, Indian and Atlantic-Mediterranean ocean in shallow, coastal waters adjacent to coral reefs (Last et al., 2016). Shark-like rays are mainly caught as bycatch in fishing gears such as trawl nets, pelagic and bottom long lines, purse seine nets, and gillnets, and are typically retained as valuable by-products of opportunistic catch (Moore, 2017, Jabado, 2018). Shark-like ray fins are considered the highest grade in the international shark fin trade, which is likely the key driver for their retention in coastal fisheries (Keong, 1996). There are few documented targeted fisheries for these species (White and Dharmadi, 2007). Wedgefish and giant guitarfish are experiencing significant declines throughout their entire ranges (Kyne et al., 2019), and

all but one species of wedgefish and giant guitarfish species are classified as Critically Endangered on

the International Union for Conservation of Nature's (IUCN) Global Red List of Threatened Species

in 2019 (Kyne et al., 2020). These species were listed on Convention of International Trade of

Endangered Species of Wild Fauna and Flora (CITES) Appendix II in 2019 (CoP18), which aims to

ensure that the international trade of products from wedgefish and giant guitarfish come from

sustainable sources (CITES, 2019).

Similar to the global decline of sawfishes and angel sharks (Lawson et al., 2020, Dulvy et al., 2016),

depletion of wedgefish and giant guitarfish likely began many decades ago, driven by the incidental

catch in fisheries and the high value of fins in the international trade (Clarke et al., 2006, Keong,

1996, Harrison and Dulvy, 2014). Quantifying the onset and extent of decline of these data-limited

species is difficult, due to depletions occurring before independent scientific monitoring, poor

fisheries and trade reporting resulting in little species-specific data, and lack of conservation

awareness (Lawson et al., 2020, Dulvy et al., 2016). Wedgefish and giant guitarfish have been

inferred to have a higher than average population productivity compared to other chondrichthyans,

and therefore can potentially recover from population declines more rapidly than other threatened

species (D'Alberto et al., 2019). However, achieving this will require significant reductions in fishing

mortality (D'Alberto et al., 2019). Yet there is little information available on shark-like ray and

stingray species' historical and contemporary interactions with fisheries, which may hinder the

development of management and enforcement efforts.

Indonesia is the world's largest contemporary elasmobranch fishing nation, accounting for ca. 13% of

reported global elasmobranch catch (Jaiteh et al., 2017, Blaber et al., 2009). It is also the third largest

exporter of shark fins in regards to quantity, with an average of 1,235 tonne and sixth largest in value,

worth an average of US$10 million per year (Dent and Clarke, 2015). Apart from the fin trade,

elasmobranch meat is also an important source of protein for communities in South East Asia (Ahmad

et al., 2016), and a large volume of elasmobranch products, particularly stingrays, continue to be

exported to regional markets such as Singapore (Clark-Shen et al., 2021). In Indonesia, wedgefishes

and guitarfishes are caught as bycatch in a variety of fisheries, but they are also specifically targeted

in the tangle net fishery, with *Rhynchobatus australiae* the main target species. The fishery also lands

stingrays and sharks as opportunistic catch (Keong, 1996, White and Dharmadi, 2007). This fishery

uses large mesh (50 – 60 cm) bottom-set nylon gillnets, to capture large elasmobranchs by

entanglement, on sandy or muddy substrates between 25 – 45 m depth (Amir, 1988). Fishing vessels

in this fishery are typically refurbished 10 – 22 meter (30 – 40 tonnes) long wooden purse seine

vessels (Amir, 1988). They have an approximate net load capacity of 15 tonnes, and are equipped

with a diesel powered inboard engine and a small freezer to hold fins (Amir, 1988). In Indonesia, this

fishery is referred to as "jaring liongbun" [= guitarfish gillnet] and "jaring cucut" [= shark gillnet].

The first records of this fishery are from Aru Island in the mid-1970s, from which the fishery rapidly

expanded throughout Indonesian waters (Keong, 1996). It reached peak fishing capacity in 1987 with

500 active vessels and began operating at other ports, before declining by 80% over a ten year period

(Amir, 1988, Dharmadi and Kasim, 2010). Declines of *Rhynchobatus* spp. began around 1992

according to local fishers that operate bottom set gillnets in the Aru-Arafura Sea (Jaiteh et al., 2016).

The reductions in vessels operating in the tangle net fishery suggests that populations of the target

species had declined and made the fishery economically unviable. The high value of wedgefish and

giant guitarfish fins was a particularly strong driver for this fishery, with the fins worth approximately

1.5 times more than those from other species (Keong, 1996). More recently, high valued leather

products from stingrays, which appear to be increasing in demand (Karthikeyan et al., 2009,

Sahubawa et al., 2018), drives the retention of large stingrays. There is strong anecdotal evidence of

declines of wedgefish and giant guitarfish in some areas of Indonesia as a result of this fishery (Amir,

1988, Keong, 1996). However, there are no catch and size composition data available for this fishery,

and the fishery as a whole is poorly defined and little understood.

To achieve sustainable use of these species, managers and conservation practitioners need to

understand their population status, risk exposure, and resilience to fishing pressure and other threats.

This requires data on fisheries catch composition, changes in relative abundance, and their

interactions with fisheries (Simpfendorfer et al., 2011, Jabado, 2018). The main aims of this paper

were to (1) examine the species, size, and sex composition of the landed catch of Indonesian tangle

net fishery in 2001 – 2005, and document changes over time in abundance and species compositions,

(2) and to examine the potential consequences of these trends for future fisheries management and

species conservation. Information of the Indonesian tangle net fishery can be used to inform the basis

for the development of local and international management plans and conservation action for

threatened rays.

2. Materials and Methods

2.1. Muara Angke landing port surveys

Between April 2001 and December 2005, elasmobranch landed catches from the tangle net fishery

were recorded at the Muara Angke landing port (North Jakarta, Indonesia) (Fig. 1) and the adjacent

village, where post-landing processing of fish occurred. Landing port surveys were conducted on 18

occasions, and for each visit the landing port was surveyed for 1 – 4 consecutive days, resulting in a

total 53 sampling days (see Supplementary Table S1). Meanwhile, data on the nature of the products

retained, their use, value, and export destination were collected by fisheries officers from the Ministry

of Marine Affairs and Fisheries (MMAF). The prices reported for these products have been converted

to Indonesian Rupiah (IDR) in 2021 values to account for inflation (www.inflationtool.com; 1 IDR

2004 = 2.54 IDR 2021; 1 IDR 2005 = 2.38 IDR 2021), and to US Dollar (USD) prices using an online

currency converter (www.xe.com/currencyconverter/; $1 USD = 13,957.10 IDR as of January 2021).

The number of each species landed from a tangle net vessel was recorded. Due to the large number of

landings and time constraints on each day surveyed, the number of specimens, biological data and

measurements could not be taken from all elasmobranches present. Only specimens that could be

accessed were surveyed, so randomised selection for sex/size was not possible. At the Muara Angke

landing port, catch composition could only be recorded for a brief period while the vessels were being

unloaded (Fig. 2a, b). After the catches were unloaded, elasmobranchs were taken to the adjacent

village processing area, located less than a kilometre from the fishing port. Here the large

elasmobranchs from the tangle net fishery were typically taken to one of the four processing 'houses'

(Fig. 2d). Similar data could be obtained at the village processing area, often from the previous day's

landings, but it was not possible to determine how many vessels they originated from if more than one

vessel had landed in the previous two days. Species and size composition data was more readily

collected during the unloading from the vessel at the fishing port. On days when catches were

recorded in Muara Angke landing port, catches were not examined again in the village processing

area. Due to the relatively low number of landings observed per trip, this issue was rarely encountered

(Table S1). On one occasion, the landings from tangle net fishing vessels operating in the Banda and

Arafura Seas that land at Benoa Harbour, Bali (Fig. 1), were observed and recorded in the Muara

Angke processing village. These catches arrived into the village processing area by freezer truck

direct from Bali.

Often individual tangle net vessels would come into Muara Angke port once a month, and on three

occasions it was possible to document the entire landed catch from three tangle net vessels. The entire

landed catch from three vessels were recorded in Muara Angke landing site in July 2004, October

2004 and two in October 2005, and referred to as MA-SKR-170704, PV-PK-061004, and PV-UK-

051005, respectively (Table 1). The landed catches from the vessels PV-KA-051005 and PV-UK-

051005 were recorded on the same day. However, not all catch was able to be examined for PV-UK-

051005, and the recorded catches for PV-KA-051005 and PV-UK-051005 were combined and

assigned the vessel identifier code of PV-CC-051005 (Table 1).

Following the methods described above, catch composition of elasmobranchs from other fisheries

were also recorded during the Muara Angke landing site surveys. This included landings from small-

mesh gillnet (<20 cm mesh size) fisheries, Java Sea and Arafura Sea trawl fisheries, southern Java

trammel net fishery, and various hand- and long-line fisheries, which were operating out of the

landing ports surveyed [see White and Dharmadi (2007)]. This allowed the comparison of the size

composition of species between the tangle net fishery and the other fisheries interacting with the same

species. Similar to the tangle net fishery landings, only landed catch that could be accessed when a

vessel was unloading was able to be surveyed, and randomised selection was not possible.

2.2. Biological data

When possible, the disc width (DW) for the Dasyatidae, Myliobatidae, Aetobatidae, Gymnuridae and

Rhinopteridae, and total length (from the tip of the snout to the tip of the upper lobe of the caudal fin;

TL) for the sharks and shark-like rays (Pristidae, Glaucostegidae and Rhinidae) were measured to the

nearest 1 cm, and sex recorded. As the shark-like rays were typically landed without fins, an estimated

TL was recorded when animals were landed without fins. Occasionally, the removal of fins from these

rays occurred following landing, and after weighing of specimens. Total weight (TW) of whole

individuals (fins attached and not gutted) was recorded to the nearest g or kg (depending on the size of

the individual), however, the vast majority of rays and sharks could not be weighed at the landing

port. When large numbers of similar sized individuals were observed, measurements were taken from

a subset of whole individuals that could be accessed, and used to estimate DW, TL and TW for the

remaining unmeasured individuals. For specimens that were not weighed but had length

measurements, the weight of individual specimens were calculated using the length to weight

conversion equations (see Supplementary Table S2). For species where a length-weight equation was

not available, the estimated weight was calculated using a conversion equation from a

morphologically-similar species (Table S2). In instances when the size of individuals for a particular

species was not recorded, the weight was estimated using the average weight of the individuals for

that species. Total weight was then determined for each species landed in the fishery. Details on the

reproductive biology of each species recorded were previously reported in White and Dharmadi

(2007).

2.3. Species identification

Species were identified using the keys in Carpenter and Niem (1998) and Last and Stevens (1994),

with taxonomic nomenclature updated using Last et al. (2016) and Last and Stevens (2009). The

identity of a subsample of *Rhynchobatus* species caught in the tangle net fishery was also further

verified by genetic analysis (Giles et al., 2016), and from images that used recently determined colour

pattern differences between species (Last et al., 2016). The key colour pattern difference used to

differentiate *R. australiae* from its closest regional congeners was the pattern of white spots around

the dark pectoral spot present in all but the largest individuals (Fig. 2d). In *R. australiae*, there is a line

of 3 white spots located adjacent to the black pectoral spot, or its usual position if faded (Last et al.,

2016). In two large pregnant females which possessed no white spots or black pectoral spots, due

mainly to their poor condition, the typical *R. australiae* spot pattern was evident in late-term embryos

allowing for confirmation of their identity. *Maculabatis macrura* was only recently recognised as

valid and distinct from *Maculabatis gerrardi* (Last et al., 2016), therefore our data could not be

retrospectively confirmed as either or both species (Last et al., 2016). Although these records could

constitute either species, herein we refer to these species as the prevailing name used for the

whitespotted ray species in Indonesia, *M. gerrardi*.

2.4. Data analysis

223 Species composition of each landing was expressed as percentage of the total number of individuals

224 by both the number and weight for each species recorded at the landing port and processing village.

225 Minimum, maximum and mean ± standard error (S.E.) for DW, TL and TW are reported for each

226 species. Size frequency histograms for the most abundant species were produced.

2.5. Ethics Statement

229 All marine life examined in this study were landed from fisheries in Indonesia and were already dead

230 upon inspection. Permission to undertake surveys in Indonesia was granted by the Research Centre for

231 Fisheries Management and Conservation in Jakarta as part of collaborative projects funded by the

232 Australian Centre for International Agricultural Research (ACIAR project codes FIS/2000/062 and

233 FIS/2003/037).

3. Results

3.1. Species and size composition of the fishery

Across 18 sampling trips, totalling 53 survey days from April 2001 – December 2005, tangle net

vessel landings were recorded in Muara Angke landing site eight times, and within the village

processing area seven times, including one frozen shipment from Benoa Harbour, Bali (Table 1; Table

S1). From discussions with a fleet manager, thirteen vessels were reported to be operational in the

tangle net industry in 2004, fishing in waters around Borneo, Sulawesi and as far as West Papua (Fig

1). Only vessels that landed at Muara Angke landing site or processing village during the 2001-2005

surveys were sampled (Table 1).

A total of 1,559 elasmobranchs were recorded from tangle net fishery landings at Muara Angke

landing port. This comprised 1,526 batoids (98.3% of the catch), of 24 species from seven families

(Table 2). The most abundant family was Dasyatidae, contributing 72.5% to the total number of

elasmobranchs recorded, followed by the family Rhinidae, and comprised 20.8% of total observed

catch. Only 25 sharks were recorded, with nine shark species from four families (Table 2). Combining

the total landed catch of the four Indonesian tangle net vessels, of which the catch was fully

documented from Muara Angke landing site, the most abundant species was *M. gerrardi*, followed by

R. australiae, P. ater, R. ancylostoma, A. ocellatus, H. uarnak, and *P. jenkinsii* (Fig. 5). In total,

eighteen other species of elasmobranchs, comprising 10 ray species and 8 shark species, were also

recorded but in low numbers. As not all individuals could be counted when landed, and in many cases

the numbers were estimated, the numbers presented represent an underestimation of the total number

of individuals caught during the survey period.

3.1.1. Family Pristidae

Two largetooth sawfish *Pristis pristis* were recorded from the Benoa Harbour, Bali landings in August

2002. It is likely the sawfish were caught in the Arafura or Banda Sea region (Fig. 1). Both

individuals were adult males and ca. 420 cm TL and had an estimated total landed weight of 220 kg

(Table 2).

3.1.2. Family Glaucostegidae

Two species of giant guitarfish were recorded in Muara Angke landing site: the clubnose guitarfish

Glaucostegus thouin, and giant guitarfish *Glaucostegus typus.* The presence of *G. thouin* in the fishery

was recorded once in April 2001. However due to the logistics of accessing these rays upon unloading

from the vessel and the decayed state of some specimens, the estimates of numbers or size were not

possible for *G. thouin,* and it was unclear if it was a regular catch in the fishery. Fourteen *G. typus*

were recorded on three occasions (Table 2). Of the subset of *G. typus* measured, seven were females,

six males and one not sexed, with an estimated total landed weight of 386 kg (Table 2).

3.1.3. Family Rhinidae

Three species of wedgefish were recorded: the bowmouth guitarfish *Rhina ancylostoma*, the eyebrow

wedgefish *Rhynchobatus palpebratus,* and *R. australiae*. *Rhina ancylostoma* were recorded on seven

occasions over the sampling period (Table S3). The landed catch of *R. ancylostoma* comprised 15

females, 10 males, and 32 specimens counted but not sexed, with an estimated total landed weight of

4.4 tonnes (Table 2). Females ranged from 139 – 250 cm TL and 22.9 – 133.6 kg, and males ranged

from 130 – 260 cm TL and 18.7 – 150.3 kg. One unsexed specimen was measured at 270 cm TL and

168 kg. The main target species of the tangle net fishery, *R. australiae* comprised the largest

component of wedgefishes and the second most abundant species recorded (Table 2; Fig. 5). On one

occasion, approximately 7.1 tonne of *R. australiae* was landed from a single tangle net vessel.

Rhynchobatus australiae was recorded on eight occasions (Table S3) and a total of 238 individuals

with an estimated total landed weight of 24 tonnes, comprising 99 females, 18 males and 121 unsexed

individuals (Table 2). A subset of 29 individuals were measured, the majority of which were females

and approximately 300 cm TL (Fig. 2a). Of the subset, 16 individuals of *R. australiae* examined

internally were pregnant as reported in White and Dharmadi (2007).

A total of 16 *Rhynchobatus* spp. landed in the tangle net fishery had their identifications confirmed by

genetic analysis [see Giles et al. (2016)]. Of these 16, two were from landings at Muara Angke and

the remaining 14 from the Benoa Harbour landings. The 14 Benoa Harbour individuals consisted of

five *R. palpebratus* [reported as *R. palpebratus/Rhynchobatus cf laevis* in Giles et al. (2016)] and nine

R. australiae. The ratio of *R. palpebratus* to *R. australiae* determined from the genetic analysis (45:6)

was used to estimate the species composition of the 100 *Rhynchobatus* individuals recorded in the

Benoa Harbour landings on the 14[th] August 2002. *Rhynchobatus palpebratus* was recorded on one

occasion, with a total of 30 individuals but not sexed, measured or weighed (Table 2).

3.1.4. Family Dasyatidae

Stingrays were present in every tangle net catch landed in Muara Angke. A total of 1130 stingrays,

with an estimated mass of 30.2 tonnes, were recorded comprised of 13 species from eight genera

(Table 2). The majority of the specimens caught of each species were near or at a larger size than their

known size at maturity (Fig. 3, 4; see Supplementary Table S4). The most abundant stingray species

were *P. fai* (9.6 tonnes; Fig. 3f), broad cowtail ray *Pastinachus ater* (7.5 tonnes; Fig. 3e), *M. gerrardi*

(2.3 tonnes; Fig. 3d; Fig. 5), Jenkin's whipray *Pateobatis jenkinsii* (2.7 tonnes; Fig. 4a), and

whitenose whipray *Pateobatis uarnacoides* (2.5 tonnes; Fig. 4b). Other species that were recorded

were the brown stingray *Bathytoshia lata,* leopard whipray *Himantura leoparda* (Fig. 3b), coach

whipray *Himantura uarnak* (Fig. 3c), honeycomb whipray *Himantura undulata*, blackspotted whipray

Maculabatis astra, smalleye stingray *Megatrygon microps*, blotched stingray *Taeniurops meyeni* (Fig.

4c) and porcupine ray *Urogymnus asperrimus*. *Maculabatis astra* was only recorded from the single

Benoa Harbour landing, this species is only present in the far eastern portion of Indonesia off West

Papua and is allopatric from *M. gerrardi* (Last et al., 2016). Specimens of *H. leoparda, H. uarnak, P.*

fai, P. uarnacoides and *T. meyeni* specimens were close to the known maximum size (Fig. 3, 4).

3.1.5. Family Aetobatidae

315   One aetobatid species, the spotted eagle ray *Aetobatus ocellatus*, was recorded in the tangle net

316   fishery on eight occasions (Table S3). A total of 45 individuals were observed, with an estimated total

317   landed weight of 1.8 tonnes (Table 2). This comprised of 21 females, 22 males, and two unsexed

318   specimens (Fig. 4d). *Aetobatus ocellatus* specimens were mainly caught close to or at a greater size

319   than the known size at maturity (Fig. 4d; Table S4).

3.1.6. Other families

Several other species from a range of families were recorded occasionally or in relatively small

quantities. This included one species of Myliobatidae, the ornate eagle ray *Aetomylaeus vespertilio*,

with an estimated landed catch of 1.1 tonnes, of which five were females, one male, and five unsexed

individuals (Table 2). These specimens were all large, including one 160 kg female, and were

recorded occasionally (Table S3), and comprised a small proportion of the total landed catch. A

single Gymnuridae species was recorded, the zonetail butterfly ray *Gymnura zonura* with seven

individuals recorded (Table 2). Sharks were a minor part of the catch and rarely observed. All of the

shark species represented less than 1% of the total catch (Table 2). Of the small amount of sharks

taken in the fishery, the species most commonly occurring in the catch were the common blacktip

shark (*Carcharhinus limbatus*; n = 6), tiger shark (*Galeocerdo cuvier*; n = 4), and hammerheads

(*Spyhrna* spp; n = 5) (Table 2; Table S3).

3.2. Products from the tangle net fishery

Data from MMAF indicated that the main products derived from the tangle net fishery catch were

elasmobranch fins, leather made from ray skins, salted elasmobranch meat, and elasmobranch

vertebrae. The most valuable product were fins from wedgefishes and giant guitarfishes. During the

surveys in 2005, the quoted price for fins from sawfish, wedgefish and guitarfish were approximately

3 million IDR kg^{-1} wet weight (~ 7,145,108 IDR / $512 USD per kg in 2021 terms). Any fins frozen

on-board the tangle net vessels did not come through the Muara Angke landing site, but instead were

directly exported through a different port to Hong Kong and Singapore.

Stingray skins, used to produce leather products, were the second most valuable product in the fishery

(Fig. 2c). The species primarily used were from the genera *Himantura, Maculabatis*, *Pastinachus*,

Pateobatis and *Urogymnus,* which together were a large component of the landed catch during the

market surveys. *Pateobatis jenkinsii* was reported to be the most sought after stingray skin due to the

row of enlarged thorns which extend down the midline of the body and tail. In 2005, the reported

value of a 13 cm and 18 cm piece of stingray leather was 25,000 (= 59,543 IDR / $4.30 USD in 2021)

and 35,000 IDR (= 83,360 IDR / $5.97 USD in 2021). Approximately 3000 – 4000 skins were

estimated to be exported per month to the Philippines and Japan. Products from the stingray leather

include wallets and belts, which were reported to be sold for approximately 290,000 IDR (= 690,694

IDR / $49.50 USD in 2021) to 500,000 IDR (= 1,190,851 IDR / $85.33 USD in 2021), as well as

bags. Estimated prices for bags were not available.

Ray meat, wedgefish in particular, was considered to be good quality. As the catch was landed in a

deteriorated condition, meat from the rays and sharks was salted and dried. In 2004, in the Muara

Angke processing village, the buying price for wedgefish and guitarfish meat was 4,000 – 5,000 IDR

kg^{-1} (= 10,143 – 12,679 IDR kg^{-1}/ $0.73 – 0.91 USD kg^{-1} in 2021). For stingrays, the meat was valued

between 2,000 – 3,500 IDR kg^{-1} (= 5,072 – 8,875 IDR kg^{-1}/ $0.36 – 0.64 USD kg^{-1} in 2021). Salted

meat was reported to be transported to West Java (Bandung, Bogor, Garut, Cianjur) and Central Java.

Salted meat from wedgefish was reported to be sold for 10,000 – 12,000 IDR kg^{-1} (= 25,358 – 30,430

IDR kg^{-1}/ \$1.82 – 2.20 USD kg^{-1} in 2021), and for stingray meat 6,000 – 8,000 IDR kg^{-1} (= 15,215 –

20,287 IDR kg^{-1}/ \$1.09 – 1.45 USD kg^{-1} in 2021).

The cartilage, such as vertebrae, comprised a small part of the products from this fishery. In 2004,

dried vertebrae were worth approximately 20,000 IDR kg^{-1} (= 50,717 IDR kg^{-1}/ \$3.63 USD kg^{-1} in

2021), which were then processed in Jakarta and exported through an undisclosed port to Korea and

Japan.

3.3. Comparison of the size range of species from tangle net fishery and other fisheries

The gill nets used in the tangle net fishery captured wedgefish and guitarfish over 130 cm TL, and

stingrays over 50 cm DW (Fig. 3), as a result of the large mesh size used (50 – 60 cm). The smallest

recorded individual caught in this fishery was a *P. uarnacoides* of 51.7 cm DW and the largest

recorded individual was a male sawfish, estimated to be 420 cm TL. Meanwhile, smaller size classes

(<50 cm DW) for many ray species were captured in a range of fishing gears used in other fisheries

operating in Indonesian waters during the 2001 – 2005 surveys, including trawl nets, hand- and long-

lines, smaller mesh gillnets, and trammel nets (Fig. 6). These other fisheries greatly increased fishing

selectivity for some species. For example, all size classes of *R. australiae* were captured; neonates

(~45 cm TL) were caught as bycatch in small mesh gillnets (Fig. 6a) while sub-adults (~90 – 130 cm

TL) were captured in the Java Sea trawl fishery, and the larger mature individuals (>170 cm TL) were

taken in hand- and long-line fisheries, likely as by-products (Fig. 6a).

Similar trends are evident for a number of dasyatid rays, with life stages from neonates to sub-adults

w landed in other fisheries during the 2001 – 2005 market surveys (Fig. 6b). Catch of *M. gerrardi* was

recorded from small mesh gillnet fishery (neonates – mature adults; ~20 – 70 cm DW), the Java Sea

trawl fishery (neonates – mature adults; ~20 – 100 cm DW), and the trammel net fishery off southern

Java (juveniles – mature adults; ~30 – 70 cm DW) (Fig. 6b). *Pastinachus ater* was recorded in the

hand- and long-line fisheries (~70 – 110 cm DW), compared to typically larger size range landed in

the tangle net fishery (~80 – 150 cm DW) (Fig. 6b). *Pateobatis fai* was recorded in the Java Sea trawl

fishery (juveniles; ~60 – 70 cm DW), in the Arafura Sea trawl fishery (sub adults; ~139 – 160 cm

DW) and hand- and long-line fisheries (juveniles to mature adults; ~65 – 160 cm DW) (Fig. 6b).

Pateobatis jenkinsii was also exposed to fishing throughout all life stages, from small mesh gillnets

(neonates – sub adults; ~25 – 75 cm DW), hand- and long-line fisheries (juveniles – mature adults;

~40 – 100 cm DW), as well as the tangle net fishery (juveniles – mature adults; ~60 – 140 cm DW)

(Fig. 5b). Catch of *P. uarnacoides* was recorded in the Java Sea trawl fishery (juveniles to mature

adults; ~30 – 60 cm DW), and in the southern Java trammel net fishery (juveniles to mature adults;

~25 – 55 cm DW) (Fig. 6b).

4. Discussion

Wedgefishes and giant guitarfishes face an extremely high risk of extinction (Kyne et al., 2020).

While the status of populations and the exact extent of declines for wedgefish and giant guitarfish are

uncertain, the current data available suggests that the global population declines are substantial (Kyne

et al., 2020, Jabado, 2018, Moore, 2017, Diop and Dossa, 2011). This study advances our

understanding of the data poor tangle net fishery in Indonesia, providing details on the species and

size composition, selectivity of the fishery for larger size classes, and the consequent declines of

shark-like ray populations in Indonesia. In doing so, the study provides important historical context

that provides indications of previous, and potentially reference population sizes, the extent of

declines, and the shift in target species. Population declines were inferred from the declining catch

rates of the target species, *R. australiae*, and the decline in the number of vessels operating in the

tangle net fishery. The number of vessels in the fishery peaked at 500 vessels in 1980s and rapidly

declining to 100 vessels left fishing in the area in 1996 (Keong, 1996), and further declined to seven

vessels in 2017/2018 (see Supplementary Table S5). The significant reduction in the number of

vessels operating in the fishery with the marked change in catch composition over the 28 year period,

suggests that target species populations have declined and the fishery became economically unviable.

Over theses decades, it seems likely that the profits from the fins, salted meat and leather, were

insufficient to maintain the tangle net fishery, over the increasing operating costs and decreases in

elasmobranch catches (Suzuki, 1997). The tangle net fishery was still operational as of July 2018

(Table S5), and the information collected from this study (i.e. the shift to target stingrays and the

values) suggests that the contemporary tangle net fishery is likely reliant on the exploitation of

stingrays for its viability and profitability, and opportunistically exploit the high valued wedgefish and

guitarfish species when they are landed. However, there is no species-specific catch information for

this fishery after 2005 and the landed catch was grouped under single labels such as 'rays', 'sharks',

'wedgefish' and 'mixed', and species diversity and catch composition are poorly understood. Given

the current state of batoids around the world and the scale of Indonesia's shark fishery, the

contemporary catch of the tangle-net fishery requires further investigation.

Opportunistic exploitation occurs when multiple species can be exploited in the same habitat (Branch

et al., 2013). The most desirable and profitable species are targeted and depleted first, before

exploitation switches to a less desirable species, leading to overexploitation and harvesting to

extinction of both the desired and less desired species (Branch et al., 2013, Burgess et al., 2017). The

apparent decline in wedgefish landings and the subsequent increase in stingray landed catches is an

indication of the opportunistic exploitation in the fishery. Both groups are common demersal species

in the Indo–West Pacific, and occupy similar habitats along inshore continental shelves waters to at

least 60 m (White et al., 2006, Compagno and Last, 1999). A decline or change in abundance of

wedgefishes in the fishing ground, could result in the stingrays that occupy the same habitat becoming

the main target catch, with opportunities to retain the higher valued and critically endangered

wedgefish and giant guitarfish species when they are encountered. The time of year may also

influence the species catch composition, however more detailed catch information from throughout

the year is needed to explore this possibility.

There has been an increasing domestic demand for stingray leather in Indonesia over the past 20

years, with the development of a commercial market for stingray leather on the north coast of Central

Java during the period 1980 – 1990 (Ibrahim, 2003). Prior to 1980, stingrays were discard as bycatch

in the tangle net fishery (Ibrahim, 2003, Sahubawa et al., 2018), now they are the main component of

the landed catch in 2018. The calcified denticles in the stingray skin, such as *P. jenkinsii,* results in a

unique and attractive finish to the leather products (e.g. bags, belts, wallets and jewellery), and have a

high economic value (Karthikeyan et al., 2009, Sahubawa et al., 2018). Additionally, stingrays remain

a major fishery export from Indonesia with stingray meat being consumed throughout the region and

conservation concerns growing for species such as *M. gerrardi* and *M. macrura* (Clark-Shen et al.,

2021). This targeted fishing pressure on the stingrays may also be resulting in population declines. It

is likely that stingrays are experiencing similar trends of declines to shark-like rays, given the

increasing pressure on stingrays as wedgefish have become less common. Similar to shark-like rays,

there is little fisheries information on tropical stingrays, and their population status is unknown (Smith

et al., 2008). The loss of large, benthic elasmobranchs may have significant social, cultural, and

economic impacts on the fishers and communities who depend on them (Jaiteh et al., 2017), as well as

ecological consequences, altering important ecological processes such as predator-prey interactions

and bioturbation from benthic feeding (Kyne and Bennett, 2002, Flowers et al., 2020). The occurrence

of large shark-like rays and stingrays and their high economic value has been used as justification for

the continuation of the tangle net fishery (Amir, 1988),but the long-term sustainability of the fishery

needs to be assessed, especially considering the impacts of other fisheries affecting different size

classes.

Length selective fishing mortality often drives a reduction in the catch length composition in heavily

fished elasmobranch populations (Walker, 1998, Stevens et al., 2000). At the time of the landing site

surveys in 2001 – 2005, a number of the abundant species were landed close to the known maximum

sizes. It can thus be inferred that the populations of wedgefishes, giant guitarfishes, and stingrays

were experiencing length selective fishing mortality. It is expected that individuals from

contemporary populations would be reaching a smaller maximum size and younger maximum age

than previous generations due to fishing pressure (Thorson and Simpfendorfer, 2009). Sexual

dimorphism is known to occur for numerous elasmobranch species, including *R. australiae, A.*

ocellatus and *U. asperrimus* (Last et al., 2016)*,* with females attaining a larger maximum size than

males (White and Dharmadi, 2007). Female elasmobranches are more likely to be captured in the

large-meshed tangle nets. The majority of *R. australiae, M. gerrardi, and P. uarnacoides* specimens

were large females, with 16 individuals of *R. australiae* examined internally being pregnant as

reported in White and Dharmadi (2007). The removal of large, breeding individuals from the

population causes a reduction in the reproductive potential, resulting in rapid declines in these

populations (Prince, 2005). The capture of large breeding females coupled with a sharp decline in

catch size suggests that recruitment overfishing may have occurred for several species in the tangle

net fishery. Recruitment overfishing occurs when the breeding stock size is reduced to a point where

future recruitment declines strongly, and/or the reduction of the number of young entering the

population (capture of juveniles), and/or through environmental degradation, which affects the size or

suitability of nurseries (Allen et al., 2013, Pauly, 1988, Walters and Martell, 2004). In fisheries where

only adults or the juveniles are caught, higher levels of fishing can be sustained (Simpfendorfer, 1999,

Prince, 2005).

The impact of other fisheries also likely contributed to these historic declines, and may continue to

affect contemporary fisheries. When all age/size classes are fished, it is much more difficult to

achieve sustainable outcomes. All life stages of these rays are experiencing fishing mortality from

multiple fisheries in Indonesia, as the smaller size classes for many of the species encountered in the

tangle net fishery are also caught as bycatch or opportunistic catch in numerous other fisheries

operating in Indonesian waters, i.e. trawls, hand- and long-lines, smaller mesh gillnets, and trammel

nets. Recruitment overfishing substantially impacts the populations' productivity and may lead to

collapse or prevent the recovery of the population (Allen et al., 2013). Some species of wedgefish and

giant guitarfish have a higher than average theoretical population productivity (i.e. recover at a faster

rate from population declines) for chondrichthyans (D'Alberto et al., 2019). However, these species

still have relatively low reproductive rates and long lifespans compared to teleost fish, and thus can

only withstand modest to low levels of fishing mortality (Dulvy et al., 2014b, Cortés, 2000, Camhi et

al., 1998, Musick, 1999). Combined with life history information, the magnitude of Indonesia

chondrichthyan catches, and the knowledge of the effects of fisheries on large elasmobranch species

that mainly takes adults (Prince, 2005, McAuley et al., 2007b, McAuley et al., 2007a), it is likely that

these populations of rays are experience unsustainable levels of exploitation and have little potential

for recovery without significant reductions in fishing mortality.

The sustainability of fisheries (both harvest and discards) and conservation related research and

management initiatives can be compromised by the misidentification of species (Garcia-Vazquez et

al., 2012). Incorrect species identification is prevalent throughout multi-species fisheries where visual

identification is challenging, and can lead to incidental overfishing, resulting in a higher risk of

extinction for the misidentified species in the fishery (Metcalfe et al., 2007). While the latest available

trade data from the Food and Agriculture Organization (FAO) does distinguish trade in rays and

skates from other elasmobranchs, there is no genera or species specific information (Ramaschiello and

Vannuccini, 2015). In South East Asia, *R. australiae* is the most commonly caught wedgefish species

(Giles et al., 2016), yet due to similarities in morphology and identification issues, it is commonly

confused with other large wedgefish species, in particular with *R. djiddensis, R. laevis* (Giles et al.,

2016) and *R. palpebratus* (Compagno and Last, 2008). Historically, all four species of wedgefish have

been referred to as 'white spotted wedgefish' as the common name and/or *R. djiddensis* as the species

name throughout the Indo-Pacific (Last et al., 2016). A recent clarification of species distributions has

recognised that *R. djiddensis* is restricted to the Western Indian Ocean (Last et al., 2016). A similar

situation has occurred for *G. typus* and *Glaucostegus granulatus,* where there are records of the

sharpnose guitarfish *Glaucostegus granulatus* from the tangle net fishery in 1987, where it constituted

4.6% of the total landed catch (Amir, 1988). This is likely to be a misidentification of *G. typus*, as

prior to 2016, the range for as *G. granulatus* was poorly described with no records to suggest that this

species occurred in Indonesia, and is now known to only occur in the northern Indian Ocean between

Myanmar and the Arabian Gulf (Last et al., 2016). Misidentification of elasmobranchs can be further

compounded by the ambiguity over the ranges of these species, as some species are rarer in landings

and possibly have more of a restricted and/or even fragmented spatial distributions. For example, the

broadnose wedgefish *Rhynchobatus springeri* distribution overlaps with *R. australiae* off Java and

Sumatra (Giles et al., 2016). The landed catches of small *Rhynchobatus* spp. during the 2001 – 2005

surveys could possibly have included this species, as maximum known size for *R. springeri* is 213 cm

TL (Last et al., 2016), compared to 300 cm TL for *R. australiae* (Last et al., 2016, Kyne et al., 2019).

The stingrays, *M. gerrardi* and *M. macrura* have overlapping distributions and differ in mostly subtle

morphological characteristics (Last et al., 2016), thus without genetic identification, the two cannot be

readily differentiated. However, our data could not be retrospectively confirmed as either or both

species. As the catch records may comprise both *M. gerrardi* and *M. macrura,* the numbers presented

may be an overestimation of *M. gerrardi* catch. This complication in species differentiation is an

ongoing issue that complicates efforts to quantify species specific catch and consumption [e.g. Clark-

Shen et al. (2021)]. The lack of species specific reporting was thought to masked the known global

declines for wedgefish and giant guitarfish throughout their distribution, where from historical and

contemporary records, there is evidence of population declines, however the extent of declines are

unknown due to the lack of species specific and time-series data (Kyne et al., 2019). Species

identification training for fisheries officials and observers, and the use of genetic identification such as

in-situ DNA barcoding field kits (Booth et al., 2018), are some methods to address the lack of species

specific information and can help with law enforcement in Indonesia and South East Asia (Tillett et

al., 2012).

Multi-species fisheries are complex social-ecological systems, and successful management will

require significant regulations in governance across local, regional and global scales (Ostrom et al.,

1999). As Indonesian government is a signatory party of CITES, it has taken important steps to

implement international obligations under CITES and FAO National Plan of Action (NPOA) for the

Conservation and Management of Sharks and Rays, with either full species protections (e.g.

sawfishes, mobulids) or export controls, and is working to regulate international trade originating

from Indonesia. Appendix II listed species are allowed to be landed and traded domestically, and there

is no oversight by CITES unless the species is being exported internationally, which requires a

positive Non Detriment Finding. These measures would not necessarily affect the harvest and

domestic use of these species, and do not cover the other 24 non – CITES elasmobranch species

caught in the tangle net fishery. Currently there are no national or regional laws in Indonesia that

specifically regulate the take and use of wedgefish, guitarfishes and stingrays (Rusandi et al., 2019).

Scientific advice and information (e.g. monitoring of fisheries landings, supply chains) will be

required to make management decisions for elasmobranch fisheries in SE Asia (Clark-Shen et al.,

2021). Indonesia has complex fisheries supply and trade chains, from fishers to buyers to exporters,

which have intricate interactions and different drivers occurring throughout the chain (Rusandi et al.,

2019). To ensure the sustainable use of wedgefish, giant guitarfish and stingrays in Indonesia, there

will be a need to target the multiple levels of supply chains, from fisheries to exporters, with strategic

and evidence based methods (Booth et al., 2020). Improved trade monitoring and traceability, as well

as awareness campaigns will be key to reduce fishing mortality, utilisation and trade (Clark-Shen et

al., 2021).

Reducing fishing mortality must be directed at managing extraction as well as increasing the

compliance and enforcement of trade regulations. For a targeted shark and ray fishery, an example of

technical management strategies that could be used to mitigate the risk of overfishing of stingrays,

wedgefish and giant guitarfish in the tangle net fishery may involve the use of gear restrictions, and

spatial and temporal closures (Harry et al., 2011, Yulianto et al., 2018). Given the size selectivity of

the tangle net gear for larger animals, modifying the gear selectivity of the tangle net through the use

of smaller mesh gill nets (> 49 cm) will likely result in a case where large and mature individuals for

R. australiae, M. gerrardi, will rarely be captured and it would likely reduce the fishing pressure on

larger size classes of the other rays e.g. *P. ater, P. fai, P. jenkinsii,* and *P. uarnacoides.* Thus adults

will be subjected to the 'gauntlet effect' and effectively excluded, while juveniles are captured by the

fishery. This harvest strategy is considered an effective method to extract long lived species, provided

that the fishing mortality on adults (Simpfendorfer, 1999, Prince, 2005) from Indonesian hook and

longline fisheries remain low. Research into remediation through post release survivability from

different gear types such as trawls and longlines will be beneficial to understand the risks of certain

gear types and reducing mortality. Spatial and temporal closures (e.g. closure of nursery grounds) may

be used to reduce fishing pressure on these rays, however this will require information on spatial

ecology of wedgefish, guitarfish and stingrays (e.g. potential sex/ habitat segregation, locations of

nursery areas). However, there is little to no such information presently available for Indonesian

waters. It would be difficult to control the level of opportunistic catch and bycatch of rays in all

fisheries in Indonesia with limited fisheries management resources and capacity. Local and regional

specific management will be required to address the overfishing of stocks, including potential

reductions in fishing effort and fishing mortality (Vincent et al., 2014). Any viable management

options and regulations will need to ensure that they are leading to noticeable conservation outcomes

(Booth et al., 2020), as well as positive social and economic outcomes for fishers (Booth et al.,

2019a). Appropriate and economically viable incentives for livelihood alternatives for fishers will be

required to address the issues of poverty, food security and lack of alternatives in resources dependent

communities (MacKeracher et al., 2020). Failure to do so will result in low compliance and illegal

fishing throughout much of their range (Crona et al., 2015, Jaiteh et al., 2017, MacKeracher et al.,

2020). Using a precautionary and holistic risk-based approach like a mitigation hierarchy framework

as proposed by Booth et al. (2019b) would be highly beneficial in Indonesia as it takes into the

account the biological and socio-economic aspect of fisheries. This method has the capability of

dealing with data paucity, and could be used for the tangle net fishery. Regional management across

multiple fisheries and life stages will be required to ensure the sustainability and conservation of

wedgefish, giant guitarfish and stingrays that are being caught across a wide range of fisheries at all

life stages.

604 5. Conclusion

Evidence of substantial and rapid declines in landings of the target species raises concerns about the

status of shark-like rays in Indonesia. The tangle net fishery was established to target wedgefish in the

1980s due to the high value of the fins, but evidence suggests populations of wedgefish have declined

dramatically in many areas. The tangle net fishery remained active in 2018, however the fishery now

appears to be mainly catching large stingrays, which are used for leather and meat, with wedgefishes

only occasionally caught. The continuing pressure on stingrays may also be resulting in population

declines, and the extent of such declines is unknown. Some ray species landed in the tangle net fishery

were also caught in other fisheries at different life stages, requiring management across multiple

fisheries and life stages, instead of single fishery management, to ensure the sustainability and

conservation of rays. There are concerns about the sustainability of the fishery and the population

status of many of ray species targeted. The results of this study emphasise the urgency for effective

management for the conservation of these rays.

Acknowledgements

The authors would like to thank the staff of the fish auction office for the Indonesian Centre for

Fisheries Research for the collection of the landed catch vessels at the Muara Angke landing port

during July 2016 – 2018. This work was funded by the Shark Conservation Fund, a philanthropic

collaborative pooling expertise and resources to meet the threats facing the world's sharks and rays

(B.M.D.). The Shark Conservation Fund is a project of Rockefeller Philanthropy Advisors. The

landing port surveys between 2001 and 2005 were funded by the Australian Centre for International

Agricultural Research (ACIAR; grants FIS/2000/062 and FIS/2003/037) and CSIRO Oceans &

Atmosphere (W.T.W.). B.M.D. was supported through an Australian Government Research Training

Program Scholarship (RTPS). The funders had no role in study design, data collection and analysis,

decision to publish, or preparation of the manuscript. The authors would like to thank Dr Stephanie

Hernandez for the design of the map. Lastly, we dedicate this paper to the memory of 'Pak'

Dharmadi, a leader and pioneer in shark and rays research and management in Indonesia.

Contributions

The manuscript was conceptualised by W.T.W., B.M.D., and C.A.S. The data was collected by

W.T.W. and D.D., and analysed by B.M.D., and W.T.W. The manuscript was written by B.M. D.,

which was reviewed and edited by W.T.W., A.C., D.D., and C.A.S.

Competing interests

The authors declare no competing interests.

ORCID

Brooke D'Alberto: 0000-0002-7834-0211

William T. White: 0000-0001-9705-2453

Andrew Chin: 0000-0003-1813-4042

Colin A. Simpfendorfer: 0000-0002-0295-2238

Data availability

All data generated or analysed during this study are included in this published article (and its

Supplementary Information files).

References

Ahmad, N. I., Wan Mahiyuddin, W. R., Tengku Mohamad, T. R., Ling, C. Y., Daud, S. F., Hussein, N. C., Abdullah, N. A., Shaharudin, R. & Sulaiman, L. H. (2016). Fish consumption pattern among adults of different ethnics in Peninsular Malaysia. *Food & Nutrition Research,* 60**,** 32697.

Allen, M. S., Ahrens, R. N. M., Hansen, M. J. & Arlinghaus, R. (2013). Dynamic angling effort influences the value of minimum length limits to prevent recruitment overfishing. *Fisheries Management and Ecology,* 20**,** 247-257.

Amir, F. (1988). *Suatu studi tentang pengoperasian shark set bottom gill net di perairan Kepulauan An, Maluku.* PhD, Hasanuddin University.

Blaber, S. J. M., Dichmont, C. M., White, W. T., Buckworth, R., Sadiyah, L., Iskandar, B., Nurhakim, S., Pillans, R., Andamari, R., Dharmadi & Fahmi (2009). Elasmobranchs in southern Indonesian fisheries: the fisheries, the status of the stocks and management options. *Reviews in Fish Biology and Fisheries,* 19**,** 367-391.

Booth, H., Muttaqin, E., Simeon, B., Ichsan, M., Siregar, U., Yulianto, I. & Kassem, K. (2018). Shark and Ray Conservation and Management in Indonesia: Status and Strategic Priorities 2018– 2023. *In:* SOCIETY, W. C. (ed.). Bogar, Indonesia: Wildlife Conservation Society.

Booth, H., Pooley, S., Clements, T., Putra, M. I. H., Lestari, W. P., Lewis, S., Warwick, L. & Milner-Gulland, E. J. (2020). Assessing the impact of regulations on the use and trade of wildlife: An operational framework, with a case study on manta rays. *Global Ecology and Conservation,* 22**,** e00953.

Booth, H., Squires, D. & Milner-Gulland, E. J. (2019a). The neglected complexities of shark fisheries, and priorities for holistic risk-based management. *Ocean & Coastal Management,* 182**,** 104994.

Booth, H., Squires, D. & Milner Gulland, E. J. (2019b). The mitigation hierarchy for sharks: A risk based framework for reconciling trade offs between shark conservation and fisheries objectives. *Fish and Fisheries*.

Branch, T. A., Lobo, A. S. & Purcell, S. W. (2013). Opportunistic exploitation: an overlooked pathway to extinction. *Trends in ecology & evolution,* 28**,** 409-413.

Burgess, M. G., Costello, C., Fredston-Hermann, A., Pinsky, M. L., Gaines, S. D., Tilman, D. & Polasky, S. (2017). Range contraction enables harvesting to extinction. *Proceedings of the National Academy of Sciences,* 114**,** 3945-3950.

Camhi, M., Fowler, S. L., Musick, J. A., Brautigam, A. & Fordham, S. V. (1998). Sharks and their relatives: ecology and conservation. *Occasional Paper of the IUCN Species Survival Commision,* 20.

Carpenter, K. E. & Niem, V. H. (1998). *FAO species identification guide for fishery purposes. Vol 2: Cephalopods, crustaceans, holothurians and sharks,* Rome, FAO.

Cites. (2019). Geneva, Switzerland: Convention on International Trade in Endangered Species Available: https://cites.org/eng/cop/18/prop/index.php [Accessed 16/01/2019 2019].

Clark-Shen, N., Tingting, K. X., Rao, M., Cosentino-Roush, S., Sandraseageren, R., Gajanur, A. R., Chapman, D. D., Ying, E. L. X., Flowers, K. I. & Feldheim, K. A. (2021). The sharks and rays at Singapore's fishery ports. *Fisheries Research,* 235**,** 105805.

Clarke, S. C., Mcallister, M. K., Milner-Gulland, E. J., Kirkwood, G. P., Michielsens, C. G., Agnew, D. J., Pikitch, E. K., Nakano, H. & Shivji, M. S. (2006). Global estimates of shark catches using trade records from commercial markets. *Ecology Letters,* 9**,** 1115-1126.

Clarke, S. C., Milner-Gulland, E. J. & Bjørndal, T. (2007). Social, economic, and regulatory drivers of the shark fin trade. *Marine Resource Economics,* 22**,** 305-327.

Compagno, L. J. & Last, P. R. (1999). Order Rhinobatiformes. *In:* CARPENTER, K. E. & VOLKER, H. N. (eds.) *The Living Marine Resources of the Western Central Pacific. Volume 3: Batoid fishes, chimaeras and bony fishes part 1 (Elopidae to Linophrynidae).* Rome, Italy: Food and Agriculture Organisation of the United Nations.

Compagno, L. J. V. & Cook, S. F. (1995). The exploitation and conservation of freshwater
elasmobranchs: status of taxa and prospects for the future. *Journal of Aquariculture and*
Aquatic Sciences, 7**,** 62-90.
Compagno, L. J. V. & Last, P. R. (2008). A new species of wedgefish, *Rhynchobatus palpebratus* sp.
nov. (Rhynchobatoide: Rhynchobatidae), from the Indo–West Pacific. *In:* LAST, P. R.,
WHITE, W. T. & POGONOSKI, J. J. (eds.) *Description of new Australian chondrichthyans.*
Hobart, Australia: CSIRO.
Cortés, E. (2000). Life History Patterns and Correlations in Sharks. *Reviews in Fisheries Science,* 8**,**
299-344.
Crona, B. I., Van Holt, T., Petersson, M., Daw, T. M. & Buchary, E. (2015). Using social–ecological
syndromes to understand impacts of international seafood trade on small-scale fisheries.
Global Environmental Change, 35**,** 162–175.
D'alberto, B. M., Carlson, J. K., Pardo, S. A. & Simpfendorfer, C. A. (2019). Population productivity
of wedgefishes, guitarfishes, and banjo rays: inferring the potential for recovery *PLoS One,*
14**,** e0225183.
Dent, F. & Clarke, S. C. (2015). State of the global market for shark products. *FAO Fisheries and*
Aquaculture Technical Paper. Rome, Italy: FAO.
Dharmadi & Kasim, K. (2010). Keragaan perikanan cucut dan pari di laut Jawa. *Jurnal Penelitian*
Perikanan Indonesia, 16**,** 205-216.
Diop, M. S. & Dossa, J. (2011). 30 Years of Shark Fishing in West Africa. *Regional Marine and*
Coastal Conservation Programme for West Africa, and the Sub-Regional Fishing
Commission. Corlet/Condé-sur-Noireau, France: Fondation Internationale Du Bassin
d'Arguin.
Dulvy, N. K., Davidson, L. N. K., Kyne, P. M., Simpfendorfer, C. A., Harrison, L. R., Carlson, J. K.
& Fordham, S. V. (2016). Ghosts of the coast: global extinction risk and conservation of
sawfishes. *Aquatic Conservation: Marine and Freshwater Ecosystems,* 26**,** 134-153.
Dulvy, N. K., Fowler, S. L., Musick, J. A., Cavanagh, R. D., Kyne, P. M., Harrison, L. R., Carlson, J.
K., Davidson, L. N., Fordham, S. V., Francis, M. P., Pollock, C. M., Simpfendorfer, C. A.,
Burgess, G. H., Carpenter, K. E., Compagno, L. J., Ebert, D. A., Gibson, C., Heupel, M. R.,
Livingstone, S. R., Sanciangco, J. C., Stevens, J. D., Valenti, S. & White, W. T. (2014a).
Extinction risk and conservation of the world's sharks and rays. *Elife,* 3**,** e00590.
Dulvy, N. K., Pardo, S. A., Simpfendorfer, C. A. & Carlson, J. K. (2014b). Diagnosing the dangerous
demography of manta rays using life history theory. *PeerJ,* 2**,** e400.
Flowers, K. I., Heithaus, M. L. R. & Papastamatiou, Y. P. (2020). Buried in the sand: Uncovering the
ecological roles and importance of rays. *Fish and Fisheries,* Early Online Version**,** 1-23.
Fowler, S. L., Reed, T. M. & Dipper, F. A. (2002). *Elasmobranch Biodiversity, Conservation and*
Management: Proceedings of the International Seminar and Workshop, Sabah, Malaysia,
July 1997, Gland, Switzerland and Cambridge, UK, IUCN

739
Garcia-Vazquez, E., Machado-Schiaffino, G., Campo, D. & Juanes, F. (2012). Species
misidentification in mixed hake fisheries may lead to overexploitation and population
bottlenecks. *Fisheries Research,* 114**,** 52-55.
Giles, J. L., Riginos, C., Naylor, G. J. P., Dharmadi & Ovenden, J. R. (2016). Genetic and phenotypic
diversity in the wedgefish *Rhynchobatus australia*e, a threatened ray of high value in the
shark fin trade. *Marine Ecology Progress Series,* 548**,** 165-180.
Harrison, L. R. & Dulvy, N. K. (2014). *Sawfish: a global strategy for conservation,* Vancouver,
Canada, IUCN Species Survival Commission's Shark Specialist Group.
Harry, A. V., Tobin, A. J., Simpfendorfer, C. A., Welch, D. J., Mapleston, A., White, J., Williams, A.
J. & Stapley, J. (2011). Evaluating catch and mitigating risk in a multispecies, tropical,
inshore shark fishery within the Great Barrier Reef World Heritage Area. *Marine and*
Freshwater Research, 62**,** 710-721.
Ibrahim, R. (2003). Earned Income for Financial Sustainability in Indonesia. *The Dian Desa.*
Phillippines: The Synergos Institude.

Jabado, R. W. (2018). The fate of the most threatened order of elasmobranchs: Shark-like batoids
(Rhinopristiformes) in the Arabian Sea and adjacent waters. *Fisheries Research,* 204**,** 448-
457.
Jaiteh, V. F., Hordyk, A. R., Braccini, M., Warren, C. & Loneragan, N. R. (2016). Shark finning in
eastern Indonesia: assessing the sustainability of a data-poor fishery. *ICES Journal of Marine*
Science: Journal du Conseil.
Jaiteh, V. F., Loneragan, N. R. & Warren, C. (2017). The end of shark finning? Impacts of declining
catches and fin demand on coastal community livelihoods. *Marine Policy,* 82**,** 224-233.
Karthikeyan, R., Babu, N. C., Mandal, A. & Sehgal, P. (2009). Soft leathers from Himantura Stingray
skins. *Journal of the Society of Leather Technologists and Chemists,* 93**,** 108.
Keong, C. H. (1996). Indonesia. *In:* KEONG, C. H. (ed.) *Shark fisheries and the trade in sharks and*
shark products of South East Asia. Selangor, Malaysia TRAFFIC.
Kyne, P. M. & Bennett, M. B. (2002). Diet of the eastern shovelnose ray, *Aptychotrema rostrata*
(Shaw & Nodder, 1794), from Moreton Bay, Queensland, Australia. *Marine and Freshwater*
Research, 53**,** 679-686.
Kyne, P. M., Jabado, R. W., Rigby, C. L., Dharmadi, Gore, M. A., Pollock, C. M., Herman, K. B.,
Cheok, J., Ebert, D. A., Simpfendorfer, C. A. & Dulvy, N. K. (2020). The thin edge of the
wedge: extremely high extinction risk in wedgefishes and giant guitarfishes. *Aquatic*
Conservation: Marine and Freshwater Ecosystems, 30**,** 1337-1361.
Kyne, P. M., Rigby, C. L., Dharmadi & Jabado, R. W. (2019). *Rhynchobatus australiae. The IUCN*
Red List of Threatened Species, e.T41853A68643043.
Lack, M. & Sant, G. (2009). Trends in global shark catch and recent developments in management.
TRAFFIC International. Cambridge, UK.
Last, P. R., Naylor, G. J., Séret, B., White, W. T., De Carvalho, M. & Stehmann, M. (2016). *Rays of*
the World, Melbourne, Australia, CSIRO Publishing.
Last, P. R. & Stevens, J. D. (1994). *Sharks and Rays of Australia,* Hobart, TAS, CSIRO
Last, P. R. & Stevens, J. D. (2009). *Sharks and rays of Australia,* Melbourne, Australia, CSIRO.
Lawson, J. M., Pollom, R. A., Gordon, C. A., Barker, J., Meyers, E. K. M., Zidowitz, H., Ellis, J. R.,
Bartolí, Á., Morey, G. & Fowler, S. L. (2020). Extinction risk and conservation of critically
endangered angel sharks in the Eastern Atlantic and Mediterranean Sea. *ICES Journal of*
Marine Science, 77**,** 12-29.
Mackeracher, T., Bergseth, B., Maung, K. M. C., Khine, Z. L., Phyu, E. T., Simpfendorfer, C. A. &
Diedrich, A. (2020). Understanding non-compliance in small-scale fisheries: Shark fishing in
Myanmar's Myeik Archipelago. *Ambio***,** 1-14.
Mcauley, R., Simpfendorfer, C. A. & Hall, N. G. (2007a). A method for evaluating the impacts of
fishing mortality and stochastic influences on the demography of two long-lived shark stocks.
ICES Journal of Marine Science, 64**,** 1710-1722.
Mcauley, R., Simpfendorfer, C. A. & Wright, I. W. (2007b). Gillnet mesh selectivity of the sandbar
shark (*Carcharhinus plumbeus*): implications for fisheries management. *ICES Journal of*
Marine Science, 64**,** 1702-1709.
Metcalfe, J. D., Pritchard, V. L., Silvestri, S. M., Jenkins, J. B., Wood, J. S., Cowley, D. E., Evans, P.
E., Shiozawa, D. K. & Martin, A. P. (2007). Across the great divide: genetic forensics reveals
misidentification of endangered cutthroat trout populations. *Molecular Ecology,* 16**,** 4445-
4454.
Moore, A. B. M. (2015). A review of sawfishes (Pristidae) in the Arabian region: diversity,
distribution, and functional extinction of large and historically abundant marine vertebrates.
Aquatic Conservation: Marine and Freshwater Ecosystems, 25**,** 656-677.
Moore, A. B. M. (2017). Are guitarfishes the next sawfishes? Extinction risk and an urgent call for
conservation action. *Endangered Species Research,* 34**,** 75-88.
Musick, J. A. (1999). Ecology and Conservation of Long-Lived Marine Animals. *American Fisheries*
Society Symposium, 23**,** 1-10.
Ostrom, E., Burger, J., Field, C. B., Norgaard, R. B. & Policansky, D. (1999). Revisiting the
commons: local lessons, global challenges. *science,* 284**,** 278-282.
Pauly, D. (1988). Some definitions of overfishing relevant to coastal zone management in Southeast
Asia.

Prince, J. D. (2005). Gauntlet Fisheries for Elasmobranchs - the Secret of Sustainable Shark Fisheries.
*Journal of Northwest Atlantic Fishery Science, 35***,** 407-416.
Ramaschiello, V. & Vannuccini, S. (2015). FAO Contribution to the Harmonized System 2017. *In:*
NATIONS, F. A. A. O. O. T. U. (ed.) *Meeting of the Expert Group on International*
Statistical Classifications. New York: Food and Agriculture Organization of the United
Nations, United Nations Department of Economic and Social Affairs Statistic Division.
Rusandi, A., Agung, F., Hadi, S., Ariansyah, W. F., Yulianto, I., Sualia, I., Muttaqin, E., Silalahi, M.,
Dewanto, H. Y. & Rahmaniar, R. (2019). Efforts and challenges in managin sharks and rays
in the CITES species in Indonesia. *In:* FISHERIES, M. O. M. A. A. (ed.). Jakarta, Indonesia:
Ministry of Marine Affairs and Fisheries.
Sahubawa, L., Pertiwiningrum, A. & Rahmadian, Y. (2018). Enhancing the economic value and
consumer preferences of commercial mondol stingray (*Himantura gerardi*) leather creative
products. *IOP Conference Series: Earth and Environmental Science, 139.*
Simpfendorfer, C. A. (Year) Published. Demographic analysis of the dusky shark fishery in
southwestern Australia. American Fisheries Society Symposium, 1999. 149-160.
Simpfendorfer, C. A., Heupel, M. R., White, W. T. & Dulvy, N. K. (2011). The importance of
research and public opinion to conservation management of sharks and rays: a synthesis
*Marine and Freshwater Research, 62***,** 518-527.
Smith, W. D., Cailliet, G. M. & Cortés, E. (2008). Demography and elasticity of the diamond
stingray, *Dasyatis dipterura*: parameter uncertainty and resilience to fishing pressure. *Marine*
*and Freshwater Research, 59***,** 575-586.
Stevens, J. D., Bonfil, R., Dulvy, N. K. & Walker, P. A. (2000). The effects of fishing on sharks, rays,
and chimaeras (chondrichthyans), and the implications for marine ecosystems. *ICES Journal*
*of Marine Science, 57***,** 476-494.
Suzuki, T. (1997). Development of shark fisheries and shark fin export in Indonesia: case study of
Karangsong Village, Indramayu, West Java. *Elasmobranch biodiversity, conservation and*
management. Sabah, Malaysia: Occasional Paper of the IUCN Species Survival Commission.
Thorson, J. T. & Simpfendorfer, C. A. (2009). Gear selectivity and sample size effects on growth
curve selection in shark age and growth studies. *Fisheries Research, 98***,** 75-84.
Tillett, B. J., Field, C. I., Bradshaw, C. J. A., Johnson, G., Buckworth, R. C., Meekan, M. G. &
Ovenden, J. R. (2012). Accuracy of species indentification by fisheries observers in a north
Australian shark fishery. *Fisheries Research, 127-128***,** 109-115.
Vincent, A. C. J., Sadovy De Mitcheson, Y. J., Fowler, S. L. & Lieberman, S. (2014). The role of
CITES in the conservation of marine fishes subject to international trade. *Fish and Fisheries,*
15**,** 563-592.
Walker, T. I. (1998). Can shark resources be harvested sustainably? A question revisited with a
review of shark fisheries. *Marine and Freshwater research, 49***,** 553-572.
Walters, C. J. & Martell, S. J. D. (2004). *Fisheries ecology and management*, Princeton University
Press.
White, J., Simpfendorfer, C. A., Tobin, A. J. & Heupel, M. R. (2013). Spatial ecology of shark-like
batoids in a large coastal embayment. *Environmental Biology of Fishes, 97***,** 773-786.
White, W. T. & Dharmadi (2007). Species and size compositions and reproductive biology of rays
(Chondrichthyes, Batoidea) caught in target and non-target fisheries in eastern Indonesia.
*Journal of Fish Biology, 70***,** 1809-1837.
White, W. T., Giles, J., Dharmadi & Potter, I. C. (2006). Data on the bycatch fishery and reproductive
biology of mobulid rays (Myliobatiformes) in Indonesia. *Fisheries Research, 82***,** 65-73.
Yulianto, I., Booth, H., Ningtias, P., Kartawijaya, T., Santos, J., Sarmintohadi, Kleinertz, S.,
Campbell, S. J., Palm, H. W. & Hammer, C. (2018). Practical measures for sustainable shark
fisheries: Lessons learned from an Indonesian targeted shark fishery. *PLOS ONE, 13***,**
e0206437.

Figure Legends

Figure 1. The location of the Muara Angke landing site and processing village in Jakarta (star), and Benoa Harbour on Bali (triangle), where the frozen shipment of landed catch was originally landed before relocated to processing village near Muara Angke, Indonesia. The map was created using ArcGIS software by Esri (Version 10.4.1; www.esri.com). Map Sources: Esri, USGS, NOAA, Garmin, NPS.

Figure 2. Tangle net fishery catches at Muara Angke landing site, Jakarta Indonesia: (a) Large bottlenose wedgefish *Rhynchobatus australiae* unloaded from tangle net vessels at the port; (b) Large stingrays being processed at the adjacent village processing area; (c) Drying ray skins which will be used to make stingray leather products such as wallets and belts; (d) Wedgefish landings from Arafura Sea at the village processing area – *Rhynchobatus australiae* in centre of image highlighting the line of three white spots (yellow circle) diagnostic in this species. Photo credits: W.T.W.

Figure 3. Size-frequency histograms of the most abundant ray species (represented by 10 or more measured individuals) in the tangle net landed catches from the Muara Angke landing site surveys 2001 – 2005: (a) *Rhynchobatus australiae*; (b) *Himantura leoparda*; (c) *Himantura uarnak*; (d) *Maculabatis gerrardi*; (e) *Pastinachus ater*; (f) *Pateobatis fai*. White bars denote females, grey bars males and black bars unsexed individuals; the total number (n) of individuals, size at birth (red solid line) and size at maturity (black dashed line; M, male; F, female) when known; the size scale bar (x-axis) extends to the maximum known size for each of the species. The species are placed in phylogenetic order from wedgefish through to eagle rays. Size at birth and size at maturity estimates are from Last et al. (2016).

Figure 4. Size-frequency histograms of the most abundant ray species (represented by 10 or more measured individuals) in the tangle net catches from the Muara Angke landing site surveys 2001 – 2005: (a) *Pateobatis jenkinsii*; (b) *Pateobatis uarnacoides*; (c) *Taeniurops meyeni*; (d) *Aetobatus ocellatus*. White bars denote females, grey bars males and black bars unsexed individuals; the total number (n) of individuals, size at birth (red solid line) and size at maturity (black dashed line; M, male; F, female) when known; the size scale bar (x-axis) extends to the maximum known size for each of the species. The species are placed in phylogenetic order from wedgefish through to eagle rays. Size at birth and size at maturity estimates are from Last et al. (2016).

Figure 5. Overall species composition and percentage (%) of catch of the four Indonesian tangle net vessels (total n = 481), of which the landed catch was able to be fully documented from Muara Angke landing site, Jakarta Indonesia. The total landed catch from the four vessels consisted of *Maculabatis*

gerrardi n = 155, *Rhynchobatus australiae* n = 112, *Rhina ancylostoma* n = 32, *Aetobatus ocellatus* n

= 25, *Pateobatis jenkinsii* n = 18, *Himantura uarnak* n = 17, and for the other species n = 90. *MA-*

SKR-170704 landed on 7[th] July 2004; PV-PK-061004 landed on 16[th] October 2004; PV-CC-051005

(combined landed catch of PV-KA-051005 and PV-UK-051005) landed on 5[th] October 2005. Photo

credit: W.T.W.

Figure 6. Comparison of the size ranges for the landed catches of (a) *Rhynchobatus australiae* using

total length (TL cm) and (b) using disk width (DW cm) for *Maculabatis gerrardi, Pateobatis ater,*

Pateobatis fai, Pateobatis jenkinsii, and *Pateobatis uarnacoides* caught in the small mesh gillnet

(blue), Java Sea trawl fishery (dark green), Arafura Sea trawl fishery (light green), trammel net

(purple), hand- and long-line (yellow) and tangle net (red) landed in Muara Angke landing site,

Jakarta Indonesia. The size at birth (red dotted line) and size at maturity (black dashed line; M, male;

F, female) are displayed from Last et al. (2016).

Tables

Table 1. Summary table of the recorded tangle net vessels landed in Muara Angke landing site, Jakarta Indonesia in April 2001 – December 2005, including a vessel identifier (Vessel ID), date of survey, type of fishing gear, type of sampling (partial landed catch, PC; total landed catch, TC), number (No.) of ray and shark species and specimens surveyed, the reported fishing grounds, and survey location (Muara Angke landing site, MA; the nearby processing village, PV).

Vessel ID	Date of Survey	Fishing Gear	Type of Sampling	No. of rays species	No. of specimens	No. of shark species	No. of specimens	Reported fishing grounds	Survey Location
MA-UK-040401	04/04/2001	Tangle net	PC	14	Unknown	3	Unknown	Unknown	MA
MA-UK-050401	05/04/2001	Tangle net	PC	15	11*	2	Unknown	Unknown	MA
MA-UK-150502	15/05/2002	Tangle net	PC	10	159	0	0	Unknown	MA
PV-BB-140802	14/08/2002	Tangle net	PC	15	197	2	3	Benoa, Bali	PV
MA-UK-181002	18/10/2002	Tangle net	PC	9	69	0	0	Unknown	MA
PV-UK-090203	09/02/2003	Tangle net	PC	10	361	0	0	Unknown	PV
MA-SKR-170704	17/07/2004	Tangle net	TC	13	147	2	4	West Sumatra, Kalimantan, Riau Islands	MA
PV-UK-200704	20/07/2004	Tangle net	PC	1	49	0	0	Unknown	PV
MA-UK-210704	21/07/2004	Tangle net	PC	13	132	2	3	Unknown	MA
PV-PK-061004	06/10/2004	Tangle net	TC	12	106	4	5	Pontianak in Kalimantan	PV
MA-UK-160705	16/07/2005	Tangle net	PC	8	51	0	0	Unknown	MA
PV-KA-051005 (PV-CC-051005)^	05/10/2005	Tangle net	TC	14	240	5	8	Kalimantan	PV
PV-UK-051005 (PV-CC-051005)^	05/10/2005	Tangle net	PC					Unknown	PV
PV-UK-081205	08/12/2005	Tangle net	PC	1	4	1	2	Unknown	PV

* Only 11 specimens were recorded from 5 species of rays, no information on the remaining 6 species of rays and sharks were documented

^Two tangle nets landed on same day, the landed catch was unable to be separate during the survey, and the landed catch was combined and assigned the ID of PV-C-2005. The number of species and specimens reflects the landed catch from both vessels.

Table 2. Species composition, number of individuals of a species observed (no.) and overall percentage (% by no.) of the catch for the elasmobranchs caught by the tangle net fishery, and landed in Muara Angke landing site, Jakarta Indonesia in April 2001–December 2005. The number of recorded specimens (No.), percentage of total specimens (% by No.), observed minimum (Min. size), observed maximum size (Max. size), observed mean (± S.E.) size (DW/TL cm), minimum (Min. weight) and maximum (Max. weight) estimated total weight (kg) are reported for each species, with the International Union for Conservation of Nature's (IUCN) Red List of Threatened Species status (as of August 2020). IUCN categories are CR, Critically Endangered; EN, Endangered; VU, Vulnerable; NT, Near Threatened; LC, Least Concern; DD, Data Deficient. Dashed lines indicate species presence was recorded in landings, but data was not able to be documented.

Family	Scientific name	Common name	IUCN Listing	Year Assessed	No.	% by No.	Min. size	Max. size	Mean Size	± S.E.	Min. weight	Max. weight	Mean Weight	± S.E.
Pristidae	*Pristis pristis*	Largetooth sawfish	CR	2013	2	0.128	--	420.0	--	--	--	220.3	--	--
Glaucostegidae	*Glaucostegus thouin*	Clubnose guitarfish	CR	2019	--	--	--	--	--	--	--	--	--	--
	Glaucostegus typus	Giant guitarfish	CR	2019	14	0.898	170.0	260.0	206.0	13.15	14.86	51.36	27.57	5.543
Rhinidae	*Rhina ancylostoma*	Bowmouth guitarfish	CR	2019	57	3.656	130.1	270.0	190.2	22.97	18.68	168.4	77.38	24.29
	Rhynchobatus australiae	Bottlenose wedgefish	CR	2019	238	15.27	190.0	300.0	282.1	4.662	30.21	118.7	100.9	4.181
	Rhynchobatus palpebratus	Eyebrow wedgefish	NT	2019	30	1.924	--	--	--	--	--	--	--	--
Dasyatidae	*Bathytoshia lata*	Brown stingray	LC	2007	1	0.064	--	202.0	--	--	--	300.0		--
	Himantura leoparda	Leopard whipray	VU	2015	31	1.988	83.20	120.0	98.72	3.016	14.37	39.45	23.61	2.061
	Himantura uarnak	Coach whipray	VU	2015	57	3.656	42.60	147.6	99.50	6.204	4.015	69.83	28.57	4.189
	Himantura undulata	Honeycomb whipray	VU	2011	1	0.064	--	112.8	112.8	--	33.27	--	33.27	--
	Maculabatis astra	Blackspotted whipray	LC	2015	4	0.257	--	79.00	79.00	--	13.26	--	13.26	--
	Maculabatis gerrardi	Whitespotted whipray	VU	2004	194	12.44	62.70	89.50	75.85	1.036	6.560	19.40	12.01	2.410
	Megatrygon microps	Smalleye stingray	DD	2015	1	0.064	--	174.8	--	--	--	111.3	--	--
	Pastinachus ater	Broad cowtail ray	LC	2015	199	12.76	86.00	149.0	114.3	2.618	15.74	71.67	37.98	2.332
	Pateobatis fai	Pink whipray	VU	2015	264	16.93	70.50	168.4	110.9	4.595	9.101	100.4	36.45	4.143
	Pateobatis jenkinsii	Jenkin's whipray	VU	2015	187	11.99	59.20	138.4	82.59	1.791	5.621	58.47	14.78	1.207
	Pateobatis uarnacoides	Whitenose whipray	VU	2004	125	8.018	51.70	118.8	91.23	4.206	3.869	38.38	20.42	2.390
	Taeniurops meyeni	Blotched stingray	VU	2015	51	3.271	62.80	164.0	116.8	5.332	6.615	93.37	40.25	4.713
	Urogymnus asperrimus	Porcupine ray	VU	2015	5	0.321	76.50	103.4	89.95	9.511	11.40	26.17	18.78	5.222
	Urogymnus granulatus	Mangrove whipray	VU	2015	10	0.641	97.20	141.0	118.8	6.300	22.07	61.55	39.72	5.728

Gymnuridae	*Gymnura zonura*	Zonetail butterfly ray	VU	2006	7	0.449	70.50	91.60	79.78	3.036	2.442	5.466	3.657	0.4433
Aetobatidae	*Aetobatus ocellatus*	Spotted eagle ray	VU	2015	45	2.886	108.9	214.4	138.6	3.994	19.37	135.4	41.43	4.151
Myliobatidae	*Aetomylaeus vespertilio*	Ornate eagle ray	EN	2015	11	0.706	146.2	240.0	187.5	17.157	45.11	187.1	100.6	26.66
Carcharhinidae	*Carcharhinus amboinensis*	Pigeye shark	DD	2005	--	--	--	--	--	--	--	--	--	--
	Carcharhinus obscurus	Dusky shark	VU	2007	2	0.128	--	--	--	--	--	--	--	--
	Carcharhinus leucas	Bull shark	NT	2005	2	0.128	--	--	--	--	--	--	--	--
	Carcharhinus limbatus	Common blacktip shark	NT	2005	6	0.385	--	--	--	--	--	--	--	--
	Galeocerdo cuvier	Tiger shark	NT	2005	4	0.257	--	--	--	--	--	--	--	--
Ginglymostomatidae	*Nebrius ferrugineus*	Tawny nurse shark	VU	2003	3	0.192	--	--	--	--	--	--	--	--
Hemigaleidae	*Hemipristis elongata*	Fossil shark	VU	2015	3	0.192	109.6	122.9	116.3	4.702	5.719	25.14	6.989	0.8984
Sphyrnidae	*Sphyrna lewini*	Scalloped hammerhead	CR	2019	3	0.192	--	175.4	175.4	--	--	--	25.14	--
	Sphyrna mokarran	Great hammerhead	CR	2019	2	0.128	--	--	--	--	--	--	--	--