Taxonomic and functional dynamics of lung microbiome in cystic fibrosis patients chronically infected with *Pseudomonas aeruginosa*

Giovanni Bacci[1], Giovanni Taccetti[2], Daniela Dolce[2], Federica Armanini[3], Nicola Segata[3], Francesca Di Cesare[1], Vincenzina Lucidi[4], Ersilia Fiscarelli[4], Patrizia Morelli[5], Rosaria Casciaro[5], Anna Negroni[6], Alessio Mengoni[1], and Annamaria Bevivino[6*]

[1]Department of Biology, University of Florence, Sesto Fiorentino, 50019, Florence, Italy,
[2]Cystic Fibrosis Center, Anna Meyer Children's University Hospital, Department of Pediatrics Medicine, Florence,
50139, Italy
[3]Centre for Integrative Biology, University of Trento, Trento, 38122, Italy
[4]Children's Hospital and Research Institute Bambino Gesù, Rome, 00165, Italy
[5]Cystic Fibrosis Center, IRCCS G. Gaslini Institute, Department of Pediatrics, Genoa, 16147, Italy
[6]Department for Sustainability, Italian National Agency for New Technologies, Energy and Sustainable Economic Development, ENEA, 00123 Rome, Italy

*Corresponding author
Annamaria Bevivino, Department for Sustainability, ENEA, Rome, Italy. Email: annamaria.bevivino@enea.it

Abstract

Background

Although the cystic fibrosis (CF) lung microbiome has been taxonomically defined, little is still known about the overall gene content and functional profiles of the resident microbiome and how it changes in relation with patient's disease status. The aim of this study was to investigate the taxonomic and functional temporal dynamics of airways microbiome in CF patients with chronic *Pseudomonas aeruginosa* infection. A shotgun metagenomic approach was used, to target the entire genomic repertoire of the microbial community and allow strain-level profiling.

Results

A cohort of 22 patients with moderate-severe lung disease carrying the ΔF_{508} mutation was enrolled and followed over 15 months. The taxonomic composition, metabolic repertoire and antibiotic resistance gene content of lung microbiome was investigated by time-resolved shotgun metagenomics. A high inter-patient taxonomic heterogeneity was found with short-term compositional changes during period of exacerbation and following antibiotic treatment. Each patient represented a unique environment showing distinct communities at the taxonomic level and high patient-specific colonization by both main and emerging CF pathogens. The extraordinary resilience of the CF microbiome was found even from a functional perspective. All samples showed a core set of antibiotic resistance genes with a limited influence of antibiotic intake.

Conclusions

 The main outcomes from this study indicate a patient-specific temporal dynamic of the microbiome and a high functional resilience following exacerbation and antibiotic treatment.

Background

Cystic Fibrosis (CF) is the most common lethal autosomal recessive disease in Caucasians, caused by mutations in the gene coding for cystic fibrosis transmembrane conductance regulator (CFTR) channel [1]. Disruption of chloride anion transport, one of the key underlying features of CF, leads to altered physiological conditions at epithelial surfaces. In the airways, CFTR mutations result in a dehydrated viscous mucus that compromises mucociliary clearance and predisposes CF patients to repeated cycles of airway infection, mucous impaction, and bronchiectasis resulting in the majority of morbidity and mortality in the patient population [2]. In particular, bacterial lung infections reduce life expectancy in most CF patients [3]. The affected individuals consistently maintain high bacterial loads in their airways also during periods of clinical stability that are punctuated by episodes of pulmonary exacerbation [4]. Such periodic episodes of acute pulmonary exacerbation strongly contribute to the irreversible decline of lung function. Though much is known about the composition of the microbial infections in CF (for a recent review see [5]), the factors leading to such exacerbations are still poorly understood. In the past years, studies employing DNA-based analyses of the airway microbiota of CF patients have shown somewhat discordant results. Indeed, some authors report a largely stable airway microbiota through periods of exacerbation and antibiotic treatment [6], while other indicate of a high inter-patients variability [5,7–9], but also suggested the possibility to identify some microbial taxa as biomarker of exacerbation [10], as well as a role of rare species in exacerbation [11]. Most of these works are targeted metagenomic surveys performed on a variable number of patients and focusing on the 16S rRNA gene sequence. However, this approach offers limited possibilities to infer strain-level and functional (meaning based on functional genes) insights [12]. These two last points are particularly relevant when host-microbiome interactions are studied. Indeed, the overall genetic repertoire of the microbiome (i.e.

the entire set of genes in all the genomes of the community members) is the main responsible of the

interaction with the host [13]. Recently, the functional interactions among members of a bacterial

community have stirred the attention of investigators for relating microbiome functionality to

human-microbe interaction [14] and as a perspective for understanding the airway microbiome

dynamics in CF [15]. In several human diseases where the microbial infection is an important factor,

such as CF, single patients harbor genomically different strains, which ultimately may lead to

explain individual differences in clinical outcomes [16–18]. Until now, few longitudinal studies,

with a limited number of patients, on CF airway microbiota have been performed [19,20].

Moreover, studies on CF microbiome are few and on a limited number of patients [9,21–23] or

specific metabolic functions [24]. Moving away from taxonomic inventories towards a better

understanding of the CF microbiome genes opens a new avenue for the identification of the

microbial gene repertoire associated with CF lung disease. An ecological perspective on

multispecies and multi-strain colonization of CF airways will permit to understand the role of

polymicrobial dynamics in lung disease progression [25] and provide the clinicians with new

biomarkers of CF progression and targets for antibiotic therapy.

In this work. we tried to fill the gap of knowledge about the temporal dynamics of the airway

microbiome in CF, paying special attention to the episodes of exacerbation, by using a shotgun

metagenomic approach [26], that is targeting the entire genomic repertoire of the microbial

community, down to the strain level [27,28]. A cohort of 22 patients with moderate-severe lung

disease, grouped according to different genotypes (homozygote and heterozygote for ΔF_{508}

mutation), and chronically infected with *Pseudomonas aeruginosa*, was selected and followed over

15 months during which 8 patients underwent exacerbation events. This offered the opportunity to

investigate the taxonomic and functional dynamics of the overall microbiome.

Results

Population and sampling

Twenty-two patients with CF were enrolled for a total of 15 females and seven males. The patients were chosen from a larger cohort of patients with moderate-severe lung disease ($30 < \%FEV_1 < 70$) and chronically infected by *Pseudomonas aeruginosa*. During the study period, they were treated with maintenance antibiotics (aerosol) and only a subset (n = 8) received clinical intervention in form of supplementary antibiotics (oral or/and intravenous) for a pulmonary exacerbation (CFPE) (Table 1 and supplementary materials Table S1). The bacterial microbiome was investigated on sputum samples obtained every 3-4 months from 22 individuals along a survey of 15 months. Within the 22 subjects monitored, 8 underwent episodes of exacerbations, which provided the opportunity to explore the microbiomes composition along the events. In total, 79 samples from these 22 subjects were collected and analyzed by a whole metagenomic sequencing approach.

Airway microbiomes are taxonomically distinct and show patient-specific strain colonization

The overall taxonomic representation of the microbiomes from the 79 samples is reported in Fig. 1a and 1b, whereas a summary of obtained reads per sample was reported in supplementary materials Table S2. Firmicutes, Proteobacteria, Bacteroidetes, and Actinobacteria were the most represented phyla. A massive presence of the "classical" CF bacterial signatures (taxa), such as *Staphylococcus aureus* and *Pseudomonas aeruginosa*, and non-traditional CF taxa, such as *Rothia mucilaginosa*, and *Prevotella melaninogenica* (all present in the top-10 species within each phylum, Fig. 1b), was found. These species, indeed, represent the 49% of all detected taxa as reported in supplementary materials Table S3.

Figure 1: Taxonomic distribution in patients enrolled in the study. a) The taxonomic distribution of all species detected using MetaPhlAn2 was reported in each row of the matrix whereas columns represent samples collected during the study. Colors from dark blue to red were used to report "copies per million" (CPM) values as obtained from HUMAnN2 with black reporting a CPM value of zero. The plot was divided according to patient status: BL, baseline; TR, treatment; RC, recovery. Species were ordered according to their mean abundance and grouped according to their Phylum. b) The mean abundance value of the top-ten species (if available) detected within each Phylum was reported together with the standard error.

Although samples can be hardly clustered based on treatment events and/or genotype (Fig. 2a), the PERMANOVA analysis (Table 2) reported a significant effect of both factors. However, the R^2 values, namely the proportion of variance explained by the factor considered, were very low (0.03 for both factors). We cannot exclude that the heterogeneity of patients (especially concerning age) could have limited the statistical power of the analysis. The interaction effect between treatment events and genotype was not significant (p-value > 0.05), meaning that different genotypes did not influence the lung microbiome during treatment (viz. exacerbation) events and vice versa. The predominant effect observed was the subject effect, reporting a R^2 value of 0.52, indicating that a

126 high fraction (more than 50%) of the total variance can be explained by subject (patient)

127 individuality. Both FEV_1 and time did not show any significant effect (Table 2).

Figure 2: Ordination analyses based on a) taxonomic assignments and b) pathway distribution
detected with MetaPhlAn2 and HUMAnN2, respectively. Ordination analyses were conducted
using the Bray-Curtis dissimilarity index and ordered following the principle coordinate
decomposition method (PCoA). The percentage of variance explained by each coordinate was
reported between round brackets. Homozygote and heterozygote refer to ΔF_{508} mutation of CFTR
gene. BL, baseline; TR, treatment; RC, recovery.

135 With the aim to evaluate the relevance of subject (patient) individuality, a strain-level analysis was

136 conducted. Results obtained revealed that strain genotypes inferred from metagenomic data are

137 highly patient-specific for many bacterial species. Indeed, samples (i.e. strains) from the same

138 patients are very closely related and tightly clustered, confirming a high patient-specific colonization

139 by strains of the above-mentioned species (Fig. 3 and supplementary materials Fig. S1).

Figure 3: Strain-level phylogenetic trees of the main CF pathogens detected in the study.
Phylogenetic trees obtained through StrainPhlAn pipeline were reported for the main pathogenic
signatures of CF disease: a) *Pseudomonas aeruginosa*; b) *Staphylococcus aureus*; c) *Rothia
mucilaginosa*; d) *Prevotella melaninogenica*. Points at the end of each clade are colored according
to patients so that two points with the same color, in the same tree, represent the same species in two
different time points, for the same patient.

Alpha diversity analysis confirmed the overall picture of results mentioned above. Different values
of bacterial diversity were found according to clinical status, genotypes, and subjects (Fig. 4a,
supplementary materials Fig. S2 and Table S4). Samples collected during clinical treatments
reported a lower biodiversity than samples collected during normal visits, highlighting the role of
clinical treatments in perturbing CF lung communities as confirmed by the Tukey's post hoc test
(supplementary materials Table S5).

Figure 4: Differences across exacerbation events. The effect of an exacerbation event on alpha diversity was inspected using both the Shannon index and the inverse Simpson index. Diversity indexes were computed for both a) taxonomic signature and b) metabolic pathways. BL, baseline; TR, treatment; RC, recovery. Each box shows the "interquartile range" (IQR) that is the differences between the third and the first quartile of data (the 75th and the 25th percentile). Horizontal bars are medians whereas whiskers represent the minimum and maximum values defined as Q1 – (1.5 x IQR) and Q3 + (1.5 x IQR), respectively. Observations that fell outside minimum and maximum values were defined as outliers and reported using white points.

Airway microbiomes are functionally consistent and show subject-specific distribution patterns

Similar results as those reported above were obtained considering the pathway distribution. Indeed, the PERMANOVA analysis (Table 2) confirmed the effect of exacerbation events and genotypes in shaping the pathway distribution of CF lung microbiome (R^2 values of 0.04 and 0.03 respectively), though less marked than the subject-specific effect ($R^2 = 0.48$). The sample distribution according to the ordination analysis (PCoA) was very heterogeneous with no sharp differences according to

genotypes or exacerbation events. Even here, alpha diversity analyses reported a significant drop of

diversity in samples collected during exacerbation events, but the drop was significant only

considering the inverse Simpson index (Fig. 4b and supplementary materials Table S4). Overall, the

pathway distribution was more consistent with respect to the taxonomic one, with biosynthetic

pathways being the most represented functional category (Fig. 5, supplementary materials Fig. S2

and Table S5). Pathways were mainly detected in members of Firmicutes and Proteobacteria phyla,

though Bacteroidetes and Actinobacteria were quite well represented. Even if these results

confirmed the results from the analysis of the taxonomic distribution, metabolic pathways showed a

more consistent distribution across samples. Indeed, the beta-diversity analysis on both taxonomic

and functional distribution showed a lower similarity based on taxonomy in respect with pathways

(Supplementary materials Table S6, Fig. 6a and 6b). These results were additionally confirmed by

the differential abundance analysis. For contrasts made within each genotype, only 40 pathways

reported significant differences across exacerbation statuses (p-values < 0.05 and |log(fold-change)|

> 5) all in the homozygote group (Supplementary materials Fig. S3 and Table S7), whereas,

considering all samples together, no pathway was found to be more abundant in one condition in

respect with another (data not shown). These results confirmed the extraordinary resilience of the

CF microbiome even from a functional perspective.

Figure 5: Pathway distribution according to exacerbation events. The pathway distribution was reported for each sample (columns) and for each pathway detected (rows). Colors from dark blue to red were used to report "copies per million" (CPM) values as obtained from HUMAnN2 with black reporting a CPM value of zero. On the left, the percentage of taxa in which each pathway was detected was reported using different colors. The main colors correspond to the Phylum whereas the different shades correspond to the genus detected (if available). BL, baseline; TR, treatment; RC, recovery.

Figure 6: Beta diversity analysis on both taxonomic and functional distribution. a) Hierarchical clustering based on UPGMA method. Clustering was performed on both pathway distribution (the upped triangle) and taxonomic composition of samples (lower triangle). The Bray-Curtis distance was used to compute distances between samples, but it was transformed into similarity value by subtracting 1 before plotting. Thus, red colors represent high similarity values whereas blue colors represent low similarity values. The shape of the points on each tip of trees refers to the hospital whereas the colors refer to the exacerbation events. b) Results of Tukey's post hoc test on beta diversity values across patient genotypes and exacerbation events. Contrasts were computed even to test differences between taxonomic distribution and pathways with taxa reporting higher level of beta diversity. Homozygote and heterozygote refer to ΔF_{508} mutation of CFTR gene. BL, baseline; TR, treatment; RC, recovery. Boxplot were computed as described in Figure 4 legend.

Antibiotic resistance genes through exacerbation events and treatments

Similar to the pathway analysis reported above, antibiotic resistance genes (ARG) were inspected in relation to treatment events. Only six genes were found to be affected by an exacerbation condition,

all regarding samples form ΔF_{508} heterozygote patients whereas, as found for metabolic pathways,

no gene was significantly impacted by antibiotic treatment when considering all samples at once

(supplementary materials Fig. S4 and Table S8). A similar approach was used to inspect the effect

of antibiotic treatment on ARG distribution. ARG were inspected in relation to the antibiotic

treatments reported in supplementary materials Table S1. The class of each antibiotic was correlated

to the presence (and the abundance) of genes that may, in principle, confer resistance to antibiotics

from the corresponding class. Differential abundance analyses were performed for each classes of

antibiotics that was used in this study and results obtained were reported in supplementary materials

Fig. S5 and Table S9. Only 11 genes were found to be affected by antibiotic intake in different

ways. Indeed, 8 out of 11 reported a reduction of abundance during the treatment whereas the

remaining 3 reported an increased abundance in respect with antibiotic intake. Results obtained

confirmed the high resilience of the gene composition of CF lung microbiome. The presence of

ARGs coupled with antibiotic intake was also explored. Results showed that the antibiotic resistance

classes of each gene corresponded to the antibiotic treatment used in each sample reporting a big

block of ARGs that were present in most of the sample considered (Supplementary materials Fig. S6

and Fig. S7).

Discussions

Longitudinal studies allow to provide important clues on stability and dynamics of microbial

ecosystems [29]. As all biotic communities, microbial communities tend to evolve towards a stable

composition, either in natural environment and in association with host (as human-associated

microbiomes). Changes in the community can be triggered by external conditions, as changes in

host physiology (e.g. inflammation status) and/or other perturbations (e.g. antibiotic treatment).

Indeed, perturbation studies help to probe community dynamics and resilience and possibly discover

new findings for accessing ways for modifying the microbiome [30,31]. Although patients with CF

experiments repeated episodes of pulmonary exacerbations during their lives, a broadly accepted

definition of these events is still missing [4]. Here, we have investigated the temporal dynamics of

CF airway microbiome by using shotgun metagenomics posing attention on exacerbation events

which usually bring to an acute decrease in lung function and an increase in respiratory symptoms

(such as: increased cough, sputum production, and shortness of breath). Key questions were i) what

was the composition and stability of the lung microbiome in patients with CF when longitudinally

sampled at stable and exacerbation events; and ii) if the clinical status influenced the metabolic

repertoire and the AR gene composition of lung bacterial community. Our results describe a unique

examination of the dynamic of the lung microbiome in patients with moderate-severe lung disease

carrying the ΔF_{508} mutation of CFTR gene and containing clinical measurements over a 15-month

period.,

The lung microbiome of CF patients seems to be a highly patient-specific environment which can be

directly conditioned by the host and its habits. Indeed, there was less variation within the same

individual at different time points than between different individuals at the same time point, proving

some degree of temporal stability of an individual's lung microbiome. This last point agrees with the

lack of a time effect on the taxonomic distribution of microbiome. The predominant taxa that

colonized the lung of CF patients showed an extraordinary resilience, as witnessed by the presence

of the same strains during the whole period of infection. These results agree with previous

observations based on 16S rRNA gene profiling, though these studies failed to report a strain-

specific overview of the whole dynamic due to the limitations intrinsic to the approach [6,8,11].

Carmody and colleagues showed a relatively stable lung community that may be altered during

period of exacerbation even in the absence of viral infection or antibiotic only in a small group of

patients [10]. Even in other pulmonary diseases, such as non-cystic fibrosis bronchiectasis, lung

bacterial communities showed a conserved structure for long period of time, as showed in the work

by Cox and colleagues where patients were followed for a six-month period [8]. A similar result was

shown in the work from Fodor and colleagues [6] where, though occasional short-term

compositional changes in the airway microbiota were found, the main taxonomic signatures of CF

disease were highly stable. A notable exception was found for *Rothia mucilaginosa*. In fact, in

contrast with other studies where it was rarely identified [15,32,33], in our samples it was detected

in high relative abundance. This finding may suggest a potential involvement of *Rothia*

mucilaginosa in CF microbiome dynamics and pathogenicity, which deserves further attention.

The antibiotic treatment used did not seem to alter this micro-environment for long period of time

since most of the main taxa linked to CF infection are still present even after exacerbation events

that are usually handled by a massive amount of antibiotic. From a taxonomic perspective, samples

coming from the same patient clustered together highlighting the role of the host in bacterial strain

selection during the baseline but even during (and after) exacerbation events. Despite this patient-

specific colonization, the taxonomic composition was very different from one subject to another

event if sampled at the same time point.

On the other hand, pathways reported a more homogeneous distribution across patients. This high

conservation could be related to the characteristic of the lung environment itself, such as mucus

compositions, nutrient availability, and oxygen levels, which can be broadly similar across patients

with a similar clinical status. This, is in line with the finding that the function of a biotic community

is more conserved than the presence of single members [34]. In fact, though the lung microbiome in

our study was populated with a relatively large set of microorganisms, the main functions detected

are similar across all patients. From this point of view the airway microbiome can be considered as

performing a similar "ecosystem service", irrespective of the taxonomy present as pointed out by

various authors in other environments [34–36]. The finding that CFTR genotypes a different

representation in some pathways, may suggest that the airways microbiome is influenced by the type

of CFTR alteration. However, this hypothesis deserves further attention to clarify the specific role of

microbial pathways with respect to CFTR genotype. Pathogenic bacteria, such as *Pseudomonas*

aeruginosa, need to colonize human tissues to grow and in this sense, even pathway that could be

related to a worsening of clinical conditions or that could be targeted by antibiotic molecules will be

part of this core set of functions. Despite a clear effect of antibiotic treatment during (and after)

exacerbation periods, the community structure is always recovered with the main pathogenic taxa

emerging again. This effect is confirmed by the correlation of ARG distribution and antibiotic

intake. Patients subjected to a given antibiotic treatment did not seem to select bacteria resistant to

the antibiotic used but the detection of a particular mechanism seems to be distributed in almost all

patients regardless of the treatment. An evidence of functional stability of the lung microbiota was

previously reported in other works not concerning CF disease [37,38]. Both works focused their

attention on the gut microbiome of obese and healthy individuals (human and mouse) reporting a

considerable metabolic redundancy. This high degree of redundancy in the gut microbiome supports

a more ecological view where subjects can be considered as different ecological niches all inhabited

by unique collections of microbial phylotypes but all sharing the same set of genes. This concept can

be extended to the lung microbiome where it is possible to define a core set of features only at the

level of metabolic functions. This functional conservation may thus be needed by the whole

community and patients can be seen as multiple micro-environments inhabited by a peculiar set of

strains, which share the same functions. This work represents a step forward toward a patient-

specific interpretation of CF microbiology [39].

Conclusions

In summary, the temporal dynamics of the airway microbiome in a large cohort of patients with CF

revealed patient-specific signatures of the airway microbiome at strain-level profiling, the lack of

variation in the microbiome across pulmonary exacerbations, and a core set of antibiotic resistance

genes that did not vary by antibiotic intake. The main conclusion of the present study is that the

management of chronic CF infection may be improved by a more patient-specific personalization of

clinical care and treatment. Longitudinal studies of CF airway microbiota will permit to tailor

therapeutic interventions and select antibiotic therapies based on the composition and relative

abundance of antibiotic resistance genes within the respiratory microbiome.

Methods

Demographic and clinical characteristics of enrolled patients

Twenty-two adolescents and adults with CF were enrolled in the study between October 2014 and

March 2015 (Table 1). The study subjects were selected based on eligibility criteria that included all

of the following: (i) a diagnosis of CF, i.e., a sweat test showing sweat Cl > 60 mmol/l and two

known CFTR mutations causing the disease with pancreatic insufficiency (elastase< 5 μg/g/feces)

[40], (ii) aged more than six years, i.e., between 11 and 55 years, (iii) chronically infected with

Pseudomonas aeruginosa according to the Leeds criteria [41] and iv) decline in %FEV_1 in the

previous three years before enrollment by measuring the difference between the best %FEV_1

registered within the previous year and the best %FEV_1 registered two-years before specimen

collection, following the criteria previously reported [42]. Patients were excluded if they were

chronically infected with *Burkholderia cepacia* complex. Using these criteria, 22 patients with

moderate-severe lung disease and carrying the ΔF_{508} mutation were included in the study for a total

of 79 shotgun metagenomic samples. The cohort was enrolled in three Italian Hospital, namely:

Bambino Gesù Children's Hospital (Rome, Italy), G. Gaslini Children's Hospital (Genoa, Italy) and

Meyer Children's Hospital (Florence, Italy). Clinical status at the time of collection was

designated as *baseline* (BL), when clinically stable and at their clinical and physiological baseline,

on treatment (TR), at exacerbation-associated antibiotic treatments, and *at recovery* (RC), upon

completion of antibiotic treatment. Subjects were treated according to current standards of care with

periodical microbiological controls [43] with at least four microbiological controls per year [4]. At

each visit, clinical data collection and microbiological status (colonizing pathogens with available

cultivation protocols) were performed according to the European CF Society standards of care [44].

Forced expiratory volume in 1 second as a percentage of predicted (%FEV$_1$) is a key outcome of

monitoring lung function in CF [45]. FEV$_1$ values were measured according to the American

Thoracic Society and European Respiratory Society standards [43]. CFTR genotype, sex, age, and

antibiotic treatment for each patient were reported in (Table 1 and supplementary materials Table

S1). During serial sampling, data (antibiotic usage and spirometry) were collected.

Sample collection, processing, DNA extraction and sequencing

Sputum samples were obtained by spontaneous expectoration at baseline, exacerbation-associated

antibiotic treatments and recovery status. Sampled were processed according to standard methods as

previously described [13,46]. Bacterial respiratory pathogens were identified using the conventional

techniques reported in the Guidelines, as previously described [46,47]. The number of samples,

microbiological status at sampling and samplings following exacerbation events are reported in

Table 1. Sputum samples were washed in 5 ml PBS and then centrifuged (3,800 g) for 15 minutes.

Resulting pellets were resuspended in 5-10 ml DNAse buffer (10 mM Tris-HCl pH 7.5; 2.5 mM

MgCl2; 0.5 mM CaCl2, pH 6.5) with 7.5 ul of DNAse I (2000 Units/ml) per 1 ml of sample

(15U/ml final), incubated for 2 hours at 37C, and washed twice by pelleting at 3,800 g for 15

minutes and resuspending in 10 ml SE buffer (75 mM NaCl, 25 mM EDTA, pH 7.5). Pellets were

then resuspended in 0.5 ml lysis buffer (20 mM Tris-HCl pH 8.0; 2 mM EDTA pH 8.0; 1% (v/v)

Triton; 20 mg/ml Lysozyme final concentration), incubated for 30 minutes at 37C before extracting

DNA with the MoBio Powersoil DNA extraction kit as per manufacturer's instructions. Libraries

were prepared with Nextera XT kit (Illumina) Sequencing was performed on an Illumina HiSeq2500

apparatus (Illumina). Raw sequence data reported in this study have been deposited in the NCBI

"Sequence Read Archive" (SRA) under the project accession PRJNA516870.

Bioinformatic analyses

Sequence quality was ensured by trimming reads using StreamingTrim 1.0 [48], with a quality

cutoff of 20. Bowtie2 [49] was used to screen out human-derived sequences from metagenomic data

with the latest version of the human genome available in the NCBI database (GRCh38) as reference.

Sequences displaying a concordant alignment (mate pair that aligns with the expected relative mate

orientation and with the expected range of distances between mates) against the human genomes

were then removed from all subsequent analyses. Metabolic and regulatory patterns were estimated

using HUMAnN2 [50] and considering only those pathways with a coverage value $\geq$ 80%, whereas

the taxonomic microbial community composition was assessed using MetaPhlAn2 [51]. Reads were

assembled into contigs using the metaSPAdes microbial assembler [52] with automatic k-mer length

selection. To establish an airway microbiome gene catalog [12] we first removed contigs smaller

than 500bp and then used prodigal in Anonymous mode [53], as suggested by the author of the tool,

to predict open reading frames (ORFs). Translated protein sequences obtained from assembled

contigs were classified using eggNOG mapper against the bactNOG database [54]. Each protein was

classified according to its best hit with an e-value lower than 0.001 as suggested in [55]. The CARD

database [56] was used in combination to the Resistance Gene Identifier (RGI, version 4.0.3) to

inspect the distribution of antibiotic resistance gene (AR genes). Genes predicted within each

metagenome were quantified using the number of reads that mapped against metagenomic contigs

obtained for each sample. Reads were mapped back to contigs using Bowtie2 [49] and the number

of reads mapping each ORF was obtained with the bedtools command "multicov" (version 2.26.0).

To quantify gene content across different samples, genes were collapsed using the bestOG given by

eggNOG mapper by summing together the number of reads that mapped genes with the same

annotation. The same approach was used to quantify AR genes predicted with RGI but this time the

unique identifier provided by CARD was used to collapse counts.

Strain characterization was performed using StrainPhlAn [27]. Sequence variants for each organism

detected were assessed against the MetaPhlAn2 [51] marker genes and a tree has been generated

including all samples in which the organism was found at least in one time point. One reference

genome per organism was downloaded from the RefSeq database and added to the tree.

Taxonomic classification of metagenomic contigs

Assembled contigs were taxonomically classified using BLAST. First, all genomes available for

each species detected with MetaPhlAn2 were downloaded from NCBI and used to build a database

for each sample. All genomes reporting an identity higher than 90% and a coverage higher than 80%

were collected and used for taxonomic classification. Contigs reporting hits with genomes coming

from a single species were assigned to that species whereas contigs reporting hits from multiple

species were flagged as unknown.

Statistical analyses

Statistical analyses were performed in R [57] version 3.4.4. The taxonomical and functional

composition on lung microbiome was explored using permutational multivariate analysis of variance

(PERMANOVA with 1000 permutations), 'adonis2' function of vegan package version 2.5-2;

whereas differences in bacterial diversity were tested using analysis of covariance (ANCOVA),

'aov' function. The model fitted for both analyses was:

$$X \sim \text{Status} + \text{Genotype} + \text{Subject} + \text{FEV}_1 + \text{days}$$

where, Exacerbation is the exacerbation event, Genotype is the CFTR genotype, Subject is the

patient, FEV_1 was the forced expiatory volume in 1 second, and days, was the number of days from

the enrollment in the study. For the ANCOVA analyses Tukey's post hoc tests were performed to

test for mean differences within each factor used to build the full model (excluding FEV_1 value and

days since they were not categorical variable). Ordination analyses were conducted on both taxa and

pathways using the function 'ordinate' of the phyloseq package (version 1.23.1) with principle

coordinate decomposition method (PCoA) and the Bray-Curtis dissimilarity index. The same index

was used to inspect the distribution of samples and compare beta diversity level in bot taxonomic

composition and pathways.

To test for differentially distributed pathways and taxa across exacerbation events and genotypes we

used a moderated t-test as implemented in the limma package [58], version 3.34.9. Data obtained

with MetaPhlAn2 (taxonomic composition) and HUMAnN2 (pathway composition) were fitted into

limma's model using subjects as blocking variable. Since both software quantify biological units

using relative counts (HUMAnN2 uses "copies per million" and MetaPhlAn2 uses percentages) we

transformed this data into logarithmic values using the formula: $\log_2(x + 0.1)$, where x are the

relative counts. Obtained p-values were corrected using the Benjamini-Hochberg correction method.

A similar approach has been used for antibiotic genes detect along assembled contigs. Here the

number of reads that mapped onto each gene was used to estimate differentially abundant gene.

Since the number of reads for each sample was variable (the ratio of the largest library size to the

smallest was more than 10-fold) we used limma's voom method [59] to fit our model, as suggested

by the author of limma.

Declarations

Ethics Statement

The study was approved by the Ethics Committees of Children's Hospital and Research Institute Bambino Gesù (Rome, Italy), Cystic Fibrosis Center, Anna Meyer Children's University Hospital (Florence, Italy) and G. Gaslini Institute (University of Genoa, Genoa, Italy) [Prot. N. 681 CM of November 2, 2012; Prot. N. 85 of February 27, 2014; Prot. N. FCC 2012 Partner 4-IGG of September 18, 2012]. All participants provided written informed consent before the enrollment in the study. All sputum specimens were produced voluntarily. All procedures were performed in agreement with the "Guidelines of the European Convention on Human Rights and Biomedicine for Research in Children" and the Ethics Committee of the three CF Centers involved. All measures were obtained and processed ensuring patient data protection and confidentiality.

Availability of data and material

All data generated or analysed during this study are included in this article. Raw sequence data reported in this study have been deposited in the NCBI "Sequence Read Archive" (SRA) under the project accession PRJNA516870.

Conflict of Interest

We have no conflict of interest to declare.

Funding

This study was supported by Italian Grants funded by the Italian Cystic Fibrosis Research Foundation (FFC) (http://www.fibrosicisticaricerca.it/) to Annamaria Bevivino: Project n. FFC#14/2015 with the contribution of "Delegazione FFC di Latina", "Latteria Montello Nonno Nanni", and "Gruppo di Sostegno FFC Valle Scrivia Alessandria"), and Project n. FFC#19/2017 with the contribution of "Delegazione FFC Lago di Garda con i Gruppi di Sostegno FFC di

Chivasso, dell'Isola Bergamasca, di Arezzo". Giovanni Bacci was supported by a postdoctoral

fellowship by Italian Cystic Fibrosis Research Foundation. The funders had no role in study design,

data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' contributions

Conceived and designed the experiments: AB VL GT EVF AM. Performed the experiments: GB

FDC. Analyzed the data: GB AM AB. Contributed reagents/materials/analysis tools: DD FA PM RS

AN. Wrote the paper: GB AM AB. Provided comments and recommendations that improved the

manuscript: NS GT VL. Supervised research: AB VL GT EVF AM.

Acknowledgments

The authors greatly acknowledge the Italian Cystic Fibrosis Research Foundation (FCC) for its

support and administrative tasks, and Ricciotti Gabriella, Tuccio Vanessa and Campana Silvia for

their technical support.

Tables

Table 1. Characteristics of patients enrolled in the study. ID, study id; Hospital, hospital in which patient has been enrolled [OPBG=Children's Hospital and Research Institute Bambino Gesù (Rome, Italy); Gaslini=G. Gaslini Institute (University of Genoa, Genoa, Italy); Meyer=Cystic Fibrosis Center, Anna Meyer Children's University Hospital (Florence, Italy)]; Genotype, CFTR genotype; Gender, gender; Age, enrollment's age; n, number of samples collected; EX, yes if an exacerbation event has occurred during the study (no otherwise) [4]; FEV1, mean value of forced expiratory volume in 1 second plus/minus the standard error on the mean; heterozygote and homozygote refers to ΔF508 genotype; %FEV1 status: S = with a rate decline lower than 1.5%, SD = with a rate decline higher than 5% [42].

ID	Hospital	Genotype	Gender	FEV_1 status	Age	n	EX	$\%FEV_1$
B01	OPBG	ΔF_{508}/2183AA->G	M	S	27	5	yes	37.0 ± 1.70
B02	OPBG	ΔF_{508}/N1303K	F	SD	26	3	no	54.7 ± 3.48
B03	OPBG	ΔF_{508}/4016insT	F	S	30	4	no	55.0 ± 1.08
B06	OPBG	ΔF_{508}/ΔF_{508}	F	SD	21	4	no	60.2 ± 3.42
G10	Gaslini	ΔF_{508}/ΔF_{508}	M	S	51	4	no	54.0 ± 3.08
G24	Gaslini	ΔF_{508}/ΔF_{508}	F	S	49	3	yes	$NA \pm NA$
G28	Gaslini	ΔF_{508}/ΔF_{508}	F	NA	38	2	no	42.5 ± 1.50
G30	Gaslini	ΔF_{508}/ΔF_{508}	F	S	50	1	no	54
G31	Gaslini	G1244E/G42X	F	SD	53	2	no	41.5 ± 1.50
G34	Gaslini	ΔF_{508}/ΔF_{508}	F	S	39	1	no	47
M05	Meyer	ΔF_{508}/ΔF_{508}	M	SD	32	4	no	34.8 ± 0.85
M19	Meyer	ΔF_{508}/ΔF_{508}	M	S	24	4	no	44.0 ± 2.04
M21	Meyer	ΔF_{508}/N1303K	M	SD	27	4	yes	51.5 ± 4.35
M22	Meyer	ΔF_{508}/2789+5G->A	F	S	29	5	yes	50.4 ± 1.03
M23	Meyer	ΔF_{508}/G542X	F	S	30	4	yes	37.0 ± 1.47
M24	Meyer	ΔF_{508}/ΔF_{508}	M	S	32	4	no	35.2 ± 0.85
M25	Meyer	ΔF_{508}/296+1G->T	F	SD	41	4	no	42.5 ± 2.02
M26	Meyer	ΔF_{508}/3849+10	F	SD	49	5	yes	39.6 ± 1.94
M28	Meyer	ΔF_{508}/N1303K	M	S	23	4	no	39.0 ± 1.08
M29	Meyer	ΔF_{508}/G542X	F	S	12	4	no	43.5 ± 3.75
M31	Meyer	ΔF_{508}/ΔF_{508}	F	SD	11	3	yes	32.7 ± 4.41
M33	Meyer	ΔF_{508}/G85E	F	SD	13	5	yes	35.4 ± 5.78
Total: 22	Gaslini:6 Meyer:12 OPBG:4	Heterozygote :47 Homozygote :29 Other:2	F:15 M:7	S:12 SD:9	32.1 ± 2.73	79	no:14 yes:8	43.5 ± 1.09

Table 2. Permutational multivariate analysis of variance (PERMANOVA) on both taxonomic

distribution and metabolic pathways. The analysis based on taxonomic distribution was reported in

the upper part of the table whereas the analysis based on metabolic pathways was reported at the

bottom. Df, degrees of freedom; SumOfSqs, sum of squares; R^2, r-squared statistic (reported as

proportion); F, F-statistic; Pr(>F), p-value associated to the F-statistic. Significant effects, namely

those reporting a p-value lower than 0.05, were reported in bold.

	Df	SumOfSqs	R^2	F	Pr(>F)
Taxonomy					
Status	**2**	**0.68**	**0.03**	**1.91**	**0.0300**
Genotype	**1**	**0.77**	**0.03**	**4.30**	**0.0020**
Sample	**18**	**11.97**	**0.52**	**3.74**	**0.0010**
FEV$_1$ value	1	0.27	0.01	1.53	0.1349
Days	1	0.28	0.01	1.58	0.1229
Status:Genotype	1	0.11	0.01	0.64	0.7642
Residual	49	8.72	0.38	-	-
Pathway					
Status	**2**	**0.20**	**0.04**	**2.37**	**0.0220**
Genotype	**1**	**0.14**	**0.03**	**3.42**	**0.0080**
Sample	**18**	**2.43**	**0.48**	**3.20**	**0.0010**
FEV$_1$ value	1	0.09	0.02	2.14	0.0989
Days	1	0.05	0.01	1.26	0.2458
Status:Genotype	1	0.08	0.02	1.96	0.1169
Residual	49	2.07	0.41	-	-

References

1. Ratjen F, Döring G. Cystic fibrosis. Lancet. 2003;361:681–9.

2. Gibson RL, Burns JL, Ramsey BW. Pathophysiology and management of pulmonary infections in cystic fibrosis. Am J Respir Crit Care Med. 2003;168:918–51.

3. Knapp EA, Fink AK, Goss CH, Sewall A, Ostrenga J, Dowd C, et al. The cystic fibrosis Foundation patient registry. Design and methods of a national observational disease registry. Ann Am Thorac Soc. Am Thoracic Soc; 2016;13:1173–9.

4. Flume PA, Mogayzel PJ, Robinson KA, Goss CH, Rosenblatt RL, Kuhn RJ, et al. Cystic fibrosis pulmonary guidelines: Treatment of pulmonary exacerbations. Am J Respir Crit Care Med. 2009;180:802–8.

5. Huang YJ, LiPuma JJ. The microbiome in cystic fibrosis. Clin Chest Med. 2016;37:59–67.

6. Fodor AA, Klem ER, Gilpin DF, Elborn JS, Boucher RC, Tunney MM, et al. The adult cystic fibrosis airway microbiota is stable over time and infection type, and highly resilient to antibiotic treatment of exacerbations. PLoS One. 2012;7:e45001.

7. Price KE, Hampton TH, Gifford AH, Dolben EL, Hogan DA, Morrison HG, et al. Unique microbial communities persist in individual cystic fibrosis patients throughout a clinical exacerbation. Microbiome. 2013;1:27.

8. Cox MJ, Turek EM, Hennessy C, Mirza GK, James PL, Coleman M, et al. Longitudinal assessment of sputum microbiome by sequencing of the 16S rRNA gene in non-cystic fibrosis bronchiectasis patients. PLoS One. 2017;12:1–17.

9. Whelan FJ, Heirali AA, Rossi L, Rabin HR, Parkins MD, Surette MG. Longitudinal sampling of the lung microbiota in individuals with cystic fibrosis. PLoS One. 2017;12:e0172811.

10. Carmody LA, Zhao J, Schloss PD, Petrosino JF, Murray S, Young VB, et al. Changes in cystic fibrosis airway microbiota at pulmonary exacerbation. Ann Am Thorac Soc. 2013;10:179–87.

11. Carmody LA, Zhao J, Kalikin LM, LeBar W, Simon RH, Venkataraman A, et al. The daily dynamics of cystic fibrosis airway microbiota during clinical stability and at exacerbation. Microbiome. 2015;3:12.

12. Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, et al. A human gut microbial gene catalogue established by metagenomic sequencing. Nature. 2010;464:59–65.

13. Bacci G, Mengoni A, Fiscarelli E, Segata N, Taccetti G, Dolce D, et al. A different microbiome gene repertoire in the airways of cystic fibrosis patients with severe lung disease. Int J Mol Sci. 2017;18:1654.

14. Franzosa EA, Morgan XC, Segata N, Waldron L, Reyes J, Earl AM, et al. Relating the metatranscriptome and metagenome of the human gut. Proc Natl Acad Sci. 2014;111:E2329–38.

15. Lim YW, Schmieder R, Haynes M, Willner D, Furlan M, Youle M, et al. Metagenomics and metatranscriptomics: Windows on CF-associated viral and microbial communities. J Cyst Fibros. 2013;12:154–64.

16. Segata N. On the Road to Strain-Resolved Comparative Metagenomics. mSystems. 2018;3:e00190-17.

17. Beghini F, Pasolli E, Truong TD, Putignani L, Cacciò SM, Segata N. Large-scale comparative
metagenomics of *Blastocystis*, a common member of the human gut microbiome. ISME J.
2017;11:2848–63.

18. Zolfo M, Tett A, Jousson O, Donati C, Segata N. MetaMLST: Multi-locus strain-level bacterial
typing from metagenomic samples. Nucleic Acids Res. 2017;45:e7.

19. Zhao J, Schloss PD, Kalikin LM, Carmody LA, Foster BK, Petrosino JF, et al. Decade-long
bacterial community dynamics in cystic fibrosis airways. Proc Natl Acad Sci. 2012;109:5809–14.

20. Delhaes L, Monchy S, Fréalle E, Hubans C, Salleron J, Leroy S, et al. The airway microbiota in
cystic fibrosis: A complex fungal and bacterial community-implications for therapeutic
management. PLoS One. 2012;7:e36313.

21. Willner D, Daly J, Whiley D, Grimwood K, Wainwright CE, Hugenholtz P. Comparison of
DNA extraction methods for microbial community profiling with an application to pediatric
bronchoalveolar lavage samples. PLoS One. 2012;7:e34605.

22. Lim YW, Evangelista JS, Schmieder R, Bailey B, Haynes M, Furlan M, et al. Clinical insights
from metagenomic analysis of sputum samples from patients with cystic fibrosis. J Clin Microbiol.
2014;52:425–37.

23. Willner D, Haynes MR, Furlan M, Hanson N, Kirby B, Lim YW, et al. Case studies of the
spatial heterogeneity of DNA viruses in the cystic fibrosis lung. Am J Respir Cell Mol Biol.
2012;46:127–31.

24. Whiteson KL, Meinardi S, Lim YW, Schmieder R, Maughan H, Quinn R, et al. Breath gas
metabolites and bacterial metagenomes from cystic fibrosis airways indicate active pH neutral 2,3-
butanedione fermentation. ISME J. 2014;8:1247–58.

25. Surette MG. The cystic fibrosis lung microbiome. Ann Am Thorac Soc. 2014;11:61–5.

26. Quince C, Walker AW, Simpson JT, Loman NJ, Segata N. Shotgun metagenomics, from
sampling to analysis. Nat Biotechnol. 2017;35:833–44.

27. Truong DT, Tett A, Pasolli E, Huttenhower C, Segata N. Microbial strain-level population
structure and genetic diversity from metagenomes. Genome Res. 2017;626–38.

28. Scholz M, Ward D V., Pasolli E, Tolio T, Zolfo M, Asnicar F, et al. Strain-level microbial
epidemiology and population genomics from shotgun metagenomics. Nat Methods. 2016;13:435–8.

29. Faust K, Lahti L, Gonze D, de Vos WM, Raes J. Metagenomics meets time series analysis:
unraveling microbial community dynamics. Curr Opin Microbiol. 2015;25:56–66.

30. Shade A, Peter H, Allison SD, Baho DL, Berga M, Bürgmann H, et al. Fundamentals of
microbial community resistance and resilience. Front Microbiol. 2012;3:417.

31. Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight R. Diversity, stability and
resilience of the human gut microbiota. Nature. 2012;489:220–30.

32. Zemanick ET, Wagner BD, Robertson CE, Ahrens RC, Chmiel JF, Clancy JP, et al. Airway
microbiota across age and disease spectrum in cystic fibrosis. Eur Respir J. 2017; 50(5). pii:
1700832

33. Carmody LA, Caverly LJ, Foster BK, Rogers MAM, Kalikin LM, Simon RH, et al. Fluctuations
in airway bacterial communities associated with clinical states and disease stages in cystic fibrosis.

PLoS One. 2018; 13(3): e0194060.

34. Doolittle WF, Booth A. It's the song, not the singer: an exploration of holobiosis and
evolutionary theory. Biol Philos. 2016;32:5–24.

35. Eng A, Borenstein E. Taxa-function robustness in microbial communities. Microbiome.
2018;6:45.

36. Louca S, Jacques SMS, Pires APF, Leal JS, Srivastava DS, Parfrey LW, et al. High taxonomic
variability despite stable functional structure across microbial communities. Nat Ecol Evol.
2016;1(1):15.

37. Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. An obesity-
associated gut microbiome with increased capacity for energy harvest. Nature. 2006;444:1027–31.

38. Thaiss CA, Itav S, Rothschild D, Meijer M, Levy M, Moresi C, et al. Persistent microbiome
alterations modulate the rate of post-dieting weight regain. Nature. 2016;540:1–27.

39. Bevivino A, Bacci G, Drevinek P, Nelson MT, Hoffman LR, Mengoni A. Deciphering the
ecology of cystic fibrosis bacterial communities: Towards systems-level integration. Trends Mol
Med. 2019; in press. DOI:10.1016/j.molmed.2019.07.008

40. Farrell PM, White TB, Ren CL, Hempstead SE, Accurso F, Derichs N, et al. Diagnosis of cystic
fibrosis: Consensus guidelines from the cystic fibrosis foundation. J Pediatr. 2017;181:S4–15.

41. Proesmans M, Balinska-Miskiewicz W, Dupont L, Bossuyt X, Verhaegen J, Høiby N, et al.
Evaluating the "Leeds criteria" for *Pseudomonas aeruginosa* infection in a cystic fibrosis centre.
Eur Respir J. 2006;27:937–43.

42. Bacci G, Paganin P, Lopez L, Vanni C, Dalmastri C, Cantale C, et al. Pyrosequencing unveils
cystic fibrosis lung microbiome differences associated with a severe lung function decline. PLoS
One. 2016;11:e0156807.

43. Kerem E, Conway S, Elborn S, Heijerman H, Committee C, others. Standards of care for
patients with cystic fibrosis: a European consensus. J Cyst Fibros. 2005;4:7–26.

44. Smyth AR, Bell SC, Bojcin S, Bryon M, Duff A, Flume P, et al. European cystic fibrosis society
standards of care: Best practice guidelines. J Cyst Fibros. 2014;13:S30–53.

45. Taylor-Robinson D, Whitehead M, Diderichsen F, Olesen H V, Pressler T, Smyth RL, et al.
Understanding the natural progression in %FEV1 decline in patients with cystic fibrosis: A
longitudinal study. Thorax. 2012;67:860–6.

46. Paganin P, Fiscarelli EV, Tuccio V, Chiancianesi M, Bacci G, Morelli P, et al. Changes in cystic
fibrosis airway microbial community associated with a severe decline in lung function. PLoS One.
2015;10:e0124348.

47. Farrell PM, Rosenstein BJ, White TB, Accurso FJ, Castellani C, Cutting GR, et al. Guidelines
for diagnosis of cystic fibrosis in newborns through older adults: Cystic fibrosis Foundation
consensus report. J Pediatr. 2008;153:S4–14.

48. Bacci G, Bazzicalupo M, Benedetti A, Mengoni A. StreamingTrim 1.0: A Java software for
dynamic trimming of 16S rRNA sequence data from metagenetic studies. Mol Ecol Resour.
2014;14:426–34.

49. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.

2012;9:357–9.

50. Abubucker S, Segata N, Goll J, Schubert AM, Izard J, Cantarel BL, et al. Metabolic
reconstruction for metagenomic data and its application to the human microbiome. PLoS Comput
Biol. 2012;8:e1002358.

51. Truong DT, Franzosa EA, Tickle TL, Scholz M, Weingart G, Pasolli E, et al. MetaPhlAn2 for
enhanced metagenomic taxonomic profiling. Nat Methods. 2015;12:902–3.

52. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: A new versatile metagenomic
assembler. Genome Res. 2017;27:824–34.

53. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: Prokaryotic gene
recognition and translation initiation site identification. BMC Bioinformatics. 2010;11:119.

54. Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC, et al. eggNOG 4.5: A
hierarchical orthology framework with improved functional annotations for eukaryotic, prokaryotic
and viral sequences. Nucleic Acids Res. 2016;44:D286–93.

55. Pearson WR. An introduction to sequence similarity ("homology") searching. Curr Protoc
Bioinformatics. 2013; Chapter 3 Unit 3.1: p. 3.1.1-3.1.8.

56. Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, et al. CARD 2017:
Expansion and model-centric curation of the comprehensive antibiotic resistance database. Nucleic
Acids Res. 2017;45:D566–73.

57. R Development Core Team. R: A language and environment for statistical computing. R
foundation for statistical computing, Vienna, Austria. http://www.R-project.org/. R Found Stat
Comput Vienna, Austria. 2012.

58. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. *limma* powers differential
expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res. 2015;43:e47.

59. Law CW, Chen Y, Shi W, Smith GK. voom: precision weights unlock linear model analysis
tools for RNA-seq read counts. Genome Biol. 2014;15(2):R29.

a 0.001

b 0.002

c 0.01

d 0.002

- ● B01
- ● B02
- ● B03
- ● B06
- ● G10
- ● G24
- ● G28
- ● G30
- ● G31
- ● G34
- ● M05
- ● M19
- ● M21
- ● M22
- ● M23
- ● M24
- ● M25
- ● M26
- ● M28
- ● M29
- ● M31
- ● M33
- ● M34
- ◇ reference

a)

Pathways	
Taxonomy	

Genotype □ heterozygote ○ homozygote △ other

Status ● Baseline ● Treatment ● Recovery

b)

Bray–Curtis dissimilarity

Exacerbation: TR, BL, RC

Genotype: heterozygote, homozygote

■ paths ■ taxa