1

The effect of symbiosis on symbiont fitness – interactions within

a simple metaorganism

Peter Deines[a]*, Katrin Hammerschmidt[b] and Thomas CG Bosch[a]

[a]Zoological Institute, Christian Albrechts University Kiel, 24118 Kiel, Germany

[b]Institute of General Microbiology, Christian Albrechts University Kiel, 24118 Kiel, Germany

*Correspondence:

Peter Deines, Zoological Institute, Christian Albrechts University Kiel, Am Botanischen Garten 1-9, 24118 Kiel, Germany

Tel: +49-431-880-4171

Email: pdeines@zoologie.uni-kiel.de

Running title: Interactions drive microbiome species composition

Abstract

Organisms and their resident microbial communities form a complex and mostly stable

ecosystem. It is known that the specific composition and abundance of certain bacterial

species affect host health and *Darwinian fitness,* but the processes that lead to these

microbial patterns are unknown. We investigate this by deconstructing the simple

microbiome of the freshwater polyp *Hydra*. We contrast the performance of its

two main bacterial associates, *Curvibacter* and *Duganella*, on germ free hosts

with two *in vitro* environments over time. We show that interactions within the

microbiome but also host modulation lead to the observed species frequencies

and abundances. More specifically, we find that rare microbiome members are

essential for achieving the observed community composition, which ultimately

sets the maximum carrying capacity. Bacterial fitness strongly depends on the

environment: while *Duganella* performs better than *Curvibacter* in a non-host

habitat, *Curvibacter* benefits through the host association. This is of particular

interest because *Curvibacter* and its host show a history of co-evolution, as

inferred from phylogenies, whereas the colonization with *Duganella* seems to be

a recent event. Our findings oppose the assumption that bacteria always benefit

through the association with the host and poses questions regarding the long-

term maintenance of such relationships.

Introduction

Eukaryotes form a distinct habitat for microbial communities (microbiomes) and these microbial associations are integral to life. The host with its associated microbial community, often dominated by bacteria but co-habited by fungi, protozoa, archaea, and viruses, is termed metaorganism. Microbiomes can contain from few up to thousands of microbial species - the human microbiome, for example, is estimated to be comprised of about 5000 bacterial species (1-3). These host-associated microbial communities have been shown to enhance host function and contribute to host fitness and health (4). Changes in microbiome diversity, function, and density have been linked to a variety of disorders in many organisms (5-8).

A major goal in host-microbe ecology is to unravel the ecological and evolutionary dynamics of microorganisms within their communities. Of particular relevance are the factors that shape the stability and resilience of such communities, despite different fitness trajectories of the microbiome members. The microbial response to stress or perturbations, e.g. exposure to a new substrate, provides a selective advantage to certain members of the community. If the system cannot tolerate the change, the microbial community dramatically shifts until a different equilibrium state is reached (9). Frequency-dependent selection forces the host to adapt to these changes and select for or against the most frequent genotypes of their associated microbiota (10). There is, for example, strong evidence that species-specific antimicrobial peptides (AMPs) shape, control, and confine host-species specific bacterial associations (11, 12). In addition, microbial communities are not evenly distributed, e.g. along the gastrointestinal tract or between the lumen and the epithelial surfaces (2, 13, 14). These significant differences in niches or micro-habitats and their occupancy is known as spatial heterogeneity and will affect community assembly rules and dynamics (15, 16). Interspecies metabolic exchange is another key biotic force acting as a major driver of species co-occurrence in diverse microbial communities (17).

One aspect that has been so far neglected in metaorganism research is why a

stable microbiome can persist in a specific host lineage for many host generations.

Central to this is the question whether microbes benefit from the association with the

host. This could either happen through direct interaction with the host or indirectly,

through interactions with another microbial strain within the microbiome. Host-

dependent advantages for the symbiont range from the availability of host-derived

nutrients, a competition-free environment to reduced predation risk (18). When

members of the microbiome benefit from the association with a particular host, one

would expect their capacity to reproduce, i.e. their *Darwinian fitness*, to be higher in the

host relative to an alternative habitat, as demonstrated for the squid-*Vibrio* system (19).

Surprisingly, this aspect has been tested less frequently than the effect of microbes on

host fitness, which has been quantified in many systems.

Methods for quantifying microbial fitness range from estimating absolute

fitness, where population growth is measured from which maximum growth rates can

be calculated, to estimates of relative fitness. The latter is of particular relevance as the

fitness of a focal microbial lineage is measured *relative* to another microbial lineage.

This provides a better representation of the turnover of strains in microbial populations

and also closely corresponds to the meaning of fitness in evolutionary theory (20). The

basic design of such an experiment involves mixing populations of the two strains

together in a particular initial ratio in the test environment and measuring their relative

contributions to future generations. Under certain ecological conditions the relative

fitness of two strains is dependent upon their relative frequencies (frequency-

dependent selection) (21). These methods of measuring microbial fitness can easily be

applied to host-microbe systems and help us in understanding the underlying dynamics.

This will yield insights into the role that microbes played in the evolution of eukaryotes

and into the ecology and evolution of host-microbe associations in general (22). Here, it

is important to unravel the extent to which the observed pattern simply result from

intrinsic properties, such as microbial growth rates, interactions of the microbes, which

could be recapitulated *in vitro* with co-culture experiments, or whether it is an emergent

property of that particular host-microbe system (23).

We here test whether microbial associates benefit from their association with

the host and their fellow community members and follow their ecological interactions

within the microbiome. We use the freshwater polyp *Hydra vulgaris* and its microbiome,

which has become a valuable experimental model in metaorganism research as it

provides an excellent bridge between the simplicity of synthetic communities and the

mouse model (24). *Hydra's* ectodermal epithelial cells are covered with a multi-layered

glycocalyx that provides a habitat for a species-specific and core microbiome of low

complexity (11, 25, 26), from which most microbes can be cultured *in vitro* (26, 27). This

allows for an integrated approach based on constructing synthetic communities of

various complexities and contrasting the host (*in vivo)* to *in vitro* habitats (24). We focus

on the two most abundant members of the microbiome that together constitute about

85% of *Hydra*'s simple microbiome, *Curvibacter* sp. AEP1.3 and *Duganella* sp. C1.2,

(hereafter called *Curvibacter* and *Duganella*) where abundances of *Curvibacter* are

several magnitudes higher as compared to *Duganella* (27). Previous work on species

interactions suggested that continued coexistence depends heavily on relative growth

rates. Using invasion-from-rare experiments we explore the potential for coexistence

under different environmental conditions (host and non-host habitat) and contribute to

a better understanding whether the host alters microbial interactions that potentially

lead to coexistence in simple microbiomes.

Results

Carrying capacity of the *Hydra* host

The carrying capacity of the *Hydra* habitat is highly stable among single *Hydra* wild-type

polyps with $1.7*10^5$ CFUs/polyp and at its maximum with the native microbiome.

Manipulations of the microbial community lead to significant changes in bacterial numbers (Welch ANOVA, $F_3=7.054$; $P<0.005$; Fig. 1A). The addition of single bacterial species (*Curvibacter* or *Duganella*) to wild-type polys does not increase the carrying capacity but, in the case of *Curvibacter*, leads to a significant reduction in microbial population size. In contrast does the addition of *Duganella* not significantly shift the bacterial numbers (Fig. 1A). Conventionalised animals (germ-free polyps incubated with tissue homogenates of wild-type animals) show a carrying capacity, which is not significantly different compared to wild-type polyps (Fig. 1A), indicating the usability of the germ-free polyps for the manipulation and construction of synthetic bacterial communities on the host/ *in vivo*.

Tracking microbial mono-colonisations in *Hydra* over time

In mono-associations *Curvibacter* and *Duganella* populations display typical bacterial growth dynamics. Following a lag and exponential phase, both strains reach a stationary phase after roughly 72 h with a stable population size/carrying capacity per host (Fig. 1B). After 72 h we find significant differences in the carrying capacity between both strains, *Curvibacter* being higher than *Duganella* (estimated by post hoc contrasts; Generalized linear model: Full model: $\chi^2=54.360$, d.f.=9, $P<0.0001$; bacterial species x days post exposure: $\chi^2=18.326$, d.f.=4, $P=0.0011$). These significant differences last until the end of the experiment. Once carrying capacity of mono-associations is reached, we find a population size of about 10^4 CFUs per host for *Curvibacter*, whereas for *Duganella* the population size reaches on average only $1.5*10^3$ CFUs per host. Further, both mono-associations do not reach the carrying capacity of wild-type polyps. The variation in bacterial density between hosts is significantly higher in *Curvibacter* than in *Duganella* (Levene: $F_1=21.496$, $P<0.0001$). No extinction events are observed in either strain after successful establishment during the mono-colonisation experiments.

Microbial growth kinetics of mono-associations *in vivo* and *in vitro*

Growth rates of *Curvibacter* did not significantly differ between the host and the microcosm environments. This is in marked contrast to *Duganella*, where significantly higher growth rates were observed in the non-host as compared to the host environment. In all environments, except for the host, *Duganella* achieved a significantly higher growth rate than *Curvibacter* (determined by post hoc t-tests; ANOVA: R^2=0.827; Full model: $F_{5,15}$ = 14.333; P<0.0001; bacterial species x environment: F_2 = 15.592; P=0.0002; Fig. 2).

Microbial di-association experiments *in vivo* and *in vitro*

Carrying capacity

Results of host and microcosm experiments (static and mixed) show that during di-associations the carrying capacity in all habitats is reached at about 72 h after inoculation. Both microcosm environments are characterized by a carrying capacity of 10^7-10^8 CFUs/ml, and so exceeding the *in vivo* carrying capacity by a factor of 10^4 (Fig. 3). Nevertheless, di-associations on the host also fail to reach the carrying capacity of wild-type polyps and reach a comparable carrying capacity as in the mono-colonisations of *Curvibacter*. Both bacterial species do not match the carrying capacities as measured in mono-colonisations on the host: whereas *Curvibacter* fails by a power of 10 to reach its density in the mono-colonisations, *Duganella* outgrows it by a power of 10.

Changes in fractions in microbial di-association experiments

When competing *Curvibacter* and *Duganella* in three different starting frequencies in the host, and the non-host environments, several interesting patterns can be observed (Fig. 4). In both non-host environments, *Duganella* outcompetes *Curvibacter* within 48 h post exposure. From then onwards, frequencies of *Curvibacter* are low, reaching a maximum of about 10%. This pattern does not depend on the initial frequency at the start of the

experiment. The host shows a different pattern. Here, a decrease in *Curvibacter* fraction

can be observed in all three initial frequencies but never to a point where it cannot be

detected in the population. From 72 h post exposure onwards the population on the host

has reached a stable state, with *Curvibacter* making up 20% of the total bacterial

population.

Microbial growth kinetics of di-associations

Overall *Curvibacter* growth rate in di-associations are lower or not different from the

mono-associations (Fig. 5; as estimated by post hoc contrasts; Generalized linear model:

Full model: χ^2=45.790, d.f.=11, P<0.0001; environment x initial frequency: χ^2=33.685,

d.f.=6, P<0.0001). *Curvibacter* grows significantly differently when inoculated in equal

densities as compared to the rare and dominant starting frequencies across the different

environments. Whereas in the host, *Curvibacter* grows better when in equal density with

Duganella, the opposite is true for both *in vitro* environments.

As observed for the growth of *Duganella* in mono-colonisations, growth rates are always

higher in the non-host environments irrespective of initial frequency (Generalized linear

model: Full model: χ^2=130.278, d.f.=11, P<0.0001; environment x initial frequency:

χ^2=59.723, d.f.=6, P<0.0001). Whereas, in di-associations, negative growth rates can be

detected only once for *Duganella*, it happens more frequently in the *Curvibacter*,

indicating a direct or indirect negative effect of *Duganella*.

Relative performance of Curvibacter

As estimation for the relative performance of *Curvibacter* in competition with *Duganella*

we determined the selection rate constant (r) for all di-association experiments (28, 29).

We observed r < 0 for all environments and frequencies, indicating that *Curvibacter*

performs worse than *Duganella* when they are in direct competition (Fig. 6). There is no

overall effect of the initial frequency - only in the mixed environment differences can be

detected with the lowest performance of *Curvibacter* at equal frequencies of the two

competitors (Generalized linear model: Full model: χ^2=51.168, d.f.=8, P<0.0001;

environment x initial frequency: χ^2=19.147, d.f.=4, P=0.0007). In general, *Curvibacter*

performs significantly better in the host environment irrespective of initial frequency.

Discussion

One of the major challenges in microbiome research is to understand the factors that

influence the dynamics and stability of host-associated microbial communities. Of

particular relevance for this are the processes governing assembly (30, 31) and

resilience (32). Insights into such processes in bacterial populations within their native

host environments can be gained through a number of ways. Relationships between

different community members in complex microbiomes, such as the human gut, can, for

example, be inferred by analysing co-occurrence data and correlation patterns from

sequencing-based metagenomic time-series experiments (33, 34). Another approach,

where ecological interactions can experimentally be dissected, allows for hypothesis

testing. For this, a relatively simple system, such as *Hydra* and its microbiome is ideal for

'deconstructing' a metaorganism and its interactions (24). This strategy is novel in

meta-organism research but has yielded exciting results regarding rules that determine

community assembly and stability in non-host associated microbial communities (35,

36). In microbiome research, the added advantage of including a host provides the

opportunity to study the performance of the individual microbiome members in their

natural environment. In addition, the comparison of the *in vivo* to *in vitro* environments

provides information on host effects in regulating its microbiome.

227   Carrying capacity is defined as the largest population size that an ecosystem can

sustainably support without degrading the ecosystem. Therefore population size is

constrained by the carrying capacity of the environment. This is also true for bacteria

populating a particular environment, including hosts. In microbiome research, the

significance of host carrying capacity has been largely overlooked until very recently

where a link between host health and microbiome density has been reported (8).

Bacterial levels have been quantified for only a few model organisms such as in the gut

of larval zebrafish (37) or the gut of *Drosophila melanogaster* (38). The host's carrying

capacity ultimately sets the upper boundary for bacterial fitness relative to any

competitors, and provides a reference for investigating the interactions within the host

microbiota at spatial and temporal scales.

We here show that wild-type *Hydra* is characterized by a specific carrying

capacity of about 10^5 bacteria per polyp that is stable in adult polyps and can be

artificially assembled through the re-population of germ-free animals with the natural

bacterial community. This is an important prerequisite for conducting the *in vivo*

experiments, where colonization patterns of single species from *Hydra's* microbiome are

individually followed. When *Curvibacter* and *Duganella* are introduced separately to the

host (in mono-associations), each bacterial species is capable to robustly colonize the

host to high abundances, which are maintained throughout the experimental period,

indicating the ability to independently utilize the resources of the *Hydra* habitat. This

confirms earlier findings from Wein *et al.* (39) that *Curvibacter* is able to robustly

colonize its host. A similar observation has been made for *Aeromonas* and *Vibrio*

colonizing patterns of the gut of larval zebrafish (23). While both, *Curvibacter* and

Duganella, fail to reach their respective carrying capacities as compared to wild type

Hydra by a factor of 10, their respective proportions of about 75% and 11% (27) are

nevertheless maintained. Results from the di-association experiments clearly show that

the abundances and relative frequencies of *Curvibacter* and *Duganella* as measured in

wild-type *Hydra* cannot be explained by their (positive) interactions in the host context

alone as this led to a frequency reversal, making *Duganella* more abundant than

Curvibacter. *Duganella* however reaches comparable carrying capacities as measured in

wild-type *Hydra* polyps. This implies that the presence of *Curvibacter* has a positive

effect on the performance of *Duganella* but *Duganella* negatively affects *Curvibacter* in

the host habitat. These observations indicate that the less frequent community members

(each 2% and less) are of central importance for reaching the full carrying capacity of

the *Hydra* microbiome. We hypothesise that two aspects are of importance here – (i) the

low abundant microbes might be able to utilize different resources as compared to

Curvibacter and *Duganella* and so inhabit different ecological niches within the

microbiome, which the two main colonizers cannot fill and (ii) they likely interact in a

positive way (either directly or indirectly) with (at least) *Curvibacter*, enabling it to

reach higher carrying capacities. It is thus important to note that *Hydra*'s carrying

capacity is not solely determined by the host (resources) alone but also by the

interactions within the microbiome.

To determine the relative importance of the host, interactions of *Curvibacter* and

Duganella, and intrinsic properties of the bacteria, such as ability to grow, in shaping the

observed patterns in the host environment, we performed the same mono- and di-

association experiments in two i*n vitro* environments. We chose to contrast static, which

closely resembles the host habitat in that it provides spatial heterogeneity facilitating

bacterial interactions and mixed, where direct interactions between individual bacteria

cannot be established but where individual bacteria have (unlimited) access to

resources and oxygen. Because community structure can be influenced by initial species

abundances (40), we performed all di-association experiments using various initial

starting frequencies. In general, both bacterial species reached much higher carrying

capacities in both *in vitro* environments as compared to the *in vivo* environment, with

Duganella reaching up to 10^8 CFUs/ml and *Curvibacter* up to 10^5 CFUs/ml. Interestingly

we were not able to replicate the *in vivo* dynamics between *Curvibacter* and *Duganella in*

vitro, suggesting that the host mediates the interaction between the two bacterial

species. Previous work has suggested that continued coexistence between bacterial

species is dependent on their relative growth. Coexistence is maintained if the slower-

growing species is released from its dependence on the other (41), which in our case

could be compensated by the host. This aspect will be subject of further investigation.

Wright and Vetsigian (42) recently demonstrated in pairwise competitions between

bacteria of the genus *Streptomyces* that the winner is often the species that starts at high

initial abundance. We find that 'survival of the common' does not apply to *Curvibacter* in

a non-host environment. Whereas pairwise competitions in the di-association

experiments resulted in coexistence of *Curvibacter* and *Duganella* in the host habitat, it

led to an overgrowth of *Duganella* in both *in vitro* habitats. More specifically, we found

that in the *Hydra* habitat *Curvibacter*, independent of its inoculation frequency, and after

the initial establishment period of 72 h, reached a constant fraction of about 20%,

whereas it was only present at very low frequencies in both *in vitro* habitats due to the

high continuous growth rates of *Duganella* (see also (43) for homogeneous

environment). This pattern seems to be more pronounced in the static environment,

fitting to the general assumption that extinctions are more likely to occur in spatially

structure environments. In terms of cross-feeding interactions one could reason that the

faster growing-species is released from its dependence on the other, resulting in the

slower-growing species being lost (a hypothesis that is termed 'feed the faster grower')

(41).

Many host-associated microbial communities live in highly competitive

surroundings, in which they struggle to persist. According to Ghoul and Mitri (44), a

strain is competitive if it shows phenotypes that cause a fitness decrease in a competitor

strain. Competition can be either passive, where strains compete for the same resources

or active, where strains directly harm one another (44). Li *et al.* (43) showed that during

co-culture experiments in a homogenous environment the presence and frequency of

Curvibacter greatly influenced the growth rate of *Duganella* (quadratic function), which

we also found. Interestingly we find the same pattern on the host, although growth rates

are drastically reduced. In the static environment we find the inverse of the quadratic

function.

When comparing the bacterial growth rates in mono-associations we observe

that *Curvibacter* performs equally well in all habitats, whereas *Duganella* shows a

reduced growth rate compared to *Curvibacter* on the host and performs best in the non-

host habitat. This suggests that *Duganella* is not particularly adapted to the host niche.

Interestingly the two species also differ in their phylogenetic association patterns with

different *Hydra* species (45). Whereas different lineages of *Curvibacter* have been shown

to be associated with the majority (N=6) of *Hydra* species analysed (N=7), *Duganella*

was detected in less than half (N=3). This indicates that *Duganella* is a more recent

member of the *Hydra* microbiome and that it is not as co-evolved as it has been

suggested for one of *the* keystone species of *Hydra's* microbiome, *Curvibacter* (43).

Nevertheless, the specific microbiome of *Hydra vulgaris*, which is studied here, has been

shown to be stable as animals living in the wild were colonized by a similar microbiome

as compared to the ones that had been maintained in the laboratory for a relatively long

time (25). One of the few metaorganisms, where symbiont performance within the host

relative to a non-host environment has been contrasted, is the squid-*Vibrio* system. For

this obligate association Wollenberg and Ruby (19) demonstrated that symbionts have

an increased reproductive capacity and higher fitness when associated with the host. In

combination with the results from this study, this indicates that more specifically

associated symbionts indeed benefit from growing in the host habitat.

Oscillatory dynamics, chaotic fluctuations and stochastic perturbations influence

species abundances and allow for the coexistence of many species and have therefore a

significant impact on species survival even in simple communities containing only a few

interacting species (46-48). Interestingly we observed such sporadic collapses in the

Curvibacter population during di-association experiments in the non-host environment

but not on the host. An earlier study following the bacterial colonization pattern of

Hydra hatchlings showed sporadic collapses of the *Curvibacter* population during the

first 15 weeks post hatching (49). In adult polyps, which were used in the experiments,

we did not observe such dynamics, which points to the fact that these sporadic collapses

might be buffered in adult polyps or are caused by the other species of the *Hydra*

microbiome. For the zebrafish host model it has been shown that *Aeromonas* during

mono-association is not resistant to disturbances related to intestinal motility and that

in the presence of *Vibrio* the magnitude of collapses are even greater (23). *Hydra* is also

characterized by spontaneous contractile activity (50) but our data suggest that this

does not have such far-reaching effects on the microbiota as found in the zebrafish gut.

Whereas in other host-microbe systems such as the zebrafish, *in vivo* dynamics

between two bacterial species can be replicated *in vitro* (51), in our system this is not

the case. We find the fitness of *Curvibacter* and *Duganella* to be highly dependent on the

specific environments tested. Whereas *Duganella* performs best *in vitro*, in terms of

carrying capacities reached, absolute fitness (growth rates) but also relative fitness

(fractions in population, selection rate constant), the opposite is true for all parameters

in *Curvibacter* which generally performs best in the host habitat, except for its carrying

capacity that is higher *in vitro*. Our experiments reveal that not only direct bacterial

interactions are important but also highlight the fact that the host plays an important

role in mediating these interactions (Fig. 7). Further, we were able to show that carrying

capacity is a fundamental property of a host-microbe system and that it is not only

determined by the physiological features of the host but also by the interactions within

the microbiome. Detecting a selection rate constant of $r < 0$ for the main *Hydra*

colonizer, *Curvibacter*, in the host environment and a reduced carrying capacity in both,

mono- and di-associations, suggests that low-abundance species of the *Hydra*

microbiome are likely playing a major role in maintaining the density of *Curvibacter* on

the host. Evidence for the importance of rare-species comes from the human intestinal

microbiota, which contains many low-abundance species (52) with some of them having

a large impact on inducing dysbiosis in the microbiome and on guaranteeing host health

(53, 54).

Materials and Methods

Animals used, culture conditions and generation of germ-free animals

Hydra vulgaris (strain AEP) was used for carrying out experiments and cultured

according to standard procedures at 18°C in standardized culture medium (Hydra

medium (HM)) (55). Animals were fed three times a week with 1st instar larvae of

Artemia salina. Germ-free (GF) polyps were obtained by treating wild-type (WT)

animals with an antibiotic cocktail solution containing 50 μ g/ml ampicillin, neomycin,

streptomycin, rifampicin and 60 μ g/ml spectinomycin as previously described (45, 50).

The antibiotic cocktail solution was exchanged every 48 h and the antibiotic treatment

lasted for two weeks, after which polyps were transferred into antibiotic-free sterile HM

for recovery (four days). The germ-free status of polyps was confirmed as previously

described (45). During antibiotic treatment and re-colonization experiments, polyps

were not fed.

Bacterial strains and media

The bacterial strains used in this study are *Curvibacter* sp. AEP1.3 and *Duganella* sp.

C1.2., which have been isolated from *Hydra vulgaris* (strain AEP) (27). These bacteria

were cultured from existing isolate stocks in R2A medium at 18°C, shaken at 250 r.p.m

for 72 h before use in the different experiments.

Carrying capacity of the host

To determine the carrying capacity of the *Hydra* habitat the microbial load of individual

Hydra polyps (N=16) was determined. In addition to wild-type polyps the carrying

capacity of conventionalized polys (N=12), obtained by incubating germ-free polyps

with tissue homogenates of wild-type animals (per germ-free polyp one wild-type polyp

was used) for 24 h was also determined. To test whether the carrying capacity can

artificially be increased or destabilises upon self-challenge we added either *Curvibacter*

or *Duganella* to wild-type polyps (N=6) (approximately 5×10^3 cells for 24 h). After

incubation all polyps were washed with and transferred to sterile HM and further

incubated at 18°C and sampled after 120 h.

Tracking microbial mono-associations in *Hydra* over time

Germ-free polys were inoculated in their aquatic environment with single bacterial

strains (mono-associations). Individual germ-free polyps were incubated with 5×10^3

cells of *Curvibacter* or *Duganella* in 1.5 ml Eppendorf tubes containing 1 ml of sterile

HM. After 24 h of incubation all polyps were washed with and transferred to sterile HM,

incubated at 18°C and followed over a period of 216 h. For each treatment 6 polys per

time point were independently analysed. Every 48 h individual polyps were

homogenized in an Eppendorf tube using a sterile pestle and serial dilutions of the

homogenate were plated on R2A agar plates to determine colony-forming units (CFUs)

per polyp.

Microbial growth kinetics of mono- and di-associations *in vivo* and *in vitro*

To study the initial phase of colonization, i.e. 96 h post inoculation (see Fig. 1B) in more

detail microbial growth of *Curvibacter* and *Duganella* was determined in different

habitats; the host habitat (*in vivo*) and two different microcosm environments (*in vitro*).

The static incubation provided a spatially structured habitat (heterogeneous), whereas

shaking of the microcosms (mixed treatment) eliminated the spatial structure

(homogenous).

Mono-associations: All germ-free polyps and microcosms were inoculated from

the same bacterial inoculation culture with approximately 5×10^3 cells of *Curvibacter* or

Duganella for 24 h, washed with and transferred to sterile HM. Samples were taken

every 12 h for 96 h. For *Hydra* six polyps were sacrificed at each time point and colony-

forming units (CFUs) were determined as described above. As microcosms 24-well

plates were used. Wells were filled with 2 ml of R2A medium, inoculated and incubated

at 18°C either under static or shaken (200 r.p.m) conditions. Each time point was

replicated four times and serial dilutions were plated on R2A agar plates to determine

CFUs. Growth rates of each strain (A and B) were determined for the exponential growth

phase (36-48 h) and were calculated as $g = \ln (A_f/A_0)$ and $g = \ln (B_f/B_0)$, where A_0, B_0 is

the starting density at time 0 and A_f, B_f is the final density at time *f*.

Di-associations: Density dependent competiveness fitness assays of the two most

dominant colonizers *Curvibacter* and *Duganella* were performed *in vivo* and *in vitro.* The

same host and microcosm experiments as described above were performed except for

using microbial di-associations of *Curvibacter* and *Duganella* with the frequency of

Curvibacter being rare (10:90), equal (50:50), or dominant (90:10). As *Curvibacter* and

Duganella form distinct colonies on R2A agar plates, their frequency can be determined

by plating serial dilutions (43). Six polyps and four microcosm replicates were assayed

per treatment (static and mixed) and time point. Also this data allowed determining the

different carrying capacities of the *in vivo* and *in vitro* habitats used. Malthusian

parameters were calculated for both bacterial strains (A and B), $m = \ln (N_f/N_0)/hour$,

where N_0 and N_f were the initial and final densities. The relative performance of the two

strains was expressed in terms of the selection rate constant (*r*). Note that this measure

of performance should be preferentially used over the more commonly used relative

fitness (*W*), when one competitor is much less fit than the other (28, 29), as was the case

in our study. The selection rate constant was calculated as $r = m_B - m_A$. Accordingly, $r = 0$

implies that there is no difference in growth rates of the two strains, $r > 0$ indicates that

strain B is advantageous in population growth ($r = 1$ indicates a 10-fold increase in the

ratio of B vs. A abundances), while r < 0 suggests that strain B fails in invading

population of strain A.

Statistical analysis

A Welch ANOVA (and subsequent Dunnett's posthoc test) was used to test for

differences in bacterial abundance patterns ('bacteria per *Hydra*') in wild-type versus

manipulated hosts as variances between the different groups were not equally

distributed.

Differences during mono-colonizations of *Curvibacter* and *Duganella* over time

were assessed using a Generalized linear model (error structure: normal; link function:

identity). The response variable was 'bacteria per *Hydra*', and explanatory variables

were 'bacterial species', 'time' and 'bacterial species' x 'time'. Differences between the

two bacterial species on each day were detected with post hoc contrasts.

Analysis of variance (ANOVA) and subsequent post hoc t-tests were used to test

for differences in growth rates of the two competitors when grown singly in the

different environments. The response variable was 'growth rate', and explanatory

variables were 'bacterial species', 'environment' and 'bacterial species' x 'environment'.

Differences in the growth rates in the di-associations of *Curvibacter* and

Duganella in the different environments and dependence on initial frequency were

assessed using a Generalized linear model (error structure: normal; link function:

identity) and post hoc contrasts. For each bacterial species, a separate model was

calculated with the response variables being either 'growth rate *Curvibacter*' or 'growth

rate *Duganella*', and the explanatory variables were 'environment', 'starting density' and

'environment' x 'starting density'.

Differences in the selection rate constant of *Curvibacter* in the different

environments and dependent on initial frequency were assessed using a Generalized

linear model (error structure: normal; link function: identity) and post hoc contrasts.

The response variable was 'selection rate constant *Curvibacter*' and the explanatory

variables were 'environment', 'starting density' and 'environment' x 'starting density'.

Sample size was chosen to maximise statistical power and ensure sufficient

replication. Assumptions of the tests, that is, normality and equal distribution of

variances, were visually evaluated. Non-significant interactions were removed from the

models. All tests were two-tailed. Effects were considered significant at the level of $P <$

0.05. All statistical analyses were performed with JMP 9. Graphs were produced with

GraphPad Prism 5.0, and Adobe Illustrator CS5.1.

Acknowledgements

PD received funding from the European Union's Framework Programme for Research and Innovation Horizon 2020 (2014–2020) under the Marie Skłodowska-Curie Grant Agreement No. 655914 and KH under the Marie Skłodowska-Curie Grant Agreement No. 657096. Both also received a Reintegration Grant from the Deutscher Akademischer Austausch Dienst (DAAD). This work was further supported by the Deutsche Forschungsgemeinschaft (DFG) Collaborative Research Center (CRC) 1182 ("Origin and Function of Metaorganisms"). TB gratefully appreciates support from the Canadian Institute for Advanced Research (CIFAR) and thanks the Wissenschaftskolleg (Institute of Advanced Studies) in Berlin for a sabbatical leave. We thank Chaitanya Gokhale for feedback on the manuscript.

Conflict of Interests

The authors declare no conflict of interest.

References

1. **Turnbaugh PJ**, **Ley RE**, **Hamady M**, **Fraser-Liggett CM**, **Knight R**, **Gordon JI**. 2007. The human microbiome project. Nature **449**:804–810.

2. **Huttenhower C**, **Gevers D**, **Knight R**, **Abubucker S**, **Badger JH**, **Chinwalla AT**, *et al.*. 2012. Structure, function and diversity of the healthy human microbiome. Nature **486**:207–214.

3. **Pasolli E**, **Asnicar F**, **Manara S**, **Zolfo M**, **Karcher N**, **Armanini F**, *et al.*. 2019. Extensive unexplored human microbiome diversity revealed by over 150,000 genomes from metagenomes spanning age, geography, and lifestyle. Cell **176**:649–662.e20.

4. **Mueller UG**, **Sachs JL**. 2015. Engineering microbiomes to improve plant and animal health. Trends Microbiol **23**:606–617.

5. **McKenney PT**, **Pamer EG**. 2015. From hype to hope: The gut microbiota in enteric infectious disease. Cell **163**:1326–1332.

6. **Gilbert JA**, **Blaser MJ**, **Caporaso JG**, **Jansson JK**, **Lynch SV**, **Knight R**. 2018. Current understanding of the human microbiome. Nat Med **24**:392–400.

7. **McKenzie VJ**, **Kueneman JG**, **Harris RN**. 2018. Probiotics as a tool for disease mitigation in wildlife: insights from food production and medicine. Ann N Y Acad Sci **287**:443.

8. **Contijoch EJ**, **Britton GJ**, **Yang C**, **Mogno I**, **Li Z**, **Ng R**, *et al.*. 2019. Gut microbiota density influences host physiology and is shaped by host and microbial factors. eLife **8**:337.

9. **Lozupone CA**, **Stombaugh JI**, **Gordon JI**, **Jansson JK**, **Knight R**. 2012. Diversity, stability and resilience of the human gut microbiota. Nature **489**:220–230.

10. **Sachs JL**, **Essenberg CJ**, **Turcotte MM**. 2011. New paradigms for the evolution of beneficial infections. Trends Ecol Evol **26**:202–209.

11. **Deines P**, **Lachnit T**, **Bosch TCG**. 2017. Competing forces maintain the Hydra metaorganism. Immunol Rev **279**:123–136.

12. **Macke E**, **Tasiemski A**, **Massol F**, **Callens M**, **Decaestecker E**. 2017. Life history and eco-evolutionary dynamics in light of the gut microbiota. Oikos **126**:508–531.

13. **Eckburg PB**, **Bik EM**, **Bernstein CN**, **Purdom E**, **Dethlefsen Les**, **Sargent M**, **Gill SR**, **Nelson KE**, **Relman DA**. 2005. Diversity of the human intestinal microbial flora. Science **308**:1635–1638.

14. **Zhang Z**, **Geng J**, **Tang X**, **Fan H**, **Xu J**, **Wen X**, *et al.*. 2014. Spatial heterogeneity and co-occurrence patterns of human mucosal-associated intestinal microbiota. ISME J **8**:881–893.

15. **Britton RA**, **Young VB**. 2012. Interaction between the intestinal microbiota and host in Clostridium difficile colonization resistance. Trends Microbiol

20:313–319.

16.   **Levy R**, **Borenstein E**. 2014. Metagenomic systems biology and metabolic
modeling of the human microbiome. Gut Microbes **5**:265–270.

17.   **Zelezniak A**, **Andrejev S**, **Ponomarova O**, **Mende DR**, **Bork P**, **Patil KR**.
2015. Metabolic dependencies drive species co-occurrence in diverse
microbial communities. Proc Natl Acad Sci USA **112**:6449–6454.

18.   **Garcia JR**, **Gerardo NM**. 2014. The symbiont side of symbiosis: do
microbes really benefit? Front Microbiol **5**:1103.

19.   **Wollenberg MS**, **Ruby EG**. 2012. Phylogeny and fitness of *Vibrio fischeri*
from the light organs of *Euprymna scolopes* in two Oahu, Hawaii
populations. ISME J **6**:352–362.

20.   **Wiser MJ**, **Lenski RE**. 2015. A comparison of methods to measure fitness in
Escherichia coli. PLoS ONE **10**:e0126210.

21.   **Lenski RE**. 1992. Relative fitness: its estimation and its significance for
environmental applications of microorganisms, pp. 183–198. *In* Levin, M,
Seidler, R, Rogul, M (eds.), Microbial Ecology: Principles, Applications and
Methods. myxo.css.msu.edu, New York.

22.   **Mushegian AA**, **Ebert D**. 2015. Rethinking "mutualism" in diverse host-
symbiont communities. Bioessays **38**:100–108.

23.   **Wiles TJ**, **Jemielita M**, **Baker RP**, **Schlomann BH**, **Logan SL**, **Ganz J**, *et al.*.
2016. Host gut motility promotes competitive exclusion within a model
intestinal microbiota. PLoS Biol **14**:e1002517.

24.   **Deines P**, **Bosch TCG**. 2016. Transitioning from microbiome composition
to microbial community interactions: The potential of the metaorganism
Hydra as an experimental model. Front Microbiol **7**:1610.

25.   **Fraune S**, **Bosch TCG**. 2007. Long-term maintenance of species-specific
bacterial microbiota in the basal metazoan *Hydra*. Proc Natl Acad Sci USA.

26.   **Bosch TCG**. 2013. Cnidarian-microbe interactions and the origin of innate
immunity in metazoans. Annu Rev Microbiol **67**:499–518.

27.   **Fraune S**, **Anton-Erxleben F**, **Augustin R**, **Franzenburg S**, **Knop M**,
Schröder K, *et al.*. 2015. Bacteria-bacteria interactions within the
microbiota of the ancestral metazoan *Hydra* contribute to fungal resistance.
ISME J **9**:1543–1556.

28.   **Lenski RE**, **Rose MR**, **Simpson SC**, **Tadler SC**. 1991. Long-term
experimental evolution in *Escherichia coli*. I. Adaptation and divergence
during 2,000 generations. Am Nat **138**:1315–1341.

29.   **Travisano M**, **Lenski RE**. 1996. Long-term experimental evolution in
Escherichia coli. IV. Targets of selection and the specificity of adaptation.
Genetics **143**:15–26.

30.   **Sonnenburg JL**, **Angenent LT**, **Gordon JI**. 2004. Getting a grip on things:

how do communities of bacterial symbionts become established in our
intestine? Nat Immunol **5**:569–573.

31. **Shafquat A**, **Joice R**, **Simmons SL**, **Huttenhower C**. 2014. Functional and
phylogenetic assembly of microbial communities in the human
microbiome. Trends Microbiol **22**:261–266.

32. **Allison SD**, **Martiny JBH**. 2008. Resistance, resilience, and redundancy in
microbial communities. Proc Natl Acad Sci USA **105**:11512–11519.

33. **Faust K**, **Raes J**. 2012. Microbial interactions: from networks to models. Nat
Rev Microbiol **10**:538–550.

34. **Trosvik P**, **de Muinck EJ**, **Stenseth NC**. 2015. Biotic interactions and
temporal dynamics of the human gastrointestinal microbiota. ISME J
9:533–541.

35. **Friedman J**, **Higgins LM**, **Gore J**. 2017. Community structure follows
simple assembly rules in microbial microcosms. Nat Ecol Evol **1**:109.

36. **Higgins LM**, **Friedman J**, **Shen H**, **Gore J**. 2017. Co-occurring soil bacteria
exhibit a robust competitive hierarchy and lack of non-transitive
interactions. bioRxiv 175737.

37. **Jemielita M**, **Taormina MJ**, **Burns AR**, **Hampton JS**, **Rolig AS**, **Guillemin
K**, ***et al.***. 2014. Spatial and temporal features of the growth of a bacterial
species colonizing the zebrafish gut. mBio, **5**:e01751-14.

38. **Pais IS**, **Valente RS**, **Sporniak M**, **Teixeira L**. 2018. *Drosophila
melanogaster* establishes a species-specific mutualistic interaction with
stable gut-colonizing bacteria. PLoS Biol **16**:e2005710.

39. **Wein T**, **Dagan T**, **Fraune S**, **Bosch TCG**, **Reusch TBH**, **Hülter NF**. 2018.
Carrying capacity and colonization dynamics of *Curvibacter* in the *Hydra*
host habitat. Front Microbiol **9**:185.

40. **Fukami T**. 2015. Historical contingency in community assembly:
Integrating niches, species pools, and priority effects. Annu Rev Ecol Evol
Syst **46**: 1-23.

41. **Hammarlund SP**, **Chacón JM**, **Harcombe WR**. 2019. A shared limiting
resource leads to competitive exclusion in a cross-feeding system. Environ
Microbiol **21**:759–771.

42. **Wright ES**, **Vetsigian KH**. 2016. Inhibitory interactions promote frequent
bistability among competing bacteria. Nat Commun **7**:11274.

43. **Li X-Y**, **Pietschke C**, **Fraune S**, **Altrock PM**, **Bosch TCG**, **Traulsen A**. 2015.
Which games are growing bacterial populations playing? J R Soc Interface
12:20150121–10.

44. **Ghoul M**, **Mitri S**. 2016. The ecology and evolution of microbial
competition. Trends Microbiol **24**:833–845.

45. **Franzenburg S**, **Walter J**, **Künzel S**, **Wang J**, **Baines JF**, **Bosch TCG**, ***et al.***.

2013. Distinct antimicrobial peptide expression determines host species-
specific bacterial associations. Proc Natl Acad Sci USA **110**:E3730–8.

46.   **Armstrong RA**, **McGehee R**. 2015. Competitive exclusion. Am Nat
115:151–170.

47.   **Sousa WP**. 1984. The role of disturbance in natural communities. Annu Rev
Ecol Syst 353–391.

48.   **Huisman J**, **Weissing FJ**. 1999. Biodiversity of plankton by species
oscillations and chaos. Nature **402**:407–410.

49.   **Franzenburg S**, **Fraune S**, **Altrock PM**, **Künzel S**, **Baines JF**, **Traulsen A**,
et al.. 2013. Bacterial colonization of Hydra hatchlings follows a robust
temporal pattern. ISME J **7**:781–790.

50.   **Murillo-Rincon AP**, **Klimovich A**, **Pemöller E**, **Taubenheim J**, **Mortzfeld**
B, **Augustin R**, *et al.*. 2017. Spontaneous body contractions are modulated
by the microbiome of *Hydra*. Sci Rep **7**:15937.

51.   **Rolig AS**, **Parthasarathy R**, **Burns AR**, **Bohannan BJM**, **Guillemin K**. 2015.
Individual members of the microbiota disproportionately modulate host
innate immune responses. Cell Host Microbe **18**:613–620.

52.   **Arumugam M**, **Raes J**, **Pelletier E**, **Le Paslier D**, **Yamada T**, **Mende DR**, *et*
al.. 2011. Enterotypes of the human gut microbiome. Nature **473**:174–180.

53.   **Hajishengallis G**, **Darveau RP**, **Curtis MA**. 2012. The keystone-pathogen
hypothesis. Nat Rev Microbiol **10**:717–725.

54.   **Sokol H**, **Pigneur B**, **Watterlot L**, **Lakhdari O**, **Bermúdez-Humarán LG**,
Gratadoux J-J, *et al.*. 2008. F*aecalibacterium prausnitzii* is an anti-
inflammatory commensal bacterium identified by gut microbiota analysis
of Crohn disease patients. Proc Natl Acad Sci USA **105**:16731–16736.

55.   **Lenhoff HM**, **Brown RD**. 1970. Mass culture of *Hydra*: an improved method
and its application to other aquatic invertebrates. Lab Anim **4**:139–154.

Figures

Fig 1. (A) Carrying capacity of the *Hydra* habitat in wild-type (WT) polyps, wild-type

polyps and the addition of either the focal species *Curvibacter* (*Curv*) or *Duganella*

(*Duga*) and germ-free (GF) animals incubated with native *Hydra* microbiota

(conventionalised polyps) (each boxplot N≥6≤16). (B) Time course analysis of microbial

abundances in mono-associations of germ-free polyps with either *Curvibacter* (red) or

Duganella (blue) (each boxplot N=6). The dashed line indicates the carrying capacity of

wild-type polys.

Fig 2. Bacterial growth rates of *Curvibacter* (red; each boxplot N≥2≤4) and *Duganella*

(blue; each boxplot N≥3≤6) in mono-associations are habitat dependent. Compared are

the host habitat (*in vivo*), and two *in vitro* environments: heterogeneous (static

microcosms), and homogenous (mixed microcosms).

Fig 3. Carrying capacity of the *in vivo* and *in vitro* habitats used in this study. Shown are

pooled total numbers of colony forming units (CFUs) from all di-association

experiments with *Curvibacter* (red) and *Duganella* (blue) (shown are s.e.m. based on

N≥11≤18 for the host, N≥9≤12 for static, and N≥4≤12 for mixed). The dashed black line

indicates the carrying capacity of WT polyps, the light grey line the carrying capacity of

polyps during *Curvibacter* mono-associations and the dark grey line the carrying

capacity of polyps during *Duganella* mono-associations.

Fig 4. Time course of fractions of *Curvibacter* (red) and *Duganella* (blue) in the three

different habitats obtained from di-association experiments. The initial inoculation

frequency of *Curvibacter* varied from being rare (R), to equal (E), to dominant (D) in

comparison to *Duganella* (each bar N≥3≤6; except for: Host, D, 24 h, where N=2, and for

Static, R and E 60 h, where N=0 due to contamination of plates).

Fig 5. Bacterial growth rates of *Curvibacter* (red) and *Duganella* (blue) from mono- and

di-association experiments across the different habitats and initial frequencies tested

(each boxplot N≥3≤4).

Fig. 6. Selection rate constants of *Curvibacter* based on competition experiments with *Duganella* for the different starting frequencies, (each boxplot N≥3≤6).

Fig. 7. Model of the microbe-microbe and host-microbe interactions occurring in

different habitats (host (A) and non-host (B and C)) that influence bacterial abundances,

which are indicated by the number of bacterial cells (*Curvibacter*: red, *Duganella*: blue).

Whereas (A) allows to study the effect of the host environment on the two microbial

species and their interactions, microcosm experiments allow to study species

interaction in a heterogeneous (B) and a homogeneous environment (C). Arrow-headed

lines indicate promotion, whereas bar-headed lines indicate inhibition.