Alternative pathway androgen biosynthesis and human fetal female virilization

Nicole Reisch[1,2*], Angela E. Taylor[1*], Edson F. Nogueira[1*], Daniel J. Asby[3*], Vivek Dhir[1],

Andrew Berry[3], Nils Krone[1,4], Richard J. Auchus[5], Cedric H.L. Shackleton[1,6], Neil A. Hanley[3,7,#],

and Wiebke Arlt[1,8,#]

* Contributed equally to this work

joint senior authors

[1] Institute of Metabolism and Systems Research, College of Medical and Dental Sciences,

University of Birmingham, Birmingham B15 2TT, United Kingdom

[2] Department of Endocrinology, Division of Internal Medicine IV, Ludwig Maximilian University

of Munich, 80336 Munich, Germany

[3] Division of Diabetes, Endocrinology & Gastroenterology, Faculty of Biology, Medicine &

Health, Manchester Academic Health Science Centre, University of Manchester, Oxford Rd,

Manchester M13 9PT, United Kingdom

[4] Department of Oncology & Metabolism, University of Sheffield, Sheffield S10 2TH, United

Kingdom

[5] Division of Metabolism, Endocrinology and Diabetes, Department of Internal Medicine,

University of Michigan, Ann Arbor, MI 48019, United States

[6] University of California at San Francisco (UCSF) Benioff Children's Hospital, Oakland, CA

94609, United States

[7] Research & Innovation, Manchester University NHS Foundation Trust, Oxford Rd, Manchester,

UK

[8] Centre of Endocrinology, Diabetes and Metabolism, Birmingham Health Partners, Birmingham

B15 2GW, United Kingdom

Abbreviated title: Human fetal androgen biosynthesis

Brief summary (25 words):

An alternative androgen synthesis pathway is active during the major period of human fetal sexual differentiation and drives female genital virilization in congenital adrenal hyperplasia.

Key words: 5α-dihydrotestosterone, 46,XX DSD, congenital adrenal hyperplasia, androgen receptor, steroid mass spectrometry

Word count (including title page, figure legends and references): 3,954

Number of figures: 4   **Number of tables:**   0

Supplemental data: 3 figures, 2 tables

Address correspondence to:

Wiebke Arlt, Institute of Metabolism and Systems Research (IMSR), Medical School IBR Tower, Rm 236, University of Birmingham, Birmingham, B15 2TT, United Kingdom

Phone: 44 121 415 8716; E-mail w.arlt@bham.ac.uk

Conflict of interest statement:

All authors have declared that no conflict of interest exists.

Role of the funding source:

The funders of the study had no role in study design, data collection, data analysis, data interpretation or writing of the manuscript. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

ABSTRACT

Androgen biosynthesis proceeds through the adrenal sex steroid precursor dehydroepiandrosterone, which is converted towards testosterone in the gonads, followed by further activation to 5α-dihydrotestosterone in genital skin, thereby facilitating male external genital differentiation. Congenital adrenal hyperplasia due to P450 oxidoreductase deficiency results in disrupted dehydroepiandrosterone biosynthesis, which explains undervirilization in affected boys. However, many affected girls are born severely virilized, despite low circulating androgens. We hypothesized that this is due to a prenatally active androgen pathway from 17α-hydroxyprogesterone to 5α-dihydrotestosterone, which bypasses dehydroepiandrosterone and testosterone, with increased activity in congenital adrenal hyperplasia. Here we employ explant cultures of human fetal organs (adrenal, gonads, genital skin) from the major period of sexual differentiation (6-10 weeks post-conception) and show that alternative pathway androgen biosynthesis is active in the fetus, and that female external genitalia respond sensitively to the androgens generated. Moreover, using steroid metabolite excretion analysis by mass spectrometry, we show that neonates with P450 oxidoreductase deficiency produce androgens through the alternative pathway during the first weeks of life. These results indicate a key role of alternative pathway androgen biosynthesis in the prenatal virilization of girls affected by congenital adrenal hyperplasia.

INTRODUCTION

Gonadal development depends on chromosomal sex, whereby the 46,XY or 46,XX karyotype, established at fertilization, dictates subsequent development of either testis or ovary (1-3). Gonadal hormones then direct differentiation of either male or female genitalia. In humans, sexual differentiation is established at 7-12 weeks *post conception* (wpc) (4).

While secretion of testosterone by fetal testis Leydig cells is thought sufficient to drive virilization of the internal genitalia in the male fetus (5), differentiation of the external genitalia requires the action of 5α-dihydrotestosterone (DHT), which is generated locally from circulating testosterone by the enzyme steroid 5α-reductase type 2 (SRD5A2) (6, 7). By contrast, differentiation of human female genitalia has been regarded as the default of a low androgen environment.

In humans, the regulation of sexual differentiation is intricately linked to early development of the adrenal cortex (4, 8). Disorders affecting adrenal steroidogenesis commonly affect sexual differentiation, as exemplified by the multiple variants of congenital adrenal hyperplasia (CAH), which result either in inappropriate or disrupted androgen biosynthesis. This consequently causes Disorder of Sex Development (DSD), which can manifest with external genital virilization in newborn girls (46,XX DSD), or undermasculinization of external genitalia in male neonates (46,XY DSD) (9). The most common variant of CAH, 21-hydroxylase (CYP21A2) deficiency, manifests with 46,XX DSD, while 17α-hydroxylase/17,20-lyase (CYP17A1) deficiency causes 46,XY DSD.

The congenital adrenal hyperplasia variant cytochrome P450 oxidoreductase (POR) deficiency can manifest with both 46,XY and 46,XX DSD (10-12). POR plays a pivotal role as the obligatory electron donor to all microsomal cytochrome P450 enzymes, including CYP21A2 and CYP17A1, the latter catalyzing the biosynthesis of dehydroepiandrosterone (DHEA), the major precursor for testosterone biosynthesis. Consequently, POR deficiency (PORD) results in

86   low circulating androgen concentrations, which readily accounts for 46,XY DSD, but fails to

87   account for the severe virilization of external genitalia observed in affected 46,XX neonates.

88   An explanation for this striking and seemingly contradictory genital phenotype paradox

in PORD has been lacking. We hypothesized that this apparent paradox could be explained by the

existence of an alternative pathway to androgen production that generates DHT from 17α-

hydroxyprogesterone (17OHP) during human fetal sexual differentiation, thereby bypassing the

classic androgen biosynthesic pathway via DHEA and testosterone (11). Elements of this

pathway have been characterized in the fetal gonad of the tammar wallaby pouch young (13-15)

and fetal opossum urogenital tract (16, 17). 17OHP accumulates in PORD but also in the most

common CAH variant, 21-hydroxylase deficiency, and thus could feed into the proposed

alternative pathway, if present in the fetus. Indirect biochemical evidence has indicated that the

proposed alternative pathway is active postnatally in individuals with CAH due to 21-hydroxylase

deficiency (18, 19) and may explain maternal virlization observed in pregnancies affected by

PORD (20, 21). However, direct delineation of the putative alternative pathway during human

fetal development, and in particular during the major period of sexual differentiation, has been

lacking.

102   Here, we present conclusive evidence for the presence and activity of the alternative

pathway in the human fetus, producing potent androgens during the major period of sex

differentiation. We show that human fetal female external genitalia respond sensitively to

androgens during the same period. In concert, these findings define an alternative pathway for

androgen biosynthesis during the critical period of sexual differentiation in the human fetus that

represents a major mechanism to explain the prenatal virilization of female infants affected by

CAH.

RESULTS AND DISCUSSION

Androgen biosynthesis in the human fetus during sexual differentiation

To ascertain the presence and activity of the hypothesized alternative androgen pathway, we performed incubations with male and female fetal adrenals, gonads and genital skin, which were collected at 6-10 wpc during the major period of human sex differentiation as previously described (8). We separately added deuterated steroid substrates for each step of the alternative pathway and employed liquid chromatography-tandem mass spectrometry (LC-MS/MS) to quantify the resulting products (**Supplemental Figure 1** and **Supplemental Tables 1 and 2**). Experiments were conducted at least in triplicate for each organ of either sex (**Supplemental Figures 2-3**).

Explant incubations with female tissue showed that some 17OHP entered the classic androgen biosynthesis pathway yielding androstenedione in adrenal, ovary and genital skin. However, testosterone was detected in only 2 of 9 female genital skin incubations, but not in female adrenals and gonads (**Figure 1A**). By contrast, all three female tissues showed conversion of 17OHP and subsequent intermediates through all steps of the proposed alternative androgen pathway, with the end product, DHT, produced from 5α-androstanediol in genital skin (3/3), ovaries (2/3) and adrenal (1/4) explant cultures (**Figure 1A**).

Male fetal tissues readily converted 17OHP along the classic androgen pathway to testosterone and and, in the alternative pathway, to 5α-17-hydroxypregnanolone (5α-17HP). In male genital skin, conversion from 17OHP proceeded until the generation of DHT (**Figure 1B**), feasibly arising from either route, as DHT represents the end product of both classic and alternative androgen pathways. However, incubations with the intermediate substrates of the alternative pathway demonstrated stepwise catalysis to DHT by male adrenal, testis and genital skin (**Figure 1B**). Consistent with studies in non-human species (22, 23), we did not identify 5α-

pregnane-17α-ol-3,20-dione as a significant intermediate of the two-step conversion of 17OHP to

5α-17HP in tissues from either sex, likely due to its immediate onward conversion.

Taken together, these data show that adrenal, gonad and genital skin are capable of

androgen biosynthesis via both classic and alternative pathway in both sexes. Our data suggest

the presence of an adreno-genital steroidogenic unit that cooperates to produce DHT during the

period of sexual differentiation in both male and female fetuses.

Steroidogenic enzyme expression during human fetal sexual differentiation

The alternative pathway conversion of 17OHP to 5α-17HP requires sequential 5α-

reductase and 3α-hydroxysteroid dehydrogenase activities.

Of the two isoforms, the first step is expected to require catalysis by steroid 5α-reductase

type 1 (SRD5A1), since SRD5A2 does not to convert 17OHP efficiently (24). While a study of

fetal tissues from 12-20 wpc did not detect SRD5A1 in adrenals, gonads or genital skin (25), we

found mRNA expression of SRD5A1 in all tissues during the major period of human sexual

differentiation (6-10 wpc) (**Figure 2A**). The second step, the conversion of 5α-pregnane-17α-ol-

3,20-dione to 5α-17HP requires 3α-hydroxysteroid dehydrogenase activity, in keeping with the

observed expression of *AKR1C1* and *AKR1C3*, both of which encode enzymes capable of

catalyzing this reaction (**Figure 2A**).

The next step in the alternative pathway requires CYP17A1 17,20 lyase activity; 5α-

17HP is the preferred substrate for this reaction and efficiently converted to 5α-androsterone (26).

CYP17A1 was robustly detected in human fetal adrenals, consistent with our previous work (8),

and also in the gonads of both sexes (**Figure 2B**). The subsequent reduction of 5α-androsterone to

5α-androstanediol requires 17β-hydroxysteroid dehydrogenase activity, which can be provided by

AKR1C3, which was expressed in all tissues investigated (**Figure 2A**).

The final step of the proposed pathway involves the conversion of 5α-androstanediol to

DHT, which requires 3β-epimerase (oxidative 3α-HSD) activity. Several enzymes have been

considered to catalyze this reaction (i.e. AKR1C2, RDH5, DHRS9, HSD17B10, HSD17B6 and

RDH16). However, only HSD17B6 and RDH16 are capable of efficient oxidation of 5α-

androstanediol to DHT, as previously demonstrated by transactivation of the androgen receptor

following cell-based overexpression (22, 27). We found expression of both HSD17B6 and

RDH16 in fetal genital skin of both sexes (**Figure 2C**).

In summary, we detected expression of all enzymes required to catalyze the alternative

androgen pathway (**Figure 2D**). Taken together with the steroid conversion studies, these data

comprehensively demonstrate that the normal adrenal, gonad and genital skin are capable of

androgen biosynthesis via both classic and alternative pathway in both sexes. Our data point to an

adreno-genital steroidogenic unit that can cooperates to produce DHT via the alternative pathway

during the major period of human sexual differentiation.

Androgen receptor in female genitalia from the start of sex differentiation

Having demonstrated the capacity for DHT production from steroidogenic precursors in

female fetuses, we corroborated its ability to function by examining the presence of the androgen

receptor (AR) in female external genitalia from the start of sex differentiation. Previously, AR

expression has been documented in four female human fetuses from 9-18 wpc (28). AR protein

was readily detected in stromal cells in the urethral folds of the external genitalia in both male

and female fetuses at the onset of sexual differentiation (**Figure 3A-C**). Its nuclear localization in

fixed tissues in both sexes implied AR was ligand-bound. To explore this further we studied the

intracellular localization of AR by immunofluorescence in female external genitalia fibroblasts

taken into primary culture from the same stage of development. In steroid-free media, the external

genitalia cells demonstrated cytoplasmic AR localization. As expected, the addition of 1 nM DHT

induced nuclear translocation of AR. Strikingly, the same translocation was observed when using

medium conditioned overnight from the corresponding female adrenal gland (**Figure 3D**). In

combination, these data show AR from the start of sex differentiation in both male and female

external genitalia and demonstrate a functional adreno-genital steroidogenic unit capable of

causing AR nuclear translocation.

Androgen biosynthesis in neonates with congenital adrenal hyperplasia

Having shown evidence of the alternative pathway during human sexual differentiation,

we next investigated whether we could demonstrate equivalent activity *in vivo*. To address our

hypothesis that excess alternative pathway androgen biosynthesis in fetal life explains female

virilization (46,XX DSD) in CAH due to PORD (12, 29), in whom the classic androgen pathway

is disrupted, we identified three patients with a mutation known to cause 46,XX DSD in affected

girls with normal male genitalia in affected boys (POR A287P) (11, 12). While postnatal

circulating androgens in PORD are low in infancy and beyond (12, 29), we hypothesized that

affected individuals would still show evidence of excess alternative pathway androgen

biosynthesis in the immediate neonatal period. We analysed urinary steroid metabolite excretion

by gas chromatography-mass spectrometry and detected significant alternative pathway activity

in the neonatal period, consistent with previous preliminary findings that argued for the adrenal

gland as a major site for alternative pathway activity (30). During the first three weeks after birth

we documented increased excretion of the 17OHP metabolite 17-HP compared to healthy

controls (**Figure 4A**), consistent with accumulation of 17OHP, the initial substrate of the

alternative androgen pathway. We found significantly increased excretion of 5α-17HP (**Figure**

4B), a key intermediate of the alternative pathway; and finally, there was significantly increased

excretion of the major DHT metabolite 5α-androsterone (**Figure 4C**). By contrast,

etiocholanolone excretion, derived from classic pathway androgen biosynthesis, did not differ

between the affected and healthy control groups (**Figure 4D**). These findings indicate increased

in vivo alternative pathway androgen biosynthesis in neonates with PORD, which ceases shortly

after birth.

Conclusions

We have provided ex vivo and in vivo evidence for the existence and activity of an

alternative pathway for the synthesis of the most potent androgen, DHT, during early human

development. Our data demonstrate that, through cooperation of an adreno-genital steroidogenic

unit, the alternative androgen pathway yields active androgen synthesis in the female fetus, with

excess activity driving female virilization, 46,XX DSD, in CAH.

METHODS

Detailed descriptions of experimental procedures are reported in the Supplemental Methods.

AUTHOR CONTRIBUTIONS

WA and NAH conceived and designed experiments; NR, AET, EFN, DJA, and CHLS

contributed to the study design. NR, EFN, DJA, VD and AB performed and analyzed the explant

culture experiments, supervised by WA and NAH. AET and NK performed and analyzed mass

spectrometry experiments, supervised by WA and CHLS. RJA provided reagents. NR wrote the

manuscript jointly with WA and NAH; all authors contributed to the discussion and

reviewed/edited the manuscript.

ACKNOWLEDGMENTS

This work was supported by the Medical Research Council UK (Program Grant 0900567,

to W.A.), the Gerald Kerkut Trust (PhD studentship to D.J.A.), the Wellcome Trust (Senior

Research Fellowship WT088566 to N.A.H.), the European Commission (Marie Curie Intra-

European Fellowship PIEF-GA-2008-221058, to N.R.), the Deutsche Forschungemeinschaft

(Heisenberg Senior Fellowship 325768017, to N.R.), and the Society for Endocrinology (Lab

visit award, to N.R.).

We thank Beverly A. Hughes, University of Birmingham, for technical support. We

thank our research nurses and study participants for tissue collection.

REFERENCES

1.　MacLaughlin DT, and Donahoe PK. Sex determination and differentiation. *N*
Engl J Med. 2004;350(4):367-78.

2.　Sekido R, and Lovell-Badge R. Sex determination involves synergistic action of
SRY and SF1 on a specific Sox9 enhancer. *Nature.* 2008;453(7197):930-4.

3.　Eggers S, Ohnesorg T, and Sinclair A. Genetic regulation of mammalian gonad
development. *Nat Rev Endocrinol.* 2014;10(11):673-83.

4.　Hanley NA, and Arlt W. The human fetal adrenal cortex and the window of
sexual differentiation. *Trends Endocrinol Metab.* 2006;17(10):391-7.

5.　Park SY, and Jameson JL. Minireview: transcriptional regulation of gonadal
development and differentiation. *Endocrinology.* 2005;146(3):1035-42.

6.　Leshin M, Griffin JE, and Wilson JD. Hereditary male pseudohermaphroditism
associated with an unstable form of 5 alpha-reductase. *J Clin Invest.*
1978;62(3):685-91.

7.　Wilson JD, Griffin JE, and Russell DW. Steroid 5 alpha-reductase 2 deficiency.
Endocr Rev. 1993;14(5):577-93.

8.　Goto M, Piper Hanley K, Marcos J, Wood PJ, Wright S, Postle AD, Cameron IT,
Mason JI, Wilson DI, and Hanley NA. In humans, early cortisol biosynthesis
provides a mechanism to safeguard female sexual development. *J Clin Invest.*
2006;116(4):953-60.

9.　Krone N, and Arlt W. Genetics of congenital adrenal hyperplasia. *Best Pract Res*
Clin Endocrinol Metab. 2009;23(2):181-92.

10.　Fluck CE, Tajima T, Pandey AV, Arlt W, Okuhara K, Verge CF, Jabs EW,
Mendonca BB, Fujieda K, and Miller WL. Mutant P450 oxidoreductase causes
disordered steroidogenesis with and without Antley-Bixler syndrome. *Nat Genet.*
2004;36(3):228-30.

11.　Arlt W, Walker EA, Draper N, Ivison HE, Ride JP, Hammer F, Chalder SM,
Borucka-Mankiewicz M, Hauffa BP, Malunowicz EM, et al. Congenital adrenal
hyperplasia caused by mutant P450 oxidoreductase and human androgen
synthesis: analytical study. *Lancet.* 2004;363(9427):2128-35.

12.　Krone N, Reisch N, Idkowiak J, Dhir V, Ivison HE, Hughes BA, Rose IT, O'Neil
DM, Vijzelaar R, Smith MJ, et al. Genotype-phenotype analysis in congenital
adrenal hyperplasia due to P450 oxidoreductase deficiency. *J Clin Endocrinol*
Metab. 2012;97(2):E257-67.

13.　Wilson JD, Auchus RJ, Leihy MW, Guryev OL, Estabrook RW, Osborn SM,
Shaw G, and Renfree MB. 5alpha-androstane-3alpha,17beta-diol is formed in
tammar wallaby pouch young testes by a pathway involving 5alpha-pregnane-
3alpha,17alpha-diol-20-one as a key intermediate. *Endocrinology.*
2003;144(2):575-80.

14.   Auchus RJ. The backdoor pathway to dihydrotestosterone. *Trends Endocrinol*
Metab. 2004;15(9):432-8.

15.   Shaw G, Fenelon J, Sichlau M, Auchus RJ, Wilson JD, and Renfree MB. Role of
the alternate pathway of dihydrotestosterone formation in virilization of the
Wolffian ducts of the tammar wallaby, Macropus eugenii. *Endocrinology.*
2006;147(5):2368-73.

16.   Wilson JD, Shaw G, Renfree MB, Auchus RJ, Leihy MW, and Eckery DC.
Ontogeny and pathway of formation of 5alpha-androstane-3alpha,17beta-diol in
the testes of the immature brushtail possum Trichosurus vulpecula. *Reprod Fertil*
Dev. 2005;17(6):603-9.

17.   Wilson JD, Renfree MB, Auchus RJ, Pask AJ, and Shaw G. Formation of 5alpha-
reduced androgens in the testes and urogenital tract of the grey short-tailed
opossum, Monodelphis domestica. *Reprod Fertil Dev.* 2009;21(5):649-54.

18.   Kamrath C, Hochberg Z, Hartmann MF, Remer T, and Wudy SA. Increased
activation of the alternative "backdoor" pathway in patients with 21-hydroxylase
deficiency: evidence from urinary steroid hormone analysis. *J Clin Endocrinol*
Metab. 2012;97(3):E367-75.

19.   Jones CM, Mallappa A, Reisch N, Nikolaou N, Krone N, Hughes BA, O'Neil
DM, Whitaker MJ, Tomlinson JW, Storbeck KH, et al. Modified-Release and
Conventional Glucocorticoids and Diurnal Androgen Excretion in Congenital
Adrenal Hyperplasia. *J Clin Endocrinol Metab.* 2017;102(6):1797-806.

20.   Shackleton C, Marcos J, Arlt W, and Hauffa BP. Prenatal diagnosis of P450
oxidoreductase deficiency (ORD): a disorder causing low pregnancy estriol,
maternal and fetal virilization, and the Antley-Bixler syndrome phenotype. *Am J*
Med Genet A. 2004;129A(2):105-12.

21.   Reisch N, Idkowiak J, Hughes BA, Ivison HE, Abdul-Rahman OA, Hendon LG,
Olney AH, Nielsen S, Harrison R, Blair EM, et al. Prenatal diagnosis of
congenital adrenal hyperplasia caused by P450 oxidoreductase deficiency. *J Clin*
Endocrinol Metab. 2013;98(3):E528-36.

22.   Bauman DR, Steckelbroeck S, Williams MV, Peehl DM, and Penning TM.
Identification of the major oxidative 3alpha-hydroxysteroid dehydrogenase in
human prostate that converts 5alpha-androstane-3alpha,17beta-diol to 5alpha-
dihydrotestosterone: a potential therapeutic target for androgen-dependent
disease. *Mol Endocrinol.* 2006;20(2):444-58.

23.   Tsujimura T, Mizutani S, and Matsumoto K. Pathway from progesterone to
5alpha-reduced c19 steroids not involving androstenedione and testosterone in
immature mouse testes in vitro. *Endocrinology.* 1975;96(2):515-8.

24.   Frederiksen DW, and Wilson JD. Partial characterization of the nuclear reduced
nicotinamide adenine dinucleotide phosphate: delta 4-3-ketosteroid 5 alpha-
oxidoreductase of rat prostate. *J Biol Chem.* 1971;246(8):2584-93.

25.   Thigpen AE, Silver RI, Guileyardo JM, Casey ML, McConnell JD, and Russell
312   DW. Tissue distribution and ontogeny of steroid 5 alpha-reductase isozyme
313   expression. *J Clin Invest.* 1993;92(2):903-10.
314   26.   Gupta MK, Guryev OL, and Auchus RJ. 5alpha-reduced C21 steroids are
315   substrates for human cytochrome P450c17. *Arch Biochem Biophys.*
316   2003;418(2):151-60.
317   27.   Bauman DR, Steckelbroeck S, Peehl DM, and Penning TM. Transcript profiling
318   of the androgen signal in normal prostate, benign prostatic hyperplasia, and
319   prostate cancer. *Endocrinology.* 2006;147(12):5806-16.
320   28.   Shapiro E, Huang HY, and Wu XR. Uroplakin and androgen receptor expression
321   in the human fetal genital tract: insights into the development of the vagina. *J
322   Urol.* 2000;164(3 Pt 2):1048-51.
323   29.   Huang N, Pandey AV, Agrawal V, Reardon W, Lapunzina PD, Mowat D, Jabs
324   EW, Van Vliet G, Sack J, Fluck CE, et al. Diversity and function of mutations in
325   p450 oxidoreductase in patients with Antley-Bixler syndrome and disordered
326   steroidogenesis. *Am J Hum Genet.* 2005;76(5):729-49.
327   30.   Homma K, Hasegawa T, Nagai T, Adachi M, Horikawa R, Fujiwara I, Tajima T,
328   Takeda R, Fukami M, and Ogata T. Urine steroid hormone profile analysis in
329   cytochrome P450 oxidoreductase deficiency: implication for the backdoor
330   pathway to dihydrotestosterone. *J Clin Endocrinol Metab.* 2006;91(7):2643-9.
331

Figure 1: Androgen biosynthesis via classic and alternative androgen pathways as observed in human fetal organ explant cultures from the major period of sexual differentiation (6-10 wpc). The steroid substrates added to the explant cultures derived from fetal adrenals, gonads and genital skin are shown in red and blue for fetal female (A) and male (B) tissues, respectively. Conversion products detected and identified by liquid chromatography-tandem mass spectrometry (LC-MS/MS) are given in black. f, female; m, male; n numbers indicate the number of biological replicates per tissue and fetal sex and how many of the cultures showed detectable synthesis of the indicated products.

Figure 2: **Steroidogenic enzyme expression in human fetal tissue from the major period of sexual differentiation (6-10 wpc) and their proposed role(s) in alternative pathway synthesis.**

(A+B) mRNA expression (mean+SEM) in human fetal tissues from the major period of sexual differentiation (6-10 wpc) as measured in by qPCR in female adrenal (n=1), gonad (n=2), genital skin (n=2) and corresponding male tissues (≥3 biological replicates for adrenal, gonads, and genital skin). Expression data were normalized to ribosomal 18S.

(C) Schematic summary of the proposed distinct roles of the identified enzymes in the classic androgen pathway (dark blue) and the alternative androgen synthesis pathway (light blue) both resulting in the synthesis of potent 5α-dihydrotestosterone. Arrows indicate observed conversions in the fetal organ explant cultures; dotted arrows represent reactions only rarely observed.

Figure 3: The androgen receptor is present in both male and female genitalia from the onset of sex differentiation. (A) Morphology of male and female external genitalia at the onset of sex differentiation. gt, genital tubercle; uf, urethral fold; lsf, labioscrotal fold; *, patency between urethral folds at 7-8 wpc that is partially sealed in males by 10-11 wpc. (B-C) Immunohistochemistry in trasverse sections through the phallus for AR in male (B) and female (C) external genitalia from the start of human sexual differentiation at 7-8 wpc counter-stained with toluidine blue. Boxes in the left-hand panel are shown at higher magnification to the right. (**D**) Immunofluorescence for AR in female external genitalia fibroblasts in the presence (+) or absence (-) of 10nM DHT (left) or medium conditioned by overnight incubation with an adrenal gland from the same female fetus (right). Size bars: (A) 500μm; (B)+(C) 100μm (low) and 20μm (high magnification); (D) 25μm.

Figure 4: Urinary steroid excretion in three male neonates with POR deficiency (closed symbols) in comparison to nine sex- and age-matched healthy controls (open symbols). POR deficient neonates harbored the A287P mutation (2 homozygous, 1 heterozygous) associated with normal male genitalia in boys and genital virilization (46,XX DSD) in girls. Longitudinal urine collections were carried out during the first three months of life and analyzed by gas chromatography-mass spectrometry. Depicted are the urinary excretion of (A) the 17OHP metabolite 17-hydroxypregnanolone (17HP) and the two alternative pathway intermediates (B) 5α-17-hydroxypregnanolone (5α-17HP) and (C) 5α-androsterone (An), in comparison to (D) etiocholanolone (Et), which is only generated via the classic androgen pathway. All steroids are shown relative to tetrahydrocortisone (THE) as the denominator; THE represents an abundant adrenal-derived steroid metabolite.