Extension of the *in vivo* haploid induction system from maize to wheat.

Chenxu Liu[1, 2#], Yu Zhong[1#], Xiaolong Qi[1#], Ming Chen[1], Zongkai Liu[1], Chen Chen[1], Xiaolong, Tian[1], Jinlong Li[1], Yanyan Jiao[1], Dong wang[1], Yuwen Wang[1], Mengran Li[1], Mingming Xin[1], Wenxin Liu[1], Weiwei Jin[1, 2], Shaojiang Chen[1, 2*]

[1]College of Agronomy and Biotechnology, China Agricultural University, No. 2, Yuan Ming Yuan West Road, Haidian District, Beijing, 100193, China.

[2]National Maize Improvement Center of China, Beijing Key Laboratory of Crop Genetic Improvement, Laboratory of Crop Heterosis and Utilization, Joint International Research Laboratory of Crop Molecular Breeding, China Agricultural University, Beijing 100193, China.

\# These authors contributed equally.

* Corresponding author: Shaojiang Chen (chen368@126.com)

Abstract

Doubled haploid breeding technology has been one of the most important techniques for
accelerating crop breeding. In compare to *in vivo* haploid induction in maize, which is efficient
and background independent, wheat haploid production by interspecific hybridization pollinated
with maize is influenced by genetic background and requires rescue of young embryos. Here, we
analyzed the homologues of maize haploid induction gene *MTL*/*ZmPLA1*/*NLD* in several crop
species systematically, the homologues are highly conserved in sorghum, millet and wheat etc.
Since wheat is a very important polyploidy crop, as a proof of concept, we demonstrated that the
in vivo haploid induction method could be extended from diploid maize to hexaploid wheat by
knocking out the wheat homologues (*TaPLAs*). Result showed that double knock-out mutation
could trigger wheat haploid induction at ~ 2%-3%, accompanied by 30% - 60% seed setting rate.
The performance of haploid wheat individual showed shorter plant, narrower leaves and male
sterile. Our results also revealed that knockout of *TaPLA*-A and *TaPLA*-D do not affect pollen
viability. This study not only confirmed the function of the induction gene and explored a new
approach for haploid production in wheat, but also provided an example that the *in vivo* haploid
induction could be applied in more crop species with different ploidy levels. Furthermore, by
combining with gene editing, it would be a fast and powerful platform for traits improvement in
polyploidy crops breeding.

Key words: Wheat, Doubled Haploid (DH), Inducer, Breeding, CRISPR/Cas9

Introduction

Doubled haploid (DH) technology substantially accelerates the breeding process for several crop species. Several different methods have been established for producing haploids in different crop species (Ishii et al., 2016), including interspecific hybridization between maize and wheat to produce wheat haploids (Laurie, 1988); *in vivo* haploid induction to produce maize haploids (E. H. Coe, 1959), and anther/male gametophyte culture method to produce haploids in rice and other crop species (Germanà, 2011). Of these, only *in vivo* haploid induction in maize has been demonstrated to be independent of genetic background and to produce haploids efficiently (> 10%), which make it possible to produce DH lines in an engineered way and laying a solid foundation for modern commercial breeding (Chen, 2013; Geiger, 2009). Recent studies of the genetics underlying *in vivo* haploid induction in maize revealed that several quantitative trait loci (QTL) contributed to haploid induction rate (HIR) (Prigge et al., 2012). Of these, *qhir1* is critical for haploid induction (Dong et al., 2013; Liu et al., 2015). Loss of function of the gene *MTL*/*ZmPLA1*/*NLD* in *qhir1* triggers haploid induction (Gilles et al., 2017; Kelliher et al., 2017; Liu et al., 2017a). Importantly, *in vivo* haploid induction system had been successfully applied to rice (Yao et al., 2018), making this method more important in plant breeding. Genetic analysis on the QTLs contributing HIR revealed that *qhir8* can improve haploid induction efficiency by 3- to 5- fold (Liu et al., 2015), this is a promising result for improving DH breeding efficiency in crop species other than maize. In addition to producing homozygous DH lines, *in vivo* haploid induction in maize has also been used for gene editing without introducing the genome of the male parent, which is meaningful for genome editing in different genetic backgrounds (Kelliher et al., 2019; Wang et al., 2019).

Compared with *in vivo* haploid induction in maize, wheat haploid production through interspecific hybridization requires the rescue of young embryos and is depend on genetic background (Niu et al., 2014), which limits its application. *In vivo* haploid induction system has been extended from maize to rice, while both are diploid crop, little is known whether it can be applied to polyploidy plants. Wheat is a very important hexaploidy crop, to extend maize *in vivo* haploid induction to wheat would create a novel protocol in speeding up wheat breeding and providing a potential method for gene editing in different genetic backgrounds (Kelliher et al.,

2019; Wang et al., 2019). Furthermore, it would also give an example for exploring the haploid
production and editing system in other polyploidy crops.

Results and Discussion

The full length amino acid sequence encoded by *MTL/ZmPLA1/NLD* was used to do a BLAST
search for homologues in different crop species (www.gramene.com), followed by amino acid
alignment and phylogenic analysis using MEGA (Knyaz et al., 2018). Results showed that the
gene is highly conserved among 19 species of Liliopsida (**Supplemental Figure 1**). Sorghum
and millet homologues had >80% amino acid sequence identity with maize, and wheat and rice
homologues were divided into two different branches (**Supplemental Figure. 1**), both which
sharing ~70%-80% amino acid sequence identity with maize. Wheat genome consists three
homologues: TraesCS4A02G018100 (*TaPLA*-A), TraesCS4B02G286000 (*TaPLA*-B) and
TraesCS4D02G284700 (*TaPLA*-D), located on chromosomes 4A, 4B and 4D, respectively. All
three shared a similar gene structure that includes four exons (**Figure1. A**). The amino acid
sequence identity between each of the three homologues and *MTL*/*ZmPLA1*/*NLD* is 75%,
representing a high level of conservation property. The amino acid sequences identity among the
three *TaPLA* genes is 96%.

Next, we designed a guide RNA sequence (**Supplemental Table 1**) targeting the first exon of all
three genes to create complete knockout lines using double-target CRISPR/Cas9 system
(**Figure1. A**) (Xing et al., 2014). After *Agrobacterium tumefaciens*-mediated transformation,
four transgenic events were obtained with mutations in both *TaPLA*-A and *TaPLA*-D. None of
them had a mutation in *TaPLA*-B in subgenome B. Of the four transgenic events obtained, two
transgenic lines, Da14-1 and ND52-6, were found to carry the same mutations: a 1bp guanine (G)
deletion in *TaPLA*-A and a GG deletion in *TaPLA*-D. Da14-2 had the same 1-bp deletion in
TaPLA-A as Da14-1 and ND52-6, but had a G to cytosine (C) mutation and a 1-bp G deletion in
the gene *TaPLA*-D (**Figure1. A**). ND52-2 had a 5-bp GGGAC deletion in *TaPLA*-A and a GG
deletion in *TaPLA*-D. Although different in sequence, all these mutations led to frame shifts and
loss of function for both *TaPLA*-A and *TaPLA*-D (**Figure1. A**). These transgenic were chosen for
subsequent analysis.

All T_0 transgenic and control plants were grown in the same greenhouse. No substantive

phenotypic difference was found between wildtype and transgenic plants (**Supplemental Table**

1), except for the seed setting rate (SSR), which ranged from ~30% to ~60% in transgenic lines,

significantly lower than the average value of 92.6% in wildtype (**Figure1. B**), implying that

TaPLAs may be involved in sexual reproduction. Haploid induction in transgenic lines was

evaluated in the self-pollination progenies of T_0 generation. Putative haploid wheat plants were

identified according to the haploid plant characteristic reported for maize and wheat (Liu et al.,

2017a; Zhang et al., 2014). Putative haploid plants were found in self-pollinated progenies of all

four transgenic lines (**Table 1**). In Da14-1 and Da14-2, two putative haploids were found among

their T_1 populations of 70 and 82 plants, respectively. In ND52-2, three putative haploids

occurred in a T_1 population of 83 plants. In ND52-6, four putative haloids were identified among

102 plants (**Figure1. C, Table 1**). Flow cytometry was used to determine the real ploidy of each

putative haploid. Compared with hexaploid controls which had an FL2A peak approximately 100,

all putative haploids had an FL2A approximately 50 (**Figure1. D**), suggesting that putative

haploids identified by phenotypic characteristic were true haploids. Compared with the hexaploid

wheat plants, these haploids had shorter plant height, narrower leaves, shorter ears and male

sterility (**Figure1. C, E**). No haploid plant was identified in a control group with 267 wildtype

individuals. Therefore, we conclude that knockout of wheat homologues of *MTL*/*ZmPLA1*/*NLD*

could trigger wheat haploids induction.

To characterize the expression pattern of the *TaPLA* genes, subcellular localization was

performed in tobacco epidermal cells infected by agrobacterium. As shown in **Supplemental**

Figure. 2, all three genes showed strong signals in the cytomembrane, that merged well with the

endoplasmic reticulum (ER) marker. This result is consistent with the cytomembrane subcellular

localization of *MTL*/*ZmPLA1*/*NLD* in maize (Gilles et al., 2017). Reciprocal crosses between the

maize haploid inducer and non-inducer lines previously revealed that the transmission of male

gametes from inducer lines is decreased significantly (Dong et al., 2013; Liu et al., 2017a),

suggesting that haploid inducer lines lead to either low pollen viability of pollen or a defect in

fusion between male and female gametes in double fertilization. Therefore, we performed

fluorescein diacetate (FDA) staining to examine pollen viability in wildtype and mutants. Pollen

from wildtype and mutant lines was divided into three classes: high-viability pollen, low-

viability pollen and dead pollen according to their pollen viability (**Figure2. A**). There was no

difference in the proportion of pollen viability classes between mutants and wildtype (**Figure2.**
B). This result demonstrates that loss of function of both *TaPLA*-A and *TaPLA*-D does not
influence pollen viability.

In summary, as a proof of concept, we have demonstrated that *in vivo* haploid induction can be
extended from diploid maize to wheat, by knocking out *TaPLA*-A and *TaPLA*-D. Previous
studies on *in vivo* haploid induction in crop species, used maize or rice, which are diploid. This
study provided the first proof that *in vivo* haploid induction is not limited to diploid crop species
but can be extended to polyploidy species, which substantially extended its potential.
Considering the potential redundancy among *TaPLA*-A, *TaPLA*-B and *TaPLA*-D, existence of
TaPLA-B may have functional complementation effect in *Tapla*-A and *Tapla*-B double mutant.
Nevertheless, the complementation effect is not sufficient to rescue the phenotype of haploid
induction and decreased SSR. On the other hand, further improvement in the efficiency of
haploid induction in wheat may be achieved by creating triple mutants.

The observed phenotype of *TaPLAs* knockout lines including a ~2-3% haploid frequency and
significantly decreased SSR, are similar to that of maize and rice (Gilles et al., 2017; Kelliher et
al., 2017; Liu et al., 2017a; Yao et al., 2018), suggesting that *MTL*/Zm*PLA1*/*NLD* homologues
are involved in the similar pathway during double-fertilization across different crop species.
Thus, our study provided an alternative crop species in exploring the biological mechanism of
haploid induction.

Several technical problems require further investigation before *in vivo* haploid induction can be
applied efficiently in wheat, including low efficiency in haploid induction and the difficulties in
wheat haploid identification in seed stage. Nonetheless, our study provides an alternative
approach other than interspecific hybridization for generating haploids in wheat. Through in-
depth study of the genetic basis and biological mechanisms of *in vivo* haploid induction in maize,
more and more genes contributing HIR will be identified (Liu et al., 2015; Prigge et al., 2012).
The efficiency of wheat haploid induction may be further improved by the knockout/over-
expression of these new identified homologous genes contributing HIR in maize. On the other
hand, recent studies enable the identification of haploids using enhanced green fluorescent
proteins (EGFP) and DsRED signals specifically expressed in the embryo and endosperm,

respectively (Dong et al., 2018). This method may provide a potential solution for haploid kernel

identification in wheat, which lacks phenotypic markers like *R1-nj* and high oil content existed in

maize (Dong et al., 2014).

Recent studies that utilized the combination of CRISPR/Cas9-mediated gene editing and haploid

induction revealed that *in vivo* haploid induction can enable gene editing in maternal haploids in

any background, without mixing the paternal genome (commonly referred to as "HI-edit" or

"IMGE") (Kelliher et al., 2019; Wang et al., 2019). This is promising for future genetic

improvement of crops. However, the "HI-edit using CRISPR/Cas9 and interspecific

hybridization has a lower editing efficiency in wheat than that in maize (Kelliher et al., 2019;

Wang et al., 2019). Because our study demonstrates the feasibility of *in vivo* haploid induction in

wheat, it also provides a putative platform for wheat haploid gene editing with CRISPR/Cas9.

Considering the conservation of gene sequence and the capacity for haploid induction among

maize, rice and wheat, this method could potentially be extended to a wider variety of crop

species.

Methods

Knockout of homologous genes in wheat

The full-length amino acid sequence of *ZmPLA1*/*MTL*/*NLD* was used to perform a BLAST

search for the homologous genes in crop species. Amino acid sequences of homologues were

downloaded (www.gramene.com) and used for cluster alignment followed by a phylogenic

analysis with the neighbor-joining method using the software MEGA X with default parameter

(Knyaz et al., 2018). A guide RNA sequence was then designed targeting the homologous gene

and then cloned into vector pBUEB411 (Liu et al., 2017b; Xing et al., 2014). The hexaploid

wheat line CB037 was used for transformation (Wang et al., 2017). Young embryos were

sampled and used for *A. tumefaciens*-mediated transformation. Positive calluses were screened

using bialaphos. Transgenic T_0 plants were transplanted in the green house and verified by

sequencing of the target site. Plants with insertion/deletion mutations that lead to frame shifts

and complete loss of function were chosen to produce the T_1 generation. Genotyping was

confirmed by sequencing in the T_1 plants by sequencing were performed again.

Haploid plant identification and verification

Putative haploids were screened in the T_1 generation. In maize, haploids have shorter plant height, narrower leaves, smaller anther and are sterile. Thus, wheat plants with the above-mentioned phenotypes were selected as putative haploids. Subsequently, ploidy of putative haploids was verified using flow cytometry (Liu et al., 2017a). Young leaves of both hexaploids (controls) and putative haploids were sampled and chopped with a razor blade in buffer (Schutte et al., 1985). Nuclei were extracted and stained with a fluorescent dye. The signal peak of standard hexaploids (controls) was set to 100, these putative haploids with signal peak at 50 were considered real haploids. HIR was calculated as follows: HIR (%) = (number of haploid plants/total number of plants derived from same individual) ×100%. For each ear, the SSR was calculated as follows: SSR (%) = (number of viable seeds/total number of embryo sacs on the ear) ×100%.

Pollen viability analysis

A fresh FDA staining buffer was made by mixing 100 μl FDA stock solution (0.5% in acetone) with 4.9 ml sucrose solution (Widholm, 1972). Fresh pollen (20 mg) was sampled from wildtype and mutant lines in the greenhouse and mixed immediately with the FDA working solution and left in dark environment for 1 h without covering to guarantee an abundant supply of oxygen for the esterase reaction. Three biological replicates were performed. Fluorescent was observed and photographed by a fluorescence microscope (Nikon, Ci-S-FL) with an excitation wavelength of 485 nm. Pollens that exhibited strong green fluorescence from the cytosol was considered to be highly viable pollen, pollen that exhibited weak fluorescence was considered as low viability pollen, and pollen with no detectable staining was considered dead.

Subcellular co-localization

RNA from mature pollen of wildtype CB037 was extracted using TRIzol reagent. Reverse transcription was performed using oligo (dT) primers to obtain full-length cDNA. The coding sequence of *TaPLA*-A, *TaPLA*-B and *TaPLA*-D without termination codons (TGA) were cloned into a vector 1305-EGFP driven by the 35S promotor. Sequencing-validated constructs of 35S::*TaPLA*-A-EGFP, 35S:: *TaPLA*-B-EGFP and 35S:: *TaPLA*-D-EGFP and ER marker were

206 used for *A. tumefaciens*–mediated transformation into tobacco epidermal cells. After culturing at

207 21 ℃ for 48 hours, EGFP signals were observed and imaged using a confocal laser-scanning

208 microscope (Zeiss LSM 710). As a control, empty 1305-EGFP vector was also transformed.

Author contributions

210 S.C. and C.L. conceive and designed the project. C.L., Y.Z. constructed plasmid. X.Q., Y.Z.,
211 M.C., Z.L., C.C., X.T., J.L., D.W., Y.W., M.L. and W.L. planted transgenic plants in greenhouse
212 and performed haploid identification and verification, phenotype investigations. Y.Z., C.L., X.Q.,
213 M.L., and W.L. performed data analysis. C.L., Y.Z., X.Q. M. X .and S.C. wrote the paper.

Acknowledgments

We thank Prof. Xingguo Ye of Institute of crop science, Chinese academy of agricultural
sciences, for his help in wheat transformation. We thank Prof. Pu Wang of College of Agronomy
and Biotechnology, China Agricultural University, for providing greenhouse. We thank Dr.
Qiguo Yu of Rutgers, The State University of New Jersey, for reading the manuscript. We thank
Prof. Jingrui Dai, Prof. Qixin Sun and Prof. Zhongfu Ni for their valuable suggestions. This
research was supported by the National Key Research and Development Program of China
(2016YFD0101200) the Modern Maize Industry Technology System (CARS–02–04) and China
Postdoctoral Science Foundation (2018M631634).

REFERENCES

Chen, S. (2013). Crop Breeding Engineering and Engineered Breeding. Crops 6:1-4.
Dong, L., Li, L., Liu, C., Liu, C., Geng, S., Li, X., Huang, C., Mao, L., Chen, S., and Xie, C. (2018). Genome
Editing and Double-Fluorescence Proteins Enable Robust Maternal Haploid Induction and
Identification in Maize. Molecular plant 11:1214-1217.
Dong, X., Xu, X., Li, L., Liu, C., Tian, X., Li, W., and Chen, S. (2014). Marker-assisted selection and
evaluation of high oil in vivo haploid inducers in maize. Molecular Breeding 34:1147-1158.
Dong, X., Xu, X., Miao, J., Li, L., Zhang, D., Mi, X., Liu, C., Tian, X., Melchinger, A.E., and Chen, S. (2013).
Fine mapping of qhir1 influencing in vivo haploid induction in maize. TAG. Theoretical and
applied genetics. Theoretische und angewandte Genetik.
E. H. Coe, J. (1959). A Line of Maize with High Haploid Frequency. 93:381-382.
Geiger, H.H., Gordillo, G. A. (2009). Doubled haploids in hybrid maize breeding. Maydica 54:485-499.
Germanà, M.A. (2011). Anther culture for haploid and doubled haploid production. 104:283-300.

Gilles, L.M., Khaled, A., Laffaire, J.B., Chaignon, S., Gendrot, G., Laplaige, J., Bergès, H., Beydon, G., Bayle,
V., Barret, P., et al. (2017). Loss of pollen-specific phospholipase NOT LIKE DAD triggers
gynogenesis in maize. The EMBO journal:e201796603.
Ishii, T., Karimi-Ashtiyani, R., and Houben, A. (2016). Haploidization via Chromosome Elimination: Means
and Mechanisms. Annual review of plant biology 67:421-438.
Kelliher, T., Starr, D., Richbourg, L., Chintamanani, S., Delzer, B., Nuccio, M.L., Green, J., Chen, Z.,
McCuiston, J., Wang, W., et al. (2017). MATRILINEAL, a sperm-specific phospholipase, triggers
maize haploid induction. Nature 542:105-109.
Kelliher, T., Starr, D., Su, X., Tang, G., Chen, Z., Carter, J., Wittich, P.E., Dong, S., Green, J., Burch, E., et al.
(2019). One-step genome editing of elite crop germplasm during haploid induction. Nature
biotechnology 37:287-292.
Knyaz, C., Stecher, G., Li, M., Kumar, S., and Tamura, K. (2018). MEGA X: Molecular Evolutionary Genetics
Analysis across Computing Platforms. Molecular Biology and Evolution 35:1547-1549.
Laurie, D.A., Bennett, M.D. (1988). The production of haploid wheat plants from wheat x maize crosses.
Theoretical and Applied Genetics 76:393-397.
Liu, C., Li, W., Zhong, Y., Dong, X., Hu, H., Tian, X., Wang, L., Chen, B., Chen, C., Melchinger, A.E., et al.
(2015). Fine mapping of qhir8 affecting in vivo haploid induction in maize. TAG. Theoretical and
applied genetics. Theoretische und angewandte Genetik 128:2507-2515.
Liu, C., Li, X., Meng, D., Zhong, Y., Chen, C., Dong, X., Xu, X., Chen, B., Li, W., Li, L., et al. (2017a). A 4bp
insertion at ZmPLA1 encoding a putative phospholipase A generates haploid induction in maize.
Molecular plant.
Liu, H., Ding, Y., Zhou, Y., Jin, W., Xie, K., and Chen, L.L. (2017b). CRISPR-P 2.0: An Improved CRISPR-Cas9
Tool for Genome Editing in Plants. Molecular plant 10:530-532.
Niu, Z., Jiang, A., Abu Hammad, W., Oladzadabbasabadi, A., Xu, S.S., Mergoum, M., and Elias, E.M. (2014).
Review of doubled haploid production in durum and common wheat through wheat × maize
hybridization. 133:313-320.
Prigge, V., Xu, X., Li, L., Babu, R., Chen, S., Atlin, G.N., and Melchinger, A.E. (2012). New insights into the
genetics of in vivo induction of maternal haploids, the backbone of doubled haploid technology
in maize. Genetics 190:781-793.
Wang, B., Zhu, L., Zhao, B., Zhao, Y., Xie, Y., Zheng, Z., Li, Y., Sun, J., and Wang, H. (2019). Development of
a Haploid-Inducer Mediated Genome Editing (IMGE) System for Accelerating Maize Breeding.
Molecular plant.
Wang, K., Liu, H., Du, L., and Ye, X. (2017). Generation of marker-free transgenic hexaploid wheat via an
Agrobacterium-mediated co-transformation strategy in commercial Chinese wheat varieties.
15:614-623.
Widholm, J.M. (1972). The Use of Fluorescein Diacetate and Phenosafranine for Determining Viability of
Cultured Plant Cells. Stain Technology 47:189-194.
Xing, H.-L., Dong, L., Wang, Z.-P., Zhang, H.-Y., Han, C.-Y., Liu, B., Wang, X.-C., and Chen, Q.-J.J.B.P.B.
(2014). A CRISPR/Cas9 toolkit for multiplex genome editing in plants. 14:327.
Yao, L., Zhang, Y., Liu, C., Liu, Y., Wang, Y., Liang, D., Liu, J., Sahoo, G., and Kelliher, T. (2018). OsMATL
mutation induces haploid seed formation in indica rice. Nat Plants 4:530-533.
Zhang, W., Wang, K., Lin, Z.-S., Du, L.-P., Ma, H.-L., Xiao, L.-L., and Ye, X.-G.J.B.S. (2014). Production and
identification of haploid dwarf male sterile wheat plants induced by corn inducer. 55:26.

LEGENDS

Figure 1. Knockout of wheat homologues of *MTL*/*ZmPLA1*/*NLD* and phenotypic analysis.

(A) Gene structure of wheat homologues *TaPLA*-A, *TaPLA*-B and *TaPLA*-D. Exons in red were target regions of guide RNAs. Shown below are knockout sequence results for of transgenic line Da14-1, Da14-2, ND52-2 and ND52-2 at *TaPLA*-A and *TaPLA*-D, respectively. Sequences in green are target sequences of *TaPLA*-A and *TaPLA*-D, the guide RNA is underlined. Black indicates the sequence of transgenic events. Deletions are shown by "-" on gray background.

(B) Ear performance of in wildtype (left) and knockout (right) plants. Statistical analysis for seed setting rate (SSR) of wildtype and four transgenic lines are shown on the right side. SSR was calculated as SSR (%) = (number of viable seeds/number of embryo sacs on an ear) ×100%. ** P < 0.01 calculated with the heteroscedastic two-tailed Student's *t*-test. Sample size of SSR for wildtype and each transgenic line was marked on the top of column.

(C) Phenotypic difference between haploid (H) and hexaploidy (HE) wheat, including whole plant, leaf, ear and anther. H, haploid; HE, Hexaploid. Bars = 5 cm, 1 cm, 1 cm and 1 mm for whole plant, flag leaf, ear and anther, respectively.

(D) Ploidy verification by flow cytometry for hexaploid and haploid plants.

(E) Statistical analysis of phenotypic differences including plant height, leaf length, leaf width and anther length between haploid and hexaploid wheat. Both stems (M) and tillers (T) of haploids and hexaploids were analyzed. A heteroscedastic two-tailed Student's *t*-test was performed for each class between wildtype and mutant. *, ** and *** P<0.05, 0.01 and 0.001, respectively.

Figure 2. Pollen viability assays with FDA staining.

(A) Comparison of pollen viability between wildtype and mutants in CB037 background (scale bar, 100μm). Pollen with a strong GFP (white triangle) and weak (yellow triangle) or no (red triangle) represented for high-viability pollen (HVP), low-viability pollen (LVP) and died pollen (DP), respectively.

(B) Ratio of the three viability classes in wildtype and mutant pollen. Statistics were conducted with three biological replicates. A heteroscedastic two-tailed Student's *t*-test was performed for each class between wildtype and mutant. NS, not significant (P≥0.05).

Table 1. Haploid induction potential of transgenic lines.

Haploid plants occurred in self-pollination progeny of transgenic lines, genotype on *TaPLA*-A and genotype *TaPLA*-D showed, H, heterozygous; M, homozygous mutant; W, wildtype. The control group comprised self-pollinated plants from the wildtype receptor background CB037.

Supplemental Figure 1. Phylogenetic analysis of *MTL/ZmPLA1/NLD* homologues among main crop species.

The green background indicates phylogenetic branch containing maize, sorghum and millet; blue background indicates wheat relative species; orange background indicates rice relative species. The scale bar indicates the proportion of the sites changing along each branch.

Supplemental Figure 2. Subcellular localization of *TaPLAs* in tobacco epidermal cells.

Confocal scanning (GFP; outside left), ER-mCherry signal (inside left), differential interference contrast (DIC; inside right), and merged (outside right) micrographs of tobacco epidermal cells transformed with a 35S::*TaPLA*-A-EGFP (row 1), 35S::*TaPLA*-B-EGFP (row 2), 35S::*TaPLA*-D-EGFP (row 3) and 35S::EGFP control (row 4). Scale bars = 100 µm.

Supplemental Table 1. Phenotypic differences between wildtype and mutant plants.

Both wildtype and double mutants are in CB037 background. PH, plant height; LW, leaf width of the flag leaf from stem; LL, leaf length of the flag leaf from stem; EL, stem ear length; NT, the number of tillers; NA, Not Applicable. Each phenotype was compared between wildtype and mutant using a heteroscedastic two-tailed Student's *t*-test.

Supplemental Table 2. Primers used in this study.

Figure. 1 knockout *TaPLA1*s leading to haploid induction in wheat.

Figure. 2

Supplemental Figure. 1

	EGFP	ER	DIC	Merged

35S::*TaPLA-A-EGFP*

35S::*TaPLA-B-EGFP*

35S::*TaPLA-D-EGFP*

35S::*EGFP*

Supplemental Figure. 2

Table. 1

Transgenic events&Control	Genotype *TaPLA*-A	Genotype *TaPLA*-D	Total plants	Haploid plants	HIR (%)
Da14-1	M	H	70	2	2.86
Da14-2	H	H	82	2	2.44
ND52-2	H	H	83	3	3.61
ND52-6	H	H	102	4	3.92
Control	W	W	267	0	0.00

Supplemental Table. 1 Phenotypic analysis between wildtype and knockout lines.

Genptype	Statistics	PH	PW	PL	EL	NT
wildtype	5	89.58	1.42	17.38	13.36	3.4
Mutant	10	84.08	1.52	18.22	13.42	2.8
T.test	NA	0.080	0.510	0.613	0.847	0.599

Supplemental Table. 2 Primers used in the study

Primer name	Left sequence (5'-3')	Functions
1305Ta4A+GFP-F	CGGAGCTAGCTCTAGAATGGCAAGCTACTGGTGCCG	
1305Ta4A+GFP-R	TGCTCACCATGGATCCGCGTCTGGATTTCAGACGGC	
1305Ta4B+GFP-F	CGGAGCTAGCTCTAGAATGATGGCAAGCTACTGGTG	Subcellular
1305Ta4B+GFP-R	TGCTCACCATGGATCCGCGGCTGGATTTCAGGCGGC	localizaiton
1305Ta4D+GFP-F	CGGAGCTAGCTCTAGAATGGCAAGCTACTGGGGCCG	
1305Ta4D+GFP-R	TGCTCACCATGGATCCGCGGCTGGATTTCAGGCGGC	
5`-A1	GAGGTGTCGTACGTCGGTTT	
3`-A1	AGCCACGCTAAGCTACAAGG	Full length gene
5`-D1	GGTGTCGTACGTTGGTTTGC	PCR
3`-D1	AGCTTTCTCTACCTATCCCAGC	
4AF	GTCAAGATCTCCAGCCGAGAC	
4AR	GGTACTTGCCGCTGTACCT	
4BF	AACTCAACATGGGGCGTCCTC	Transgenic plant
4BR	ACGTCGTATGTGGAGAAGATGATG	sequencing
4DF	TTCGGGTCCGGATTCTATTGTG	
4DR	GCAGGTACTTGCCGTTGTACC	
Target1	GCCCCTACATCTTCCCGCAAAGG	
Target2	GAAGCGGTTGACGTCCCTGGCGG	Crispr Cas9 target