Title

MLKS2 is an ARM domain and F-actin-associated KASH protein that functions in

stomatal complex development and meiotic chromosome segregation

Authors

Gumber, Hardeep K.[1]; McKenna, Joseph F.[2]; Tolmie, Andrea F.[2]; Jalovec, Alexis M.[1];

Kartick, Andre C.[1]; Graumann, Katja[2]; Bass, Hank W.[1]

Affiliations

1. Department of Biological Science, Florida State University, Tallahassee, FL, USA,

32306-4295.

2. Department of Biological and Medical Sciences, Faculty of Health and Life Sciences,

Oxford Brookes University, Oxford, UK, OX30BP.

Corresponding Author:

Hank W. Bass, bass@bio.fsu.edu

Abstract: The linker of nucleoskeleton and cytoskeleton (LINC) complex is an

essential multi-protein structure spanning the eukaryotic nuclear envelope. The LINC

complex functions to maintain nuclear architecture, positioning, and mobility, along with

specialized functions in meiotic prophase and chromosome segregation. Members of

the LINC complex were recently identified in maize, an important scientific and

agricultural grass species. Here we characterized *Maize LINC KASH AtSINE-like2*,

MLKS2, which encodes a highly conserved SINE-group plant KASH protein with

characteristic N-terminal armadillo repeats (ARM). Using a heterologous expression

system, we showed that actively expressed GFP-MLKS2 is targeted to the nuclear

periphery and colocalizes with F-actin and the endoplasmic reticulum, but not

microtubules in the cell cortex. Expression of GFP-MLKS2, but not GFP-MLKS2ΔARM,

resulted in nuclear anchoring. Genetic analysis of transposon-insertion mutations,

mlks2-1 and *mlks2-2*, showed that the mutant phenotypes were pleiotropic, affecting

root hair nuclear morphology, stomatal complex development, multiple aspects of

meiosis, and pollen viability. In male meiosis, the mutants showed defects for bouquet-

stage telomere clustering, nuclear repositioning, perinuclear actin accumulation,

dispersal of late prophase bivalents, and meiotic chromosome segregation. These

findings support a model in which the nucleus is connected to cytoskeletal F-actin

through the ARM-domain, predicted alpha solenoid structure of MLKS2. Functional

conservation of MLKS2 was demonstrated through genetic rescue of the misshapen

nuclear phenotype of an Arabidopsis (triple-*WIP*) KASH mutant. This study establishes

a role for the SINE-type KASH proteins in affecting the dynamic nuclear phenomena

required for normal plant growth and fertility.

Keywords

48   KASH, LINC, nuclear envelope, actin, FRAP, maize, MLKS2, meiosis, telomere,

49   bouquet

List of abbreviations

FRAP; Fluorescence recovery after photobleaching

DPI; Days post infiltration

OD; Optical density

MLKS2; Maize LINC KASH AtSINE-like2

LINC; Linker of nucleoskeleton and cytoskeleton

NE; Nuclear envelope

INM; Inner nuclear membrane

ONM; Outer nuclear membrane

Introduction

The cell nucleus is a eukaryotic organelle that houses, organizes and expresses the bi-parentally inherited genetic material. As a cellular compartment, the nucleus is separated from the cytoplasm by the multifunctional double-membraned nuclear envelope (NE). The NE has embedded macromolecular protein complexes, such as the nuclear pore complex (NPC) and the linker of nucleoskeleton and cytoskeleton (LINC) complex, which coordinate fundamental cell processes such as signaling and gene regulation (reviewed by Hetzer, 2010; Van de Vosse et al., 2011; D'Angelo, 2018).

The LINC complex was discovered relatively recently, but has been shown to be highly conserved in eukaryotes, with myriad functions as described in numerous reviews (Burke and Roux, 2009; Chang et al., 2015; Hieda, 2017; Kim et al., 2015; Luxton and Starr, 2014; Meier et al., 2017). These LINC-mediated functions include regulation of nuclear shape and position (Dittmer and Richards, 2008; Sakamoto and Takagi, 2013), mechanosensory signal transduction (Uzer et al., 2016), control of cell division, intranuclear architecture and gene regulation (Alam et al., 2016; Poulet et al., 2017b; Wang et al., 2018), and the specialized behavior of meiotic chromosomes to ensure segregation and fertility (Link et al., 2014; Murphy et al., 2014; Varas et al., 2015). The two core LINC components are the inner nuclear membrane (INM) SUN-domain proteins (Hagan and Yanagida, 1995; Malone et al., 1999) and outer nuclear membrane (ONM) KASH-domain proteins (Starr and Han, 2002). The SUN domain proteins are relatively well conserved across taxa and can be recognized by sequence homology and domain composition which includes transmembrane domains, the SUN domain, and a coiled-coil domain. In contrast, KASH proteins are functionally

conserved, with a C-terminal region that includes a transmembrane domain and

diagnostic terminal residues, but otherwise can be quite divergent with limited apparent

homology (Zhou and Meier, 2013; Evans et al., 2014; Kim et al., 2015; Meier, 2016;

Poulet et al., 2017a).

Structural studies reveal that SUN proteins form trimers that can bind to three

KASH proteins (Sosa et al., 2012). The cytoplasmic N-terminal regions of KASH domain

proteins interact directly or indirectly with a variety of cytoskeletal structures including

motor proteins, microfilaments, microtubules, or intermediate filaments (Starr and

Fridolfsson, 2010; Tamura et al., 2013; Luxton and Starr, 2014; Varas et al., 2015). The

forces generated by the cytoskeletal elements result in LINC-dependent movements

known or thought to be required for specialized nuclear morphology associated with

mitotic and meiotic cell division, karyogamy at fertilization, cell polarity, and response to

biotic and abiotic interactions (Starr and Fridolfsson, 2010; Gundersen and Worman,

2013; Griffis et al., 2014; Fernández-Álvarez et al., 2016; Groves et al., 2018).

Most of our knowledge of the LINC complex comes from studies of metazoans

and yeast. However, given the global challenges in agriculture and the need for plant-

based biorenewable resources (Godfray et al., 2010), it is important to better

understand the plant NE and its fundamental components. Recently, progress towards

understanding plant LINC biology has expanded beyond Arabidopsis to include

Medicago and maize (Gumber et al., 2019; Newman-Griffis et al., 2019).

The first SUN domain protein identified in plants was OsSad1, found in a nuclear

proteomic study in rice (Moriguchi et al., 2005). Subsequently, SUN domain proteins

have been identified in many plant species (reviewed by Meier, 2016; Meier et al., 2017;

Poulet et al., 2017a) with roles in the maintenance of nuclear shape and size

(Graumann et al., 2010; Oda and Fukuda, 2011; Graumann et al., 2014) and meiotic

chromosome behavior (Murphy et al., 2014; Varas et al., 2015). In maize, SUN2 has

been shown to form a novel structure called the meiotic SUN belt during prophase of

meiosis I (Murphy et al., 2014). The SUN belt region of the NE includes the telomere

cluster which defines the zygotene bouquet (Murphy et al., 2014).

The first plant KASH proteins were originally discovered in Arabidopsis through

analysis of RanGAP-NE anchoring proteins, and implicated in NE-associated signaling

and maintenance of nuclear shape (Xu et al., 2007; Zhou et al., 2012). They were later

shown to bind to SUN (Meier et al., 2010; Zhou et al., 2012) and possess a single near-

C-terminal TMD and a C-terminal amino acid sequence of V/P-P-T. These features are

diagnostic of plant KASH-like proteins and were collectively used as criteria for

identification of ten KASH candidates in maize (Gumber et al., 2019). From that study,

four groups of Maize LINC KASH (MLK) related genes were identified: the *AtWIP*-like

group (*MLKP1 - MLKP4*), the *AtSINE*-like group (*MLKS1* and *MLKS2*), the grass-

specific group (*MLKG1* and *MLKG2*), and the KASH-associated *AtWIT*-like group

(*MLKT1* and *MLKT2*).

A particularly interesting and distinct group of plant KASH-like proteins are the

SUN-interacting nuclear envelope (SINE) proteins, which contain armadillo domains

(Armadillo-type fold, InterPro ID: IPR000225). The armadillo (ARM) type fold is a

protein-protein interaction surface formed by twisting of multiple alpha helices around a

central axis, and has been reported in a variety of proteins involved in intracellular

signalling and cytoskeletal regulation (Coates, 2003; Hatzfeld, 1998). Genetic analysis

of the SINE-group KASH genes in Arabidopsis reveals non-redundant functions ranging

from actin-dependent nuclear positioning in guard cells for *AtSINE1* to innate immunity

against a fungal pathogen for *AtSINE2* (Zhou et al., 2014). Despite their deep

conservation, the biological functions of plant SINE-group KASH genes in monocot

grass species remain largely uncharacterized. Maize contains two *AtSINE1* homologs,

MLKS1 (Zm00001d031134) and *MLKS2* (Zm00001d052955). Both of these contain an

ARM domain towards their N-terminus (Gumber et al., 2019). Given the fundamental

importance of connecting the nucleus to the cytoplasm, these maize SINE-group KASH

genes were predicted to participate in a broad array of processes requiring coordinated

actions of the NE with the cytoskeleton.

In this study, we have undertaken an investigation of a single maize LINC gene,

MLKS2, for several reasons. First of all, the MLKS2 protein was detected in CoIP

assays using ZmSUN2 antisera with earshoot nuclei (Gumber et al., 2019). Secondly,

MLKS2 is expressed in actively dividing cells, including meiosis-stage tassels,

according to transcriptomic studies (Stelpflug et al., 2016). Thirdly, *MLKS2* orthologs are

conserved across land plants, increasing the potential translational impact of the study

(Gumber et al., 2019). Finally, two independent Mu transposon insertion alleles of

MLKS2 were available (https://maizegdb.org/uniformmu) allowing for genetic and

phenotypic analyses. Here we characterize *MLKS2* using genetic and cellular

experiments in order to: (1) test for canonical KASH properties of NE localization and

SUN-binding; (2) investigate colocalization with cytoplasmic structures; (3) address its

role in meiosis and nuclear shape and positioning via mutant phenotyping; (4) gain

mechanistic insight into its role in nuclear positioning and actin interaction using domain

deletion assays; and (5) test for functional conservation through cross-species genetic

rescue of nuclear shape phenotype.

Results

The structure of the *MLKS2* gene and protein are summarized in Figure 1. The

MLKS2 gene Zm00001d052955_T001 (from B73 v4) has three exons and is predicted

to encode a protein with 637 amino acids (Fig. 1A-B). The secondary and domain

structure of the MLKS2 protein includes the canonical KASH features of a single

transmembrane domain near the C-terminus, followed by a 14-residue C-terminal KASH

domain and terminal sequence of LVPT (Fig. 1C). Nearly half of the protein (8-294) has

an armadillo-type fold domain (InterPro IPR016024). Another large segment (301-598)

is composed of three segments classified as disordered regions, designated here as

DR1 (301-401), DR2 (414-478), and DR3 (488-598). Tertiary structure prediction using

I-TASSER revealed that the ARM and disordered regions could adopt a single large

structure similar to that of human protein phosphatase 2A, PP2A, a structural

scaffolding subunit. PP2A (pdb 1b3uA) was the top ranked structural analog as shown

separately and threaded together with MLKS2 (Fig. 1D). This overall structure is

referred to as an alpha solenoid, broadly deployed in biological systems as a binding

structure (Fournier et al., 2013).

Nuclear envelope localization of MLKS2 is disrupted by C-terminal but not ARM

domain deletions

We first examined the cellular localization of MLKS2 and its potential to interact

with SUN or the cytoskeleton. GFP-tagged MLKS2 was transiently expressed and

imaged as shown in Figure 2, using the tobacco expression system previously

described (Gumber et al., 2019). The GFP-MLKS2 localized to the nuclear periphery as

expected for a KASH protein (Fig. 2B). The GFP-MLKS2 was also seen at the cell

periphery in a conspicuous pattern resembling that of a cytoskeletal network (Fig. 2B).

In order to map the protein regions responsible for these patterns, we made a series of

domain deletion constructs (Fig. 2A). The deletions were MLKS2Δ4, with removal of

the terminal four amino acid residues; MLKS2ΔKASH, with replacement of the terminal

KASH domain with three alanine residues; MLKS2ΔTM, with removal of the

transmembrane domain; or MLKS2ΔARM, with removal of the entire Armadillo fold

region, DR1, DR2, and part of DR3.

We found that the terminal Δ4 and ΔKASH constructs reduced the relative

abundance of MLKS2 at the nuclear periphery but retained the network-like pattern in

cytoplasm as clearly seen at the cell periphery (arrows, Fig. 2B). In contrast, removal of

the TM domain completely changed the localization pattern at both the nuclear and the

cell periphery to one resembling that of a soluble protein showing nucleoplasmic and

cytoplasmic labelling. The N-terminal deletion construct, ΔARM, retained its NE

localization but with a more uniform and less patchy signal. However, the ΔARM variant

had a dramatic effect on the cell peripheral labelling, which changed from a filamentous

cytoskeleton-like pattern to a more endoplasmic reticulum-like pattern (arrow heads,

Fig. 2B), including both sheets and tubules. These results demonstrate that when

expressed under the control of a constitutive promoter, MLKS2 can accumulate at two

places in the cell, the NE requiring C-terminal domains and the cytoplasm requiring the

N-terminal ARM domains. It remains a formal possibility that the non-nuclear

cytoplasmic MLKS2 may be an artifact of experimental over- or heterologous

expression and may not, therefore, reflect its primary biological function.

The KASH and ARM domains immobilize the NE-associated MLKS2

We next tested for interaction of MLKS2 with ZmSUN2 using fluorescence

recovery after photobleaching (FRAP). The mobility of GFP-MLKS2 expressed alone

was compared to that co-expressed with full length mCherry-ZmSUN2, or deletion

constructs lacking the SUN domain or coiled-coil regions of ZmSUN2 (Fig. 2C, D, S1).

GFP-MLKS2 has a large immobile fraction to start with, showing only 25% fluorescence

recovery 30 s after photobleaching (Fig. 2D). When co-expressed with full length

SUN2, the immobile fraction of GFP-MLKS2 showed a slight but statistically insignificant

increase compared to GFP-MLKS2 alone (Fig. 2C, D). When co-expressed with SUN2

protein lacking the coiled-coil domain (SUN2ΔCC), the immobile fraction of GFP-MLKS2

was significantly increased compared to GFP-MLKS2 alone or when co-expressed with

full length ZmSUN2. When co-expressed with ZmSUN2 protein lacking the SUN

domain (SUN2ΔSUN), the immobile fraction of GFP-MLKS2 was significantly and

markedly decreased compared to GFP-MLKS2 alone or when co-expressed with full

length ZmSUN2.

Given the large immobile fraction of MLKS2 to begin with, we did not expect to

observe a marked increase in its immobile fraction when co-expressed with full length

SUN. However, the drastic reduction in the immobile fraction when co-expressed with

SUN2ΔSUN, compared to co-expression with full-length SUN, was unexpected but

consistent with the interpretation that MLKS2 interacts with SUN2 via its SUN domain.

One possible explanation for the unexpected effect of the SUN2ΔSUN is that its

expression could have a dominant negative effect on the tobacco nuclear envelope

components and their ability to interact with GFP-MLKS2. We conclude that two

regions of MLKS2 each play a role in anchoring it within the NE, the C-terminal KASH

domain and the N-terminal ARM domain.

We used the FRAP mobility assay to decipher the contributions of the KASH and

ARM domains to the relatively large immobile fraction of MLKS2. Removal of the KASH

domain decreased the immobile fraction (Fig. 2E, F, S1), consistent with a KASH-SUN

binding interaction in the NE. Remarkably, removal of the ARM domain decreased the

immobile fraction even more (Fig. 2E, F) making over 60% of the protein population

mobile. This indicates that SINE2-type KASH proteins are likely docked in the NE

through interactions at both ends, one via SUN-KASH binding in the NE and one via

ARM domain binding to the cytoskeleton.

Genetic analysis of two transposon-disrupted alleles of *MLKS2*

For genetic analysis, we searched the UniformMu transposon mutagenesis

records (McCarty and Meeley, 2009) at maizeGDB and found two alleles for *MLKS2*.

The mutator element mobilized in UniformMu is MuDR, a 1.4 kbp non-autonomous *Mu1*

element (Barker et al., 1984). The two insertion alleles are designated here as *mlks2-1*

and *mlks2-2*, as diagrammed in Figure 3. Both alleles involved insertions into the

protein coding region in Exon 2. PCR primers were selected to confirm the location and

sequences surrounding the transposon insertion sites (Fig. 3 A,B). For each allele, a 9-

bp target site duplication was detected (underlined, Fig. 3B), characteristic for *Mu*

insertions. The PCR reactions used for genotyping of individual plants are shown for

representative individuals from families segregating for the insertions (Fig. 3C)

We examined RNA from immature, meiosis-staged tassels of wild-type and

mutant plants to check if the transposon insertions reduced the mRNA levels.

Quantitative RT-PCR showed that both alleles, *mlks2-1* and *mlks2-2*, resulted in

decreased relative RNA levels when compared to those from wild-type siblings (*Mu -/-*)

of *mlks2-1* and *mlks2-2* or those from parental W22 line of maize (Fig. 3D). Of the two

alleles, *mlks2-2* showed the greatest reduction in transcript levels.

MLKS2 is required for normal male meiosis and pollen development.

Given the role of the LINC complex and NE in meiosis, we asked whether

genetic disruption of *MLKS2* resulted in meiotic or post-meiotic phenotypes. In maize,

like most eukaryotes, the NE and the LINC complex have highly specialized functions

including a role in telomere bouquet formation, pairing, synapsis, and recombination, all

of which must occur with high fidelity to ensure proper genome reduction and euploid

transmission. We therefore used 3D cytology to look for several hallmarks of normal

male meiosis, including the telomere bouquet, the eccentric positioning of the nucleus at

early prophase I, 10 bivalents at late prophase I indicative of complete pairing, chiasma

formation, and equal distribution of chromosomes after meiosis I and II as evidenced by

daughter nuclei with matching sizes and shapes.

The *mlks2-2* mutants showed deviations from all of the cytological meiotic

hallmarks inspected as summarized in Figure 4. At early meiotic prophase, 3D

telomere FISH revealed that the *mlks2-2* plants showed partial telomere clustering on

the NE at the bouquet stage (Fig. 4A-C) with variable numbers (4-12) of telomere FISH

signals far removed from the primary cluster (bracketed "bq" in Fig. 4A, B). The

magnitude of telomere dispersion was measured using pairwise telomere FISH signal

distances per nucleus. The average pairwise telomere distance increased by 50% in

the mutant, from 4.3 microns in normal nuclei (n=6) to 6.5 microns in *mlks2-2* nuclei

(n=11). The tendency for multiple telomeres to reside far from the bouquet is also seen

in the fold change of inter-telomere pairwise distances for the long distance bins (Fig.

4C). At early prophase I, meiotic nuclei tend to relocate towards one edge of the cell

(Cowan et al., 2002), but *mlks2-2* nuclei remained closer to the cell center (Fig. 4D,E),

according to measures of nuclear eccentricity. The cell-to-nucleus centroid offset at

early prophase in normal cells decreased from 15 microns in normal (n=6) plants to 7

microns in mutant (n=11) plants (Fig. 4F).

In the late prophase I stages of diplotene or diakinesis, chromosomes continue to

condense while successful disomic pairing and recombination becomes cytologically

evident in the form of well-separated bivalents. These bivalents total up to the haploid

chromosome number, 10 in the case of normal maize (Fig. 4G). Failure to properly pair

or recombine typically results univalents or multivalents, which often fail to segregate

properly at anaphase during meiosis I division. The *mlks2-2* mutation caused

conspicuous and obvious departure from these normal late prophase I patterns (Fig.

4H-J). The meiotic chromosomes from mutant plants showed aggregation and

pronounced clumping of multiple chromosomes at late prophase I, with relatively few

clear examples of well separated bivalents (compare Fig. 4G to 4H-I). The tendency

and degree of aggregation varied in the mutant plants, even for cells from the same

anther. The majority of the mutant cells showed these late prophase I phenotypes.

This late prophase I clumping phenotype is intriguing given that telomere-NE tethering

normally persists even after the bouquet stage. These results place MLKS2 in the

poorly understood pathway that ensures uniform dispersal of bivalents after pachytene.

Anaphase and telophase can be particularly informative stages for detection of

abnormal chromosome segregation. We found that the *mlks2-2* mutants showed

numerous such segregation defects, including chromosomes not in the metaphase I

plate bundle (arrows, Fig 4L-N). The mutant plants frequently showed chromosome

laggards (Fig. 4P-Q) and micronuclei (Fig. 4R), both of which are indicative of

missegregation. Frequency of cells with one or more aberrant patterns at meiosis I

(metaphase, anaphase, telophase) were low for wild-type (5%, n = 200) but high for

mlks2-2 (46%, n = 303) Specifically, the percentages of cells with abnormal

chromosomes per stage were 61% in diakinesis (n=109), 37% in metaphase I (n=123),

60% in anaphase I (n=10), and 34% in the combined stages of telophase I, telophase

II, and 4-nucleate tetrad (n=61).

We next checked for post-meiotic phenotypes associated with pollen viability.

Acid fuchsin and malachite green differential pollen staining revealed that both *mlks2-1*

and *mlks2-2* plants showed reduced pollen viability compared to normal plants (Fig. 4S-

V, n>1,000 for each genotype). The pollen sterility phenotype was only partially

penetrant and, like the other phenotypes, more severe for *mlks2-2*. Taken together,

these phenotypes show that MLKS2 is required for proper male reproduction, with

cytological phenotypes appearing as early as the zygotene bouquet stage of meiosis.

MLKS2 plays a conserved role in nuclear shape maintenance in root hair cells

Root hair nuclei in plants are pleomorphic, and in Arabidopsis their shape and positioning are disrupted by mutations in LINC or LINC-associated genes (Dittmer and Richards, 2008; Oda and Fukuda, 2011; Tamura et al., 2013; Zhou et al., 2012). Interestingly, this rounding up or loss of elongation nuclear shape phenotype appears to be generally diagnostic for genetic disruption of the LINC complex. We measured nucleus lengths in maize root hairs 5 days after imbibition, from normal and *mlks2* mutants. Normal (W22, also designated W22+ or wild-type) nuclei averaged a length of 37 microns, but mutant nuclei were only half that long on average (Fig. 5A, B). A related measure, circularity index from edge-traced nuclei in 2D images, also changed from 0.4 in normal plants to a more circular value of 0.7 in the more "rounded up" mutants (Fig. 5C). Although the biological role of root hair nuclear elongation remains to be determined, these shape measures provide a quick and quantitative phenotypic readout for genetic disruption of the LINC complex at the seedling stage.

MLKS2 is required for normal stomatal complex development

While screening seedling leaf tissues for nuclear shape phenotypes, we found an unexpected and dramatic effect on stomatal complex development in *mlks2* mutants, as summarized in Figure 6. Normal stomatal development in maize leaves (reviewed by Smith, 2001) results in a bilaterally symmetric 4-cell complex composed of 2 dumbbell-shaped guard cells (GC in Fig. 6A) flanked by two subsidiary cells (SC in Fig. 6A). In the *mlks2* mutants, the subsidiary cells, but not the guard cells, exhibited dramatic and

variable deviations (denoted by arrows, Fig. 6D-I) in size, shape, and number per

stomatal complex. Given the known role of asymmetric cell division in the normal

development of the guard cell complex, we checked for cell polarity phenotypes at

earlier stages of development. Images from the meristematic base of the fourth leaf

showed that the *mlks2-2* mutant exhibited irregular cell polarities and division planes

(Fig. 6L), abnormal shaped subsidiary cells (Fig. 6O), or extra inter-stomatal and

subsidiary mother cells (Fig. 6K, N). These phenotypes closely resemble those of the

pangloss or *brick* mutants of maize, known to perturb nuclear positioning and the actin

cytoskeleton, culminating in irregular subsidiary cell development. These findings

uncover an unanticipated developmental role for *MLKS2* that likely reflects its function in

cell polarization and nuclear positioning for specialized, highly asymmetric cell divisions

leading to the differentiation of the stomatal complex.

MLKS2 interacts with F-actin via its ARM domain and is required for recruitment

of meiotic perinuclear actin

Given that MLKS2 has ARM domains, localizes to a cytoskeletal-like network,

and shows mutant phenotypes suggestive of cytoskeletal interactions, we wanted to

more directly explore the interaction of MLKS2 with actin. First, GFP-MLKS2 was co-

expressed with markers for the endoplasmic reticulum (RFP-HDEL), the actin

cytoskeleton (RFP-Lifeact) or the microtubule cytoskeleton (mCherry-TUA5) as shown

in Figures 7, S2. Colocalization of GFP-MLKS2 was seen for both ER and F-actin but

not microtubule markers (Fig. 7A, S2). The actin, but not the ER colocalization with

MLKS2, requires the presence of the MLKS2 ARM domain region (Fig. 7B). To further

characterize the interaction of MLKS2 with F-actin, we examined the localization of

MLKS2 in cells treated with Latrunculin-B (LatB) to depolymerize the F-actin. This

treatment caused the GFP-MLKS2 staining pattern to change from fibrous to a more

ER-like network, similar to that of the ARM domain deletion of MLKS2 (compare 2B to

7C). These findings show that when expressed in tobacco, MLKS2 appears to induce

an actin-ER colocalization. MLKS2 appears to connect the ER to F-actin through its

ARM domain. Consistent with this, the mobile fraction of MLKS2 in the NE was

increased after F-actin depletion by LatB (Fig. 7D, E). This result provides additional

evidence that MLKS2 binds to F-actin in an ARM domain-dependent manner.

Given these results together with the previously established roles for some KASH

proteins' actin-based nuclear anchoring (Gundersen and Worman, 2013; Huelsmann

and Brown, 2014; Zhou et al., 2014), we next asked if GFP-MLKS2 could affect nuclear

mobility in a live cell assay. The nuclear mobility baseline was first established in

N. benthamiana using GFP-LBR (Graumann et al., 2010) as a general NE marker (Fig.

7F-H). For these experiments, nuclear movement was tracked in live cells for five min

by imaging at 10 s intervals to obtain total distance migrated. Compared to the GFP-

LBR-expressing control, the nuclei expressing GFP-MLKS2 were dramatically

immobilized as shown by temporal color-coded projections, kymograph analysis (Fig.

7F-G), and measurements of total distance of nuclear movement over the 30-frame

time-lapse (Fig. 7H). Expression of MLKS2 and MLKS2ΔKASH resulted in significant

decreases in nuclear movement compared to the LBR control. In contrast, the nuclear

anchoring activity of MLKS2 could be abolished by deletion of the ARM domains (Fig.

7H). These live nuclear mobility assays define an ARM-dependent nuclear anchoring

activity for MLKS2, consistent with the core function of the LINC complex in connecting

the nucleus to the cytoskeleton.

Given the role of the NE dynamics in meiotic prophase and our findings that

MLKS2 interacts with F-actin in somatic cells, we returned to maize to check genetic

evidence of MLKS2-actin interactions in pollen mother cells at meiotic prophase (Fig. 7I-

P). The overall appearance of phalloidin-stained F-actin in W22 was similar to that of

mlks2-2 meiocytes, with two notable exceptions. First, the F-actin in the mutant plants

appeared to exhibit a subtle phenotype of somewhat discontinuous or fragmented F-

actin structures (compare fiber lengths of Fig. 7 panels J and L versus N and P).

Secondly, and most importantly, we observed that the *mlks2-2* meiocytes exhibited a

conspicuous loss of perinuclear F-actin staining. Perinuclear actin is common in maize

meiocytes (Murphy et al., 2014) and intensity line profile tracings show clear evidence of

F-actin concentration spikes around the nucleus (yellow arrows, Fig. 7M, N). However,

we were unable to observe by inspection (Fig. 7I-L) or line profile analysis (Fig. 7M-P)

any such perinuclear actin staining in *mlks2-2* mutant meiocytes. These findings

implicate MLKS2 in the formation of perinuclear F-actin to achieve proper meiotic

chromosome behavior.

MLKS2 can rescue Arabidopsis triple *wip* nuclear shape phenotype

Finally, the functional conservation of MLKS2 in plants was addressed using a

genetic rescue assay. Previously, it has been shown that deletion of certain KASH

proteins, such as AtWIP proteins, causes nuclei to become more circular (Zhou et al.,

2012). GFP-MLKS2 was expressed in Arabidopsis harboring mutations in three *WIP*

genes to determine if the maize SINE-group KASH protein could correct the nuclear

shape defect phenotype. For this experiment, stable transformants expressing GFP-

MLKS2 were grown and nuclei were imaged and analyzed from two different tissues, as

shown in Figure 8. Circularity index measurements of leaf and root nuclei showed that

maize GFP-MLKS2 complemented the nuclear shape phenotype of the triple-*WIP*

mutant in Arabidopsis by restoring the nuclei to wild-type length in two different cell

types. These experiments demonstrate that a GFP-tagged maize SINE-family KASH

protein can substitute for Arabidopsis *WIP*-family KASH proteins in nuclear shape

control. In summary, the findings from this study establish that the maize KASH protein

MLKS2 plays a role in multiple biological pathways including stomatal complex

development, meiotic nuclear architecture and positioning, meiotic chromosome

segregation, and production of viable pollen.

Discussion

In this study we characterized a core LINC component, MLKS2, in maize using genetic and cellular assays. We showed that MLKS2 localizes to two cellular compartments, the NE as expected and cytoplasmic ER where it colocalizes with actin via the ARM domain region. Genetic analysis uncovered mutant phenotypes in both vegetative and reproductive organs while documenting the first case of a plant KASH required for proper telomere clustering and chromosome behavior in meiotic prophase. Interactions of MLKS2 with F-actin via the ARM domains of MLKS2 may represent a general mechanistic principle that explains how MLKS2 might control nuclear shape and positioning. The pleiotropic nature of the mutants and MLKS2's deep conservation in plants may be important for a variety of tissue-specific biological functions that require specialized control of nuclear morphology and dynamics. For instance, MLKS2 may play a role in female meiosis and ovule development where highly organized nuclear positioning occurs in the mature ovary, pollen development and nuclear migration of sperm and vegetative nuclei, and fertilization itself. Interestingly, the overall plant morphology of both the wild type (WT) and homozygous mutant plants was largely normal.

At the onset of this study, the diagnostic ARM domains of the SINE group of plant KASH and previous findings from Arabidopsis AtSINE1 (Zhou et al., 2014) pointed to a possible MLKS2 interaction with F-actin. Interestingly, GFP-AtSINE1, but not GFP-AtSINE2, colocalizes with F-actin when expressed in tobacco or Arabidopsis (Zhou et al., 2014). Medicago SINE-group KASH genes, *MtSINE1* and *MtSINE2*, also encode proteins that colocalize with F-actin (Newman-Griffis et al., 2019). The ARM domains

alone may not, therefore, be predictive of F-actin binding. Consistent with this is the fact

that a variety of proteins have ARM domains, but their binding partners vary (Coates,

2003).   In this study, there exists a clear preponderance of evidence, direct and

indirect, that MLKS2 has F-actin binding activities. In the heterologous expression

system, MLKS2 localizes to the NE, as expected for a KASH domain protein, and to a

cytoskeletal-like reticulate network that colocalizes with F-actin and ER markers at the

cell periphery (Fig. 7A). Two complementary cell localization experiments, latrunculin-B

depletion of F-actin and ARM domain deletions, uncover a possible cytoplasmic function

that does not involve the nucleus, but rather deploys MLKS2 to tether the ER to

cytoplasmic F-actin. In this regard, MLKS2 resembles members of the Networked

(NET) proteins, as defined by Hussey and colleagues (Deeks et al., 2012). Although

MLKS2 does appear to fit the definition of membrane-bound plant-specific actin-binding

adapter protein, it lacks the tryptophan-rich NET actin-binding (NAB) domain sequence,

common to NET families 1-4 (Hawkins et al., 2014). The NET NAB domain is thought

to confer actin-binding activity, whereas for MLKS2 that activity may come from the

ARM domain. We note that the non-nuclear cytoplasmic MLKS2 may result from

overexpression or compromised targeting in the heterologous expression system used.

It nonetheless predicts that misregulation by overexpression may present cellular

dysfunction phenotypes generally associated with the cytoplasm and specifically

associated with the ER and cortical F-actin.

The actin interaction with core LINC NE components provides mechanistic

insights into several cellular functions of MLKS2 in both vegetative and reproductive

parts of the maize plant. We focused on three major nuclear phenotypes assayed

primarily in somatic cells: (1) positioning within the cell, (2) shape, and (3) mobility.

Nuclear positioning was implicated by genetic analysis of stomatal complex

development, nuclear shape was examined via rescue of root hair nuclei rounding up,

and nuclear mobility was directly measured in live cells.

Nuclear positioning defines cell polarity. One well-characterized example of

highly polarized cells associated with asymmetric cell divisions is the normal stomatal

complex development in leaves. In wild-type maize, guard mother cells (GMC) signal to

adjacent subsidiary mother cells (SMC) to commence a series of cellular events

involving nuclear migrations and localized actin patches that coordinate the eventual

production of a mature stomatal complex (Müller et al., 2009). The mature stomatal

complex is characterized by two guard cells flanked by subsidiary cells, displaying a

bilateral symmetry. Mutations known to disrupt stomatal complex development in maize

include the *discordia* (*dcd1*), the *panglass* (*pan1*, *pan2*), and the *brick* (*brk1*, *brk2*, and

brk3) mutants (Facette et al., 2015). The brick proteins are part of the SCAR complex

which interacts with ARP to form actin patches on the GMC side of the SMC. How the

nucleus moves towards the actin patches is not known. The KASH phenotypes

reported here bear a striking resemblance to those of *pan1* and *brk1*, placing *MLKS2* in

the stomatal complex development pathway. Given that MLKS2 ARM domains appear

to interact with F-actin, it is tempting to speculate that the developmental defect seen in

mlks2 reflects a failure to coordinate nuclear positioning with the cortical-ER-patches

known to occur at the region adjacent to the inducing GMC (Giannoutsou et al., 2011).

Nuclear shape maintenance is one of the fundamental functions performed by

LINC complex proteins. Nuclear morphology is commonly altered in LINC mutants,

some of which present pathological conditions such as envelopathies (Burke and

Stewart, 2006; Janin et al., 2017). In eudicot plants, disruption of LINC components

cause alterations in root hair nuclear shape phenotypes (Newman-Griffis et al., 2019;

Oda and Fukuda, 2011; Tamura et al., 2013; Zhou et al., 2015a). In maize, we similarly

observed that *mlks2* nuclei change to more compact and spherical shapes than those

from normal cells (Fig. 5A-C). There is growing evidence that disruption of any

component of the core LINC components or even the LINC-interacting nucleoskeletal

proteins can cause a "rounded up" nuclear phenotype. The finding that maize GFP-

MLKS2 was able to rescue the nuclear shape phenotype in Arabidopsis plants lacking

three WIP proteins is important for two reasons. First, it represents the first

demonstration of cross-species genetic rescue of a LINC component. Second, it

suggests that even diverged LINC components may possess some degree of structural

redundancy in establishing a balance of forces required for maintaining nuclear shape.

The triple *WIP* rescue by a maize SINE-group KASH may reflect, therefore, the capacity

of MLKS2 to replace a NE-cytoskeletal tether normally formed by the WIP--WIT--

MyosinXI-i-actin bridge (Tamura et al., 2013).

Nuclear mobility in plants is associated with a wide variety of cellular processes

including symbiosis, pathogenesis, fertilization, and division planes in development

(Griffis et al., 2014). Nuclear mobility can either be inferred from variable positioning in

fixed cells, such as the eccentric nucleus stage of early meiotic prophase (Fig. 4), or by

direct time-lapse tracking in live cells (Fig. 7F-H). Here we observed both. Our results

show that the mobility of tobacco nuclei is reduced or stopped by heterologous

expression of GFP-MLKS2, and this immobilization is reversed by removing the ARM

domain region of MLKS2. These findings, consistent with our genetic analyses,

implicate MLKS2 in nuclear anchoring, which may further explain, for example, the

appearance of wild-type elongated nuclei in root hair cells (Fig. 5). The lack of

perinuclear actin around the meiotic nucleus in *mlks2-2* plants further suggests that

MLKS2 may function to bind or recruit filamentous actin to the perinuclear cytoplasm. In

this way, MLKS2 may play an organizational or mechanical role whereby

microfilaments, microtubules, or their associated motor proteins may direct the

dynamics of meiotic nuclei (Sheehan and Pawlowski, 2009).

The role of the NE in meiotic chromosome behavior has been recognized for

over a century. In recent decades, the LINC complex has been shown to be required for

telomere clustering at the bouquet stage, as first described in fission yeast (Chikashige

et al., 2009). Here we show that MLKS2 is required for normal bouquet formation and

proper chromosome segregation. Our findings represent the first genetic evidence that

a plant KASH interacts with the actin cytoskeleton during male meiosis. These results

suggest that actin may be more important in plant meiosis than previously recognized.

This idea is consistent with live imaging of maize meiotic prophase chromosome and

nuclear movements, both of which were stopped following MF depletion by latrunculin-B

treatment (Sheehan and Pawlowski, 2009). In our mutants, the earliest cytological

meiotic defect is seen at the zygotene stage, when the telomere bouquet is normally

formed. The *mlks2* mutants exhibit defects in telomere clustering and nuclear

positioning. These early stage problems may later manifest in the failure of homologs to

synapse and recombine with high fidelity. MLKS2 may also have post-bouquet

functions such as interlock resolution and telomere dispersal, ideas consistent with

detection of irregular and pronounced clumping at late prophase (Fig. 4G-J). The

chromosome segregation laggards, micronuclei, and pollen sterility phenotypes may

therefore reflect multiple upstream failures of disomic pairing and recombination.

Interestingly, several of our *mlks2* phenotypes resemble those described for mutations

in AtSUN genes (Varas et al., 2015) or the meiotic kinesin1-like gene *AtPSS1 (Duroc et*

al., 2014). *AtSUN1-1* and *AtSUN2* double mutants also show partial polarization of

telomeres, absence of full synapsis, presence of unresolved interlock-like structures,

and reduction in the mean cell chiasma frequency which manifests in the formation of

laggard chromosomes during anaphase and severely reduced fertility (Varas et al

2015). Several of these phenotypes are the same or similar to what we observed with

mlks2, which is not surprising given that SUN and KASH are core components of the

LINC complex. Similarly, mutations in *AtPSS1* leads to phenotypes shared with mlks2,

including production of univalents and aneuploid cells after meiosis (Duroc et al 2014).

Findings such as these, together with the large body of evidence implicating

microtubules in plant meiotic chromosome behavior (Cowan and Cande, 2002; Cowan

et al., 2002; Driscoll and Darvey, 1970; Higgins et al., 2016; Shamina, 2003; Thomas

and Kaltsikes, 1977) suggest that meiotic chromosome behavior likely requires both

actin and tubulin cytoskeletal systems. Indeed, the LINC complex in general, and

MLKS2 in particular, may provide a coordinating structure that integrates cytoskeletal

systems to affect nuclear architecture.

In conclusion, this study identifies MLKS2 as a core LINC component with

multiple functions in fundamental plant vegetative and reproductive processes. Analysis

of the ARM domain sheds light on how the SINE group of plant KASH proteins can

interact with the actin cytoskeleton. In Figure 9, we present a working model for MLKS2

where we propose that MLKS2 interacts with F-actin via its N-terminal ARM domains in

wild-type (Fig. 9A) but not mutant plants (Fig. 9B). The MLKS2 protein could extend a

Velcro-type hook into the cytoplasm to bind generally or specifically with cytoskeletal

linear polymers, most likely F-actin. This interaction between MLKS2 and F-actin could

be achieved by direct or indirect contact (Fig. 9B, compare left and right). In tobacco

(Fig. 9C,D), the expression of GFP-MLKS2 or GFP-MLKS2ΔKASH results in nuclear

anchorage according to kymograph analysis, suggesting the ARM domain can anchor

the nucleus even if the KASH domain is absent. The anchorage is relieved by removal

of the ARM domains (Fig. 9D, right half). The models based on heterologous

expression in *N. benthamiana* include GFP-MLKS2 in the core LINC complex via a

presumed interaction with tobacco SUN (NbSUN, grey), or a demonstrated interaction

with maize SUN2 (red, ZmSUN2). These findings highlight the importance of

investigating individual components of the LINC complex in order to understand how the

NE coordinates cytoplasmic and nuclear phenomena to carry out fundamental biological

processes at the cellular level.

Materials and Methods:

Microscopy with tobacco and Arabidopsis

Nicotiana benthamiana (*Nb*) transformation was performed as described (Sparkes et al.,

2006). Agrobacterium cultures were used at an OD of 0.05 and plants imaged 3 days

post-infiltration. All imaging was performed on a Zeiss LSM 800 confocal system

equipped with a 100X 1.4NA lens. Bi-directional scanning was used for all imaging. For

DAPI excitation the 405nm laser was used and emission collected at 400-491 nm. For

GFP excitation the 488 nm laser was used and emission collected between 491-617

nm. For mCherry excitation the 561 nm laser was used and emission collected from

561-620 nm. Colocalization imaging was performed with multi-track line switching to

avoid fluorescence bleedthrough, and 4x frame averaging was used. For FRAP

acquisition a 5X digital zoom was used with a frame time of 0.15s. Fluorescence

recovery was monitored by consecutive imaging of 240 frames. Bleaching was

performed after 5 control frames were imaged using the 488 nm laser at 100X power for

20 iterations in a 160x160 pixel ROI containing the NE. For nuclear movement assays,

nuclei were imaged at 10 s intervals for 30 frames (5 minutes). We used LBR-GFP as a

control for baseline nuclear movement as described in (Graumann et al., 2010). For

DAPI staining, *N. benthamiana* or *A. thaliana* tissue was incubated in 5μg/ml DAPI and

0.001% Triton X-100 for 10 min before imaging.

For image analysis *ZEN* (Blue edition, v2.3.69.1000, Zeiss) was used and figures

compiled in *Illustrator* (v22.1, Adobe). For FRAP analysis, normalized intensity values

were calculated as described (Martinière et al., 2012), and one-phase association for

each replicate and combined curves plotted in *GraphPad* v7.04. For nuclear movement

assays, manual tracking was used to track the center of nuclear movement for each

experimental condition, and FIJI (Image J version 1.52h) was used to generate temporal

color-coded projections and kymographs over a 25 μm line. All whisker plots showing

data points were generated in *GraphPad* v7.04. Student's T-test or one-way ANOVA

statistical tests were used depending on the suitability of the data, as indicated in each

figure legend. The number of replicates used for each experiment can be found in the

figure legends. Circularity index analysis was performed as described (Zhou et al.,

2012).

Gene constructs

Gene constructs, nomenclature, and sequence information for the clones used in this

study are listed in Table S1.   GFP-MLKS2 constructs were synthesized (GenScript

Biotech Corp.) using the full-length ORF of MLKS2 fused to the C-terminal end of

eGFP-FLAG-HA. The synthetic clone (GFP-MLKS2) was obtained in a pUC18 vector,

inserted at the *Bam*HI restriction site to produce a plasmid designated pHWBF07. From

this vector, the eGFP-FLAG-HA-MLKS2 sequence was PCR amplified with *att* flanking

primers (Table S2) using high fidelity Q5 polymerase (NEB) and cloned into pDONR221

vector by BP cloning (Cat. No. 1235019, Invitrogen) to generate the entry clone

designated pHWBF07EC. The eGFP-FLAG-HA-MLKS2 sequence from the entry clone

was transferred to vector pH7WG2 (Karimi et al., 2002) by Gateway LR recombination

(Cat. No. 1235019, Invitrogen) to obtain the expression vector designated pPK1Fexp.

Deletion constructs were generated from the full-length entry clone as a template to

PCR-amplify required regions with internal primers containing flanking *att* sequences

and cloned into pDONR211 by Gateway BP cloning method. These deletion constructs

were also subcloned into pH7WG2 destination vector by Gateway LR recombination.

Full length and domain deletion constructs for ZmSUN2 were generated as described in

(Gumber et al., 2019).

Maize plant material and genotyping

The transposon insertion alleles of *MLKS2* (unique insertion IDs: mu1038603 and

mu1058535) were identified through MaizeGDB (http://www.maizegdb.org/) and derived

from the UniformMu transposon mutagenesis project (McCarty and Meeley, 2009;

Settles et al., 2007). Seeds for stock number UF-Mu04133 carrying allele *mlks2-1* or

stock number UF-Mu07312 carrying allele *mlks2-2* were obtained from the Maize

Genetics Cooperation Stock Center (http://maizecoop.cropsci.uiuc.edu/). For "wild-

type" maize referred to here as W22, we used color-converted W22 obtained from Hugo

Dooner (Waksman Inst., Rutgers, New Jersey, USA) derived by Brink (Brink, 1956).

The seeds were planted at the Florida State University Mission Road Research Facility

(Tallahassee, FL, USA) and propagated by out-crossing to W22 in summer 2014. In the

fall, the progeny seeds were grown in a greenhouse in the King Life Sciences building

(Biological Science Dept, Florida State University, Tallahassee, FL). The segregating

outcross plants were self-crossed to obtain mutant plants from among the progeny.

For PCR genotyping, leaf samples from 4-week-old maize W22 and plants from

families segregating for MLKS2 alleles were harvested, frozen, and stored at -80 C.

Genomic DNA was isolated using a modified mini-CTAB method from approximately 2

cm^2 leaf tissue in 96 well plates in a Mixer Mill (Retsch®) as described (Labonne et al.,

2013). Plants were genotyped using three different primer pair combinations to

accurately predict the genotype as described by (McCarty et al., 2013). For the -*mlks2-1*

allele, DNA extracted from each of the UF-Mu04133 stock plants was individually

amplified with 603F1-603R1 gene specific primer pair, as well as a gene specific primer

651 in combination with the Tir6 primer corresponding to terminal inverted repeats of the

652 Mu1 transposable element (603F1-Tir6 and 603R1-Tir6). Similarly, for the *mlks2-2*

653 allele, DNA extracted from each of the UF-Mu07312 stock plants was individually

654 amplified with 535F2-535R2, 535F2-Tir6 and 535R2-Tir6 primer pairs. Normal W22

655 DNA was also amplified with all of these primer pairs. All the primer sequences used for

656 genotyping are listed in Table S3. The PCR amplification products were resolved by

657 agarose gel electrophoresis followed by ethidium bromide staining. The PCR products

658 were sequence-verified using TA cloning in pCR™4Blunt-TOPO® Vector (Invitrogen cat

659 # K2875-20). M13F and M13R vector primers were used for sequencing (Molecular

660 Cloning Facility, Department of Biological Sciences, Florida State University) and the

661 resulting sequences were aligned with the W22 reference genome sequence to validate

662 the mu transposon element insertion sites.

RNA extraction and qRT-PCR

665 After one round of outcrossing to W22 and two rounds of self crossing, seeds were

666 grown in the greenhouse to harvest meiotic tassels from V13 -stage plants. The tassels

667 were frozen in liquid nitrogen and stored at -80°C until further use. The tissue was

668 ground in liquid nitrogen and 100 mg of the resulting powder used for RNA extraction as

669 per manufacturer's instructions (Qiagen RNeasy kit, Cat # 74904). The extracted RNA

670 was treated with amplification-grade DNase I (ThermoFisher, Cat # 18068015) for 15

671 min at room temperature, then treated with RNA clean and concentrator-25 kit (Zymo

672 Research, Cat # R1017). The resulting RNA was inspected for quantity and quality by

673 UV spectroscopy and gel electrophoresis, respectively. Superscript III reverse

transcriptase (ThermoFisher, Cat# 18080093) was used to reverse transcribe 1μg of

high quality RNA into cDNA according to the manufacturer's instructions. The resulting

cDNA was used as a template to amplify ~ 100 bp products downstream of Mu1

insertion sites in *mlks2-1* and *mlks2-2* using three gene-specific primers. Two

housekeeping genes CDK1 and RPN2 were amplified as internal controls (Lin et al.,

2014). The products were amplified using SYBR Green Master Mix (Applied

Biosystems) on a 7500 Fast Real-Time PCR System (Applied Biosystems). Relative

expression change was calculated using the 2ΔΔCt method (Livak and Schmittgen,

2001). Two technical replicates were carried out to ascertain the accuracy of the

procedure.

Microscopy in maize

For 3D telomere FISH analysis, meiotic stage tassels were harvested from W22

and *mlks2-2* plants grown in the greenhouse. Flowers were microdissected and fixed in

1X Buffer A with 4% paraformaldehyde for 1 hour on rotary shaker at room temperature

as described (Bass et al., 1997). The fixed anthers were microdissected and embedded

in acrylamide pads as described in (Howe et al., 2013) and subjected to telomere FISH

using a 0.13 μM fluorescein-conjugated telomere-specific oligonucleotide probe, MTLF.

The meiocytes were counterstained with 3 μg/mL DAPI then mounted in vectashield

(Vector labs) and imaged using a 3D-deconvolution microscope equipped with a 60X

lens and CCD camera (Deltavision, GE Healthcare). Telomere coordinates in X, Y, and

Z were identified using the 3D MODEL tool of SoftWorx program (DeltaVision, GE

Healthcare) and pairwise telomere distances were calculated using R software for

Euclidean distances.

For actin staining, meiotic-staged anthers were fixed in PHEMS buffer

supplemented with 8% paraformaldehyde for 2 hours with shaking at room temperature

as described (Chan and Zacheus Cande, 1998). The fixed anthers were microdissected

on a glass slide in PBS (pH 7.0) and stained with 3.3 µM Alexa Fluor™ 594 Phalloidin

(Thermo Fisher Scientific) and DAPI (1.5 µg/mL) for 20 min at room temperature in the

dark. The cells were imaged using a Deltavision 3D deconvolution microscope.

For pollen viability, anthers from male flowers just before dehiscence were fixed

in Carnoy's fixative (6 alcohol:3 chloroform:1 acetic acid) for a minimum of 2 hours at

room temperature. The fixed anthers were placed on glass slides and stained with

modified Alexander's stain containing Malachite green (0.01%), Acid Fuchsin (0.05%)

and Orange G (0.005%) as described (Peterson et al., 2010) to differentiate viable

(magenta) pollen grains from aborted (green) pollen grains. Bright field images of pollen

grains were collected on Revolve microscope (Echo Labs). At least 300 pollen grains

each from 3 plants of every genotype were counted to calculate pollen viability.

For maize root hair imaging, seeds were sterilized with 1% bleach, 0.01% Triton

X-100 in deionized water for 10 min, rinsed and soaked in water overnight. The next

day seeds were spread on wet paper towels, kept dark and moist for 5-6 days, then

roots were cut off and fixed in Buffer A (Howe et al., 2013) plus 4% final concentration

paraformaldehyde for 1 hour, with rotation at room temperature. Small pieces of roots

with root hair from the elongation zone were placed on glass slides and chopped to

small pieces with a razor blade. The tissue was stained with 3 µg/mL DAPI for 20 min

at room temperature, mounted with VECTASHIELD, and imaged on an EVOS

fluorescence microscope (Thermo Fisher Scientific). The images were processed using

the *Analyze particle* function of *ImageJ* to measure the longest diameter and circularity

of the nuclei.

For leaf and stomatal complex imaging, plants were grown in the greenhouse

and the 4th leaf was harvested at its first appearance. For developing stomatal complex,

up to 2 cm tissue from the base of the leaf was harvested, and for mature stomatal

complex, the remainder of the leaf was harvested and fixed in FAA fixative (3.7%, 10%

Formalin, 50% ethanol and 5% acetic acid in water) for at least 15 min at room

temperature as described in (Facette et al., 2015). The leaf tissue was cut in thin

longitudinal strips and treated with PBS supplemented with 1% Triton for 10 min to

permeabilize the tissue. The tissue was then rinsed and incubated in a solution of

propidium iodide (Life technologies) and DAPI for 30 min. The tissue was rinsed with

PBS to get rid for excess dye and mounted in water. The tissue was imaged on a 3D

deconvolution microscope at 60X magnification.

Figure Legends

Figure 1: Gene model and protein domain diagram of *MLKS2*.

A) Gene model diagram for *MLKS2* with gene features drawn to scale. Total length of

the primary transcript (transcript model for B73v4 is indicated above) is 3007 bp. Three

exons are shown as black boxes. The 5'UTR of the gene is interrupted by "Intron 1."

Gene features are indicated: TSS, transcription start site; UTR, untranslated region;

ORF, open reading frame. B) The amino acid sequence of MLKS2 with domains

underlined with lines of different styles, or boxed. The protein sequence is identical to

that from genotype W22 (not shown) used for transposon mutagenesis. The key is

given under the sequence. C) Domain diagram for MLKS2 showing the names and

locations of secondary structural features (alpha helices) or domains are indicated. The

conserved terminal four amino residues (LVPT) at the end of the KASH domain are

shown . D) The I-TASSER protein structure prediction software was used with MLKS2

as input sequence. The top-ranked structural homolog is shown for a predicted tertiary

structure of MLKS2 (1st panel) next to the known structure of PP2A structural subunit

(pdb 1b3uA, 2nd panel), followed by an overlay of the two (3rd panel). The rainbow

annotation denotes the protein polarity from the N-terminus (blue) to the C-terminus

(red). The transmembrane (TMD) and KASH domains of MLKS2 are annotated.

Figure 2: MLKS2 cellular localization and ZmSUN2-interaction.

Subcellular localization of full length and deletion mutants of GFP-MLKS2 in transiently

expressing *N. benthamiana* leaf cells as previously described (Gumber et al., 2019). A)

Domain deletion diagram of all constructs used showing the ARM domain (yellow),

disordered regions (blue), transmembrane domain (TM), KASH domain (purple),

terminal AA residues, and stop codon (*). B) Full length GFP-MLKS2 (green) and all but

one deletion mutant localize to the NE around DAPI-stained chromatin (magenta). GFP-

MLKS2ΔTM appears soluble and distributed throughout the nucleoplasm. A cell

periphery network-like pattern was observed for all constructs, with GFP-MLKS2ΔTM

appearing as soluble in the cytoplasm and GFP-MLKS2ΔARM appearing as associated

with the ER. C) Fluorescence recovery curve from MLKS2 FRAP alone or when co-

expressed with mCherry-ZmSUN2, mCherry-ZmSUN2ΔCC (without coiled coil domain),

or mCherry-ZmSUN2ΔSUN (without SUN domain). D) FRAP recovery plateau values

for fluorescence recovery curves shown in C). E) Fluorescence recovery curves

showing increased mobile fraction in the NE for KASH or ARM domain deletion variants

of GFP-MLKS2. F) FRAP recovery plateau values of MLKS2, MLKS2ΔKASH, and

MLKS2ΔARM shown in D). Scale bar denotes 10 μm. For whisker plots, blue lines

denote SD error bars, red lines denote mean. Nuclei (N = ≥30) imaged across three

experimental repeats per treatment. ANOVA statistical test used where Ns = P≥0.05, ***

= P≤0.001 and **** = P≤0.0001.

Figure 3. Genotypic characterization of *MLKS2* alleles. Two transposon-tagged

alleles of *MLKS2, mlks2-1* and *mlks2-2* are depicted. A) *MLKS2* gene model showing

the location of *Mu1* transposon insertion sites (triangles) for two alleles. The positions of

allele-specific primer pairs (gF1, gR1 and gF2, gR2) and Tir6 primer (T6) at the Mu1

transposon terminal inverted repeat sequence are marked with arrows. The positions of

primer pairs (qF1, qR1; qF2, qR2; or qF3, qR3) used for qRT-PCR are also marked with

arrows. The PCR products (21FR, 21FT, 21TR, 22FR, 22FT, and 22TF) used for

sequence verification, genotyping, and quantitative RT-PCR are indicated below the

gene model. B) Sequences aligned around the insertion site include the published

parental sequence (Springer et al., 2018), the wild-type allele from PCR products ("a"

and "d") from W22, and the mutator-flanking sequences from PCR products ("b","c", "e",

and "f") using one gene primer and one mutator-specific primer (T6). In both alleles, a

9-bp duplication (underlined) was detected. C) Agarose gels showing PCR products

amplified from W22 and plants from families segregating for *mlks2-1* (top gel) or *mlks2-*

2 (bottom gel) allele. The PCR products were amplified using gene-specific primer pairs

(lanes/PCR products a, d) or primer pairs from one gene-specific primer and one

mutator (T6) primer (lanes/PCR products b, c, e, f). Plant genotypes are shown on the

top of the gels, primer pairs and band sizes are indicated on the right. The lanes "M"

contain 100 bp DNA marker fragments at the lengths indicated. D) Fold change in the

transcript levels of *MLKS2* in families segregating for *mlks2-1* or *mlks2-2*; homozygous

WT siblings (+/+) or homozygous mutant plants (-/-) were quantified relative to W22

using an average of 3 primer pairs (qF1-qR1, qF2-qR2, qF3-qR3) as measured by qRT-

PCR.

Figure 4: Multiple meiotic and post-meiotic defects of *mlks2* mutants.

Cytological defects of *mlks2-2* during and after male meiosis. A) Early prophase I stage

W22 meiocyte nucleus showing a typical bouquet (green dots, "bq") of NE-associated

telomeres visualized in the nucleus (DAPI, shown in magenta) using the 3D acrylamide

oligo FISH method (Bass et al., 1997). B) Partial telomere bouquet in *mlks2-2* mutant at

meiotic prophase, showing unusually distant telomeres (arrows) relative to the main

bouquet (bq) telomere cluster region. C) Histogram showing bouquet-stage telomere

pairwise distance distributions plotted as log2 fold change of mutant/wild-type per 1

micron distance bins (n=6 W22, n=11 *mlks2-2*). The mutants show a pronounced

increase in the longer telomere-to-telomere distance bins. Nuclear position phenotypes

for normal D) or mutant E) cells shown as projections from the middle-most 1/5 of the

optical sections through the nuclei stained with DAPI, including traces around the cell

and nuclear peripheries to ascertain 2D centroid locations. F) Eccentricity plots

showing the distribution of distances of between the pairwise centroids of nuclei and

cells for normal (W22) and mutant (*mlks2*) meiocytes at at the bouquet stage, using the

same nuclei as those analyzed in A-C. G) Late prophase I stage W22 meiocyte showing

bivalents spread throughout the volume of the nucleus. H-J) Late prophase I stage

mlks2-2 meiocytes showing clumping of bivalents. K) W22 meiocyte showing bivalents

on a normal meiosis I metaphase plate. L-N) Mutant *mlks2-2* meiocytes showing one or

more chromosomes (arrows) not located in the meiosis I metaphase plate. O) W22

meiocyte at late anaphase I or early telophase I. P-Q) Mutant *mlks2-2* late anaphase I

or early telophase I showing irregularly positioned "laggard" chromosomes (arrows). R)

Mutant mlks2-2 at telophase after meiosis I, before meiosis II, and showing micronuclei

(arrows) that are associated with failure of chromosomes or chromosomal fragments to

reach the spindle poles. S-U) Pollen viability stains for wild-type S) or mutant T, U)

pollen. Dark purple indicates viable pollen, light blue indicates inviable pollen. V)

Quantification of pollen viability with n=1,000 or more for each genotype.

Figure 5. MLKS2 is required for the elongated nuclear shape phenotype in root

hair cells.

A) Root hair nuclei stained with DAPI in W22, *mlks2-1* and *mlks2-2* maize plants. B)

Quantitation of longest dimension of W22, *mlks2-1*, or *mlks2-2* root hair nuclei. C)

Circularity index measurements of nuclei from W22, *mlks2-1* and *mlks2-2* in which 1.0

represents a perfectly round shape.

Figure 6. MLKS2 is required for normal stomatal complex development.

Mature stomatal complex in DAPI stained leaf from wild-type W22 (A-C), *mlks2-1* (D-F)

or *mlks2-2* (G-I) plants. For comparison, a typical stomatal complex has bilateral

symmetry with two elongated central dumbbell-shaped guard cells (GC) flanked by two

outer subsidiary cells (SC). In the *mlks2-1* D-F) or *mlks2-2* G-I) mutants, subsidiary

cells appear abnormal (arrows) in their number, shape, or nuclear position relative to

the guard cells. J-L) Representative images from early stages of stomatal development

where subsidiary mother cells (SMC) are polarizing, with nuclei migrating towards the

guard mother cells (GMC). Interstomatal cells (ISC) are annotated for W22. M-O)

Representative images of developing stomatal complex after SMC polarization.

Boundaries of abnormally shaped cells or cells with abnormal nuclear positioning are

marked with dotted lines. Scale bars denote 15 µm.

Figure 7. MLKS2 interaction with actin.

Co-localization of ER marker (RFP-HDEL, magenta) or actin marker (RFP-LifeAct,

magenta) with A) GFP-MLKS2 or B) GFP-MLKS2ΔARM. C) GFP-MLKS2 localization in

F-actin-depleted cells (25 uM LatB) phenocopies GFP-MLKS2ΔARM with an ER-like

staining pattern. D) Depolymerization of the actin cytoskeleton with LatB results in

increased GFP-MLKS2 FRAP recovery compared to mock (MK) controls. E) FRAP

Plateau value of MLKS2 mock (MK) and LatB treated as shown in D). For whisker plots,

blue lines denote SD error bars, red lines denote mean. Student's T-test used to test

statistical significance. **** = P≤0.0001. Scale bar denotes 10 µm. F) Temporal color-

coded projections of randomly selected LBR, MLKS2, MLKS2ΔKASH and

MLKS2ΔARM nuclei imaged every 10 seconds over 5 minutes. G) Kymographs of

nuclear movement shown in F) for a different nucleus. H) Quantification of total nuclear

movement over time, imaged for LBR, MLKS2, MLKS2ΔKASH and MLKS2ΔARM. N =

at least 30 nuclei imaged across three experimental repeats per treatment; ANOVA

statistical test used. Ns = P≥0.05, * = P≤0.05, *** = P≤0.001 and **** = P≤0.0001. (I-L)

DAPI (magenta) and phalloidin (green) stained early prophase maize meiocytes. (M-P)

Gray-scale images of phalloidin staining of cells shown in panels I-L. Line trace plots

show intensity of phalloidin in the middle of the cell marked by horizontal band,

illustrating the spike in perinuclear actin in wild-type but not mutant nuclei. Scale bar

denotes 15 μm.

Figure 8. Arabidopsis triple *wip* mutant phenotype rescued with MLKS2.

DAPI stained representative images of A) leaf and B) root nuclei from three

Arabidopsis genotypes: *Arabidopsis thaliana* WT (Col0); (Columbia WT strain), *wip123*

(*AtWIP*-type KASH triple mutant where *wip123* refers to genotype *wip1-1*, *wip2-1*, *wip3-*

1) (Zhou et al., 2015b), or *wip123* plus GFP-MLKS2 (WIP triple mutant transformed with

GFP-MLKS2). (C, D) Nuclear circularity index summaries for these same tissues and

genotypes are shown below each tissue/genotype combination, where 1 is a perfectly

round nucleus and values lower than 1 are a measure of nuclear elongation.

Arabidopsis WIP triple mutant nuclei are significantly more rounded (leaf CI=0.81±0.03;

root CI=0.72±0.02) than WT (leaf CI=0.56±0.03; root CI=0.39±0.02) whereas GFP-

MLKS2 complemented nuclei are similar to WT (leaf CI=0.52±0.03, root CI=0.32±0.02);

p>0.05=ns, p<0.0001=****.

Figure 9. Summary diagrams and models of ZmMLKS2.

Summary diagram illustrates how MLKS2 may interact with F-actin to produce the

genetic and heterologous expression phenotypes reported in this study. A) MLKS2 is

presumed to be arranged with an ARM domain-containing alpha solenoid structure in

the cytoplasm, where it interacts with F-actin directly (left half) or indirectly through a

hypothetical connector depicted by the boxed question mark (right half). B) The *mks2-2*

mutant has lost the ability to bind or contribute to the recruitment of F-actin. Not

depicted here are data from MLKS2 in vegetative (leaf, root) organs. Results from

kymograph analysis of GFP-MLKS2 expressed in *N. benthamiana* are summarized for

experiments that showed C) anchored nuclei in cells expressing full length MLKS2 or

MLKS2ΔKASH, or D) mobile nuclei in cells expressing a control NE marker (lbr) or

MLKSΔARM. In these diagrams, evidence for interaction with ZmSUN2 co-expression

is indicated (red) on the basis of FRAP assays or depicted from presumed interactions

with tobacco SUN, NbSUN (grey). In these diagrams (C,D), the MLKS2 interaction with

F-actin is depicted as direct for convenience, but the alternative indirect mode (A, right

half) of MLKS2 interaction with F-actin remains a formal possibility.

Figure S1. Individual nucleus data from FRAP experiments presented throughout

the main figures. A) FRAP recovery curves from all individual nuclei imaged with GFP-

MLKS2 for comparison to co-expression with SUN2. B) FRAP recovery curves from all

individual nuclei imaged with GFP-MLKS2 and co-expressing mCherry-SUN2. C) FRAP

recovery curves from all individual nuclei imaged with GFP-MLKS2 and co-expressing

mCherry-SUN2ΔCC. D) FRAP recovery curves from all individual nuclei imaged with

GFP-MLKS2 and co-expressing mCherry-SUN2ΔSUN. E) Halftime values from FRAP

recovery experiments shown in A-D. F) FRAP recovery curves from all individual nuclei

imaged with GFP-MLKS2 as control during GFP-MLKS2ΔKASH and GFP-

MLKS2ΔARM experiments. G) FRAP recovery curves from all individual nuclei imaged

with GFP-MLKS2ΔKASH. H) FRAP recovery curves from all individual nuclei imaged

with GFP-MLKS2ΔARM. I) Halftime values from FRAP recovery experiments shown in

F-H. J) FRAP recovery curves from all individual nuclei imaged with GFP-MLKS2 after 1

hour mock treatment, for comparison to LatB treatment. K) FRAP recovery curves from

all individual nuclei imaged with GFP-MLKS2 after 1 hour 25µM LatB treatment. L)

Halftime values from FRAP recovery experiments shown in J and K. Recovery curves

blue dots show individual data points, black lines individual fluorescence recovery

curves, and red curves average recovery curves for all nuclei. For whisker plots, blue

lines denote error bars showing standard deviation, red lines show mean values.

Statistical tests performed were either one-way ANOVA (E and I) or Student's t-test (L)

as appropriate. ns = P≥0.05, ** = P = ≤ 0.01.

Figure S2. Co-expression of GFP-MLKS2 and tubulin alpha-5 (TUA5)

Representative images of *N. benthamiana* leaves transiently co-transformed with GFP-

MLKS2 and mcherry-TUA5. MLKS2 and microtubules do not seem to colocalize. Scale

bar denotes 10 µm.

References

Alam, S. G., Zhang, Q., Prasad, N., Li, Y., Chamala, S., Kuchibhotla, R., Kc, B., Aggarwal, V., Shrestha, S., Jones, A. L., et al. (2016). The mammalian LINC complex regulates genome transcriptional responses to substrate rigidity. *Sci. Rep.* **6**, 38063.

Barker, R. F., Thompson, D. V., Talbot, D. R., Swanson, J. and Bennetzen, J. L. (1984). Nucleotide sequence of the maize transposable element Mul. *Nucleic Acids Res.* **12**, 6924–6924.

Bass, H. W., Marshall, W. F., Sedat, J. W., Agard, D. A. and Cande, W. Z. (1997). Telomeres cluster de novo before the initiation of synapsis: a three-dimensional spatial analysis of telomere positions before and during meiotic prophase. *J. Cell Biol.* **137**, 5–18.

Brink, R. A. (1956). A Genetic Change Associated with the R Locus in Maize Which Is Directed and Potentially Reversible. *Genetics* **41**, 872–889.

Burke, B. and Roux, K. J. (2009). Nuclei take a position: managing nuclear location. *Dev. Cell* **17**, 587–597.

Burke, B. and Stewart, C. L. (2006). The laminopathies: the functional architecture of the nucleus and its contribution to disease. *Annu. Rev. Genomics Hum. Genet.* **7**, 369–405.

Chan, A. and Zacheus Cande, W. (1998). Maize meiotic spindles assemble around chromatin and do not require paired chromosomes. *J. Cell Sci.* **111**, 3507–3515.

Chang, W., Worman, H. J. and Gundersen, G. G. (2015). Accessorizing and anchoring the LINC complex for multifunctionality. *J. Cell Biol.* **208**, 11–22.

Chikashige, Y., Yamane, M., Okamasa, K., Tsutsumi, C., Kojidani, T., Sato, M., Haraguchi, T. and Hiraoka, Y. (2009). Membrane proteins Bqt3 and -4 anchor telomeres to the nuclear envelope to ensure chromosomal bouquet formation. *J. Cell Biol.* **187**, 413–427.

Coates, J. C. (2003). Armadillo repeat proteins: beyond the animal kingdom. *Trends Cell Biol.* **13**, 463–471.

Cowan, C. R. and Cande, W. Z. (2002). Meiotic telomere clustering is inhibited by colchicine but does not require cytoplasmic microtubules. *J. Cell Sci.* **115**, 3747–3756.

Cowan, C. R., Carlton, P. M. and Cande, W. Z. (2002). Reorganization and

polarization of the meiotic bouquet-stage cell can be uncoupled from telomere
clustering. *J. Cell Sci.* **115**, 3757–3766.

D'Angelo, M. A. (2018). Nuclear pore complexes as hubs for gene regulation. *Nucleus*
9, 142–148.

Deeks, M. J., Calcutt, J. R., Ingle, E. K. S., Hawkins, T. J., Chapman, S.,
Richardson, A. C., Mentlak, D. A., Dixon, M. R., Cartwright, F., Smertenko, A.
P., et al. (2012). A Superfamily of Actin-Binding Proteins at the Actin-Membrane
Nexus of Higher Plants. *Curr. Biol.* **22**, 1595–1600.

Dittmer, T. A. and Richards, E. J. (2008). Role of LINC proteins in plant nuclear
morphology. *Plant Signal. Behav.* **3**, 485–487.

Driscoll, C. J. and Darvey, N. L. (1970). Chromosome pairing: effect of colchicine on
an isochromosome. *Science* **169**, 290–291.

Duroc, Y., Lemhemdi, A., Larchevêque, C., Hurel, A., Cuacos, M., Cromer, L.,
Horlow, C., Armstrong, S. J., Chelysheva, L. and Mercier, R. (2014). The
kinesin AtPSS1 promotes synapsis and is required for proper crossover distribution
in meiosis. *PLoS Genet.* **10**, e1004674.

Evans, D. E., Pawar, V., Smith, S. J. and Graumann, K. (2014). Protein interactions at
the higher plant nuclear envelope: evidence for a linker of nucleoskeleton and
cytoskeleton complex. *Front. Plant Sci.* **5**, 183.

Facette, M. R., Park, Y., Sutimantanapi, D., Luo, A., Cartwright, H. N., Yang, B.,
Bennett, E. J., Sylvester, A. W. and Smith, L. G. (2015). The SCAR/WAVE
complex polarizes PAN receptors and promotes division asymmetry in maize. *Nat*
Plants **1**, 14024.

Fernández-Álvarez, A., Bez, C., O'Toole, E. T., Morphew, M. and Cooper, J. P.
(2016). Mitotic Nuclear Envelope Breakdown and Spindle Nucleation Are
Controlled by Interphase Contacts between Centromeres and the Nuclear
Envelope. *Dev. Cell* **39**, 544–559.

Fournier, D., Palidwor, G. A., Shcherbinin, S., Szengel, A., Schaefer, M. H., Perez-
Iratxeta, C. and Andrade-Navarro, M. A. (2013). Functional and genomic
analyses of alpha-solenoid proteins. *PLoS One* **8**, e79894.

Giannoutsou, E. P., Apostolakos, P. and Galatis, B. (2011). Actin filament-organized
local cortical endoplasmic reticulum aggregations in developing stomatal
complexes of grasses. *Protoplasma* **248**, 373–390.

Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J.
F., Pretty, J., Robinson, S., Thomas, S. M. and Toulmin, C. (2010). Food
security: the challenge of feeding 9 billion people. *Science* **327**, 812–818.

Graumann, K., Runions, J. and Evans, D. E. (2010). Characterization of SUN-domain
proteins at the higher plant nuclear envelope. *Plant J.* **61**, 134–144.

Graumann, K., Vanrobays, E., Tutois, S., Probst, A. V., Evans, D. E. and Tatout, C.
(2014). Characterization of two distinct subfamilies of SUN-domain proteins in
Arabidopsis and their interactions with the novel KASH-domain protein AtTIK. *J.*
Exp. Bot. **65**, 6499–6512.

Griffis, A. H. N., Groves, N. R., Zhou, X. and Meier, I. (2014). Nuclei in motion:
movement and positioning of plant nuclei in development, signaling, symbiosis, and
disease. *Front. Plant Sci.* **5**, 129.

Groves, N. R., Biel, A. M., Newman-Griffis, A. H. and Meier, I. (2018). Dynamic
Changes in Plant Nuclear Organization in Response to Environmental and
Developmental Signals. *Plant Physiol.* **176**, 230–241.

Gumber, H. K., McKenna, J. F., Estrada, A. L., Tolmie, A. F., Graumann, K. and
Bass, H. W. (2019). Identification and characterization of genes encoding the
nuclear envelope LINC complex in the monocot species Zea mays. *J. Cell Sci.* **132**,
1–16.

Gundersen, G. G. and Worman, H. J. (2013). Nuclear positioning. *Cell* **152**, 1376–
1389.

Hagan, I. and Yanagida, M. (1995). The product of the spindle formation gene sad1+
associates with the fission yeast spindle pole body and is essential for viability. *J.*
Cell Biol. **129**, 1033–1047.

Hatzfeld, M. (1998). The Armadillo Family of Structural Proteins. In *International*
Review of Cytology (ed. Jeon, K. W.), pp. 179–224. Academic Press.

Hawkins, T. J., Deeks, M. J., Wang, P. and Hussey, P. J. (2014). The evolution of the
actin binding NET superfamily. *Front. Plant Sci.* **5**, 254.

Hetzer, M. W. (2010). The nuclear envelope. *Cold Spring Harb. Perspect. Biol.* **2**,
a000539.

Hieda, M. (2017). Implications for Diverse Functions of the LINC Complexes Based on
the Structure. *Cells* **6**,.

Higgins, D. M., Nannas, N. J. and Dawe, R. K. (2016). The Maize Divergent spindle-1
(dv1) Gene Encodes a Kinesin-14A Motor Protein Required for Meiotic Spindle
Pole Organization. *Front. Plant Sci.* **7**, 1277.

Howe, E. S., Murphy, S. P. and Bass, H. W. (2013). Three-Dimensional Acrylamide
Fluorescence In Situ Hybridization for Plant Cells. In *Plant Meiosis: Methods and*
Protocols (ed. Pawlowski, W. P.), Grelon, M.), and Armstrong, S.), pp. 53–66.
Totowa, NJ: Humana Press.

Huelsmann, S. and Brown, N. H. (2014). Nuclear positioning by actin cables and
perinuclear actin: Special and general? *Nucleus* **5**, 219–223.

Janin, A., Bauer, D., Ratti, F., Millat, G. and Méjat, A. (2017). Nuclear envelopathies:
a complex LINC between nuclear envelope and pathology. *Orphanet J. Rare Dis.*
12, 147.

Karimi, M., Inzé, D. and Depicker, A. (2002). GATEWAY^TM vectors for Agrobacterium-
mediated plant transformation. *Trends Plant Sci.* **7**, 193–195.

Kim, D. I., Birendra, K. C. and Roux, K. J. (2015). Making the LINC: SUN and KASH
protein interactions. *Biol. Chem.* **396**, 295–310.

Labonne, J. D. J., Dorweiler, J. E. and McGinnis, K. M. (2013). Changes in
nucleosome position at transcriptional start sites of specific genes in Zea mays
mediator of paramutation1 mutants. *Epigenetics* **8**, 398–408.

Lin, F., Jiang, L., Liu, Y., Lv, Y., Dai, H. and Zhao, H. (2014). Genome-wide
identification of housekeeping genes in maize. *Plant Mol. Biol.* **86**, 543–554.

Link, J., Leubner, M., Schmitt, J., Göb, E., Benavente, R., Jeang, K.-T., Xu, R. and
Alsheimer, M. (2014). Analysis of meiosis in SUN1 deficient mice reveals a distinct
role of SUN2 in mammalian meiotic LINC complex formation and function. *PLoS*
Genet. **10**, e1004099.

Livak, K. J. and Schmittgen, T. D. (2001). Analysis of relative gene expression data
using real-time quantitative PCR and the 2- ΔΔCT method. *Methods* **25**, 402–408.

Luxton, G. W. G. and Starr, D. A. (2014). KASHing up with the nucleus: novel
functional roles of KASH proteins at the cytoplasmic surface of the nucleus. *Curr.*
Opin. Cell Biol. **28**, 69–75.

Malone, C. J., Fixsen, W. D., Horvitz, H. R. and Han, M. (1999). UNC-84 localizes to
the nuclear envelope and is required for nuclear migration and anchoring during C.
elegans development. *Development* **126**, 3171–3181.

Martinière, A., Lavagi, I., Nageswaran, G., Rolfe, D. J., Maneta-Peyret, L., Luu, D.-
T., Botchway, S. W., Webb, S. E. D., Mongrand, S., Maurel, C., et al. (2012). Cell
wall constrains lateral diffusion of plant plasma-membrane proteins. *Proc. Natl.*
Acad. Sci. U. S. A. **109**, 12805–12810.

McCarty, D. R. and Meeley, R. B. (2009). Transposon Resources for Forward and
Reverse Genetics in Maize. In *Handbook of Maize: Genetics and Genomics* (ed.
Bennetzen, J. L.) and Hake, S.), pp. 561–584. New York, NY: Springer New York.

McCarty, D. R., Suzuki, M., Hunter, C., Collins, J., Avigne, W. T. and Koch, K. E.
(2013). Genetic and Molecular Analyses of UniformMu Transposon Insertion Lines.
In *Plant Transposable Elements: Methods and Protocols* (ed. Peterson, T.), pp.

157–166. Totowa, NJ: Humana Press.

Meier, I. (2016). LINCing the eukaryotic tree of life - towards a broad evolutionary
comparison of nucleocytoplasmic bridging complexes. *J. Cell Sci.* **129**, 3523–3531.

Meier, I., Zhou, X., Brkljacić, J., Rose, A., Zhao, Q. and Xu, X. M. (2010). Targeting
proteins to the plant nuclear envelope. *Biochem. Soc. Trans.* **38**, 733–740.

Meier, I., Richards, E. J. and Evans, D. E. (2017). Cell Biology of the Plant Nucleus.
Annu. Rev. Plant Biol. **68**, 139–172.

Moriguchi, K., Suzuki, T., Ito, Y., Yamazaki, Y., Niwa, Y. and Kurata, N. (2005).
Functional isolation of novel nuclear proteins showing a variety of subnuclear
localizations. *Plant Cell* **17**, 389–403.

Müller, S., Wright, A. J. and Smith, L. G. (2009). Division plane control in plants: new
players in the band. *Trends Cell Biol.* **19**, 180–188.

Murphy, S. P., Gumber, H. K., Mao, Y. and Bass, H. W. (2014). A dynamic meiotic
SUN belt includes the zygotene-stage telomere bouquet and is disrupted in
chromosome segregation mutants of maize (Zea mays L.). *Front. Plant Sci.* **5**, 314.

Newman-Griffis, A. H., Del Cerro, P., Charpentier, M. and Meier, I. (2019). Medicago
LINC Complexes Function in Nuclear Morphology, Nuclear Movement, and Root
Nodule Symbiosis. *Plant Physiol.* **179**, 491–506.

Oda, Y. and Fukuda, H. (2011). Dynamics of Arabidopsis SUN proteins during mitosis
and their involvement in nuclear shaping. *Plant J.* **66**, 629–641.

Peterson, R., Slovin, J. P. and Chen, C. (2010). A simplified method for differential
staining of aborted and non-aborted pollen grains. *1* **1**, e13–e13.

Poulet, A., Probst, A. V., Graumann, K., Tatout, C. and Evans, D. (2017a). Exploring
the evolution of the proteins of the plant nuclear envelope. *Nucleus* **8**, 46–59.

Poulet, A., Duc, C., Voisin, M., Desset, S., Tutois, S., Vanrobays, E., Benoit, M.,
Evans, D. E., Probst, A. V. and Tatout, C. (2017b). The LINC complex contributes
to heterochromatin organisation and transcriptional gene silencing in plants. *J. Cell*
Sci. **130**, 590–601.

Sakamoto, Y. and Takagi, S. (2013). LITTLE NUCLEI 1 and 4 regulate nuclear
morphology in Arabidopsis thaliana. *Plant Cell Physiol.* **54**, 622–633.

Settles, A. M., Holding, D. R., Tan, B. C., Latshaw, S. P., Liu, J., Suzuki, M., Li, L.,
O'Brien, B. A., Fajardo, D. S., Wroclawska, E., et al. (2007). Sequence-indexed
mutations in maize using the UniformMu transposon-tagging population. *BMC*
Genomics **8**, 116.

Shamina, N. V. (2003). [Dynamics of cytoskeleton microtubules in higher plant meiosis.
I. The perinuclear band of microtubules and the meiotic spindle formation].
Tsitologiia **45**, 650–654.

Sheehan, M. J. and Pawlowski, W. P. (2009). Live imaging of rapid chromosome
movements in meiotic prophase I in maize. *Proc. Natl. Acad. Sci. U. S. A.* **106**,
20989–20994.

Smith, L. G. (2001). Plant cell division: building walls in the right places. *Nat. Rev. Mol.*
Cell Biol. **2**, 33–39.

Sosa, B. A., Rothballer, A., Kutay, U. and Schwartz, T. U. (2012). LINC complexes
form by binding of three KASH peptides to domain interfaces of trimeric SUN
proteins. *Cell* **149**, 1035–1047.

Sparkes, I. A., Runions, J., Kearns, A. and Hawes, C. (2006). Rapid, transient
expression of fluorescent fusion proteins in tobacco plants and generation of stably
transformed plants. *Nat. Protoc.* **1**, 2019–2025.

Springer, N. M., Anderson, S. N., Andorf, C. M., Ahern, K. R., Bai, F., Barad, O.,
Barbazuk, W. B., Bass, H. W., Baruch, K., Ben-Zvi, G., et al. (2018). The maize
W22 genome provides a foundation for functional genomics and transposon
biology. *Nat. Genet.* **50**, 1282–1288.

Starr, D. A. and Fridolfsson, H. N. (2010). Interactions between nuclei and the
cytoskeleton are mediated by SUN-KASH nuclear-envelope bridges. *Annu. Rev.*
Cell Dev. Biol. **26**, 421–444.

Starr, D. A. and Han, M. (2002). Role of ANC-1 in tethering nuclei to the actin
cytoskeleton. *Science* **298**, 406–409.

Stelpflug, S. C., Sekhon, R. S., Vaillancourt, B., Hirsch, C. N., Buell, C. R., de Leon,
N. and Kaeppler, S. M. (2016). An Expanded Maize Gene Expression Atlas based
on RNA Sequencing and its Use to Explore Root Development. *Plant Genome* **9**,.

Tamura, K., Iwabuchi, K., Fukao, Y., Kondo, M., Okamoto, K., Ueda, H., Nishimura,
M. and Hara-Nishimura, I. (2013). Myosin XI-i links the nuclear membrane to the
cytoskeleton to control nuclear movement and shape in Arabidopsis. *Curr. Biol.* **23**,
1776–1781.

Thomas, J. B. and Kaltsikes, P. J. (1977). THE EFFECT OF COLCHICINE ON
CHROMOSOME PAIRING. *Can. J. Genet. Cytol.* **19**, 231–249.

Uzer, G., Rubin, C. T. and Rubin, J. (2016). Cell Mechanosensitivity is Enabled by the
LINC Nuclear Complex. *Curr Mol Biol Rep* **2**, 36–47.

Van de Vosse, D. W., Wan, Y., Wozniak, R. W. and Aitchison, J. D. (2011). Role of
the nuclear envelope in genome organization and gene expression. *Wiley*

Interdiscip. Rev. Syst. Biol. Med. **3**, 147–166.

Varas, J., Graumann, K., Osman, K., Pradillo, M., Evans, D. E., Santos, J. L. and
Armstrong, S. J. (2015). Absence of SUN1 and SUN2 proteins in Arabidopsis
thaliana leads to a delay in meiotic progression and defects in synapsis and
recombination. *Plant J.* **81**, 329–346.

Wang, S., Stoops, E., Cp, U., Markus, B., Reuveny, A., Ordan, E. and Volk, T.
(2018). Mechanotransduction via the LINC complex regulates DNA replication in
myonuclei. *J. Cell Biol.* **217**, 2005–2018.

Xu, X. M., Meulia, T. and Meier, I. (2007). Anchorage of plant RanGAP to the nuclear
envelope involves novel nuclear-pore-associated proteins. *Curr. Biol.* **17**, 1157–
1163.

Zhou, X. and Meier, I. (2013). How plants LINC the SUN to KASH. *Nucleus* **4**, 206–
215.

Zhou, X., Graumann, K., Evans, D. E. and Meier, I. (2012). Novel plant SUN–KASH
bridges are involved in RanGAP anchoring and nuclear shape determination. *J.*
Cell Biol. **196**, 203–211.

Zhou, X., Graumann, K., Wirthmueller, L., Jones, J. D. G. and Meier, I. (2014).
Identification of unique SUN-interacting nuclear envelope proteins with diverse
functions in plants. *J. Cell Biol.* **205**, 677–692.

Zhou, X., Groves, N. R. and Meier, I. (2015a). Plant nuclear shape is independently
determined by the SUN-WIP-WIT2-myosin XI-i complex and CRWN1. *Nucleus* **6**,
144–153.

Zhou, X., Groves, N. R. and Meier, I. (2015b). SUN anchors pollen WIP-WIT
complexes at the vegetative nuclear envelope and is necessary for pollen tube
targeting and fertility. *J. Exp. Bot.* **66**, 7299–7307.

1163

A

Zm00001d052955_T001 (B73v4)

TSS   5' UTR   ORF   3' UTR

Intron 1   200 bp   Intron 2

B

```
  1 MGRSLSPLLR QELDNLDKDA DSRRAAMKAL KSYARHLDSK SIPHFLAEVS DTTAAGAGAG

 61 SGMPTGGEFT ISLYEVLARV HGRNIVPQIG NIMATIMCTL SSSGGSFPLH QACSKVVPAI

121 ARHGIDPSAP DGEKAGIIAS LCRPLCSALM GGSQDGGAAS GAALCLKALV ESSNWRFASG

181 ELVNEVCLKV AGAMHDSATR SNAHMGLAMA LVKHNGLIAE AYARSIVRSG LQILDGDTAD

241 SSSQKRLSAI QMINFFMKFV DPRCLSSELG RVIDVMERCQ NDRMPFVRGA AFEASQSARN

301 IAAQKGSRHE VGTSPMVGSN FHKRREKSPC RSLWGAKRSP AASTMAAASP VQFRSPESQV

361 VDSSIMNGST LHDSPVSVGQ SSCNFDQSRR TNRRLWSNDA VDVSLKDGLF IQLCSNSNSY

421 EDDLGEVCDS EVTDANFECT NTFAGFVSPS HNGSISRDKT PSPRASDRPI NIDDVKIYST

481 PRKLLRLLQS SYDSDSASNI GQSTAKLRGL SSPDQEHKGL VISSEQMQSL HSDNKGDEMK

541 DENETIYMQN SKGRTQTLSN ADASVLSTTE AENTSIKTSP EIELNGDDVC ITSSIGKTRK

601 YRAKFTLFLS VIVIFLAIIT MLIRIENDDD SAYLVPT*
```

<u>α-helix</u>   <u>Transmembrane domain</u>   KASH domain

C

Predicted alpha solenoid scaffold structure   TM domain   KASH

-LVPT

DR1   DR2   DR3

0   50   100   150   200   250   300   350   400   450   500   550   600   650

armadillo type fold   α-helix   disordered region   TM domain   KASH domain

D

MLKS2   PP2A   Overlay

TMD

KASH domain

Figure 1

Figure 2

A

B

```
MLKS2 published      GCCAGAATGACCGCATGCCTTTCGTCCGAGGTGCTGCGTTTGAGGCATC
MLKS2 PCR 21FR(a)     GCCAGAATGACCGCATGCCTTTCGTCCGAGGTGCTGCGTTTGAGGCATC
mlks2-1 PCR 21FT(b)   GCCAGAATGACCGCATGCCTTTCGTCCGAgagataa...
mlks2-1 PCR 21TR(c)          ...ttatctcTTCGTCCGAGGTGCTGCGTTTGAGGCATC

MLKS2 published      AGCAATTCCTATGAAGATGACTTGGGCGAGGTCTGTGACAGTGAAGTGA
MLKS2 PCR 22FR(d)     AGCAATTCCTATGAAGATGACTTGGGCGAGGTCTGTGACAGTGAAGTGA
mlks2-2 PCR 22FT(e)   AGCAATTCCTATGAAGATGACTTGGGCGAgagataatt...
mlks2-2 PCR 22TR(f)          ...ttatctcCTTGGGCGAGGTCTGTGACAGTGAAGTGA
```

C

D

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Maize pollen mother cell

A MLKS2 (normal)

F-actin
MLKS2
SUN2

NE:
ONM
INM

direct   indirect

B *mlks2-2 (mutant)*

GFP-ZmKASH in eudicot leaf

C ANCHORED

MLKS2   MLKS2ΔKASH

GFP
MLKS2
ZmSUN2
NbSUN

NE:
ONM
INM

D MOBILE

Control (lbr)   MLKS2ΔARM

Figure 9

Figure S1

GFP-MLKS2	mCherry-TUA5	Overlay

Figure S2

Table S1. Plasmid information (Gumber et al.)

	Plasmid name	Alternate lab name	Construct	Vector backbone	Antibiotic marker	Deletion
Synthetic construct Genscript	pHWBF08		eGFP-FLAG-HA-MLKS2	pUC18	Amp	none
Full length MLKS2 entry clone	MLKS2ec	pPK1F-EC	eGFP-FLAG-HA-MLKS2	pDONR221	Kan	none
Full length MLKS2 expression clone	MLKS2exp	pPK1Fexp	eGFP-FLAG-HA-MLKS2	pH7WG2	Sm/Sp	none
	MLKS2del4ec		eGFP-FLAG-HA-MLKS2del4	pDONR221	Kan	terminal 4 amino acid residues
Deletion construct entry clones	MLKS2delKASHec	pPK1FdelKASHec	eGFP-FLAG-HA-MLKS2delKASH-AA/	pDONR221	Kan	KASH domain
	MLKS2delTMec	pMLKS2delTMec	eGFP-FLAG-HA-MLKS2delTM	pDONR221	Kan	transmembrane domain onwards
	MLKS2delARMec	pMLKS2delARMec	eGFP-FLAG-HA-MLKS2delARM	pDONR221	Kan	ARM domain
	MLKS2del4exp	pPK1Fdel4exp	eGFP-FLAG-HA-MLKS2del4	pH7WG2	Sm/Sp	terminal 4 amino acid residues
Deletion construct expression clones	MLKS2delKASHexp	pPK1FdelKASH-AAAexp	eGFP-FLAG-HA-MLKS2delKASH-AA/	pH7WG2	Sm/Sp	KASH domain
	MLKS2delTMexp	pPK1FdelTMexp	eGFP-FLAG-HA-MLKS2delTM	pH7WG2	Sm/Sp	transmembrane domain
	MLKS2delARMexp	pPK1FdelARMexp	eGFP-FLAG-HA-MLKS2delARM	pH7WG2	Sm/Sp	ARM domain

Table S2. List of primers used for construct cloning (Gumber et al.)

Plasmid genera	Name	Sequence (5'-3')	Purpose
			att primers to amplify eGFP-MLKS2 for
MLKS2-EC	BPHWB8F	GGGGACAAGTTTGTACAAAAAAGCAGGCTTCATGGTCTCCAAGGGCGAGG	creating entry clone
	BPHWB8R	GGGGACCACTTTGTACAAGAAAGCTGGGTCTCAGGTCGGCACCAGGTACG	"
MLKS2ΔARM	1Fdel4F	AAGGGCCTCGTCATTTCTTCAGAGCAGATGC	To generate MLKS2delARM entry clone
	1Fdel4R	CCCCATGCCAGCCCCGGC	
			To delete terminal nucleotide bases
MLKS2Δ4*	1Fdel1F	TGAGACCCAGCTTTCTTGTAC	corresponding to LVPT
	1Fdel1R	GTACGCGCTGTCATCGTC	
			To delete terminal 42 bases corresponding to TMD and add 9 bases corresponding
MLKS2ΔKASH*	1Fdel2F	CGCATGAGACCCAGCTTTCTTG	to 3 alanines
	1Fdel2R	GCAGCGATCAGCATCGTAATGATAG	
			To delete bases TMD
MLKS2ΔTM*	1Fdel3F	TGAGACCCAGCTTTCTTGTACAAAG	onwards
	1Fdel3R	CTTCGCGCGGTACTTCCG	

Table S3. List of genotyping primers (Gumber et al.)

Primers used for genotyping

Primer name	Alternate lab name	Sequence 5'-3'
gF1	mu1038603F1	GGAGGCGTACGCGAGGTCCA
gR1	mu1038603R1	GAGGCTCCGGCATGGGCTCT
gF2	mu1056535F2	ATTCATCCATCATGAACGGCAGCAC
gR2	mu1056535R2	CTGGGACTGGGAGTCTTGTCCCTTG

Primers used for qRT-PCR

qF1	qS2F1	AAGGGACAAGACTCCCAGTCC
qR1	qS2R1	GAGCTTTGAAGCAATCGGAGA
qF2	qS2F2	CCACTACCGAGGCTGAGAACA
qR2	qS2R2	TGGTCTTCCCTATGCTGCTTG
qF3	qS2F3	AAGATGACTTGGGCGAGGTCT
qR3	qS2R3	GAACCATTGTGGCTTGGTGAC