Understanding requires tracking: noise and knowledge interact in bilingual comprehension

Authors: Esti Blanco-Elorrieta[1,2] *, Nai Ding[3], Liina Pylkkänen[1,2,4], David Poeppel[1,5,6]

Affiliations:

[1]Department of Psychology, New York University, New York, NY, USA.

[2]NYUAD Institute, New York University Abu Dhabi, Abu Dhabi, UAE.

[3]Speech, Cognition and Neuroscience Department, Zhejiang University, Zhejiang Sheng, China.

[4]Department of Linguistics, New York University, New York, NY, USA.

[5]Department of Neuroscience, Max Planck Institute for Empirical Aesthetics, Frankfurt, Germany.

[6]Center for Neural Science, New York University, New York, NY, USA.

*Correspondence to: eb134@nyu.edu.

Abstract: Understanding speech in noise is a fundamental challenge for speech comprehension. This perceptual demand is amplified in a second language: it is a common experience in bars, train stations, and other noisy environments that degraded signal quality severely compromises second language comprehension. Through a novel design, paired with a carefully selected participant profile, we independently assessed signal-driven and knowledge-driven contributions to the brain bases of first versus second language processing. The neurophysiological data show that the locus of bilinguals' difficulty in understanding second language speech in noisy conditions arises from a failure to successfully perform a basic, automatic, low-level process: cortical entrainment to speech signals above the syllabic level.

Main Text: Speaking more than one language is the norm for the majority of the world's population (*1, 2*), and multilingualism is increasing notably (*3*). Although language proficiency can improve remarkably through exposure over the years, even to the point of reaching native-like proficiency, there is a familiar phenomenon that remains challenging throughout the life of a bilingual individual: in noisy environments, comprehension is hard in a second language - but seems relatively effortless in a first. Our understanding of the computational and neural foundations of this ubiquitous phenomenon is rather limited. A few hypotheses have attempted to account for this experience (*4, 5, 6*), and although somewhat different in scope, they have all proposed post-lexical sources for this effect. Here we tested an alternative possibility that shifts the locus of the impairment from post-lexical to pre-lexical processes. Specifically, we investigated whether the source of this persistent difficulty may lie in an automatic, lower level process: the neurophysiologically well-established concept of neural entrainment to speech (*7, 8*).

Working with bilingual (Mandarin Chinese, American English) participants with varying proficiency levels, we recorded electrophysiological (MEG) responses to speech at different signal-to-noise (SNR) ratios. We discovered that the neural responses that track the physical speech rhythm are affected by noise – but not by language proficiency. In contrast, responses tracking linguistic structure reflect the interaction between noise and knowledge. Hence, contrary to previous research suggesting that greater availability of top-down linguistic information may account for the difference in L1 versus L2 comprehension (*9*), the data show that an automatic lower-level mechanism, oscillatory tracking of speech, also contributes to the prevalent effect of impoverished comprehension of L2 speech in noise.

We collected MEG data from 40 Mandarin-English bilinguals. Participants listened to four-syllable sentences, concatenated and isochronously presented, in English and Mandarin (cf. *10*) at 4 different levels of noise, ranging from clear speech (+15 dB SNR) to unintelligible speech in a noisy background (-15 dB SNR), in 7.5 dB SNR intervals. The presentation of these sentences was fully randomized, and each sentence consisted of 4 monosyllabic words combined to form two-word noun phrases (adjective + noun) and two-word

verb phrases (verb + noun). The combination of these two phrases resulted in four-word sentences (e.g., "big rocks block roads"; Fig. 1AB, left panels). English syllables were between 250 ms and 347 ms, Mandarin syllables between 75 and 354 ms, and were adjusted to 320 ms and 250 ms, respectively, by truncation or padding silence at the end (Fig. 1AB, top right panels). Crucially, the spectral content of the envelope of the stimuli showed clear syllabic rhythm at all levels of noise, but no phrasal or sentential modulations in the signal (Fig. 1AB, bottom right panels, Supplementary Figure 1 for stimulus spectrograms).

Fig. 1. Sample English (A) and Mandarin (B) stimuli. Monosyllabic words were presented isochronously, forming phrases and sentences. Left panels: presentation rate and syntactic structure of the stimuli. In each of A and B, top right: waveform of a sample stimulus. Bottom right: spectral intensity of the stimulus at each tested level of noise, revealing a syllabic-scale rhythm at all levels of noise, but no phrasal-rate or sentential-rate modulation.

In order to characterize quantitatively the effect of noise across L2 proficiency levels, we tested bilinguals who were (i) Mandarin Chinese native speakers with low English proficiency, (ii) Mandarin native speakers with high English proficiency (these participants lived in China until adulthood and had learned English since they were young, but only in an academic setting), (iii) balanced bilinguals (these participants had been born in Mandarin speaking countries but attended English speaking international educational institutions, and they had at least one English speaking caregiver), and (iv) native speakers of Mandarin who were English dominant (born to at least one Mandarin speaking caregiver in an English speaking country). In the last group, participants reported that their life unfolded fully in English except for at home, where they spoke Mandarin, and they reported their English to be significantly better than their Chinese. Thus, our carefully selected and tested participant sample covered the full spectrum of possible language proficiency combinations in both languages. Participants listened to sets of 10 randomly presented 4-syllable sentences in

either language and were then presented with a 1 – 4 scale onscreen. Listeners had to indicate via button press how much they understood (1 = "nothing at all" and 4 = "everything"). Note that methodological restrictions prevented the use of more standard recall tests to evaluate comprehension, since these tasks would elicit motor artifacts that would contaminate our brain signals of interest. However, the validity of this comprehension measure has been previously established by showing qualitatively comparable results between this subjective measure of comprehension and participants' performance on recall tests (*11, 12*).

Behavioral results show that the comprehension of speech as a function of SNR varied depending on the language background of the participants. Specifically, we found that for Mandarin-dominant participants with low English proficiency (Fig. 2A), the lower comprehension of English compared to Mandarin was constant across all levels of noise ($p < .001$). However, Mandarin speakers with high English proficiency understood both languages equally in clear speech (15 dB $p = .09$; 7.5 dB $p = .14$), but their comprehension of English was severely impaired in noisy conditions (-7.5 dB and -15 dB $p < .001$; Fig. 2B). Balanced bilinguals were equally influenced by noise in both of their languages, as demonstrated by the fact that their comprehension of English and Mandarin did not diverge at any level of noise (all levels $p > .1$ Fig. 2C). Finally, we found that English dominant bilinguals showed an overall impaired comprehension of Mandarin, except at the highest level of noise; in that case, participants did not report understanding in either language (15 dB; 7.5 dB, -7.5 dB $p < .001$; -15 dB $p = .12$; Fig. 2D). These behavioral results reveal distinct interactions between noise and comprehension, contingent on language knowledge.

Fig. 2. Comprehension performance for all participant groups as a function of noise, averaged across subjects (shaded area: 95% C.I.). Below each graph, we report the average age of acquisition (AoA), self-reported proficiency (Prof.$_{self}$) and English proficiency score in the Woodcock language questionnaire (Prof.$_{test}$) for the participants in each group.

While participants listened to the sentences, we recorded neural activity with MEG (recordings for balanced bilinguals had to be interrupted due to environmental noise and could not be analyzed). Cortical oscillations have been proposed to be likely candidates for segmentation of continuous speech (*13, 14, 15*). We performed analyses aimed to quantify how entrainment to different linguistic levels (words versus phrasal structures) was disrupted by noise, on the one hand, and language proficiency, on the other. The MEG responses were transformed into the frequency domain, and we retained for subsequent analysis the 5 neural response components that were the most consistent over trials, as identified by spatial filters (see Methods). Results revealed the distinct influence of noise on the cortical tracking of different linguistic levels. In particular, participants of all proficiency combinations tracked the syllabic rhythm at all levels of noise, and the amplitude of this tracking response decreased as noise increased (Fig. 3; 3.2 Hz response in upper panels, 4 Hz response in lower panels). However, the tracking of phrasal structure was heavily dependent on language proficiency and comprehension (Fig. 3; 1.6 Hz response in upper panels, 2 Hz response lower panels).

Specifically, with regard to speech rhythm tracking, Mandarin speakers with low English proficiency tracked syllabic rhythm in Mandarin (15 dB $p < .001$; 7.5 dB $p = .002$; -7.5 dB $p = .001$; -15 dB $p = .008$) and

English (15 dB *p* < .001; 7.5 dB *p* < .001; -7.5 dB *p* = .003; -15 dB *p* = .002); as did Mandarin native speakers with high English proficiency (Mandarin: 15 dB *p* = .04; 7.5 dB *p* = .04; -7.5 dB *p* = .04; -15 dB *p* = .03; English: 15 dB *p* < .04; 7.5 dB *p* = .04; -7.5 dB *p* = .03; -15 dB *p* = .04) and English dominant speakers (English: all noise-levels *p* < .001; Mandarin: 15 dB and 7.5 dB *p* < .001; -7.5 dB *p* = .021; -15 dB *p* = .01).

In contrast, there was a clear disparity in the tracking response to phrase-level structure across languages depending on the proficiency combinations of the participants. Mandarin dominant speakers with low English proficiency did not track English phrases (15 dB *p* = .33; 7.5 dB *p* = .52; -7.5 dB *p* < .1 and -15 dB *p* = .21; Fig. 3A) although they did track phrases in Mandarin at all levels of noise (15 dB, 7.5 dB and -7.5 dB *p* < .001) except during pure noise (-15 dB *p* = .53; Fig. 3D). As proficiency in English increased, so did the tracking of the English phrases. Mandarin speakers with high English proficiency did track phrases at the clearest levels of speech in English (15 dB *p* < .04; 7.5 dB *p* = .02) although not at the two noisier levels (-7.5 dB *p* = .16 and -15 dB *p* = .22; Fig. 3B); while also being able to track Mandarin phrases at all levels of noise (15 dB *p* = .04; 7.5 dB *p* = .04; and -7.5 dB *p* = .02) except during pure noise (-15 dB *p* = .15; Fig. 3E). Finally, English dominant speakers showed phrasal tracking of English phrases at 15 and 7.5 dB, as did the Mandarin dominant speakers with high English proficiency (*p* = .04 and *p* = .002, respectively). But crucially, the speakers with higher English proficiency were additionally able to track phrases at -7.5 dB (*p* = .04; Fig. 3C). Hence the increase in English proficiency was accompanied by tracking of phrasal structures at higher levels of noise. In contrast, these same English dominant speakers whose Mandarin was mildly worse were only able to track phrases in Mandarin at the two clearest levels of speech (15 dB (*p* = .03) and 7.5 dB (*p* = .01)) but not at -7.5 dB (*p* = .09) or -15 dB (*p* = .1) like Mandarin dominant speakers had done (Fig. 3F).

English stimuli

Mandarin stimuli

Fig. 3. MEG-derived neural response spectra for each language group (Mandarin dominant – low English (n=16), Mandarin dominant – high English (n=12) and English dominant – high Mandarin (n=12). Solid lines indicate average response; shading is the 95% C.I. Spectral peaks at corresponding frequencies reflect whether there was neural tracking of syllabic or phrasal rhythms at a given level of noise. Frequency bins with significantly stronger power than two neighbors on each side are marked with a * of the corresponding color (p < 0.05, paired one-sided t-test, FDR corrected).

These results build on previous findings reporting an effect of intelligibility on cortical tracking of speech (*12, 16, 17*), but reveal a more complex and informative pattern than previously known. Specifically, we show (i) that not all levels of entrainment are affected equally by noise and (ii) that not only the physical properties of the stimuli but also the language proficiency of the listener affects the degree of entrainment to different linguistic structures.

Finally, we performed analyses to identify from where in the cortex the reported activation patterns were emerging. We source-localized the same 5 MEG components submitted to the frequency-domain analysis and compared neural responses to English and Mandarin at -7.5 dB, as this was the SNR at which language proficiency clearly determined the presence or absence of phrasal entrainment. The analysis revealed that activity in areas surrounding the auditory cortex elicited increased activity in response to the better-understood language. Furthermore, this increase widened commensurate with the imbalance between languages (Fig. 4). This is consistent with previous research that localized intelligibility effects (*17, 18*).

Fig. 4. Whole brain source localization of the 5 neural response components most consistent over trials, identified with a Denoising Source Separation (DSS) technique. The whole brain images show the result of subtracting the average activity elicited by English sentences (at -7.5 dB) from the average activity elicited by Mandarin sentences (at -7.5 dB), averaged across subjects. Activity is displayed in noise-normalized statistical parametric maps (dSPM; blue indicates higher activity for English stimuli, red for Mandarin stimuli).

The combined behavioral and neurophysiological data we present capitalize on recent findings and illuminate speech and language processing in new mechanistic ways. In particular, it has been shown that

neural entrainment to speech signals enables both tracking of the acoustic input, principally the amplitude envelope, but also higher order structure building operations that are not accessible from the physical input alone (Fig.1, Fig.3; *10, 14, 18*). We advance current understanding in two important ways, deriving from the parametric nature of the experimental design, in which we concurrently vary the quality of the speech signal (Fig.1) and the language proficiency of the listeners (Fig.2).

First, the paradigm allows us to establish that the language knowledge of the listener determines the required spectral information required for speech recognition (cf. *19*). Next, we show at which processing level (cf. Fig. 1) differential knowledge of language is especially influential. There is a minimum level of SNR (-7.5 dB in native language and 7.5 dB in the non-native language; Fig. 3) necessary to facilitate phrase-level tracking, which in turn forms the basis for structure building. Listeners who track their L1 well at -7.5 dB fail at tracking their L2 at that same SNR, suggesting that the tracking impairment is not due to peripheral causes but implicates higher-order limitations. Second, our data shed new light on the conceptualization of multilingual language comprehension. Contrary to previous accounts positing that the source of the prevalent bilingual impairment to comprehend speech in noise emerges from lexical or syntactic deficiencies in this language (*4, 5, 6*), our results suggest that the impairment emerges at a much lower-level, and perhaps largely automatic, process. Mechanistically, comprehension may be enhanced by a feedback loop such that bottom-up rhythmic structure and top-down information mutually aid the prediction and processing of upcoming signals (*17*). In bilinguals, both of these processes may be impaired: While previous research has focused on the top-down aspect (*9*), we show that the impoverished comprehension of speech in noise by L2 learners is *also* determined by a bottom-up disruption in the oscillatory entrainment to linguistic structures.

Our results characterize the minimum SNR requirements for neural entrainment to different linguistic structures, specify the differential influence of noise and knowledge on syllabic and phrasal tracking, and reveal a neurophysiological effect that plausibly underlies the widely experienced phenomenon of compromised comprehension of second language speech in noisy environments.

References and Notes:

Craik, F. I., & Bialystok, E. (2006). Cognition through the lifespan: mechanisms of change. Trends in Cognitive Sciences, 10(3), 131-138.

U. S. Census Bureau (2010) The 2011 Statistical Abstract. Languages Spoken at Home by Language: 2008, Table 53.

Cenoz, J., Nunes, P., Riganti, P., Onofri, L., Puzzo, B., & Sachdeva, R. Benefits of linguistic diversity and multilingualism (2006). Sustainable Development in a Diverse World (SUS. DIV) position paper, Research Task, 1. European Comission: Community Research and Development Information Service.

Ferreira, F., Engelhardt, P. E., & Jones, M. W. (2009). Good enough language processing: A satisficing approach. In *Proceedings of the 31st Annual conference of the Cognitive Science Society. Austin: Cognitive Science Society*.

Hahne, A., & Friederici, A. D. (2001). Processing a second language: Late learners' comprehension mechanisms as revealed by event-related brain potentials. Bilingualism: Language and Cognition, 4(2), 123-141.

Hervais-Adelman, A., Pefkou, M., & Golestani, N. (2014). Bilingual speech-in-noise: Neural bases of semantic context use in the native language. Brain and language, 132, 1-6.

Buzsáki, G., & Draguhn, A. (2004). Neuronal oscillations in cortical networks. Science, 304(5679), 1926-1929.

Lakatos P, Karmos G, Mehta AD, Ulbert I, Schroeder CE. Entrainment of neuronal oscillations as a mechanism of attentional selection, Science, 2008, vol. 320 (pg. 110-113)

Hervais-Adelman, A., Pefkou, M., & Golestani, N. (2014). Bilingual speech-in-noise: Neural bases of semantic context use in the native language. Brain and Language, 132, 1-6.

Ding, N., Melloni, L., Zhang, H., Tian, X., & Poeppel, D. (2016). Cortical tracking of hierarchical linguistic structures in connected speech. Nature Neuroscience, 19(1), 158.

Ghitza, O. (2012). On the role of theta-driven syllabic parsing in decoding speech: intelligibility of speech with a manipulated modulation spectrum. Frontiers in psychology, 3, 238.

Doelling, K. B., Arnal, L. H., Ghitza, O., & Poeppel, D. (2014). Acoustic landmarks drive delta–theta oscillations to enable speech comprehension by facilitating perceptual parsing. Neuroimage, 85, 761-768.

Gross, J., Hoogenboom, N., Thut, G., Schyns, P., Panzeri, S., Belin, P., & Garrod, S. (2013). Speech rhythms and multiplexed oscillatory sensory coding in the human brain. PLoS Biology, 11(12), e1001752.

Luo, H., & Poeppel, D. (2007). Phase patterns of neuronal responses reliably discriminate speech in human auditory cortex. Neuron, 54(6), 1001-1010.

Zoefel, B., Archer-Boyd, A., & Davis, M. H. (2018). Phase Entrainment of Brain Oscillations Causally Modulates Neural Responses to Intelligible Speech. Current Biology.

Park, H., Ince, R. A., Schyns, P. G., Thut, G., & Gross, J. (2015). Frontal top-down signals increase coupling of auditory low-frequency oscillations to continuous speech in human listeners. Current Biology, 25(12), 1649-1653.

Peelle, J. E., Gross, J., & Davis, M. H. (2012). Phase-locked responses to speech in human auditory cortex are enhanced during comprehension. Cerebral Cortex, 23(6), 1378-1387.

Keitel, A., Gross, J., & Kayser, C. (2018). Perceptually relevant speech tracking in auditory and motor cortex reflects distinct linguistic features. PLoS Biology, 16(3), e2004473.

Shannon, R. V., Zeng, F. G., Kamath, V., Wygonski, J., & Ekelid, M. (1995). Speech recognition with primarily temporal cues. Science, 270(5234), 303-304.

Acknowledgments: Funding: National Institute of Health grant 5R01DC05660 and Max Planck Society funding to D.P, NYUAD Institute grant G1001 to L.P., Woodcock Institute Research Award to E.B.E.

Author contributions: E.B.E and D.P designed the experiment; E.B.E collected and performed data analyses, N.D. supported data analyses; E.B.E, D.P and L.P wrote the manuscript.

Competing interests: Authors declare no competing interests. **Data and materials availability:** All data and analysis code is available upon request.

Supplementary Materials:

Participants

Fourty-three right-handed Mandarin-English bilingual individuals participated in the experiment (15 male, 28 female, mean 20; SD ± 2.45 years). In order to meaningfully characterize the effect of noise across varied second language proficiency levels, we selected Mandarin-English bilinguals with diverse language backgrounds. 16 of the participants were native speakers of Mandarin who had acquired English later in life and had always lived in a Mandarin-speaking environment (age of acquisition Mandarin = 1.31 (1.53 SD), English = 7.62 (3.5 SD)). Their self-reported oral (speaking and understanding) proficiency was 96.8% in Mandarin (4.7 SD) and 68.5% in English (7.1 SD), and their score in the Woodcock-Muñoz English language survey was 51.8%. English had never been their language for socializing, and they had rarely used it outside of

a classroom context. 12 participants were native speakers of Mandarin with high proficiency in English. They had acquired English earlier in life at international schools but had grown up in a Mandarin-speaking environment (age of acquisition Mandarin = 1.33 (0.88 SD), English = 5.91 (4.42 SD)). They had moved to the US for undergraduate education at some point in the past 3 years and had since been in an English-speaking environment. Their self-reported oral proficiency was 94.1% in Mandarin (3.2 SD) and 82.9% in English (6.4 SD), and their score in the Woodcock-Muñoz English language survey was 68.8%. The third group of participants were balanced bilinguals (n=3), who had one parent that spoke each of the languages and had learned both languages simultaneously (age of acquisition Mandarin = 2.40 (2.59 SD), English = 3.01 (2.74 SD)). Their self-reported oral proficiency was 75.7% in Mandarin (5.3 SD) and 95.5% in English (2.29 SD), and their English proficiency score was 80%. Lastly, we tested a group of English dominant speakers (n=12), who were born to at least one Mandarin speaking parent in the US (age of acquisition Mandarin = 1.83 (0.70 SD), English = 2.25 (4.24 SD)). Hence, they had learned Mandarin from birth but the dominant language in their environment and everyday use had always been English. Their self-reported oral proficiency was 76.4% in Mandarin (6.1 SD) and 95.6% in English (2.41 SD), and their score in the English language survey was 91.4%. Information about their language use and proficiency level was gathered with a modified version of the language background questionnaire of Marian and colleagues (2007; see additional materials for full language background information). All subjects were neurologically intact with normal or corrected-to-normal vision and all provided informed written consent following NYU Institutional Review Board protocols.

Stimuli

Mandarin stimuli were 50 four-syllable sentences taken from Ding, Melloni, Zhang, Tian, and Poeppel (2016; MandarinMaterials, Supplementary Table 1, *Four-syllable sentences*), in which the first two syllables formed a noun phrase and the last two formed a verb phrase. The combination of the noun and the verb phrase formed a complete sentence. All syllables were presented isochronously, lasted between 75 and 354 ms (mean duration 224 ms), and were adjusted to 250 ms by truncation or padding silence at the end. Each trial consisted of the sequential presentation of 10 of these sentences, and no acoustic gaps were inserted between sentences. For this reason, the intensity of the stimulus as shown by the sound envelope, only fluctuated at the syllabic rate (Fig. 1, bottom right panel).

English materials consisted of 60 four-syllable sentences. Each syllable was a monosyllabic word, and all sentences had the following syntactic structure: adjective/pronoun + noun + verb + noun. All syllables were between 250 and 347 ms in duration, and were adjusted to 320 ms by padding or truncation. For both English and Mandarin, a 25-ms cosine window smoothed the offset of each syllable. All sentences in English and Mandarin are displayed in Supplementary materials.

We embedded all sentences in 4 different levels of white noise. We first measured the power of the sentences in isolation and then added white noise to reach the desired output signal-to-noise ratio in dB. The SNR levels ranged from +15 dB (clear speech) to -15 dB (unintelligible speech in noise) in 7.5 dB intervals. Although ideally (and eventually) babbling noise may be a better background noise to mirror the type of noise experienced in real life, white noise was selected for a first characterization to avoid confounding semantic, phonological, or language interference effects that went beyond the effects of noise qua noise. Participants heard 160 sentences at each noise level for each language, and the experiment took approximately one hour to complete. Across participants, each four-syllable sentence was presented an equal number of times at each level of noise.

Procedure

Before the MEG recording, each subject's head shape was digitized using a Polhemus dual-source handheld FastSCAN laser scanner. MEG data were collected in the Neuroscience of Language Laboratory at NYU using a whole-head 157 channel axial gradiometer system (Kanazawa Institute of Technology, Kanazawa, Japan) as subjects lay in a dimly lit, magnetically shielded room. Trials began with the binaural auditory presentation of the stimuli. After listening to a set of 10 sentences participants were presented with a scale from 1 – 4 onscreen, and they were asked to indicate via button press how much they had understood (1 being "nothing at all" and 4 being "everything"). After the button press the next trial began. Following the

MEG recording, all participants completed the Woodcock-Muñoz Language Survey to evaluate their proficiency in English. Subjects completed the first four parts of this survey, aimed to assess their oral, listening, reading, and writing skills. The completion of this test took around 45 minutes.

Data acquisition and preprocessing

MEG data were recorded at 1000 Hz (200 Hz low-pass filter), noise reduced via the continuously adjusted least-squares method (Adachi et al., 2001) in MEG Laboratory software (Yokogawa Electric and Eagle Technology) and epoched from beginning to end of the auditory stimulus. The MEG responses were decomposed into components using a Denoising Source Separation technique (DSS; de Cheveigné & Simon, 2008; for a detailed explanation see Ding et al., 2016), and the first 5 components were kept for analysis and projected back into sensor space. This technique decomposes MEG recordings to extract the neural response components that are consistent over trials, and it was applied to accurately estimate the strength of neural activity phase-locked to the stimulus. To avoid the transient response at the beginning of each trial, data were only analyzed from the beginning of the second sentence of each 10-sentence trial. Single trial responses per noise level and participant were Fourier transformed (DFT) into the frequency domain, and subsequently averaged within condition to obtain an evoked response per condition per participant.

Data were source localized with MNE-Python (Gramfort et al., 2013, 2014). To estimate the distributed electrical current image in the brain at each time sample, we used the Minimum Norm Approach (Hämäläinen and Ilmoniemi, 1994) via MNE (MGH/HMS/MIT Athinoula A. Martinos Center for Biomedical Imaging, Charleston, MA). The cortical surfaces were constructed using an icosahedron subdivision of 4 and mapping an average brain from FreeSurfer (CorTech and MGH/HMS/MIT Athinoula A. Martinos Center for Biomedical Imaging) to the head-shape data gathered from the headscanning process. This generated a source space of 5124 points for each reconstructed surface, leaving ~6.2 mm of spacing within sources (cortical area per source = ~39 mm^2). Then, the boundary-element model method was used to calculate the forward solution. The 100 ms pre-stimulus period was used to construct the noise covariance matrix and to apply as a baseline correction. The inverse solution for each subject was then computed from the noise-covariance matrix, the forward solution, and the source covariance matrix, and was applied to the evoked response for each condition. The application of the inverse solution determined the most likely distribution of neural activity in source space. Minimum norm current estimates were computed for three orthogonal dipoles, of which the root mean square was retained as a measure of activation at that source (thus, the orientation of the dipole was free unsigned). The resulting minimum norm estimates of neural activity were transformed into normalized estimates of noise at each spatial location using the default regularization factor (SNR = 3). Hence, we obtained noise-normalized statistical parametric maps (SPMs), which provide information about the statistical reliability of the estimated signal at each location in the map with millisecond accuracy. Then, those SPMs were converted to dynamic maps (dSPMs). To quantify the spatial resolution of these maps, the point-spread function for different locations on the cortical surface was computed. The point spread is defined as the minimum norm estimate resulting from the signals coming from a current dipole located at a certain point on the cortex. The calculation of the point-spread function following the approach of Dale et al. (2000) reduces the location bias of the estimates, in particular, the tendency of the minimum norm estimates to prefer superficial currents (i.e., their tendency to misattribute focal, deep activations to extended, superficial patterns). Hence, by transforming our minimum norm estimates to dSPM, we obtained an accurate spatial blurring of the true activity patterns in the spatiotemporal maps (Dale et al., 2000).

Analyses

Behavioral data: Responses for each trial were averaged within participant for each noise level. We subsequently conducted a related samples t-test across participants to assess whether their comprehension in English and Mandarin at each noise level significantly differed.

Oscillation analyses: MEG activity for each trial was averaged within participant at each noise level in the frequency domain. For spectral peaks, a one-tailed paired t-test was used to test if the neural response across all sensors in a frequency bin was significantly stronger across participants than the average of the neighboring

four frequency bins (two bins on each side), FDR-corrected for multiple comparisons. The application of a 1000 permutation test in lieu of the original t-test revealed the same significant results.

Source localized analysis: Having obtained distinct tracking effects across languages for each participant group at -7.5 dB, we turned to the source localized data to identify from where in the cortex this activity was emerging. For this purpose, we subtracted the response to the English sentences at -7.5 dB from the response to Mandarin sentences at -7.5 dB (averaged across participants). This analysis hence revealed the localization of the oscillatory analysis effect.

Supplementary Figure 1:

English stimuli

| +15 dB | +7.5 dB | -7.5 dB | -15 dB |

Mandarin stimuli

| +15 dB | +7.5 dB | -7.5 dB | -15 dB |

Supplementary Figure 1. Spectrograms for a sample stimuli sentence at each SNR. Top row: spectrograms for English stimuli. Bottom row: spectrograms for Mandarin stimuli.