Title:

Longevity Relatives Count score identifies heritable longevity carriers and suggests case

improvement in genetic studies.

Authors:

Niels van den Berg[1,2,*] , n.m.a.van_den_berg@lumc.nl

Mar Rodríguez-Girondo[3]

Kees Mandemakers[4]

Angelique A.P.O. Janssens[2]

Marian Beekman[1]

P. Eline Slagboom[1,5]

*: corresponding author

Affiliations:

[1]. Department of Biomedical Data Sciences, section of Molecular Epidemiology, Leiden

University Medical Center, Albinusdreef 2, 2333 ZA Leiden the Netherlands

[2]. Radboud Group for Historical Demography and Family History, Radboud University,

Erasmusplein 1, 6525 HT Nijmegen, the Netherlands

[3]. Department of Biomedical Data Sciences, section of Medical Statistics, Leiden University Medical

Center, Albinusdreef 2, 2333 ZA Leiden the Netherlands

[4]. International Institute of Social History, Cruquiusweg 31, 1019 AT Amsterdam

[5]. Max Planck Institute for Biology of Ageing, Joseph-Stelzmann-Str. 9b, D-50931 Cologne, Germany

*Corresponding author:

Name: Niels van den Berg

Email: n.m.a.van_den_berg@lumc.nl

Address: Albinusdreef 2, 2333 ZA Leiden the Netherlands

Phone: +31 (0) 71 5269738

Abstract

Longevity loci represent key mechanisms of a life-long decreased mortality and decreased/compressed morbidity. However, identifying such loci is challenging. One of the most plausible reasons is the uncertainty in defining long-lived cases with the heritable longevity trait amongst long-living phenocopies. To avoid phenocopies, family selection scores have been constructed but these have not yet been adopted as state of the art in longevity research. Here we aim to identify individuals with the heritable longevity trait by using current insights and a novel family score based on these insights. We use a unique dataset connecting living study participants to their deceased ancestors covering 37,825 persons from 1,326 five-generational families, living between 1788 and 2019. Our main finding suggests that longevity is transmitted for at least 2 subsequent generations only when at least 20% of all relatives are long-lived. This proves the importance of family data to avoid phenocopies in genetic studies.

Main

In contrast to the low heritability of human lifespan[1–4], human longevity is strongly heritable as illustrated by the familial clustering of survival into extreme ages[5–17]. Identifying longevity loci is important because these loci likely represent key mechanisms of a life-long decreased mortality[15,16], decreased morbidity[9,12,18] and compression of morbidity towards the end of the lifespan[19–21]. Currently, genome wide linkage and association studies (GWAS) identified a limited number of loci promoting longevity[22–31], for example the *APOE* and *FOXO3A* genes (more details can be found in current review papers[22,23,30]). However, many of the identified loci could not be replicated in independent studies as yet. In addition, the largest and most recent longevity GWAS, based on cases belonging to the top 10% oldest survivors, again only replicated association of the APOE locus[32].

One of the main reasons for the limited success of longevity genetic studies[24–26,31–34] is the uncertainty in defining the heritable longevity trait itself[1,16]. Given the increased life expectancy of the past 200 years due to non-genetic factors (improved hygiene, nutrition and medication) there are likely many phenocopies among the long-lived cases selected for our genetic studies[35,36]. The presence of phenocopies is illustrated by the increase of centenarians in the United States between 1994 and 2012 from 1 in 10,000 to 1 in 5,000[37]. To avoid phenocopies, family selection scores, such as the Family Longevity Selection Score (FLoSS) and the Family Excess Longevity (FEL) score have been constructed[38,39]. The use of such scores is substantiated by novel studies which showed that that including family history information can provide valuable information about an individual's genetic liability for a trait and is likely to increase the power to detect genetic[40–42]. The scores focus, in different ways,

on selecting multiple family members with the same trait[15,38,39,43,44] and usually focus on a

single group of relatives, such as parents[15,43] or siblings[39] of cases.

As the definition of heritable longevity was not yet established, the construction and

application of the family selection scores have not yet been adopted as state of the art in

longevity research. As such, the majority of genealogical[5,6,10–14,45] and genetic studies[24–26,31–

34] focus only on single, and thus including sporadic, long-lived individuals (singletons), with

some exceptions focusing for example on parental age[28,29] or multiple siblings[7,25]. In previous

work, we showed that longevity defined as top 10% survivors or more extreme is

transmitted to subsequent generations[16]. With this, a consistent definition of longevity was

provided that is also adopted in the largest longevity GWAS up to now[32]. In addition, we

showed that every additional long-lived relative independently contributes to the survival

advantage of study participants, according to their genetic distance[16]. As such, there is room

to incorporate these novel insights into family selection scores to gain knowledge about the

extent that longevity needs to cluster in families in order to include individuals with the

heritable longevity trait and increase the power of genetic studies.

Here, we aim to establish the proportion of ancestral blood relatives that should be long-

lived (top 10% survivors of their birth cohort or more extreme) in order to observe a survival

advantage in their descendants and incorporate these insights into a novel family score to

define cases with the heritable longevity trait for inclusion in genetic studies. For our

analyses we use the data available in the Historical Sample of the Netherlands (HSN) for the

period between 1860 and 1875 which is based on Dutch citizens[46–48]. We primarily identify

cases who died beyond 80 years (N=884, on average top 10% survivors of their birth cohort),

allowing us to select on more extreme ages at death, and controls who died between 40 and

59 years (N=442). We extend this filial (F) 1 generation data with a parental and 3

descendant generations of individual life course and mortality data and refer to the data as

the HSN case/control dataset. We subsequently exclude groups with high rates of missing

mortality information and where the majority was still alive (Supplementary Figure 4). This

study covers 37,825 persons from 1,326 three-generational families (F1-F3) and contains F1

index persons (IPs), 2 consecutive generations of descendants (F2-F3) and 2 generations of

spouses (F2-F3) (Table 1). The dataset is unique in that it covers multiple generations and

connects   alive   persons   to   at   least   two   generations   of   deceased   ancestors.

Results

Outline

We analyzed the data across multiple steps (Supplementary Figure 5) in two phases. In the first phase, we used Standardized Mortality Ratios (SMRs) to compare the transmission of longevity for cases (died beyond 80 years) and controls (died between 40 and 59 years) as defined in the original approach (Figure 1A), focusing on the F1 index persons (IPs) and two generations of descendants.

In the second phase of our study (the combined approach), we combined original cases and controls and their descendants into one combined group and focused on the survival of the F3 descendants in relation to their F2 and F1 ancestral family members (Figure 1B). First, we constructed the Longevity Relatives Count (LRC) score. We used the LRC score to investigate the proportion of long-lived (top 10% survivors of their birth cohort) F1 and F2 ancestors required for F3 descendants to express a survival advantage compared to members of the same birth cohort and sex (family method, Figure 1B). On the basis of these observations we defined a new case and control group in F3, where we labeled F3 descendants with ≥30% long-lived ancestors as family cases and those without long-lived ancestors as family controls. Subsequently, these F3 family cases and controls were compared for their survival, that of their spouses (to investigate environmental influences), and for survival differences with the F3 descendants, selected to have at least one (singleton) long-lived ancestor or at least one average-lived ancestor. This means that they could have more than 1 long or average lived ancestor but we actively selected for the presence of only 1 such ancestor. Supplementary Figure 3A provides a conceptual overview of this selection. To this end, we

selected either F3 descendants with at least one top 10% grandparent, at least one top 10%

parent, or with grandparents who died between 40 and 59 years (their children (parents)

resembled the general population). In a final step, we focused on the F3 descendants with at

least one long-lived parent and calculated LRC scores within this F3 group to determine if

parents transmitted their longevity more frequently if they were part of a long-lived

(LRC≥0.30) family (Figure 1B). The analysis steps are summarized in Supplementary Figure 5

and an overview of the available data per group and generation is shown in Table 1.

Longevity is transmitted in the case group and not in the control group

Focusing on the original approach (Figure 1A), we determined to what extent longevity is

transmitted in the original case and the control group by estimating SMRs per generation for

all cases and controls separately. Table 2 shows that F1 cases had a similar survival pattern

to birth cohort members of the same sex, indicating that they resemble a representative

group of random Dutch persons aged ≥ 80 years and born between 1860 and 1875. The SMR

for the descendants of the cases (F2 case descendants) was 0.87 (95%CI=0.84-0.89),

indicating 13% less deaths than expected based on individuals from a similar birth cohort

and sex. From here we refer to this as 13% excess survival (or, if appropriate, excess

mortality) compared to the general population. The descendants of controls (F2 control

descendants) had a similar survival pattern to the general population (SMR=1.01

(95%CI=0.96-1.05)). The spouses of the F2 case and control descendants surprisingly also

showed a pattern of excess survival (SMR$_{case\_F2spouses}$=0.89 (95%CI=0.85-0.94) and

SMR$_{control\_F2spouses}$ =0.9 (95%CI=0.83-0.97)). Next we observed 14% (95%CI=11%-16%) excess

survival compared to the general population for F3 descendants of the F1 cases, whereas F3

control descendants resembled the general population (SMR=0.96 (95%CI=0.93-1.00)) just as

observed in the F2 generation. The spouses of both F3 groups resembled the general

population (SMR$_{case\_F3spouses}$=1.00 (95%CI=0.95-1.05) & SMR$_{control\_F3spouses}$=1.07 (95%CI=0.99-

1.15)). We conclude that two descendant generations of cases, who belong on average to

the top 10% survivors, have 13-14% excess survival compared to the general populations

and that the descendants of controls resemble the general population.

To explore to what extent the survival of F2 and F3 descendants depends on the extremity of

the longevity of their parents, we calculated SMRs for F2 and F3 case and control

descendants with increasing parental longevity (for example, a parent belonged to the top

10%, 5%, or 1% survivors). We observed that the SMR decreased in descendants when

defining parental longevity in terms of more extreme survival percentiles. This was the case

for descendants of both the IP cases and controls although the effects were stronger in the

descendants of the cases, especially in F3, since this group is now selected to have long-lived

parents and grandparents (Supplementary table 1). This illustrates that selection on single

long-lived persons belonging on average to the top 10% survivors, as we did for the IP

selection, leads only to a modest transmission of longevity in two generations (max 14%).

Likely, the control group includes misclassified persons of which the descendants do live

longer, whereas the case group includes long-lived persons that do not transmit longevity to

their descendants (potentially these are phenocopies). Such misclassification can jeopardize

genetic studies immensely. To be able to evaluate living persons as potential carriers of the

heritable longevity trait in genetic studies, we constructed and validated a familial longevity

score.

Constructing the Longevity Relatives Count score

We now look at the HSN data from a different perspective, the combined approach (Figure 1B). In the combined approach we consider the F3 generation as the focal point of the pedigree, instead of the F1 generation, as was the case in the original approach. To identify individuals with the heritable longevity trait, we constructed the LRC score.

$$LRC_i = \frac{\text{weighted number of top 10\% ancestors}}{\text{weighted total number of ancestors}} = \frac{\sum_{k=1}^{N_i} w_k \cdot I(P_k \geq 0.9)_i}{\sum_{k=1}^{N_i} w_k}$$

Where k=1,…,N_i are all the available ancestral blood relatives (from here: ancestors) of F3 descendant i used to build the score (parents, aunts and uncles and grandparent of the F3 descendants, Figure 1B), P_k is the sex and birth year-specific survival percentile, based on lifetables, of ancestor k, and I($P_k \geq 0.9$) indicates if ancestor k belongs to the top 10% survivors. $\sum_{k=1}^{N_i} w_k$ is the weighted total number of ancestors of F3 descendant i. The relationship coefficients are used as weights w_k. The LRC score indicates the proportion of ancestors that has become long-lived. For example, an LRC of 0.5 indicates 50% long-lived ancestors (see methods for a more detailed and general description of the LRC score).

Longevity is transmitted when at least 20% of all ancestors are long-lived

To determine what proportion of long-lived ancestors could be associated with the survival of F3 descendants, we calculated LRC scores for all F3 descendants and subsequently defined 9 mutually exclusive LRC groups (g) of F3 descendants: LRC_g1=0, LRC_g2=[>0 &

<0.1], LRC_g3=[≥0.1 & <0.2], LRC_g4=[≥0.2 & <0.3], LRC_g5=[≥0.3 & <0.4], LRC_g6=[≥0.4 &

<0.5], LRC_g7=[≥0.5 & <0.6], LRC_g8=[≥0.6 & <0.7], LRC_g9=[≥0.7 & ≥1.0]. For each group of

F3 descendants we explored whether they have a survival benefit compared to the general

population by estimating SMRs (Figure 2). F3 descendants without any long-lived ancestors

(LRC score of 0) had a survival pattern that resembled the general population (SMR=0.97

(95%CI=0.93-1.01)). Similarly, we observed a survival pattern that resembled the general

population for F3 descendants with up to 20% long-lived ancestors (group 2 and 3,

SMR=0.97 (95%CI=0.91-1.04) and SMR=0.95 (95%CI=0.91-1.00) respectively). This shows

that the long-lived ancestors of group 2 and 3 F3 descendants were likely phenocopies

instead of genetically enriched long-lived persons. We observed a pattern of excess survival

for F3 descendants with more than 20% long-lived ancestors. The weakest significant effect

was observed for group 3, with an SMR of 0.84 (95%CI=0.80-0.89) which is comparable to

the excess survival of the F3 descendants of the singleton F1 cases in the original approach

(first part of the results). The strongest significant effect was observed for group 8, with an

SMR of 0.56 (95%CI=0.45-0.69). Hence, the higher the degree of long-lived ancestors, the

lower the SMR. This indicates that the more long-lived ancestors an F3 descendant has, the

higher the level of excess survival of these F3 descendants is compared to the general

population, and the more likely that genetic effects drive the transmission of longevity.

Using the LRC score family method we defined a new case and control group in the F3

generation, which is based on the presence or absence of longevity among the ancestors of

the F3 generation and potential excess survival or mortality in the F3 generation itself

(Figure 1B). The F3 family controls include all F3 descendants without any long-lived

ancestors (LRC score of 0, N=4,166). To define the F3 family cases we chose an LRC cutoff

based on a trade-off between the size and the uncertainty, given by the sample size, of the

SMR. The F3 family cases include all F3 descendants with at least 30% long-lived ancestors

(LRC score ≥ 0.30 (N=2,526)). Even if F3 family cases are not long-lived themselves, their

survival reflects the presence of longevity of their ancestors, which is transmitted by their

parents. Similarly, F3 controls reflect the absence of longevity of their ancestors.

Supplementary Figure 1 shows the variation in lifespan of the F3 family case and control

descendants. F3 descendants with more than 0% and up to 20% long-lived ancestors (LRC

score >0 and < 0.2) did not express excess survival (N=5,340). The F3 descendants with an

LRC score ≥0.2 and < 0.30 showed some excess survival compared to the general population,

but the size of the SMR was considered too low to enter our family case definition. Hence,

we denoted them as non-classified (N=2,639).

Strong survival advantage and genetic enrichment for F3 family cases

To validate the LRC score, we investigate survival differences, measured as age at death or

last observation, between the F3 family cases and controls and used a Cox-type random

effects (frailty) regression model to adjust for within-family relations of the F3 descendants.

Figure 4 and table 3A show that F3 cases have a 25% (95%CI=18-31%) lower hazard of dying

than F3 controls, even after adjustment for sibship size, birth year, and sex. The difference

between the cases and controls became increasingly more pronounced when confining the

cases to a higher proportion of long-lived ancestors, for example an LRC score of 0.40, 0.50,

or 0.60, reflecting 40%, 50%, or 60% long-lived ancestors (Supplementary figure 2). The

strongest effect was observed for those with an LRC score ≥ 0.60 (hazard ratio (HR) of 0.62

(95%CI=0.50-0.77)). The mortality pattern for the spouses of these F3 cases resembled that

of the F3 controls (HR=0.94 (95%CI=0.82-1.07),Table 3B) and the general population

(SMR=0.92 (95%CI=0.83-1.02)). The survival of the spouses, equal to the F3 controls and the

general population, in addition to the absence of effects of environmental covariate

adjustment, indicates that environmental factors were likely of limited influence to the

observed survival benefit of the F3 cases as defined by our novel family based definition.

Hence, the observed survival benefit of F3 cases likely represents a genetic longevity

component.

Family cases live longer than those with one long-lived parent or grandparent

Next, we test if the F3 descendants with 30% long-lived ancestors (the family cases) have a

stronger survival advantage than F3 descendants with at least 1 long-lived (top 10%) parent

or grandparent. We actively selected this group of F3 descendants to have 1 long-lived

parent or grandparent, meaning that other ancestors could also be long-lived but there was

no active selection on the presence of their longevity (Supplementary Figure 3A and 3B),

hence the designation 'at least' for this group. Subsequently, we tested if F3 descendants

without long-lived ancestors (the family controls) had a similar survival pattern to the F3

descendants with parents resembling the general population (those with a grandparent who

died between 40 and 59 years). Table 4 shows that we observed 14% (95%CI=11%-17%)

excess survival compared to the general population for F3 descendants with at least one

long-lived grandparent (F1). When identifying F3 descendants with at least one long-lived

parent (F2), we observed 16% (95%CI=8%-24%) excess survival compared to the general

population. Using the family method at 30% long-lived family members to identify F3 family

cases, we observed 26% (95%CI=22%-30%) excess survival compared to the general

population and this increased to 38% (95%CI=31%-45%) when applying a 50% threshold to

the family method. For the identification of controls both methods seem to preform equally

well, with almost identical SMRs of around 1. This indicates that the F3 controls, whether

defined by having no long-lived ancestors or by grandparents dying between 40 and 50

years, have a similar survival pattern to the general population. We conclude that, at least

for cases, the family method provides a better contrast in excess survival compared to the

general population and seems to better represent the heritable longevity trait.

Since the F3 descendants with ≥ 30% long-lived ancestors have a stronger survival advantage

than those with at least one long-lived parent, it is possible to get an indication of how many

F3 descendants did not appear to have a survival advantage compared to the general

population, even though at least one parent was long-lived. This is relevant in view of case

definitions used in large genetic studies into longevity. Figure 3 and Supplementary Figure 3

show that 919 F3 descendants had a long-lived parent. Out of those 919 F3 descendants,

247 (27%) had more than 0% but less than 20% long-lived ancestors (LRC > 0 and < 0.20) and

thus as a group had an SMR that resembled the general population (Supplementary Figure

3D). The other 672 (73%) had exactly, or more than 20% long-lived ancestors (LRC ≥ 0.20)

and thus, as a group, showed excess survival compared to the general population

(Supplementary Figure 3B and C). These results suggest that if living persons are selected as

case in genetic studies on the basis of one long-lived parent, 27% of these persons is unlikely

to be a carrier of the longevity trait. Persons defined as 30% long-lived ancestors, on the

other hand would be potential carriers.

Discussion

Human longevity is heritable and clusters in specific families. Studying the familial clustering of longevity in these families is important to improve our understanding of genetic factors promoting longevity and healthy aging. The main observations supporting this are (1) In the original approach, we observed 14% excess survival of the cases compared to their birth cohort for two subsequent generations (F2-F3) while in the controls no such benefit was observed, (2) in the combined approach, the excess survival of the F3 cases compared to the general population was 26-38% depending on the proportion of long-lived family members being 30-50% and these estimates strongly overlap to the survival difference between the F3 family cases and controls based on the Cox models, (3) no excess survival as compared to the birth cohort and general population was observed for F3 controls, spouses of cases or controls and neither for F3 cases with up to 20% long-lived ancestors. The analyses in the HSN case/control dataset provides strong evidence that longevity is transmitted for at least 2 subsequent generations and only when at least 20% of all ancestors are long-lived. Moreover, the family cases seem to be genetically enriched for longevity while the controls resemble the general population. Finally, 27% of the F3 descendants showed a survival pattern similar to the general population even though they had at least one long-lived parent.

Previous family studies, usually focusing on 2 generations and single individuals, showed that siblings and children of long-lived persons lived longer than first degree ancestors of non-long-lived persons or population controls[5–7,9–15,45,49]. This knowledge about the familial clustering of longevity was utilized to construct longevity ranking scores such as the Family

Mortality History Score (FMHS)[43], the est(SE) which subsequently was developed into the

FLOSS[39,44], the Longevity Family Score (LFS) which is an adaptation to the est(SE) and the

FMHS[15], and finally a method was developed to rank individuals by the survival of their

ancestors, the Familial Excess Longevity (FEL) score[38]. The FMHS, FLOSS, and LFS all resemble

excess survival of a family (FMHS focus on parents and FLOSS and LFS focus on siblings)

compared to the general population. The FEL score focuses on excess survival, defined as the

difference between a person's attained and expected age, derived from an accelerated

failure time model. This excess survival was estimated for ancestors and from this a score

was created for individuals. Although these scores all resemble a continuous familial

estimate of a lifespan advantage and not necessarily longevity, they might be used as an

inclusion tool for cases in genetic (association) studies[39]. However, these scores are not

based on a clear longevity definition that represents the heritable longevity trait and they

always require an arbitrary and difficult to interpret decision to make a cutoff in the scores

so that they resemble longevity. In addition, the majority of the scores are not based on

ancestors and thus do not capture the full family history of longevity. As such, the scores are

not suitable to establish the proportion of family members that should be long-lived in order

to properly define long-lived cases with a heritable longevity trait and thus, increase the

power of genetic longevity studies.

To overcome these issues, we developed a novel tool based on mapping the longevity of a

person's ancestors, the LRC score. The LRC score can be used to select carriers of the

heritable longevity trait (cases) and controls who resemble the general population. Another

interesting group, which we did not address in this article, is composed of persons without

any long-lived ancestors who themselves are long-lived. It may be interesting to study

environmental factors contributing to a long and healthy life in this group. Here we used the

LRC score to construct a novel family case and control group and observed a survival

advantage for F3 case descendants, even when their parents were not necessarily long-lived,

supporting the idea that a beneficial genetic component was transmitted. Likewise, the

increase in the LRC score ≥ 20% associated with an increase in survival advantage for F3

descendants. This indicates that every additional ancestor contributes to the survival

advantage of F3 descendants and confirms our previous findings in the LINKing System for

historical demography (LINKS) data and the Utah Population Database (UPDB)[16]. This

additive pattern is not readily expected if the observations are due to non-genetic factors,

such as wealth, that cluster in families. The fact that none of the environmental confounders

(sex, birth year, and sibship size) affected the survival differences between the family cases

and controls provided additional evidence for the transmission of a genetic component. A

final indication for the genetic enrichment of the family cases is based on the observed

mortality pattern for the spouses of the family cases and controls which resembled the

family controls themselves and the general population.

We observed that F3 descendants with at least one long-lived parent had less excess survival

than a subset of these F3 descendants who had at least 30% long-lived ancestors and this

difference increased when at least 50% of their ancestors were long-lived. These results

indicate that some parents were long-lived but might not have transmitted their longevity to

the subsequent F3 generation. In fact, 27% of the F3 descendants with at least one long-

lived parent did not have an LRC ≥0.20 and, as a group, did not express excess survival.

Hence the parents of theses 27% F3 descendants were sporadically long-lived as they did not

transmit their longevity. Thus, genetic studies may benefit from a case definition, where

cases are long-lived and have at least 30% long-lived ancestors, as current genetic studies, based on long-lived cases, often not include ancestral longevity in their case selection. Even though our data did not allow for an exact misclassification analysis, studies showed that the level of phenotypic misclassification in case and control annotation has a strong inhibiting effect on the power to identify variants in genetic association studies, including GWAS[42,50–58]. Moreover, it was shown that the power to identify genetic variants decreases at an equal rate to the level of misclassification[42]. For example, a study with 95% power to detect an association based on a sample of 100 cases and controls when there are no phenotypic errors may actually have only 75% power when 20% of the cases are misclassified as controls and vice versa[42]. Interestingly, when known, methods exist to adjust for the level of phenotypic misclassification[51–53,55,59], providing opportunities for specific application in genetic longevity research.

Due to the nature of the HSN data we could not use the mortality data for the parents (F0), siblings (F1), and spouses (F1) of the F1 IPs. Mortality data was less incomplete for the F2 and F3 spouses (table 1A) but there was still a relatively large number of missing mortality data. Thus, for future studies with this dataset it might be interesting to extend the mortality information for these groups. Furthermore, life course data was only present for persons with an identified personal card or personal list (details in the methods section). Consequently, socio-economic status and religion was only available for a small part (around 15%) of the F3 descendants with an unequal share of availability between men and women. This led to the exclusion of these environmental factors from our analyses. Even though we could not adjust our models for socio-economic status and religion, it is known from other studies that those factors are not influencing the association between parental longevity and

offspring survival[16]. Similarly, previous studies showed only a minor[60] or no[16,61] influence of

early and mid-life environmental covariates, such as farm ownership, parental literacy,

parental and own occupation, and birth intervals, on the association between parental

longevity and offspring survival. We, however, cannot completely rule out that other,

unobserved non-genetic familial effects may affect our results. The observed excess survival

of F2 case and control group spouses in the original approach seem to be an exception, as

we observed a survival advantage for both groups. This is likely a form of ascertainment bias

because mortality data for this group was difficult to obtain in the Dutch Personal Records

Database, leading to an overrepresentation of high ages at death. These observations add to

the mixed results about whether spouses married to a long-lived person have a survival

advantage themselves[7,11,15–17,62].

Our results have two important implications. First, existing studies based on living study

participants who have not yet reached the ages to express longevity, but have ancestral

survival data, such as UK Biobank, can now better distinguish cases by incorporating a

liability based on the LRC score. Second, new studies would obtain a maximum power to

identify loci that promote survival to the highest ages in the population when cases are

included with at least 30% (LRC≥0.30) ancestors who belong at least to the top 10% survivors

of their birth cohort and are themselves among the 10% longest lived. More extreme

selections can be made on the survival percentile by for example focusing on the top 5% or

1% survivors, and/or on the proportion of long-lived family members, for example 50%.

However, this is not strictly necessary and might unnecessarily lead to limited sample sizes[16].

In addition, controls without any ancestors living to the top 10% survivors of their birth

cohort should be included, as their mortality pattern resembles that of the general

population. Finally, for future research it may be interesting to study the environmental

factors causing the longevity in those individuals who were long-lived but had no long-lived

ancestors. If our proposed method is consistently applied across studies, the comparative

nature of longevity studies may improve and facilitate the discovery of novel genetic

variants.

Methods

Historical Sample of the Netherlands

The Historical Sample of the Netherlands (HSN) Dataset Life Courses, Release 2010.01 is based on a sample of birth certificates and contains complete life course information for 37,137 Dutch individuals (index persons (IPs)) born in and between 1850 and 1922[46–48]. These 37,137 persons were subsequently identified in the Dutch population registers and followed in the registers throughout their entire life course[47,48,63]. The database includes information about the IPs' household, including their siblings, parents, and children, occupation at several points in time and religion. Households were only followed as long as the IP was present in that household meaning that information on kin was only partly covered[48,63]. For this study we selected 884 IPs who died at 80 years or beyond (case group) and 442 IPs who died between 40 and 59 years (control group), representing 1,326 disjoint families. IPs from both groups were born between 1860 and 1875. The case group was defined so that we would obtain a sample with overrepresentation of long-lived individuals. This was interesting since it would potentially allow to select on more extreme ages at death and still guarantee numbers reasonably large. The control group was selected to represent the mortality pattern of the general population of that time as best as possible. Individuals from both groups were selected to have an available date of birth, date of death, and at least one child should be identified. In conclusion, we identified 1,326 IPs (cases and controls), their F0 parents (N=2,652), F1 siblings (N=5,179), F2 descendants (N=7,404) and F1 spouses (N=1,409), covering 3 filial generations (F0 - F2) spanning from 1788 to 1941 (Figure 1A and Table 1). The underlying data for this specific study were released as Kees

Mandemakers and Cor Munnik, Historical Sample of the Netherlands. Project Genes, Germs

and Resources. Dataset LongLives. Release 2016.01.

Extending the HSN study

For this study we extended the pedigrees until we identified the living descendants for all

1326 families. From the population registers we know the names of all F2 descendants and

we subsequently identified the F2 descendants on personal cards (PCs) and personal lists

(PLs) which were obtained from the Dutch central bureau of genealogy (CBG). These PLs and

PCs were respectively introduced in 1939 and 1994 as the individualized and subsequently,

digitized form of the population register[48]. The cards contain similar information to the

population registers and because of privacy legislation could only be obtained for deceased

persons,   one   year   after   they   passed   away   (https://cbg.nl/bronnen/cbg-

verzamelingen/persoons kaarten-en-lijsten). Hence, from these cards we obtained similar

life course and mortality information for the F2 descendants as for the F1 IPs and we

obtained the names of their descendants (F3). We repeated this procedure until no cards

could be obtained anymore, which was at the F3 generation. Thus the F4 generation was not

identified on the PCs of PLs anymore. In conclusion, we identified and obtained information

for the F2 descendants, F2 spouses, F3 descendants, F3 spouses, and F4 descendants

(Figure1A and Table1). We will refer to this database as the HSN case/control database.

Obtaining information for the living descendants

In a final step we obtained as much mortality information as possible for the relatives of the

identified persons and we obtained addresses, as contact information for the living

descendants. This information was obtained through the Personal Records Database (PRD)

which is managed by Dutch governmental service for identity information.

https://www.government.nl/topics/personal-data/personal-records-database-brp. The PRD

contains PL information on all Dutch citizens (alive and death) and PC information is

continuously added. We were granted permission (permission number: 2016-0000364875)

to obtain the date of death, date of last observation, current living address, and identifying

information such as names of a person's father and mother to double check if the person

identified in the PRD was identical to the person in our HSN case/control database. Using the

PRD we were able to obtain addresses for F3 and F4 descendants and additional mortality

information for F2 descendants, F2 spouses, F3 descendants, F3 spouses, and F4

descendants (Figure1A and Table1). The final database covers 57,337 persons from 1,326

five-generational families (F0-F4) and contains F1 index persons (IPs), their parents (F0),

siblings (F1), spouses (F1), and 3 consecutive generations of descendants (F2-F4) and

spouses (F2-F4), connecting deceased persons to their living descendants.

Exclusion criteria and study population

Due to the nature of the source data there is a high rate of missing mortality information for

F0 parents, F1 spouses and F1 siblings, which we therefore excluded from analyses. We

further excluded F4 descendants because 92% is still alive (Table 1 and Figure 1B). The final

study population covers 37,825 persons from 1,326 three-generational families (F1-F3) and

contains F1 index persons (IPs), 2 consecutive generations of descendants (F2-F3) and 2

generations of spouses (F2-F3).

Statistical analyses

Statistical analyses were conducted using R version 3.4.1[64]. We reported 95% confidence

intervals (CIs) and considered p-values statistically significant at the 5% level (α = 0.05).

Lifetables

In the Netherlands, population based cohort lifetables are available from 1850 until

2019[65,66]. These lifetables contain, for each birth year and sex, an estimate of the hazard of

dying between ages x and x + n (hx) based on yearly intervals (n=1) up to 99 years of age.

Conditional cumulative hazards (Hx) and survival probabilities (Sx) can be derived using

these hazards. In turn, we can determine to which sex and birth year based survival

percentile each person of our study belonged to. For example: a person was born in 1876,

was a female, and died at age 92. According to the lifetable information this person

belonged to the top three percent survivors of her birth cohort, meaning that only three

percent of the women born in 1876 reached a higher age. We used the lifetables to calculate

the birth cohort and sex specific survival percentiles for all persons in the HSN case/control

study. This approach prevents against the effects of secular mortality trends over the last

centuries and enables comparisons across study populations[1,14]. Supplementary Figure 6

shows the ages at death corresponding to the top 10, 5, and 1 percent survivors of their

birth cohorts for the period 1850-1935.

Standardized Mortality Ratios

To indicate excess mortality or excess survival of groups, such as F2 case or control group

descendants in the HSN case/control study compared to Dutch birth cohort members of the

same sex, we used Standardized Mortality Ratios (SMRs). An SMR is estimated by dividing

the observed number of deaths by the expected number of deaths. The expected number of

deaths are given by the sum of all individual cumulative hazards based on the birth cohort

and sex specific lifetables of the Dutch population. An SMR between 1 and 0 indicates excess

survival, an SMR of 1 indicates that the study population shows a similar survival to the

reference population, and an SMR above 1 indicates excess mortality. The SMR can be

estimated conditional on the specific age at which an individual starts to be observed in the

study (correction for left truncation). This was necessary to avoid selection bias if individuals

in a study population were not at risk of dying before a specific age of entry.

$$SMR = \frac{observed\ number\ of\ deaths}{expected\ number\ of\ deaths} = \frac{\sum_{i=1}^{N} d_i}{\sum_{i=1}^{N} H_{t0i}(t_i|t_{0i})}$$

Where d_i=dead status (1=dead, 0=alive), H_{t0i}=sex and birth year specific cumulative hazard

based on lifetable, t_i=timing, referring to age at death or last observation, t_{0i}=liftable age

conditioning, for example from birth (t_{0i}=0), N= group sample size. Exact CIs were derived [67]

and compared to bootstrap CIs for family data [15]. Both methods provided identical CIs and

thus, to reduce the amount of computational time necessary to estimate bootstrap CIs, we

estimated exact CIs.

Longevity Relatives Count score

Based on the results of a recent study which shows that longevity is heritable beyond the

10% survivors of their birth cohort and that multiple family members, such as parents and/or

aunts and uncles, should belong to the top 10% survivors[16] we constructed a novel score

that summarizes the familial history of longevity, the Longevity Relatives Count score (LRC).

$$\text{LRC}_i = \frac{\text{weighted number of top x percentile relatives}}{\text{weighted total number of relatives}} = \frac{\sum_{k=1}^{N_i} w_k \cdot I(P_k \geq 0.9)_i}{\sum_{k=1}^{N_i} w_k}$$

521 Where k=1,…,N_i are the available relatives of individual i used to build the score, P_k is the sex

522 and birth year–specific survival percentile based on lifetables of relative k and I($P_k \geq 0.9$)

523 indicates if relative k belongs to the top 10% survivors $\sum_{k=1}^{N_i} w_k$ is the weighted total

524 number of relatives of person i. The relationship coefficients are used as weights w_k. For

525 example, persons share on average 50% of their nuclear DNA with their parents and siblings

526 and this is 25% for aunts, uncles or grandparents. Hence, in the LRC, each parent and sibling

527 contributes 0.5 to the score while each aunt, uncle or grandparent contributes only 0.25.

528 This is consistent to a previous study of us, which shows that distant longevous relatives

529 associate significantly, but less strong to a person's survival than a close long-lived relative[16].

530 The higher the score, the higher the familial aggregation level of longevity. For example, a

531 score of 0.5 indicates that 50% of a person's relatives were long-lived. We utilized the LRC

532 score to map the proportion of long-lived ancestors for all F3 descendants, select cases with

533 the heritable longevity trait and controls resembling the general population, and compare

534 the survival advantage of F3 descendants who had at least one long-lived parent to those

535 who had at least 30% long-lived descendants. The LRC scores were based on all identified

536 relatives of F3 descendants with sufficient data quality (Supplementary Figure 4 and 5).

538 **Survival analysis (Cox-type random effects regression model)**

539 To investigate the extent of a survival difference between the family F3 case and control

540 group we use a Cox-type random effects model:

$$\lambda(t_{ij}) = u_i \lambda_0(t_{ij}) \exp{(\beta Z_{ij} + \gamma X_{ij})}$$

where t_{ij} is the age at death for person j in family i. $\lambda_0(t_{ij})$ refers to the baseline hazard,

which is left unspecified in a Cox-type model. β is the vector of regression coefficients for

the main effects of interest (Z). γ is a vector of regression coefficients for the effects of

covariates and possible confounders (X). $u_i > 0$ refers to an unobserved random effect

(frailty). In all Cox models we adjust for sibship size, birth year, and sex.

Code availability

The scripts containing the code for data pre-processing and data analyses can be freely

downloaded at: https://git.lumc.nl/molepi/PUBLIC/LRCscore.

Data availability

Currently all data is cleaned and we are constructing a data description file. As soon as the

data description file is completed the data will be made freely available in a data repository

on DANS (https://dans.knaw.nl/en/front-page?set_language=en).

Competing interests

The authors declare no competing interests.

References

561   1.   van den Berg, N., Beekman, M., Smith, K. R., Janssens, A. & Slagboom, P. E. Historical

562   demography and longevity genetics: Back to the future. *Ageing Res. Rev.* **38**, 28–39

563   (2017).

564   2.   Herskind, A. M. *et al.* The heritability of human longevity: A population-based study of

565   2872 Danish twin pairs born 1870–1900. *Hum. Genet.* **97**, 319–323 (1996).

566   3.   Kaplanis, J. *et al.* Quantitative analysis of population-scale family trees with millions of

567   relatives. *Science* **360**, 171–175 (2018).

568   4.   Ruby, J. G. *et al.* Estimates of the Heritability of Human Longevity Are Substantially

569   Inflated due to Assortative Mating. *Genetics* **210**, 1109–1124 (2018).

570   5.   Houde, L., Tremblay, M. & Vézina, H. Intergenerational and Genealogical Approaches

571   for the Study of Longevity in the Saguenay-Lac-St-Jean Population. *Hum. Nat.* **19**, 70–

572   86 (2008).

573   6.   Perls, T. T. *et al.* Life-long sustained mortality advantage of siblings of centenarians.

574   *Proc. Natl. Acad. Sci.* **99**, 8442–8447 (2002).

575   7.   Schoenmaker, M. *et al.* Evidence of genetic enrichment for exceptional survival using

576   a family approach: the Leiden Longevity Study. *Eur. J. Hum. Genet.* **14**, 79–84 (2006).

577   8.   Ljungquist, B., Berg, S., Lanke, J., McClearn, G. E. & Pedersen, N. L. The Effect of

578   Genetic Factors for Longevity: A Comparison of Identical and Fraternal Twins in the

579   Swedish Twin Registry. *Journals Gerontol. Ser. A Biol. Sci. Med. Sci.* **53A**, 441–446

580   (1998).

581   9.   Terry, D. F. *et al.* Lower All-Cause, Cardiovascular, and Cancer Mortality in

582   Centenarians' Offspring. *J. Am. Geriatr. Soc.* **52**, 2074–2076 (2004).

583   10.   Willcox, B. J., Willcox, D. C., He, Q., Curb, J. D. & Suzuki, M. Siblings of Okinawan

584   centenarians share lifelong mortality advantages. *J. Gerontol. A. Biol. Sci. Med. Sci.* **61**,

585   345–54 (2006).

586   11.   Jarry, V., Gagnon, A. & Bourbeau, R. Survival advantage of siblings and spouses of

587   centenarians in 20th-century Quebec. *Can. Stud. Popul.* **39**, 67 (2013).

588   12.   Dutta, A. *et al.* Longer lived parents: Protective associations with cancer incidence and

589   overall mortality. *Journals Gerontol. - Ser. A Biol. Sci. Med. Sci.* **68**, 1409–1418 (2013).

590   13.   Gudmundsson, H., Gudbjartsson, D. F. & Kong, A. Inheritance of human longevity in

591   Iceland. *Eur. J. Hum. Genet. EJHG* **8**, 743–749 (2000).

592   14.   Sebastiani, P., Nussbaum, L., Andersen, S. L., Black, M. J. & Perls, T. T. Increasing

593   Sibling Relative Risk of Survival to Older and Older Ages and the Importance of Precise

594   Definitions of "Aging," "Life Span," and "Longevity". *Journals Gerontol. Ser. A Biol. Sci.*

595   *Med. Sci.* **71**, 340–346 (2016).

596   15.   Berg, N. van den *et al.* Longevity Around the Turn of the 20th Century: Life-Long

597   Sustained Survival Advantage for Parents of Today's Nonagenarians. *Journals*

598   *Gerontol. Ser. A* **73**, 1295–1302 (2018).

599   16.   van den Berg, N. *et al.* Longevity defined as top 10% survivors and beyond is

600   transmitted as a quantitative genetic trait. *Nat. Commun.* **10**, 35 (2019).

601   17.   Pedersen, J. K. *et al.* The Survival of Spouses Marrying Into Longevity-Enriched

602   Families. *Journals Gerontol. Ser. A Biol. Sci. Med. Sci.* **72**, 109–114 (2017).

603   18.   Westendorp, R. G. J. *et al.* Nonagenarian Siblings and Their Offspring Display Lower

604   Risk of Mortality and Morbidity than Sporadic Nonagenarians: The Leiden Longevity

605   Study. *J. Am. Geriatr. Soc.* **57**, 1634–1637 (2009).

19.   Christensen, K. *et al.* Physical and cognitive functioning of people older than 90 years:

a comparison of two Danish cohorts born 10 years apart. *Lancet* **382**, 1507–1513

(2013).

20.   Christensen, K., McGue, M., Petersen, I., Jeune, B. & Vaupel, J. W. Exceptional

longevity does not result in excessive levels of disability. *Proc. Natl. Acad. Sci.* **105**,

13274–13279 (2008).

21.   Andersen, S. L., Sebastiani, P., Dworkis, D. a., Feldman, L. & Perls, T. T. Health Span

Approximates Life Span Among Many Supercentenarians: Compression of Morbidity

at the Approximate Limit of Life Span. *Journals Gerontol. Ser. A Biol. Sci. Med. Sci.* **67A**,

395–405 (2012).

22.   Shadyab, A. H. & LaCroix, A. Z. Genetic factors associated with longevity: A review of

recent findings. *Ageing Res. Rev.* **19**, 1–7 (2015).

23.   Eline Slagboom, P., van den Berg, N. & Deelen, J. Phenome and genome based studies

into human ageing and longevity: An overview. *Biochim. Biophys. Acta - Mol. Basis Dis.*

1864, 2742–2751 (2018).

24.   Deelen, J. *et al.* Genome-wide association meta-analysis of human longevity identifies

a novel locus conferring survival beyond 90 years of age. *Hum. Mol. Genet.* **23**, 4420–

4432 (2014).

25.   Broer, L. *et al.* GWAS of longevity in CHARGE consortium confirms APOE and FOXO3

candidacy. *Journals Gerontol. - Ser. A Biol. Sci. Med. Sci.* **70**, 110–118 (2015).

26.   Willcox, B. J. *et al.* FOXO3A genotype is strongly associated with human longevity.

Proc. Natl. Acad. Sci. **105**, 13987–13992 (2008).

27.   Flachsbart, F. *et al.* Association of FOXO3A variation with human longevity confirmed

in German centenarians. *Proc. Natl. Acad. Sci.* **106**, 2700–2705 (2009).

28.   Pilling, L. C. *et al.* Human longevity: 25 genetic loci associated in 389,166 UK biobank

participants. *Aging (Albany. NY).* **9**, 2504–2520 (2017).

29.   Joshi, P. K. *et al.* Genome-wide meta-analysis associates HLA-DQA1/DRB1 and LPA and

lifestyle factors with human longevity. *Nat. Commun.* **8**, 1–13 (2017).

30.   Partridge, L., Deelen, J. & Slagboom, P. E. Facing up to the global challenges of ageing.

Nature **561**, 45–56 (2018).

31.   Sebastiani, P. *et al.* Four Genome-Wide Association Studies Identify New Extreme

Longevity Variants. *J. Gerontol. A. Biol. Sci. Med. Sci.* **72**, 1453–1464 (2017).

32.   Deelen, J. *et al.* A meta-analysis of genome-wide association studies identifies

multiple longevity genes. *Nat. Commun.* **10**, 3669 (2019).

33.   Flachsbart, F. *et al.* Immunochip analysis identifies association of the RAD50/IL13

region with human longevity. *Aging Cell* **15**, 585–588 (2016).

34.   Zeng, Y. *et al.* Novel loci and pathways significantly associated with longevity. *Sci. Rep.*

6, 21243 (2016).

35.   Oeppen, J. & Vaupel, J. W. Broken limits to life expectancy. *Science* **296**, 1029–1031

(2002).

36.   Vaupel, J. W. Biodemographic Trajectories of Longevity. *Science (80-.).* **280**, 855–860

(1998).

37.   Sebastiani, P. & Perls, T. T. The Genetics of Extreme Longevity: Lessons from the New

England Centenarian Study. *Front. Genet.* **3**, 1–7 (2012).

38.   Kerber, R. A., Brien, E. O., Smith, K. R. & Cawthon, R. M. Familial Excess Longevity in

Utah Genealogies. *Journals Gerontol. Ser. A Biol. Sci. Med. Sci.* **56**, 130–139 (2001).

39.   Sebastiani, P. *et al.* A Family Longevity Selection Score: Ranking Sibships by Their

Longevity, Size, and Availability for Study. *Am. J. Epidemiol.* **170**, 1555–1562 (2009).

40.   Hujoel, M. L. A., Gazal, S., Loh, P., Patterson, N. & Alkes, L. Combining case-control

status and family history of disease increases association power. 1–20 (2019).

41.   Liu, J. Z., Erlich, Y. & Pickrell, J. K. Case–control association mapping by proxy using

family history of disease. *Nat. Genet.* **49**, 325–331 (2017).

42.   Gordon, D. Factors affecting statistical power in the detection of genetic association. *J.*

Clin. Invest. **115**, 1408–1418 (2005).

43.   Rozing, M. P. *et al.* Familial longevity is associated with decreased thyroid function. *J.*

Clin. Endocrinol. Metab. **95**, 4979–4984 (2010).

44.   Arbeeva, L. S. *et al.* How Well Does the Family Longevity Selection Score Work: A

Validation Test Using the Utah Population Database. *Front. Public Heal.* **6**, (2018).

45.   Deluty, J. A., Atzmon, G., Crandall, J., Barzilai, N. & Milman, S. The influence of gender

on inheritance of exceptional longevity. *Aging (Albany. NY).* **7**, 412–418 (2015).

46.   Mandemakers, K. Historical sample of the Netherlands. In P. K. Hall, R. McCaa, & G.

Thorvaldsen (Eds.). in *Handbook of International Historical Microdata for Population*

Research 149–177 (2000).

47.   Mandemakers, K. https://socialhistory.org/en/hsn/hsn-releases. *HSN 2010.01 release*

(2010).

48.   van den Berg, N. *et al.* Families in Comparison: An individual-level comparison of life

course and family reconstructions between population and vital event registers.

SocArXiv (2018).

49.   Kemkes-Grottenthaler, A. Parental effects on offspring longevity—evidence from 17th

to 19th century reproductive histories. *Ann. Hum. Biol.* **31**, 139–158 (2004).

50.   Buyske, S., Yang, G., Matise, T. C. & Gordon, D. When a Case Is Not a Case: Effects of

Phenotype Misclassification on Power and Sample Size Requirements for the

Transmission Disequilibrium Test with Affected Child Trios. *Hum. Hered.* **67**, 287–292

679  (2009).

680 51. Gordon, D., Haynes, C., Yang, Y., Kramer, P. L. & Finch, S. J. Linear trend tests for case–

681   control genetic association that incorporate random phenotype and genotype

682   misclassification error. *Genet. Epidemiol.* **31**, 853–870 (2007).

683 52. Barral, S., Haynes, C., Stone, M. & Gordon, D. LRTae: improving statistical power for

684   genetic association with case/control data when phenotype and/or genotype

685   misclassification errors are present. *BMC Genet.* **7**, 24 (2006).

686 53. Rekaya, R., Smith, S., Hay, E. H., Farhat, N. & Aggrey, S. Analysis of binary responses

687   with outcome-specific misclassification probability in genome-wide association

688   studies. *Appl. Clin. Genet.* **Volume 9**, 169–177 (2016).

689 54. Ji, F., Yang, Y., Haynes, C., Finch, S. J. & Gordon, D. Computing Asymptotic Power and

690   Sample Size for Case-Control Genetic Association Studies in the Presence of

691   Phenotype and/or Genotype Misclassification Errors. *Stat. Appl. Genet. Mol. Biol.* **4**,

692   (2006).

693 55. Edwards, B. J., Haynes, C., Levenstien, M. A., Finch, S. J. & Gordon, D. Power and

694   sample size calculations in the presence of phenotype errors for case/control genetic

695   association studies. *BMC Genet.* **6**, 18 (2005).

696 56. Royall, D. R., Chiodo, L. K. & Polk, M. J. Misclassification Is Likely in the Assessment of

697   Mild Cognitive Impairment. *Neuroepidemiology* **23**, 185–191 (2004).

698 57. Bross, I. Misclassification in 2 X 2 Tables. *Biometrics* **10**, 478 (1954).

699 58. Platz, E. A., De Marzo, A. M. & Giovannucci, E. Prostate cancer association studies:

700   Pitfalls and solutions to cancer misclassification in the PSA era. *J. Cell. Biochem.* **91**,

701   553–571 (2004).

702 59. Smith, S., Hay, E., Farhat, N. & Rekaya, R. Genome wide association studies in

presence of misclassified binary responses. *BMC Genet.* **14**, 124 (2013).

60.   Danzhen You, Danan Gu & Zeng Yi. Familial Transmission of Human Longevity Among

the Oldest-Old in China. *J. Appl. Gerontol.* **29**, 308–332 (2010).

61.   Gavrilov, L. A. & Gavrilova, N. S. Predictors of Exceptional Longevity: Effects of Early-

Life and Midlife Conditions, and Familial Longevity. *North Am. Actuar. J.* **19**, 174–186

(2015).

62.   Montesanto, A. *et al.* The genetic component of human longevity: analysis of the

survival advantage of parents and siblings of Italian nonagenarians. *Eur. J. Hum.*

Genet. **19**, 882–886 (2011).

63.   Mandemakers, K. Building life courses datasets from population registers by the

historical sample of the Netherlands. in *History and Computing* **14**, 87–107 (2002).

64.   R Core Team. R: A language and environment for statistical computing. (2016).

65.   Van Der Meulen, A. *Life tables and Survival analysis.* (2012).

66.   Carolina, T., Uijvenhoven, L. & van der Laan, J. Overlevingstafels en longitudinale

analyse. *CBS* 1–25 (2009).

67.   Ulm, K. Simple method to calculate the confidence interal of a standardized mortality

ratio (SMR). *Am. J. Epidemiol.* **131**, 373–375 (1990).

Figure legends

Figure 1: Pedigree overview of the data structure

This figure illustrates the two approaches; 1. the original approach and 2. the combined

approach. The original approach refers to the case and control group based on the F1 IPs

where cases died at 80 years or older and controls died between 40 and 59 years (panel A).

Panel B shows a pedigree of the data from the perspective of F3 children (combined

approach). The combined approach refers to the dataset where we combined the cases and

controls form the original design and constructed a new case and control group in the F3

descendants. To this end, F3 descendants with ≥30% long-lived ancestors were labeled as

family cases and those without long-lived ancestors as family controls. F3 spouses were left

out of this figure but this group was used to confirm a genetic enrichment in the F3

descendants.

Figure 2: LRC score in mutually exclusive F3 descendant groups

The figure shows Standardized Mortality Ratios for all F3 descendants without missing

mortality information. The F3 descendants are grouped into mutually exclusive groups based

on the Longevity Relatives Count (LRC) score. The LRC score represents the family approach

as illustrated in figure 1B. The dark red color of group one represents F3 descendants

without any long-lived (top 10%) ancestors and are denoted as family controls. The light red

represents F3 descendants who had more than 0 and less than 20% long-lived ancestors. The

light blue colors represent the F3 descendants with 20% or more long-lived ancestors. The

dark blue color represent our cut-off point for the family case definition. Hence all F3

descendants with 30% or more long-lived ancestors were considered family cases. The beige

color of group 9 shows that this bar represents all F3 ancestors with more than 70% long-

lived ancestors as their sample size was very low, we grouped them into one group.

Figure 3: LRC score for F3 descendants with at least one long-lived parent

This center of this doughnut figure shows all F3 descendants (N=919) with at least one long-

lived (top 10%) parent, ignoring the rest of the ancestors. Thus, at least means that they

could have more than 1 long-lived ancestor but we actively selected for the presence of only

1 such ancestor. The edges of the doughnut illustrate the number and proportion of these

919 F3 descendants with at least one long-lived parent who had 1. 30% or more long-lived

ancestors (LRC ≥ 0.30) and excess survival compared to the general population (SMR < 1),

N=335 (36%) 2. between 20% and 30% long-lived ancestors (LRC ≥ 0.20 and < 0.30) and

excess survival compared to the general population (SMR < 1), N=337 (37%) and 3. between

0% and 20% long-lived ancestors (LRC > 0.20 and < 0.20) and a similar survival pattern to the

general population (SMR ~ 1), N=247(27%).

Figure 4: Survival differences between family based cases and their spouses

This figure shows the survival curve for the difference in survival between the F3 family

cases and controls. The figure is connected to Table 3A which shows the Hazard Ratios

corresponding to the difference between the two curves. Blue color represent the cases, red

color represents the controls.

Table 1: Overview study sample for groups in all generations based on the proband and F3 perspective

Role	Number	Deceased (%)	Alive (%)	Female (%)	Range Birth cohort	Mean age (sd)	Median age (sd)	missing_age (%)
Cases (Original design)								
F1 IPs	884	884 (100)	0 (0)	422 (50)	1860-1875	85.79 (4.59)	84.99 (4.95)	0 (0)
F2 descendants	4916	4405 (90)	11 (1)	2435 (50)	1879-1941	63.04 (31.11)	75.51 (17.72)	500 (9)
F2 spouses	3899	1500 (38)	16 (1)	1504 (38)	1873-1934	76.2 (15.09)	78.78 (12.83)	2383 (61)
F3 descendants	9910	4869 (49)	4146 (42)	4733 (48)	1901-1973	70.35 (19.54)	74.77 (11.38)	895 (9)
F3 spouses	3431	1289 (38)	792 (23)	1963 (57)	1900-1959	77.14 (11.31)	79.25 (10.1)	1350 (39)
F4 descendants*	9001	746 (8)	7172 (80)	3937 (44)	1922-1995	57.7 (10.68)	58.21 (9)	1083 (12)
Controls (Original design)								
F1 IPs	442	442 (100)	0 (0)	214 (48)	1860-1875	51.71 (5.71)	52.88 (6.21)	0 (0)
F2 descendants	2488	2202 (89)	1 (<1)	1217 (49)	1881-1925	58.17 (32.49)	71.72 (21.37)	285 (11)
F2 spouses	1877	690 (37)	7 (<1)	734 (39)	1875-1935	76.02 (14.77)	78.34 (13.76)	1180 (63)
F3 descendants	4761	2540 (53)	1813 (38)	2265 (48)	1904-1966	69.39 (20.38)	74.49 (11.36)	408 (9)
F3 spouses	1778	721 (41)	376 (21)	972 (55)	1893-1965	76.54 (11.5)	78.66 (10.47)	681 (38)
F4 descendants*	4710	387 (8)	3744 (80)	2099 (45)	1871-1992	57.72 (11.17)	58.37 (9.35)	579 (12)
F3 perspective (Combined design)								
F3 descendants	14671	7409 (51)	5959 (41)	6998 (48)	1901-1973	70.03 (19.82)	74.68 (11.38)	1303 (8)
F3 spouses	5209	2010 (38)	1168 (22)	2935 (55)	1893-1965	76.93 (11.38)	79.07 (10.24)	2031 (40)
F2 parents	9728	6139 (63)	23 (1)	4137 (43)	1873-1935	76.8 (13.4)	78.9 (12.31)	3566 (36)
F2 aunts & uncles	7036	6382 (91)	10 (1)	3456 (49)	1879-1941	61.81 (31.47)	74.4 (18.67)	644 (8)
F1 grandparents	1181	1181 (100)	0 (0)	560 (47)	1860-1875	74.88 (16.6)	81.94 (9.72)	0 (0)

The Cases and Controls rows provide an overview of the groups of persons from the original case/control perspective of the data, described as part a. The F3 perspective rows provide an overview of the groups of persons from the perspective of F3 descendants, described as part b. mean and missing age refer to an unknown age at death or an unknown age at last observation. For the F0 and F1 groups we assume everyone is dead because the birth cohorts date back further than 120 years. From the F2 generations we requested Personal Records Data indicating if a person was still alive or not and if not, what the date of death was. The F1 IPs are the focal persons in the pedigrees as they are selected to be 80 years or older (cases) or to have died between 40 and 59 years (controls). * indicates that the group is excluded for this study, sd refers to standard deviation.

Table 2: Standardized mortality ratios for original case and control group individuals

Role	Case group SMRs	Number (N)	Control group SMRs	Number (N)	Adjustment for right truncation
F1 IPs	1.06 (0.99-1.13)	884	NA	NA	80 years
F2 descendants	0.87 (0.84-0.89)	4416	1.01 (0.96-1.05)	2203	No adjustment
F2 spouses	0.89 (0.85-0.94)	1516	0.9 (0.83-0.97)	697	20 years
F3 descendants	0.86 (0.84-0.89)	9015	0.96 (0.93-1.00)	4353	No adjustment
F3 spouses	1.00 (0.95-1.05)	2081	1.07 (0.99-1.15)	1097	20 years

Original cases (F1 IPs) died at 80 years or older, original controls (F1 IPs) died between 50 and 69 years. If persons could not die before a specific age due to direct or indirect selection, due to for example that all persons in a group were selected to have a child an adjustment for right truncation was applied so that a fair comparison could be made with their birth cohort members. An SMR for F1 control IPs could not be estimated due to a combination of left and right truncation in the data. The lifetables can only be adjusted for right or left truncation, but not a combination between the two.

Table 3: Mortality difference between family cases and controls and their spouses

	A			B		
	N (mean)	HR (95% CI)	P-value	N (mean)	HR (95% CI)	P-value
Family based case/control group						
Control group (ref)	3714 (0.62)			3714 (0.50)		
Case group	2282 (0.38)	0.75 (0.69-0.82)	1.75e-10	2282 (0.30)	0.74 (0.68-0.80)	4.08e-12
Spouses of cases				541 (0.07)	0.94 (0.82-1.07)	3.44e-01
Spouses of controls				937 (0.13)	1.12 (1.00-1.25)	4.07e-02
Birth year	5996 (1933)	0.99 (0.98-0.99)	1.99e-05	7474 (1932)	0.98 (0.98-0.99)	1.39e-12
Sex						
Males (ref)	3133 (0.52)			3364 (0.45)		
Females	2863 (0.48)	0.56 (0.52-0.61)	<1.00e-15	4110 (0.55)	0.49 (0.46-0.53)	<1.00e-15
Sibship size						
Small - 1-2 sibs (ref)	1531 (0.26)					
Medium - 3-5 sibs	1770 (0.30)	1.17 (1.04-1.32)	8.51e-03			
Large - 6-8 sibs	927 (0.15)	1.22 (1.04-1.43)	1.21e-02			
Exceptional - 9-15 sibs	441 (0.07)	1.36 (1.09-1.68)	5.84e-03			
Single child - 0 sibs	1327 (0.22)	1.81 (1.62-2.02)	<1.00e-15			

Table 3A corresponds to the CH curves of panel a of figure 4. Means represent a mean for a continuous variable and a proportion for a categorical variable. When the p-value was lower than 1.00e-15 we indicated the P-value as <1.00*10-15. SES = socio-economic status, OCC = occupational coding scheme of 1950, CI = confidence interval, CH = cumulative hazard. P-values are estimated with cox regression. F3 children with relatives who were still alive and had no last moment of observation ≥ 100 years were removed to assure an equal comparison between cases and controls. In table 3B the spouses of cases and controls are adjusted for the fact that they could not die before the birth of at least their first child (left truncation). We adjusted for this left truncation by entering the spouses of cases and controls in the model based on the first observed death in the groups (cases: 30 years and controls: 25 years). In model A no adjustment for left truncation was necessary. In both models we adjusted for right censoring by including a censoring indicator in the cox model.

Table 4: Standardized Mortality Ratio for different F3 descendant groups

Group	SMR	N
Cases		
F3 descendant with at least one long-lived grandparent	0.86 (95%CI=0.83-0.89)	4986
F3 descendant with at least one long-lived parent	0.84 (95%CI=0.76-0.92)	852
F3 descendant with ≥ 30% long-lived ancestors (LRC ≥ 30%)	0.74 (95%CI=0.70-0.78)	2304
F3 descendant with ≥ 50% long-lived ancestors (LRC ≥ 50%)	0.62 (95%CI=0.55-0.96)	565
Controls		
F3 descendant with grandparent who died between 40 and 59 years	0.96 (95%CI=0.93-1.00)	4353
F3 descendant with no long-lived ancestors (LRC = 0)	0.97 (95%CI=0.93-1.01)	3782

Long-lived is defined as belonging to the top 10% survivors of their birth cohort. Note that the group size (N) reflects only those with a known age at death as this was necessary to estimate a standardized mortality ratio.

A

F1 Cases Controls F1

F2 Spouse Child Child Spouse Child Spouse Child Child Spouse Child F2

F3 Child Child Child Child Child Child Child Child Child Child Child Child F3

B

Combined approach

Family approach

F1 Grand parent

F2 Father Mother Father Mother Uncle Aunt

F3 IP IP IP IP IP IP

Proband line   Spouse line

N=247
(27%)

N=335
(36%)

F3 descendants with at least
one long-lived parent
N=919

N=337
(37%)

F3 descendants with 30% long-lived ancestors
(LRC ≥ 0.30) and an SMR < 1

F3 descendants with ≥ 20% and less than 30% long-lived ancestors
(LRC ≥ 0.20 and < 0.30) and an SMR <1

F3 descendants with ≥ 0% and less than 20% long-lived ancestors
(LRC ≥ 0 and < 0.20) and an SMR = 1