Competition and diversity determine vaccine impact on antibiotic resistance

evolution

Nicholas G. Davies[1*], Stefan Flasche[1], Mark Jit[1,2], Katherine E. Atkins[1,3]

1. Centre for Mathematical Modelling of Infectious Diseases; Vaccine Centre; and

Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical

Medicine, London, UK.

2. Modelling and Economics Unit, Public Health England, London, UK.

3. Centre for Global Health, Usher Institute of Population Health Sciences and

Informatics, The University of Edinburgh, Edinburgh, UK.

* Corresponding author. Address: Department of Infectious Disease Epidemiology,

Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical

Medicine, Keppel Street, London WC1E 7HT, United Kingdom. Phone: +44 20 7927

2880. E-mail: nicholas.davies@lshtm.ac.uk.

One sentence summary: Competition and diversity are key to antibiotic resistance

evolution and determine whether vaccines will prevent or increase resistant infections.

Bacterial vaccines can protect recipients from contracting potentially antibiotic-resistant infections. But by altering the selective balance between sensitive and resistant strains, vaccines may also help suppress—or spread—antibiotic resistance among unvaccinated individuals. Predicting the outcome requires knowing the drivers of resistance evolution. Using mathematical modelling, we identify competition and diversity as key mediators of resistance evolution. Specifically, we show that the frequency of penicillin resistance in _Streptococcus pneumoniae_ (pneumococcus) across 27 European countries can be explained by between-host diversity in antibiotic use, heritable diversity in pneumococcal carriage duration, or within-host competition. We use our calibrated model to predict the impact of universal pneumococcal vaccination upon the prevalence of carriage, incidence of disease, and frequency of resistance for _S. pneumoniae_. The relative strength and directionality of competition between resistant and sensitive pneumococcal strains determines whether vaccination promotes, inhibits, or has little effect on the evolution of antibiotic resistance. Finally, we find that differences in overall bacterial transmission and carriage alter predictions, suggesting that evidence-based policies for managing resistance with vaccines must be tailored to both pathogen and setting.

In an age of widespread antibiotic resistance, there is growing interest in using vaccines to prevent bacterial infections that would otherwise call for treatment with antibiotics (_1–4_). This interest arises for two main reasons: first, vaccines are effective against both antibiotic-resistant and antibiotic-sensitive bacteria; and second, successful prophylaxis removes the need for a course of antibiotic therapy that might promote more resistance (_2–5_). Over the past two decades, the use of pneumococcal conjugate vaccines (PCVs) has seemingly borne out these advantages. Administering PCVs to young children has substantially reduced disease caused by _S. pneumoniae_ (_5–8_)—a common asymptomatic coloniser of the nasopharynx which can cause pneumonia, meningitis and other infections when invasive—and has decreased demand for antibiotic therapy, largely by reducing cases of otitis media (_5, 9_). But because PCV formulations target only a fraction of the ~100 known pneumococcal serotypes, the niche vacated by PCV-targeted serotypes has been filled by non-vaccine serotypes, and overall pneumococcal carriage has rebounded to pre-vaccine levels (_10, 11_). Concomitantly, the incidence of infections

attributed to non-vaccine serotypes (*12*) and the proportion of non-vaccine-type

infections exhibiting antibiotic resistance (*5*, *13*) have risen in many settings. Concern

over serotype replacement—along with the high cost of PCV manufacturing—has

spurred the development of "universal" whole-cell or protein-based pneumococcal

vaccines protecting against all serotypes, some of which are now in early-stage clinical

trials (*14*).

However, it is unclear how universal vaccination may itself impact upon the evolution of

antibiotic resistance in *S. pneumoniae*, which is a concern given that vaccination is

unlikely to eliminate pneumococcal disease (*15*). Mathematical models can be used to

generate predictions from nonlinear transmission dynamics (*16*, *17*), but existing

models focus on serotype-specific vaccines and, even then, disagree over the expected

impact of vaccination on resistance evolution (*18–24*). Comparing and interpreting the

results of these models is hampered by the fact that none starts from a position of

recapitulating large-scale empirical patterns of antibiotic resistance. The main challenge

in replicating these patterns lies in identifying the mechanisms that maintain long-term

coexistence between sensitive and resistant pneumococcal strains across a wide range

of antibiotic treatment rates, like those seen across Europe and the United States (*25*,

26). Robust predictions of the long-term impact of vaccination on resistant

pneumococcal disease require a mechanistic understanding of these patterns.

<div align="center">**Results**</div>

Competition and diversity maintain stability in resistance evolution. A model must

be able to explain the current burden of an infectious disease before it can be used to

robustly predict the impact of interventions for managing that disease. Across Europe,

the frequency of antibiotic resistance among isolates from pneumococcal infections

shows two salient features for models to recapitulate (Supplementary Fig. 1). One is

spatial: the frequency of penicillin non-susceptibility varies between countries, and is

higher in countries where more penicillin is consumed (*27*). The other is temporal:

although in individual countries resistance fluctuates from year to year, the overall

frequency across Europe of penicillin non-susceptibility in pneumococcal isolates has

remained steady at roughly 12% since consolidated records began in 2005 (*28*). These

observations contradict simple models of resistance evolution, which predict that

intermediate frequencies of resistance cannot be stably maintained in the long term:

that is, either sensitive strains will competitively exclude resistant strains, or resistant

strains will competitively exclude sensitive strains, unless there is some mechanism

maintaining coexistence between them (*25*, *29*).

By conducting a literature search, we identify nine such mechanisms (*25*, *26*, *30–41*)

that fall into two broad classes. In one class, coexistence is maintained by

environmental or genetic diversity that prevents resistant and sensitive strains from

completely overlapping in competition. In the other class, competition between

resistant and sensitive strains is itself the stabilising factor that maintains coexistence,

because resistant and sensitive strains exhibit alternative competitive phenotypes that

afford strains a competitive advantage when rare, thus promoting negative frequency-

dependent selection for resistance. Thus, diversity and competition are two key forces

maintaining stability in resistance evolution. Of the nine identified mechanisms, we find

that four are biologically plausible for maintaining coexistence in *S. pneumoniae* (Table

1).

Four models of resistance evolution. To compare these four mechanisms, we embed

each in a shared model framework of person-to-person transmission of nasopharyngeal

pneumococcal carriage. This framework tracks the country-specific frequency of

resistance in pneumococci circulating among children under five years old, the age

group that drives the majority of pneumococcal transmission and disease (*42*, *43*). We

assume that each individual makes effective contact with another random individual at

rate β, thereby potentially acquiring a strain (either sensitive or resistant) carried by

the contacted person. With probability c, resistant strains fail to transmit, where c

represents the transmission cost of resistance (*44*, *45*). A carrier naturally clears all

strains at rate u, and is exposed to antibiotic therapy at a country-specific rate τ, which

clears the host of sensitive strains only. We assume this treatment rate is independent

of carriage status (*46*) and we do not explicitly track disease progression in hosts.

Under the "Treatment diversity" and "Pathogen diversity" models, coexistence is

maintained because diversity among hosts or among pathogens prevents resistant and

sensitive strains from fully overlapping in competition. In the "Treatment diversity" model (Fig. 1a), heterogeneity in the consumption of antibiotics between host subpopulations within a country maintains coexistence (*25*, *34*, *35*). These subpopulations could correspond to geographical regions, socioeconomic strata, host age and risk classes, or a combination of these. Provided that transmission between high-consumption (resistance-promoting) and low-consumption (resistance-inhibiting) subpopulations is not too frequent, an intermediate frequency of resistance can be maintained across the whole population. The key parameters governing coexistence in this model are κ, the variability in antibiotic consumption between subpopulations, and *g*, the relative rate at which within-country contact is made within subpopulations rather than between them.

In the "Pathogen diversity" model (Fig. 1b), pneumococci are divided into subtypes ("D-types"(*38*)) that vary in their mean duration of natural carriage. All else equal, the D-type with the longest carriage duration would be expected to competitively exclude all other strains; the model assumes that diversifying selection acting on the D-type locus keeps all subtypes in circulation. What D-types correspond to is not explicitly specified by this model, but one candidate is serotype variation. For example, if antigenic diversity is promoted by host acquired immunity to capsular serotypes, and serotypes tend to differ in their intrinsic ability to evade clearance by the immune system, then intermediate resistance can be maintained because selection for resistance tends to be greater in strains that have a longer duration of carriage (*38*). Long-lasting serotypes will tend to evolve resistance, while shorter-lived serotypes will tend not to—a pattern observed in *S. pneumoniae* (*38*) and reproduced by this model (Supplementary Fig. 2). The parameters governing coexistence in this model are *a*, the strength of diversifying selection on the D-type locus, and δ, the variability between subtypes in clearance rate.

Under the "Treatment competition" and "Growth competition" models, coexistence is maintained because of competition between sensitive and resistant strains. In these models, hosts can be co-colonised by multiple strains, but which strain a co-colonised host transmits to other potential hosts is determined by within-host competition between strains (*26*). The "Treatment competition" model (Fig. 1c) assumes that antibiotic therapy mediates within-host competition, such that when a co-colonised

host takes antibiotics (*i.e.*, at rate τ), the sensitive strains are cleared and only the

resistant strains are transmitted to other hosts. The "Growth competition" model (Fig.

1d) has both treatment-mediated and growth-mediated competition: while in the

presence of antibiotics, resistant strains still outcompete co-colonising sensitive strains,

in the absence of antibiotics, sensitive strains gradually outcompete co-colonising

resistant strains at rate *b*. We assume that there is no transmission cost of resistance in

this latter model (*i.e.*, *c* = 0); instead, the within-host growth advantage *b* of sensitive

strains accounts for the cost of resistance. In these competition models, resistant strains

have an advantage in antibiotic-mediated competition, while sensitive strains have an

advantage in growth-mediated competition. These alternative forms of within-host

competition can both promote coexistence because rare strains can more consistently

exploit a competitive advantage over common strains, thus creating negative frequency-

dependent selection for resistance (*26*). The key parameter governing coexistence in

these two models is *k*, the relative rate of co-colonisation compared to primary

colonisation.

In all four models, we assume that contact between individuals is assortative by

country, such that with probability *f*, contact is with a random person from the same

country, and with probability 1 – *f*, contact is with a random person from any country.

We implement these models using systems of ordinary differential equations. All four

models (*25*, *26*, *38*) are structurally neutral (*25*, *29*), meaning that any coexistence

exhibited by the models is accounted for by the specified biological mechanism rather

than by any bias in the logical structure of the model that generates coexistence "for

free". Additionally, while the within-host competition models capture co-colonisation

using a simplified subset of only 2 "mixed-carriage" states (S_R and R_S, Fig. 1a&b), we

have previously shown (*26*) that this is equivalent to a more complex individual-based

model with an arbitrary number of mixed-carriage states.

All four models reproduce observed patterns of resistance. The European Centre

for Disease Prevention and Control (ECDC) monitors antibiotic consumption and

resistance evolution across European countries (*13*, *28*). These data capture a natural

experiment in resistance evolution: for each monitored drug and pathogen, each

country reports a different rate of antibiotic consumption in the community and

exhibits a different frequency of resistance among invasive bacterial isolates. By fitting

models to this multi-country data set, we can potentially rule out models that cannot

reproduce the large-scale patterns that are observed. We use Bayesian inference to fit

the model-predicted equilibrium frequency of resistance to the reported frequency of

penicillin non-susceptibility in *S. pneumoniae* across 27 European countries, assuming a

50% carriage prevalence (*11*, *42*) and a carriage duration of 47 days (*47*, *48*) in children

under five years old. We begin by assuming that countries only differ by their reported

treatment rate—where we define a treatment course as equivalent to $z = 5$ defined daily

doses of penicillin—with other model parameters shared across countries.

Strikingly, each model fits equally well to the empirical relationship between resistance

and antibiotic use (all model WAICs are similar; Fig. 2a) and recovers plausible

posterior parameter distributions (Fig. 2b). That is, the empirical data do not

distinguish between the four alternative mechanisms of resistance evolution we have

identified. Later, we relax the assumption that only the treatment rate varies between

countries, allowing us to capture additional between-country variation in resistance not

explained by population-wide penicillin consumption.

Competition and diversity determine the impact of vaccination on resistant

disease. To determine the impact of universal vaccination on pneumococcal disease, we

consider three outcomes. The first is the impact of the vaccine upon the prevalence of

pneumococcal carriage. The second is the vaccine impact upon the frequency of

penicillin resistance among circulating pneumococcal strains remaining after

vaccination. The third is the impact of the vaccine upon the prevalence of resistant

pneumococcal carriage—*i.e.*, the prevalence of carriage multiplied by the frequency of

penicillin resistance. Since all four models are equally capable of recapitulating

observed patterns of penicillin resistance in *S. pneumoniae*, our aim is to determine

whether the mechanism of resistance evolution—competition or diversity—matters

when forecasting the impact of interventions for managing resistance.

We consider two alternative vaccines: an "acquisition-blocking" vaccine, which prevents

carriage from being established with probability ε_a, and a "clearance-accelerating"

vaccine, which shortens the duration of carriage by a fraction ε_c. Both vaccines reduce

pneumococcal transmission through alternative modes of host immunity that might be elicited by a whole-cell or protein-based universal pneumococcal vaccine. Analogously to naturally-acquired serotype-independent pneumococcal immunity (*49*), the protective effect of whole-cell vaccines manifests as accelerated clearance (*50*); it is unclear whether protein-based vaccines would block pneumococcal acquisition, like PCVs, or accelerate clearance (*51*). We refer to ε_a or ε_c as the vaccine efficacy, and for simplicity, we assume that all children under five years old have vaccine protection, as would be established by an infant vaccination programme rolled out across Europe. In order to compare these vaccines with an alternative intervention of antibiotic stewardship, we also evaluate the impact of reducing the rate of penicillin prescribing by a fraction ε_s.

We find that both vaccines have a similar impact upon carriage prevalence, regardless of whether competition or diversity maintains stability in resistance evolution (Fig. 3a). Specifically, as the vaccine efficacy ε_a or ε_c increases, carriage decreases, with the elimination of pneumococcal carriage occurring at a vaccine efficacy between 50 and 60%. Reducing antibiotic prescribing moderately increases pneumococcal carriage, such that carriage prevalence increases to approximately 54% across all countries when penicillin prescribing is eliminated completely.

However, the mechanism of resistance evolution has a substantial impact upon whether vaccines increase or decrease the frequency of resistance in *S. pneumoniae* in the long term (Fig. 3b). In the "Treatment diversity" and "Pathogen diversity" models, the acquisition-blocking vaccine has relatively little impact upon the frequency of resistance, because administering a universal pneumococcal vaccine to all individuals does not substantially alter the distribution of antibiotic use or of heritable variation in clearance rates. By contrast, in the within-host competition models, vaccination has a substantial impact upon resistance evolution because by reducing pneumococcal circulation, vaccines decrease the rate at which strains encounter each other within hosts, and hence strongly decrease competition between pneumococcal strains. Specifically, the acquisition-blocking vaccine selects strongly against resistance in the "Treatment competition" model: since antibiotic-mediated within-host competition benefits the resistant strain in this model, the vaccine works against this competitive

advantage and therefore inhibits resistance. Conversely, in the "Growth competition"

model, growth-mediated competition benefits the sensitive strain, and so by reducing

competition, vaccination tends to promote resistance. These results expand upon our

previous finding that the rate of co-colonisation modulates resistance evolution through

its impact upon within-host competition (*26*).

The clearance-accelerating vaccine exhibits similarly divergent impacts across

mechanisms of resistance evolution. However, compared with the acquisition-blocking

vaccine, it also has an additional resistance-inhibiting effect across all models, because a

shorter duration of carriage—whether natural or vaccine-induced—selects against

resistance (*38*). This suggests that vaccines that accelerate natural clearance have a

particular potential for managing resistant infections.

Reducing the rate of penicillin prescribing selects against resistance, as expected,

exhibiting a similar impact across all four models.

The impact on resistant carriage (Fig. 3c), which combines changes in the prevalence of

carriage and changes in the frequency of resistance, can be treated as a proxy for the

incidence of resistant infections. Overall, under the "Growth competition" model,

vaccination at intermediate efficacy is expected to increase the rate of resistant carriage,

and hence the number of cases of resistant disease. In other models, vaccination always

reduces resistant carriage, particularly under the "Treatment competition" model. A

summary of the strongest vaccine impacts is shown in Fig. 3d.

Evidence to inform policy and vaccine trials. For vaccines to be considered an

efficient means of controlling resistant infections, they must compare favourably to

existing interventions, such as reducing inappropriate antibiotic use (*52*). The UK

government has recently announced an initiative to reduce antibiotic consumption by

15% by the year 2020 (*52*). Our models predict that a 15% reduction in primary-care

penicillin consumption would reduce carriage of penicillin-non-susceptible

pneumococci from 6% to 3%. The vaccine efficacy required to yield the same effect

varies considerably depending upon the mechanism of resistance evolution (Fig 4a); for

example, the required vaccine efficacy is lowest under the "Treatment competition"

model (ε_a = 11%; ε_c = 7%), and highest under the "Growth competition" model (ε_a =

52%; ε_c = 50%). A full comparison of vaccine and stewardship interventions would

require accounting for the economic cost of vaccines versus antibiotics, the wider range

of resistant pathogens that would be targeted by restrictions on antibiotic use, and any

potential increase in pathogen circulation that might be brought about by inadvertent

decreases in appropriate antibiotic use.

In randomized controlled trials of pneumococcal conjugate vaccines, resistance-related

endpoints have routinely been evaluated over a follow-up period of between 6 months

and 3.5 years after vaccination (*53*, *54*). If vaccine-induced changes in resistance

evolution unfold over a considerably longer timescale, similarly-designed trials may not

appropriately capture vaccine impact on resistance. Indeed, we find that it can take 5–

10 years for the full effects of resistance evolution to be seen (Fig. 4b), and that short-

term drops in resistance can be reversed—or even give way to increased resistance—in

the long term. Moreover, a trial in which vaccination is not offered to a substantial

fraction of the population would not capture the full impact of reduced pneumococcal

circulation, which is what drives competition-mediated changes in resistance in our

models. Finally, our analysis assumes that vaccines are administered to all recipients

simultaneously. In a real-world setting where vaccination would be rolled out gradually,

the full effect of vaccination could take even longer to observe.

The impact of vaccination at a national level varies depending upon the treatment rate

in a given country. Focusing on the specific outcome of childhood pneumococcal

pneumonia cases, we find that while interventions have a consistent impact from

country to country on the total pneumonia case rate, the impact on resistant pneumonia

cases is greatest in those countries where antibiotic use, and hence resistance, is highest

(Fig. 4c). We focus on resistant carriage, but the realised health benefits of any

intervention targeting both resistant and sensitive strains will depend upon the relative

health burdens of susceptible versus non-susceptible *S. pneumoniae* infections, which is

an area of ongoing research (*55*).

Vaccination in a high-burden setting. High prevalences of carriage, disease, and

resistance are often observed in low-income settings, and this may substantially alter

predictions of vaccine impact. As an illustrative example, a 90% pneumococcal carriage rate, with 81% of isolates resistant to penicillin, has been observed among children under five years old in western Kenya (*56*). This may be partly attributable to a longer average duration of carriage in this setting, as a 71-day mean duration of natural pneumococcal carriage has been measured in Kilifi, eastern Kenya (*57*).

To model a similar high-burden setting, we adjust model parameters estimated from European data: increasing the mean natural carriage duration, transmission rate, and treatment rate to match observed data, and ignoring mixing with any other countries (f = 1), while keeping other parameters the same. We find that a comparatively greater vaccine efficacy is needed to reduce the prevalence of resistant carriage in a high-burden, high-resistance setting (Fig. 5). This is particularly true under the "Growth competition" model, because in this model resistant carriage only declines as total pneumococcal carriage declines, and it is particularly difficult to reduce overall carriage in a high-transmission setting. Simultaneously, vaccination may have a comparatively greater impact in high-burden settings because of a comparatively higher incidence of disease: for example, Kenya is estimated to have an 8.8-fold higher incidence of severe pneumococcal pneumonia than the average in Europe (*58*).

Accounting for additional between-country variation does not substantially alter predictions. Our focus thus far has been on the impact of the four identified mechanisms *per se* upon resistance evolution, and accordingly we have focused on reproducing the positive association between treatment rate and resistance frequency rather than attempting to capture the additional variability in resistance frequency between countries not accounted for by the reported treatment rate alone (Fig. 2a). This additional variability may partially stem from differences in national testing and reporting practices, or between-country differences in the distribution of pneumococcal serotypes among invasive isolates (*59*). However, another possibility is that this additional variability in resistance results from systematic differences in pathogen biology or host behaviour across countries which can be captured by our modelling framework.

To help identify which model parameters could account for this variability, we relax the

assumption that only the treatment rate varies across countries, and perform Bayesian

maximum *a posteriori* fitting, assuming one additional parameter (c, b, β, u, f, z, g, κ, a, δ,

or k) is free to vary between countries while other parameters are held constant. We

find that additional variation in resistance between countries can be explained by

variation in certain other parameters, depending upon which model is used (Fig. 6a–b).

Importantly, among those parameters for which additional variation between countries

can explain the variation in resistance (Fig. 6c), predictions for the overall impact of

vaccination remain similar with the major differences still attributable to the underlying

mechanism of resistance evolution (Fig. 6d; Supplementary Figs. 3–14). Models that

could make more accurate country-specific predictions would need to account for the

effects of demographic structure, differences in carriage prevalence and disease rates

between settings, and variable vaccine protection among individuals.

Discussion

We have identified four mechanisms of resistance evolution that are capable of

recapitulating the observed relationship between penicillin consumption and penicillin

non-susceptibility in *S. pneumoniae* across Europe. These mechanisms are not mutually

exclusive, but the relative importance of each is predicted to have a substantial impact

upon predictions for resistance evolution under vaccination. In particular, the

"directionality" of within-host competition—that is, whether, on average, within-host

competition tends to benefit resistant or sensitive strains—strongly determines

whether vaccination selects for a decrease or an increase in antibiotic resistance in the

long term. This directionality may vary between pathogens, but is also sensitive to the

antibiotic treatment rate, and so may also vary between settings. Although we have

focused on competition between sensitive and resistant strains of *S. pneumoniae* only,

competition between serotypes (*24*) and with other bacteria colonizing the

nasopharynx will also impact upon resistance evolution, and determining the

importance of these other sources of within-host competition is crucial.

A key result of our models is that the mode of vaccine protection—whether acquisition-

blocking or clearance-accelerating—has an appreciable impact upon resistance

evolution. Whole-cell and purified-protein pneumococcal vaccines may induce

antibody-mediated humoral immunity, CD4+ T helper-17 cell-mediated immunity, or

both, with the type of immunity mediating pneumococcal acquisition, carriage, and

disease in ways that are still not fully understood (*49–51*). By modelling both modes of

vaccine action, we have highlighted that clearance-accelerating vaccines have increased

potential for preventing the spread of resistance, because in shortening the duration of

asymptomatic carriage they limit the fitness advantage of resistant pathogens under

selection pressure from antibiotic use.

The prevalence of penicillin non-susceptibility in *S. pneumoniae* has remained largely

stable in Europe between 2005–2017, a period which saw the incorporation of PCV into

the routine immunization schedules of most European countries (*60*). However,

because serotype replacement has largely negated any vaccine impact on the

prevalence of nasopharyngeal pneumococcal carriage (*10*, *11*), it is not clear that we

should expect to see any effects of competition-mediated resistance evolution following

a serotype-specific vaccine such as PCV—particularly given the complexity of detecting

vaccine-attributable changes in resistance in a population-level associational study that

would be confounded both by serotype replacement and by other changes in resistance

evolution that might be expected to occur at a national level over the course of multiple

years.

Under the "Treatment diversity" and "Pathogen diversity" models, we have argued that

universal pneumococcal vaccination will have little impact upon the long-term

evolution of antibiotic resistance because it does not change the sources of diversity

that modulate resistance evolution. Nonetheless, it is possible to target vaccines such

that this diversity is harnessed to manage resistance: high-resistance serotypes could

be targeted with a serotype-specific vaccine, or high-treatment subpopulations could be

targeted for vaccination in order to more effectively manage resistance. Indeed,

vaccination does have an additional inhibiting effect upon resistance in our models

because of the latter effect. This inhibition occurs because the vaccine has a relatively

greater impact upon transmission in populations where the prevalence of carriage is

already low, which in our models occur in countries or subpopulations with more

antibiotic consumption. Since these populations drive resistance more strongly, the

vaccine's comparatively greater impact in these populations tends to moderately inhibit
resistance overall. We note that while previous work (*38*) has suggested that resistance
evolution under a "Pathogen diversity" model results in a "stepped" resistance pattern
in which D-types are either fully sensitive or fully resistant at equilibrium, we find that
small amounts of mixing between populations can smooth out this pattern and allow
intermediate rates of resistance within subtypes (Supplementary Fig. 2). Finally, while
we have framed "Treatment competition" and "Growth competition" as two distinct
alternatives, they can instead be viewed as endpoints on a continuum, with possible
models of resistance evolution for which both $c > 0$ and $b > 0$ lying between them. The
impact of vaccination on resistance in such a model would depend upon the relative
importance of treatment-mediated and growth-mediated competition.
425
This analysis has necessarily made simplifying assumptions. We have focused on
prevalence (the fraction of individuals who are carriers) rather than incidence (the rate
of new carriage episodes) of nasopharyngeal carriage in presenting our findings. There
is evidence that pneumococcal disease progression is more likely to occur shortly after
nasopharyngeal acquisition (*61*), suggesting that incidence may be more relevant than
prevalence for predicting disease outcomes. Of particular note, recent modelling work
has suggested that clearance-accelerating vaccines can increase rates of pneumococcal
acquisition, if extended carriage is protective against new acquisition (*62*). However, it
is not obvious how to compare rates of carriage acquisition across the models examined
in this paper, particularly because co-colonisation is explicitly tracked—and fitted to
data—in some but not all models. More work is required to clarify the links between
acquisition, carriage, and disease across competing models of pneumococcal
transmission. Additionally, we have assumed that antibiotic treatment rates among
pneumococcal carriers remains constant after the introduction of a vaccine, even
though treatment rates dropped in many settings following PCV introduction (*5*, *9*).
However, for a universal pneumococcal vaccine that reduces antibiotic treatment rates
because it reduces carriage and thereby prevents antibiotic-treatable disease, any
reduction in treatment will only occur among individuals who, because of vaccine
protection, are not pneumococcal carriers, all else being equal. It might then be
expected that treatment rates in carriers would remain equally high among those
individuals for whom vaccine protection has failed.

Our work helps resolve the question: What explains the persistent coexistence between

resistant and sensitive strains of *S. pneumoniae*? (*25*) by demonstrating that multiple

mechanisms are capable of explaining trends of resistance across European countries.

Since there is empirical support for within-host competition between sensitive and

resistant pathogen strains (*63–66*), heritable differences in the propensity for resistance

within species (*38*), and within-country heterogeneity in antibiotic consumption rates

(*67–69*), all of these mechanisms likely contribute to this pattern. Our results

contextualize previous mathematical studies which have variously suggested that

serotype-specific vaccination may increase (*24*), decrease (*22*) or have no impact upon

(*18*) the frequency of resistance in *S. pneumoniae*. While the potential for vaccination to

promote resistance because of competition between sensitive and resistant strains has

been described previously (*24*), we have shown that vaccination can either promote or

inhibit resistance depending upon the directionality of within-host competition. While

vaccines targeting highly-resistant serotypes can decrease resistance (*22*), we have

shown that a serotype-independent vaccine promoting accelerated natural clearance

can decrease resistance across all circulating subtypes. And where single-population

models have found no long-term impact of vaccination on resistance frequency (*18*), we

have shown that in multi-population models, vaccination can inhibit resistance if it has

a larger impact in subpopulations that consume more antibiotics. The direction and

magnitude of this effect would depend upon variation in vaccine uptake, vaccine

efficacy, and pathogen transmission among subpopulations, and we have not

systematically explored this variation here.

A highly efficacious serotype-independent pneumococcal vaccine can indeed reduce the

overall burden of antibiotic-resistant pneumococcal infections. However, the long-term

effect upon resistance of a vaccine with intermediate efficacy is less certain, as vaccine

impact depends crucially upon the mechanisms that drive resistance evolution. Thus,

empirical investigation of pathogen competitive dynamics—and the impact of setting-

specific factors on these dynamics—is needed to make accurate predictions of vaccine

impact on resistant infections.

Methods

Study design. This study comprises four parts: a literature search used to identify plausible mechanisms through which coexistence can be maintained between sensitive and resistant pneumococcal strains across a range of antibiotic treatment rates; a mathematical modelling study embedding these mechanisms of resistance evolution in four models of pneumococcal transmission; a Bayesian statistical analysis to fit these models to empirically observed frequencies of penicillin non-susceptibility and community penicillin consumption across 27 European countries for the year 2007; and a vaccine impact analysis using these fitted models to forecast the impact of a universal pneumococcal vaccine. We use data from 2007 because changes in pneumococcal resistance reporting standards for some countries after this year hamper the between-country comparability of data (*70*). Our objectives were to identify the mechanisms potentially responsible for maintaining coexistence between resistant and sensitive pneumococci in Europe, and to determine whether the impact of vaccination on the evolution of resistance depends upon which mechanism is assumed to operate.

Mechanisms driving resistance. We searched PubMed using the terms: (AMR OR ABR OR ((antimicrobial OR antibiotic) AND resist*)) AND ((model OR modelling OR modeling) AND (dynamic* OR transmi* OR mathematical)) AND (coexist* OR intermediate). This yielded 93 papers (Supplementary Table 1). We included all papers containing a dynamic host-to-host pathogen transmission model analysing both sensitive and resistant strains with stable coexistence as an outcome of the model. From the 11 studies meeting these criteria, we identified nine unique mechanisms, two of which correspond to alternative parameterisations of a within-host competition model. We ruled out four mechanisms because of implausibility or because previous work shows that the mechanism does not bring about substantial coexistence, leaving four mechanisms (Table 1).

Model framework. We analyse the evolution of antibiotic resistance by tracking the transmission of resistant and sensitive bacterial strains among hosts in a set of M countries indexed by $m \in \{1, 2, \dots, M\}$ using systems of ordinary differential equations.

In a simple model, hosts can either be non-carriers (X), carriers of the sensitive strain

(S), or carriers of the resistant strain (R). Omitting country-specific subscripts m for

concision, model dynamics within a country are captured by

$\qquad dS/dt = \lambda_S X - (u + \tau)S$

$\qquad dR/dt = (1 - c)\lambda_R X - uR$

$\qquad X = 1 - S - R$,   (1)

where λ_S is the force of infection of the sensitive strain, λ_R is the force of infection of the

resistant strain, c is the transmission cost of resistance, u is the rate of natural

clearance, and τ is the treatment rate. In this model, in a given country, the total

carriage of the sensitive strain is S and the total carriage of the resistant strain is R.

Force of infection terms are defined below.

The "Treatment diversity" model extends the simple model (eq. 1) by structuring each

country into multiple subpopulations that exhibit different rates of antibiotic treatment

and make contact with each other at unequal rates (*25, 34, 35, 71*). In each country, we

model N equally-sized representative subpopulations indexed by $i \in \{1, 2, ..., N\}$, where

we assume $N = 10$. Dynamics within a country are

$\qquad dS_i/dt = \lambda_{S,i} X - (u + \tau_i)S$

$\qquad dR_i/dt = (1 - c)\lambda_{R,i} X - uR$

$\qquad X_i = 1 - S_i - R_i$ (2)

where we assume that treatment rates of subpopulations within a country

approximately follow a gamma distribution with shape parameter κ and mean

treatment rate τ. Accordingly, the rate of antibiotic consumption in subpopulation i is

$\tau_i = \int_{Q_\Gamma\left(\frac{i-1}{N}|\kappa\right)}^{Q_\Gamma\left(\frac{i}{N}|\kappa\right)} t\, P_\Gamma(t|\kappa)\, dt$, where $Q_\Gamma(q|\kappa)$ is the quantile q of the gamma distribution

with shape κ and $P_\Gamma(t|\kappa)$ is the probability density at t of the same gamma distribution.

The "Pathogen diversity" model extends the simple model (eq. 1) by structuring the

pathogen population into D different "D-types" (we assume $D = 25$), each with a

different natural clearance rate, where each type is kept circulating by diversifying

selection acting on D-type (38). Dynamics within a country are

$dS_d/dt = q_d\lambda_{S,d}X - (u_d + \tau)S_d$

$dR_d/dt = q_d(1 - c)\lambda_{R,d}X - u_d R_d$

$X = 1 - \Sigma_d (S_d + R_d)$ (3)

where $q_d = (1 - \frac{S_d + R_d}{\Sigma_{j=1}^{D}(S_j + R_j)} + \frac{1}{D})^a$ is the strength of diversifying selection for D-type $d \in$

$\{1, 2, \dots, D\}$ and $u_d = u\left(1 + \delta\left(2\frac{d-1}{D-1} - 1\right)\right)$ is the clearance rate for D-type d, where

model parameter a is the power of diversifying selection and model parameter δ is the

range of clearance rates (38). In a given country, the total carriage of the type-d

sensitive strain is S_d and the total carriage of the type-d resistant strain is R_d.

Finally, the within-host competition models (26) allow hosts to carry a mix of both

strains. Hosts can carry the sensitive strain with a small complement of the resistant

strain (S_R) or the resistant strain with a small complement of the sensitive strain (R_S).

Dynamics within a country are

$dS/dt = \lambda_S X - (u + \tau)S - k(1 - c)\lambda_R S + b_0 S_R$

$dS_R/dt = k(1 - c)\lambda_R S - (u + \tau)S_R + bR_S - b_0 S_R$

$dR_S/dt = k\lambda_S R - (u + \tau)R_S - bR_S$

$dR/dt = (1 - c)\lambda_R X - uR - k\lambda_S R + \tau(S_R + R_S)$

$X = 1 - S - R - S_R - R_S$,   (4)

where k is the rate of co-colonisation relative to primary colonisation, b is the within-

host growth benefit of sensitivity (i.e. the rate of the $R_S \rightarrow S_R$ transition), and $b_0 = 4b$ is

the rate of the $S_R \rightarrow S$ transition as a function of this growth benefit. In a given country,

the total carriage of the sensitive strain is $S + S_R$ and the total carriage of the resistant

strain is $R + R_S$. "Treatment competition" assumes the cost of resistance is incurred by

reduced transmission potential ($b = 0$ and $c > 0$), while "Growth competition" assumes

that the cost of resistance is incurred through decreased within-host growth ($b > 0$ and

$c = 0$).

In equations 1, 3 and 4, the force of infection of a particular strain A in country m is $\lambda_A =$

$\beta\left(f A_{\text{tot}|m} + (1 - f) \sum_{\ell=1}^{M} h_\ell A_{\text{tot}|\ell}\right)$, where β is the transmission rate, f is the between-

country assortativity, h_ℓ is the relative population size of country m (such that $\sum_\ell h_\ell =$

1), and $A_{\text{tot}|\ell}$ is the total carriage of strain A in country ℓ. The probability with which

individuals contact an individual from another country, $1 - f$, captures those contacts

made with individuals from another country in either one's home country or a foreign

country. In equation 2, the force of infection of a particular strain A in subpopulation i of

country m is $\lambda_{A,i} = \beta\left(f\left(g A_{\text{tot}|m,i} + (1 - g) \sum_{j=1}^{N} \frac{1}{N} A_{\text{tot}|m,j}\right) + (1 -$

$f) \sum_{\ell=1}^{M} \sum_{j=1}^{N} \frac{h_\ell}{N} A_{\text{tot}|\ell,j}\right)$, where g is the within-country assortativity and $A_{\text{tot}|\ell,j}$ is the

total carriage of strain A in subpopulation j of country ℓ.

Data and model fitting. We extracted community penicillin consumption and penicillin

non-susceptibility in *S. pneumoniae* invasive isolates from databases made available by

the ECDC (*13, 28*). We assume that community penicillin consumption drives penicillin

resistance, that antibiotic consumption is independent of whether an individual is

colonised by pneumococcus, and that resistance among invasive bacterial isolates is

representative of resistance among circulating strains more broadly. Countries report

community penicillin consumption in defined daily doses (DDD) per thousand

individuals per day. To transform this bulk consumption rate into the rate at which

individuals undertake a course of antibiotic therapy, we analysed prescribing data from

eight European countries, estimating that, on average, 5 DDD in the population at large

correspond to one treatment course for a child under 5 years of age. This conversion

rate varies between countries (Supplementary Table 4), but since the data are

incomplete (8 of 27 countries) we have not explicitly accounted for this variability in

our main model fitting results.

Our model framework tracks carriage of *S. pneumoniae* among children aged 0–5 years,

the age group driving both transmission and disease. In European countries, we assume

that the prevalence of pneumococcal carriage in under-5s is 50% (*11, 42*) and the

average duration of carriage is 47 days (*47, 48*). We calculate the average incidence of *S.*

pneumoniae-caused severe pneumonia requiring hospitalisation as 610 per million

children under 5 per year (*58*) across the European countries in our data set. See

Supplementary Tables 3, 5, and 6 for details of calculations relating to pneumococcal

carriage duration and disease incidence.

We use Bayesian inference via differential evolution Markov chain Monte Carlo (*72*) to

identify model parameters that are consistent with empirical data. Country m has

antibiotic treatment rate τ_m and reports r_m of n_m isolates are resistant. Over all M

countries, these data are denoted $\tau = (\tau_1, \tau_2, \ldots, \tau_M)$, $r = (r_1, r_2, \ldots, r_M)$, and $n =$

$(n_1, n_2, \ldots, n_M)$, respectively. The probability of a given set of model parameters θ is

then

618
$$P(\theta|\tau, r, n) \propto P(\tau, r, n|\theta)P(\theta),$$

620   where $P(\theta)$ is the prior probability of parameters θ and

621
$$P(\tau, r, n|\theta) = C\big(Y = Y(\tau|\theta)\big) \prod_{m=1}^{M} R\big(r = r_m, n = n_m, \rho = \rho(\tau_m|\theta)\big)^{N_m/\bar{N}}$$

is the likelihood of data τ, r, n given model parameters θ. Above, $Y(\theta)$ is the average

model-predicted prevalence of carriage across all countries and $\rho(\tau_m|\theta)$ is the model-

predicted resistance prevalence for country m. $C(Y)$ is the credibility of prevalence of

carriage Y and $R(r,n,\rho)$ is the credibility of r out of n isolates being resistant when the

model-predicted resistance prevalence is ρ. For $C(Y)$, we use a normal distribution with

mean 0.5 and standard deviation 0.002. For $R(r,n,\rho)$, we use $R(r, n, \rho) =$

$\int_0^1 T\big(x|\mu = \rho, \sigma = \sigma(\theta)\big)\binom{n}{r}x^r(1-x)^{n-r}\,dx$, a binomial distribution where the

probability of success is modelled as a [0,1]-truncated normal distribution centred on ρ

and with standard deviation σ. The parameter σ captures the unexplained between-

country variation in resistance frequency. Here, $T(x|\mu, \sigma) = \frac{\varphi(x|\mu,\sigma)}{(\Phi(1|\mu,\sigma) - \Phi(0|\mu,\sigma))}$, where

$\varphi(\mu, \sigma) = \frac{1}{\sqrt{2\pi\sigma^2}} exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right)$ is the untruncated normal PDF and $\Phi(\mu, \sigma) = \frac{1}{2}(1 +$

$erf\left(\frac{x-\mu}{\sigma\sqrt{2}}\right))$ is the untruncated normal cumulative distribution function. Finally, N_m is the

population size of country m and $\bar{N}$ is the average population size across all countries;

the exponent $N_m/\bar{N}$ allows us to weight the importance of each country by its

population size, which allows a closer fit with the overall resistance prevalence across

all countries.

We adopt $c \sim \text{Beta}(\alpha = 1.5, \beta = 8.5)$, $b \sim \text{Gamma}(\kappa = 2, \theta = 0.5)$, $\beta \sim \text{Gamma}(\kappa =

5, \theta = 0.35)$, $g \sim \text{Beta}(\alpha = 10, \beta = 1.5)$, $\kappa \sim \text{Gamma}(\kappa = 4, \theta = 2)$, $a \sim \text{Gamma}(\kappa =

2, \theta = 5)$, $\delta \sim \text{Beta}(\alpha = 20, \beta = 25)$, and $k \sim \text{Normal}(\mu = 1, \sigma = 0.5)$ as weakly

informative prior distributions for model fitting. We set the unexplained between-

country variation in resistance prevalence σ to 0.06 across all models based on a

preliminary round of model fitting with σ as a free parameter, and set the between-

country assortativity *f* to 0.985 (*i.e.*, 1.5% of contacts occur with individuals from a

different country) based on rates of travel within the EU. See Supplementary Table 7 for

MCMC diagnostics.

To match model predictions to a high-burden setting, we increase the duration of

carriage to 71.4 days; increase the transmission rate by a factor of 3.49 (Treatment

diversity), 3.62 (Pathogen diversity), 3.61 (Treatment competition), or 3.20 (Growth

competition), so that carriage prevalence reaches 90.0%; and increase the antibiotic

consumption rate to 1.670, 1.458, 1.138, or 5.887 courses per person per year,

respectively, so that resistance prevalence reaches 81.4%.

Interventions. Interventions have the following impact on model parameters: for the

acquisition-blocking vaccine, the transmission rate becomes $\beta' = (1-\varepsilon_a)\,\beta$; for the

clearance-accelerating vaccine, the clearance rate becomes $u' = u/(1-\varepsilon_c)$; and under

antibiotic stewardship, the average treatment rate in each country *m* becomes $\tau_m' = \tau_m$

$(1-\varepsilon_s)$.

Capturing additional between-country variation in resistance frequency. We begin by

finding the maximum *a posteriori* model fits according to the likelihood and prior

distributions for each of the four models of resistance evolution. This identifies the

following parameter values for each model. "Treatment diversity": $\beta = 1.41$, $c = 0.124$, $g

= 0.976$, and $\kappa = 2.22$. "Pathogen diversity": $\beta = 1.33$, $c = 0.191$, $a = 10.8$, and $\delta = 0.608$.

"Treatment competition": $\beta = 1.42$, $c = 0.191$, and $k = 1.64$. "Growth competition": $\beta =

1.39$, $b = 0.195$, and $k = 1.61$. Then, we perform maximum *a posteriori* model fits for each

potentially-varying parameter under each model, allowing the varying parameter to

take on a different value for each country and fixing other parameters at their maximum

a posteriori values as determined in the previous step, or at specific assumed values for

$u = 0.65$, $f = 0.985$, and $z = 5$. For the second step, we use a modified likelihood function

675
$$P(\tau, r, n | \theta) = C\big(Y = Y(\tau | \theta)\big) \prod_{m=1}^{M} \phi\left(\mu = \frac{r_m + 1}{n_m + 2}, \sigma = 0.001 \middle| x = \rho(\tau_m | \theta)\right)^{N_m / \bar{N}},$$

where $\phi(\mu, \sigma | x)$ is the normal probability density function. This modified likelihood

function ensures that the model-predicted resistance frequency for each country is

matched as closely as possible to the maximum-likelihood resistance prevalence $\frac{r_m + 1}{n_m + 2}$

(*i.e.*, assuming a uniform prior on resistance frequency) for each country m, so that

model fits are comparable across different varying parameters. We use the Nelder-Mead

algorithm to maximize the posterior probability in both steps.

Supplementary Figs. 3–6 show maximum *a posteriori* fits when allowing an additional

parameter to vary freely between countries, along with the parameter values identified

by model fitting. Supplementary Figs. 7–10 show the impact of vaccination, focusing on

those parameters for which model fitting was able to capture the observed variability in

resistance frequency between countries (*i.e.*, those parameters plotted to the left of the

dashed line in Fig. 6b of the main text). Supplementary Figs. 11–14 show the impact of

vaccination for the remaining parameters.

List of Supplementary Materials

Supplementary Table 1. Literature review.

Supplementary Table 2. Summary of model parameters.

Supplementary Table 3. Carriage duration.

Supplementary Table 4. Penicillin consumption.

Supplementary Table 5. Pneumococcal morbidity.

Supplementary Table 6. Carriage duration (Kilifi).

Supplementary Table 7. MCMC diagnostics.

Supplementary Fig. 1. Patterns of penicillin non-susceptibility across European

countries, 2005–2017.

Supplementary Fig. 2. Carriage and resistance of D-types in "Pathogen variability"

model.

Supplementary Fig. 3. Varying-parameter fits for "Treatment diversity" model.

Supplementary Fig. 4. Varying-parameter fits for "Pathogen diversity" model.

Supplementary Fig. 5. Varying-parameter fits for "Treatment competition" model.

Supplementary Fig. 6. Varying-parameter fits for "Growth competition" model.

Supplementary Fig. 7. Vaccine impact for "Treatment diversity" model, varying

parameters c and z**.**

Supplementary Fig. 8. Vaccine impact for "Pathogen diversity" model, varying

parameters c, δ, and z.

Supplementary Fig. 9. Vaccine impact for "Treatment competition" model, varying

parameters β, c, u, k, and z.

Supplementary Fig. 10. Vaccine impact for "Growth competition" model, varying

parameters b and z.

Supplementary Fig. 11. Vaccine impact for "Treatment diversity" model, varying other

parameters.

Supplementary Fig. 12. Vaccine impact for "Pathogen diversity" model, varying other

parameters.

Supplementary Fig. 13. Vaccine impact for "Treatment competition" model, varying

other parameters.

Supplementary Fig. 14. Vaccine impact for "Growth competition" model, varying other

parameters.

References

1.   J. O'Neill, "Vaccines and alternative approaches: reducing our dependence on
antimicrobials" (2016), (available at http://amr-
review.org/sites/default/files/Vaccines and alternatives_v4_LR.pdf).

2.   M. Lipsitch, R. Siber, How Can Vaccines Contribute to Solving the Antimicrobial
Resistance Problem ? **7**, 1–8 (2016).

3.   K. E. Atkins, M. Lipsitch, Can antibiotic resistance be reduced by vaccinating
against respiratory disease? *Lancet Respir. Med.* **6**, 820–821 (2018).

4.   J. P. Sevilla, D. E. Bloom, D. Cadarette, M. Jit, M. Lipsitch, Toward economic
evaluation of the value of vaccines and other health technologies in addressing
AMR. *Proc. Natl. Acad. Sci.* **115**, 12911–12919 (2018).

5.   K. P. Klugman, S. Black, Impact of existing vaccines in reducing antibiotic
resistance: Primary and secondary effects. *Proc. Natl. Acad. Sci.* **115**, 12896–
12901 (2018).

6.   M. H. Kyaw, R. Lynfield, W. Schaffner, A. S. Craig, J. Hadler, A. Reingold, A. R.
Thomas, L. H. Harrison, N. M. Bennett, M. M. Farley, R. R. Facklam, J. H. Jorgensen,
J. Besser, E. R. Zell, A. Schuchat, C. G. Whitney, Effect of Introduction of the
Pneumococcal Conjugate Vaccine on Drug-Resistant Streptococcus pneumoniae.
N. Engl. J. Med. **354**, 1455–1463 (2006).

7.   D. Thorrington, N. Andrews, J. Stowe, E. Miller, A. J. van Hoek, Elucidating the
impact of the pneumococcal conjugate vaccine programme on pneumonia, sepsis
and otitis media hospital admissions in England using a composite control. *BMC*
Med. **16**, 1–14 (2018).

8.   C. Chen, F. Cervero Liceras, S. Flasche, S. Sidharta, J. Yoong, N. Sundaram, M. Jit,
Effect and cost-effectiveness of pneumococcal conjugate vaccination: a global
modelling analysis. *Lancet Glob. Heal.* **7**, e58–e67 (2019).

9.   K. J. Wilby, D. Werry, A review of the effect of immunization programs on
antimicrobial utilization. *Vaccine*. **30**, 6509–6514 (2012).

10. W. P. Hanage, J. A. Finkelstein, S. S. Huang, S. I. Pelton, A. E. Stevenson, K.
Kleinman, V. L. Hinrichsen, C. Fraser, Evidence that pneumococcal serotype
replacement in Massachusetts following conjugate vaccination is now complete.
Epidemics. **2**, 80–84 (2010).

11.   S. Flasche, A. J. Van Hoek, E. Sheasby, P. Waight, N. Andrews, C. Sheppard, R.
George, E. Miller, Effect of pneumococcal conjugate vaccination on serotype-
specific carriage and invasive disease in England: a cross-sectional study. *PLoS*
Med. **8**, e1001017 (2011).

12.   J. A. Lewnard, W. P. Hanage, Making sense of differences in pneumococcal
serotype replacement. *Lancet Infect. Dis.* **3099** (2019), doi:10.1016/s1473-
3099(18)30660-1.

13.   European Centre for Disease Prevention and Control, Antimicrobial consumption
rates by country (2018), (available at
http://ecdc.europa.eu/en/healthtopics/antimicrobial_resistance/esac-net-
database/Pages/Antimicrobial-consumption-rates-by-country.aspx).

14.   World Health Organization, WHO vaccine pipeline tracker (2019), (available at
https://www.who.int/immunization/research/vaccine_pipeline_tracker_spreads
heet/en/).

15.   G. L. Masala, M. Lipsitch, C. Bottomley, S. Flasche, Exploring the role of
competition induced by non-vaccine serotypes for herd protection following
pneumococcal vaccination. *J. R. Soc. Interface.* **14** (2017),
doi:10.1098/rsif.2017.0620.

16.   K. E. Atkins, E. I. Lafferty, S. R. Deeny, N. G. Davies, J. V. Robotham, M. Jit, Use of
mathematical modelling to assess the impact of vaccines on antibiotic resistance.
Lancet Infect. Dis. **18**, e204–e213 (2018).

17.   K. E. Atkins, S. Flasche, Vaccination to reduce antimicrobial resistance. *Lancet*
Glob. Heal. **6**, e252 (2018).

18.   L. Temime, D. Guillemot, P. Y. Boëlle, Short- and long-term effects of
pneumococcal conjugate vaccination of children on penicillin resistance.
Antimicrob. Agents Chemother. **48**, 2206–2213 (2004).

19.   L. Temime, P. Y. Boëlle, A. J. Valleron, D. Guillemot, Penicillin-resistant
pneumococcal meningitis: High antibiotic exposure impedes new vaccine
protection. *Epidemiol. Infect.* **133**, 493–501 (2005).

20.   L. Opatowski, L. Temime, E. Varon, R. Leclerc, H. Drugeon, P.-Y. Boëlle, D.
Guillemot, Antibiotic Innovation May Contribute to Slowing the Dissemination of
Multiresistant Streptococcus pneumoniae: The Example of Ketolides. *PLoS One.* **3**,
e2089 (2008).

21.   T. Van Effelterre, M. R. Moore, F. Fierens, C. G. Whitney, L. White, S. I. Pelton, W. P.
Hausdorff, A dynamic model of pneumococcal infection in the United States:
Implications for prevention through vaccination. *Vaccine*. **28**, 3650–3660 (2010).

22.   M. D. De Cellès, M. Pons-Salort, E. Varon, M. A. Vibet, C. Ligier, V. Letort, L.
Opatowski, D. Guillemot, Interaction of vaccination and reduction of antibiotic use
drives unexpected increase of pneumococcal meningitis. *Sci. Rep.* **5**, 1–11 (2015).

23.   P. K. Mitchell, M. Lipsitch, W. P. Hanage, Carriage burden, multiple colonization
and antibiotic pressure promote emergence of resistant vaccine escape
pneumococci. *Philos. Trans. R. Soc. B Biol. Sci.* **370**, 3–9 (2015).

24.   U. Obolski, J. Lourenço, C. Thompson, R. Thompson, A. Gori, S. Gupta, Vaccination
can drive an increase in frequencies of antibiotic resistance among nonvaccine
serotypes of Streptococcus pneumoniae. *Proc. Natl. Acad. Sci.* **115**, 3102–3107
(2018).

25.   C. Colijn, T. Cohen, C. Fraser, W. Hanage, E. Goldstein, N. Givon-Lavi, R. Dagan, M.
Lipsitch, What is the mechanism for persistent coexistence of drug-susceptible
and drug-resistant strains of *Streptococcus pneumoniae*? *J. R. Soc. Interface*. **7**,
905–919 (2010).

26.   N. G. Davies, S. Flasche, M. Jit, K. E. Atkins, Within-host dynamics shape antibiotic
resistance in commensal bacteria. *Nat. Ecol. Evol.* (2019), doi:10.1038/s41559-
018-0786-x.

27.   H. Goossens, M. Ferech, R. Vander Stichele, M. Elseviers, Outpatient antibiotic use
in Europe and association with resistance: A cross-national database study.
Lancet. **365**, 579–587 (2005).

28.   European Centre for Disease Prevention and Control, Data from the ECDC
Surveillance Atlas - Antimicrobial resistance (2016), (available at
https://ecdc.europa.eu/en/antimicrobial-resistance/surveillance-and-disease-
data/data-ecdc).

29.   M. Lipsitch, C. Colijn, T. Cohen, W. P. Hanage, C. Fraser, No coexistence for free:
Neutral null models for multistrain pathogens. *Epidemics*. **1**, 2–13 (2009).

30.   P. Rodrigues, M. G. M. Gomes, C. Rebelo, Drug resistance in tuberculosis-a
reinfection model. *Theor. Popul. Biol.* **71**, 196–212 (2007).

31.   O. Patterson-Lomba, B. M. Althouse, G. M. Goerg, L. Hébert-Dufresne, Optimizing
Treatment Regimes to Hinder Antiviral Resistance in Influenza across Time

Scales. *PLoS One.* **8**, 1–11 (2013).

32.   I. M. Hastings, Complex dynamics and stability of resistance to antimalarial drugs.
Parasitology. **132**, 615–624 (2006).

33.   R. Sergeev, C. Colijn, T. Cohen, Models to understand the population-level impact
of mixed strain M. tuberculosis infections. *J. Theor. Biol.* **280**, 88–100 (2011).

34.   F. Blanquart, S. Lehtinen, M. Lipsitch, C. Fraser, The evolution of antibiotic
resistance in a structured host population. *J. R. Soc. Interface.* **15** (2018),
doi:10.1098/rsif.2018.0040.

35.   M. S. Krieger, A. L. Hill, Long-term coexistence and regional heterogeneity of
antibiotic-resistant infections reproduced by a simple spatial model. *bioRxiv.* **14**,
469171 (2018).

36.   R. Kouyos, E. Klein, B. Grenfell, Hospital-Community Interactions Foster
Coexistence between Methicillin-Resistant Strains of Staphylococcus aureus. *PLoS*
Pathog. **9** (2013), doi:10.1371/journal.ppat.1003134.

37.   M. F. Boni, M. W. Feldman, Evolution of Antibiotic Resistance By Human and
Bacterial Niche Construction. *Evolution (N. Y).* **59**, 477 (2006).

38.   S. Lehtinen, F. Blanquart, N. J. Croucher, P. Turner, M. Lipsitch, C. Fraser, Evolution
of antibiotic resistance is linked to any genetic mechanism affecting bacterial
duration of carriage. *Proc. Natl. Acad. Sci.* **114**, 1075–1080 (2017).

39.   D. J. Austin, K. G. Kristinsson, R. M. Anderson, The relationship between the
volume of antimicrobial consumption in human communities and the frequency
of resistance. *Proc. Natl. Acad. Sci.* **96**, 1152–1156 (1999).

40.   A. B. Beams, D. J. A. Toth, K. Khader, F. R. Adler, Harnessing intra-host strain
competition to limit antibiotic resistance: mathematical model results. *Bull. Math.*
Biol. **78**, 1828–1846 (2016).

41.   K. Græsbøll, S. S. Nielsen, N. Toft, L. E. Christiansen, How fitness reduced,
antimicrobial resistant bacteria survive and spread: A multiple pig - multiple
bacterial strain model. *PLoS One.* **9** (2014), doi:10.1371/journal.pone.0100458.

42.   D. Bogaert, A. Van Belkum, M. Sluijter, A. Luijendijk, R. De Groot, H. C. Rümke, H. A.
Verbrugh, P. W. M. Hermans, Colonisation by *Streptococcus pneumoniae* and
Staphylococcus aureus in healthy children. *Lancet.* **363**, 1871–1872 (2004).

43.   D. M. Weinberger, V. E. Pitzer, G. Regev-Yochay, N. Givon-Lavi, R. Dagan,
Association between the Decline in Pneumococcal Disease in Unimmunized

Adults and Vaccine-Derived Protection Against Colonization in Toddlers and

Preschool-Aged Children. *Am. J. Epidemiol.* **188**, 160–168 (2019).

44.   D. I. Andersson, The biological cost of mutational antibiotic resistance: any

practical conclusions? *Curr. Opin. Microbiol.* **9** (2006), pp. 461–465.

45.   D. I. Andersson, D. Hughes, Antibiotic resistance and its cost: Is it possible to

reverse resistance? *Nat. Rev. Microbiol.* **8**, 260–271 (2010).

46.   C. Tedijanto, S. Olesen, Y. Grad, M. Lipsitch, Estimating the proportion of

bystander selection for antibiotic resistance among potentially pathogenic

bacterial flora. *Proc Natl Acad Sci USA*. **115**, E11988–E11995 (2018).

47.   L. Högberg, P. Geli, H. Ringberg, E. Melander, M. Lipsitch, K. Ekdahl, Age- and

serogroup-related differences in observed durations of nasopharyngeal carriage

of penicillin-resistant pneumococci. *J. Clin. Microbiol.* **45**, 948–952 (2007).

48.   A. Melegaro, N. J. Gay, G. F. Medley, Estimating the transmission parameters of

pneumococcal carriage in households. *Epidemiol. Infect.* **132**, 433–441 (2004).

49.   S. Cobey, M. Lipsitch, Niche and neutral effects of acquired immunity permit

coexistence of pneumococcal serotypes. *Science (80-.).* **335**, 1376–1380 (2012).

50.   R. Malley, P. W. Anderson, Serotype-independent pneumococcal experimental

vaccines that induce cellular as well as humoral immunity. *Proc. Natl. Acad. Sci. U.*

S. A. **109**, 3623–3627 (2012).

51.   S. P. Jochems, J. N. Weiser, R. Malley, D. M. Ferreira, The immunological

mechanisms that control pneumococcal carriage. *PLoS Pathog.* **13**, 1–14 (2017).

52.   UK Department of Health and Social Care, "Tackling antimicrobial resistance 2019

to 2024: the UK's 5-year national action plan" (2019), (available at

https://www.gov.uk/government/publications/uk-5-year-action-plan-for-

antimicrobial-resistance-2019-to-2024).

53.   B. S. Buckley, N. Henschke, H. Bergman, B. Skidmore, E. J. Klemm, G. Villanueva, C.

Garritty, M. Paul, Impact of vaccination on antibiotic usage: a systematic review

and meta-analysis. *Clin. Microbiol. Infect.* **In press** (2019),

doi:10.1016/j.cmi.2019.06.030.

54.   A. Von Gottberg, L. De Gouveia, S. Tempia, V. Quan, S. Meiring, C. Von Mollendorf,

S. A. Madhi, E. R. Zell, M. Stat, J. R. Verani, K. L. O'Brien, C. G. Whitney, K. P.

Klugman, C. Cohen, Effects of vaccination on invasive pneumococcal disease in

South Africa. *N. Engl. J. Med.* **371**, 1889–1899 (2014).

889 55. A. Cassini, L. D. Högberg, D. Plachouras, A. Quattrocchi, A. Hoxha, G. S. Simonsen,

890   M. Colomb-Cotinat, M. E. Kretzschmar, B. Devleesschauwer, M. Cecchini, D. A.

891   Ouakrim, T. C. Oliveira, M. J. Struelens, C. Suetens, D. L. Monnet, R. Strauss, K.

892   Mertens, T. Struyf, B. Catry, K. Latour, I. N. Ivanov, E. G. Dobreva, A. Tambic

893   Andraševic, S. Soprek, A. Budimir, N. Paphitou, H. Žemlicková, S. Schytte Olsen, U.

894   Wolff Sönksen, P. Märtin, M. Ivanova, O. Lyytikäinen, J. Jalava, B. Coignard, T.

895   Eckmanns, M. Abu Sin, S. Haller, G. L. Daikos, A. Gikas, S. Tsiodras, F. Kontopidou,

896   Á. Tóth, Á. Hajdu, Ó. Guólaugsson, K. G. Kristinsson, S. Murchan, K. Burns, P.

897   Pezzotti, C. Gagliotti, U. Dumpis, A. Liuimiene, M. Perrin, M. A. Borg, S. C. de Greeff,

898   J. C. Monen, M. B. Koek, P. Elstrøm, D. Zabicka, A. Deptula, W. Hryniewicz, M.

899   Caniça, P. J. Nogueira, P. A. Fernandes, V. Manageiro, G. A. Popescu, R. I. Serban, E.

900   Schréterová, S. Litvová, M. Štefkovicová, J. Kolman, I. Klavs, A. Korošec, B. Aracil, A.

901   Asensio, M. Pérez-Vázquez, H. Billström, S. Larsson, J. S. Reilly, A. Johnson, S.

902   Hopkins, Attributable deaths and disability-adjusted life-years caused by

903   infections with antibiotic-resistant bacteria in the EU and the European Economic

904   Area in 2015: a population-level modelling analysis. *Lancet Infect. Dis.* **19**, 56–66

905   (2019).

906 56. M. Kobayashi, L. M. Conklin, G. Bigogo, G. Jagero, L. Hampton, K. E. Fleming-Dutra,

907   M. Junghae, M. da G. Carvalho, F. Pimenta, B. Beall, T. Taylor, K. F. Laserson, J.

908   Vulule, C. Van Beneden, L. Kim, D. R. Feikin, C. G. Whitney, R. F. Breiman,

909   Pneumococcal carriage and antibiotic susceptibility patterns from two cross-

910   sectional colonization surveys among children aged <5 years prior to the

911   introduction of 10-valent pneumococcal conjugate vaccine - Kenya, 2009-2010.

912   *BMC Infect. Dis.* **17**, 1–12 (2017).

913 57. M. Lipsitch, O. Abdullahi, A. D'Amour, W. Xie, D. M. Weinberger, E. Tchetgen

914   Tchetgen, J. A. G. Scott, Estimating rates of carriage acquisition and clearance and

915   competitive ability for pneumococcal serotypes in Kenya with a Markov

916   transition model. *Epidemiology*. **23**, 510–519 (2012).

917 58. I. Rudan, K. L. O'Brien, H. Nair, L. Liu, E. Theodoratou, S. Qazi, I. Luksic, C. L. F.

918   Walker, R. E. Black, H. Campbell, Child Health Epidemiology Reference Group,

919   Epidemiology and etiology of childhood pneumonia in 2010: estimates of

920   incidence, severe morbidity, mortality, underlying risk factors and causative

921   pathogens for 192 countries. *J. Glob. Health*. **3**, S114-010401 (2013).

59.   A. N. Torné, J. G. Dias, C. Quinten, F. Hruba, M. C. Busana, P. L. Lopalco, A. J. A.
Gauci, L. Pastore-Celentano, M. Sabbe, J. Verhaegen, M. Koliou, D. Pieridou-
Bagkatzouni, P. Křižovà, J. Kozakova, J. Motlova, P. Valentiner-Branth, L.
Lambertsen, T. Georgakopoulou, H. Humphreys, T. Melillo, P. Caruana, M. Knol, H.
de Merkel, K. Elberse, D. Frimann, A. Skoczynska, W. Hryniewicz, A. Kuch, I.
Paradowska-Stankiewicz, M. Pana, M. Vitek, V. Učakar, B. H. Normark, T. Lepp, M.
Slack, P. A. Waight, European enhanced surveillance of invasive pneumococcal
disease in 2010: Data from 26 European countries in the post-heptavalent
conjugate vaccine era. *Vaccine*. **32**, 3644–3650 (2014).

60.   M. Tin Tin Htar, D. Christopoulou, H. J. Schmitt, Pneumococcal serotype evolution
in Western Europe. *BMC Infect. Dis.* **15**, 1–10 (2015).

61.   B. Simell, K. Auranen, H. Käyhty, D. Goldblatt, R. Dagan, K. L. O'Brien, The
fundamental link between pneumococcal carriage and disease. *Expert Rev.*
Vaccines. **11**, 841–855 (2012).

62.   S. Flasche, A. Guilder, Using a mathematical model to assess the potential effects
of a new vaccine approach that targets enhanced clearance of pneumococcal
carriage. (2018).

63.   M. C. Negri, M. Lipsitch, J. Blázquez, B. R. Levin, F. Baquero, Concentration-
dependent selection of small phenotypic differences in TEM beta-lactamase-
mediated antibiotic resistance. *Antimicrob. Agents Chemother.* **44**, 2485–91
(2000).

64.   A. R. Wargo, S. Huijben, J. C. de Roode, J. Shepherd, A. F. Read, Competitive release
and facilitation of drug-resistant parasites after therapeutic chemotherapy in a
rodent malaria model. *Proc. Natl. Acad. Sci.* **104**, 19914–19919 (2007).

65.   N. Wale, D. G. Sim, M. J. Jones, R. Salathe, T. Day, A. F. Read, Resource limitation
prevents the emergence of drug resistance by intensifying within-host
competition. *Proc. Natl. Acad. Sci.*, 201715874 (2017).

66.   J. A. Lewnard, P. A. Tähtinen, M. K. Laine, L. Lindholm, J. Jalava, P. Huovinen, M.
Lipsitch, A. Ruohola, Impact of antimicrobial treatment for acute otitis media on
carriage dynamics of penicillin-susceptible and penicillin–non-susceptible
Streptococcus pneumoniae. *J. Infect. Dis.*, jiy343 (2018).

67.   M. Filippini, G. Masiero, K. Moschetti, Socioeconomic determinants of regional
differences in outpatient antibiotic consumption: Evidence from Switzerland.

Health Policy (New. York). **78**, 77–92 (2006).

68.   D. Koller, F. Hoffmann, W. Maier, K. Tholen, R. Windt, G. Glaeske, Variation in

antibiotic prescriptions: Is area deprivation an explanation? Analysis of 1.2

million children in Germany. *Infection.* **41**, 121–127 (2013).

69.   M. Di Martino, A. Lallo, U. Kirchmayer, M. Davoli, D. Fusco, Prevalence of antibiotic

prescription in pediatric outpatients in Italy: The role of local health districts and

primary care physicians in determining variation. A multilevel design for

healthcare decision support. *BMC Public Health.* **17**, 1–8 (2017).

70.   M. C. Goossens, B. Catry, J. Verhaegen, Antimicrobial resistance to benzylpenicillin

in invasive pneumococcal disease in Belgium, 2003-2010: The effect of altering

clinical breakpoints. *Epidemiol. Infect.* **141**, 490–495 (2013).

71.   S. Cobey, E. B. Baskerville, C. Colijn, W. Hanage, C. Fraser, M. Lipsitch, Host

population structure and treatment frequency maintain balancing selection on

drug resistance. *J. R. Soc. Interface.* **14**, 20170295 (2017).

72.   C. Ter Braak, A Markov Chain Monte Carlo version of the genetic algorithm

Differential Evolution: Easy Bayesian computing for real parameter spaces. *Stat.*

Comput. **16**, 239–249 (2006).

73.   T. Smieszek, K. B. Pouwels, F. C. K. Dolk, D. R. M. Smith, S. Hopkins, M. Sharland, A.

D. Hay, M. V Moore, J. V Robotham, Potential for reducing inappropriate antibiotic

prescribing in English primary care. *J. Antimicrob. Chemother.* **73**, ii36–ii43

(2018).

Acknowledgements

N.G.D., M.J. and K.E.A. were funded by the National Institute for Health Research Health Protection Research Unit in Immunisation at the London School of Hygiene and Tropical Medicine in partnership with Public Health England. The views expressed are those of the authors and not necessarily those of the NHS, National Institute for Health Research, Department of Health or Public Health England. S.F. was supported by a Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and Royal Society (grant number 208812/Z/17/Z).

Table 1. Mechanisms for maintaining coexistence

	Mechanism	Mode of action	Plausible mechanism for coexistence in *S. pneumoniae*?	Consistent with empirical patterns?
DIVERSITY	**Treatment diversity**	Assortatively-mixing subpopulations differ in treatment rates (*25, 34–37*)	✓ Yes	✓ Yes
	Pathogen diversity	Subtypes maintained by diversifying selection differ in propensity for resistance (*38*)	✓ Yes	✓ Yes
	Treated class	Individuals currently in treatment maintain resistant strains (*25, 34, 39, 40*)	✗ No: Only supports a small amount of coexistence (*25*)	▇ N/A
	Within-host niches	Sensitive and resistant strains exploit separate niches within the host (*30, 41*)	✗ No: Resistant and sensitive strains are known to occupy the same niches (*29*)	▇ N/A
	Mutation pressure	Mutation-selection balance maintains intermediate resistance frequency (*30, 31, 37*)	✗ No: De novo acquisition of resistance in *S. pneumoniae* is not frequent enough (*25*)	▇ N/A
	Prescription feedback	Doctors reduce prescribing of a drug as resistance to it increases (*37, 39*)	✗ No: Does not explain how coexistence is maintained over a range of treatment rates	▇ N/A
COMPETITION	**Within-host competition: Treatment competition**	Within-host competition creates frequency-dependent selection for resistance (*25, 26, 32, 33, 40*)	✓ Yes	✓ Yes
	Within-host competition: Growth competition	"	✓ Yes	✓ Yes
	Superinfection	Superinfection creates frequency-dependent selection for resistance (*30*)	✗ No: Requires resistant strain to transmit better than sensitive strain in absence of antibiotics	▇ N/A

Fig. 1. Four models of resistance evolution. X hosts are uncolonised, S hosts are
colonised with the sensitive strain and R hosts are colonised with the resistant strain. λ_S
and λ_R are the force of infection of the sensitive and resistant strain, respectively; c is
the transmission cost of resistance; u is the natural clearance rate; and τ is the rate of
antibiotic treatment. **(a)** "Treatment diversity": each country is split into
subpopulations varying in treatment rate τ_i. Assortative mixing between
subpopulations maintains coexistence. **(b)** "Pathogen diversity": the pathogen comes in
multiple subtypes maintained by diversifying selection, each with its own clearance rate
u_d. Subtypes with a longer carriage duration experience stronger selection for
resistance than those with a shorter carriage duration, which maintains circulation of
both sensitive and resistant strains overall. **(c)** "Treatment competition": singly-
colonised hosts can acquire a small amount of another strain at relative rate k (host
states S_R and R_S). Population-level coexistence is maintained by treatment-mediated
within-host competition between co-colonising strains. **(d)** "Growth competition": as in
(c), but the transmission cost of resistance is removed and sensitive strains now
outgrow resistant strains within co-colonised hosts at rate b. Coexistence is maintained
by both treatment-mediated and growth-mediated within-host competition.

1007

1008

Fig 2. Four models reproduce patterns of resistance in *S. pneumoniae* in Europe.

(a) Model fits with associated WAIC (± standard error). Vertical lines show the 95%

highest density intervals (HDIs) for the reported proportion of invasive *S. pneumoniae*

isolates that are resistant to penicillin plotted against the penicillin consumption rate in

under-5s. Ribbons show the 50% and 95% HDIs for resistance prevalence from each

fitted model. **(b)** The top row shows estimated posterior distributions for the free

parameters in each model; the bottom row shows model outputs associated with these

parameters to aid interpretation.

1017

1018

1019 **Fig. 3. Impact of interventions.** Impact of vaccine and treatment interventions on **(a)**
1020 carriage prevalence, **(b)** resistance frequency, and **(c)** resistant carriage (mean and
1021 95% HDI). **(d)** Illustration of the strongest forces selecting for greater or lesser
1022 resistance across models.

Fig. 4. Policy considerations. (a) Median equivalent reduction in prescribing across
four models of resistance evolution, in terms of vaccine efficacy at reducing the
prevalence of resistant pneumococcal carriage. This demonstrates the vaccine efficacy
required to achieve a similar decrease in resistant carriage to a given reduction in
antibiotic prescription rates. The impact on overall pneumococcal carriage is not
considered here. The shaded bar shows an 8.8–23.1% reduction in prescriptions, an
estimate of the percentage of prescriptions which are clinically inappropriate in the UK
(*73*). The dashed line shows a 15% reduction in prescriptions, which has recently been
announced as a target by the UK government (*52*). **(b)** The full impact of vaccination,
illustrated here with 30% vaccine efficacy, can take 5–20 years to play out (mean and
95% HDI). **(c)** Per-country impact of vaccination at 30% efficacy. Countries reporting to
ECDC are ordered from lowest (NL) to highest (CY) reported rate of penicillin
consumption. Diamonds show the estimated change in all pneumococcal pneumonia
cases, while filled distributions show the change in resistant cases.

1039
1040
1041 **Fig. 5. Vaccine impact in a high-burden setting.** Adjusting fitted models to be
1042 consistent with a high-burden setting yields different predictions for vaccine impact,
1043 highlighting both increased challenges and greater opportunities for resistance
1044 management via vaccination.

1045
1046

Fig 6. Explaining additional between-country variation in resistance frequency.
Allowing model parameters to vary across countries captures additional between-country variation in resistance frequency not captured by variation in the treatment rate. For example, **(a)** allowing the transmission rate β to vary across countries can explain the variation in some but not all models. **(b)** Depending upon which parameter is allowed to vary, models differ in how well they explain all additional between-country variation, with a clear separation (dashed line) between flexible and inflexible models. **(c)** Model-specific predictions for the impact of vaccination among those parameters that do fully capture the observed variation remain similar. Solid grey lines show "base" model; dashed lines correspond with colours in panel b.

1057   Supplementary Materials for

1058

1059   Competition and diversity determine vaccine impact

1060   on antibiotic resistance evolution

1061

1062   Nicholas G. Davies, Stefan Flasche, Mark Jit, Katherine E. Atkins

Supplementary Tables 1–7

These tables can be found in an Excel spreadsheet accompanying the article.

Literature review — Details of the literature review used to identify mechanisms for maintaining coexistence between sensitive and resistant bacterial strains.

Summary of model parameters — Table describing model parameters and assumed values or prior distributions for model fitting.

Carriage duration — Calculation of mean pneumococcal carriage duration for children under 5 years old in European settings.

Penicillin consumption — Calculation of the mean number of defined daily doses of penicillin corresponding to a single treatment course for children under 5 years old in European countries.

Pneumococcal morbidity — Calculation of the annual number of pneumococcal pneumonia cases in children under 5 in Europe and Kenya.

Carriage duration (Kilifi) — Calculation of mean pneumococcal carriage duration for children under 5 years old in Kilifi, Kenya.

MCMC diagnostics — Widely Applicable Information Criteria (WAIC), Leave-One-Out Information Criteria (LOOIC), effective posterior sample size and Gelman-Rubin diagnostics for Bayesian inference model fitting using Markov chain Monte Carlo.

1089

Supplementary Fig. 1. Patterns of penicillin non-susceptibility across European
countries, 2005–2017. **(a)** The frequency of penicillin non-susceptibility in
pneumococcal isolates (mean and 95% HDI) increases roughly linearly with the
primary care consumption of penicillins (ATC class J01C). Data from Belgium after 2007
are excluded because of changes to the definition of non-susceptibility after this year.
(b) The frequency of penicillin non-susceptibility has fluctuated in individual countries
between 2005 and 2017, but the European population-weighted average (thick dashed
line) has remained stable at roughly 12%.

1098

Supplementary Fig. 2. Distribution of carriage prevalence and resistance frequency in the "Pathogen diversity" model. This verifies that the D-types with the highest prevalence of carriage (averaged over all countries, above) also exhibit the highest resistance frequency (separated by country, below), and shows that at equilibrium, D-types can exhibit intermediate frequencies of resistance.

1104

1105

1106 **Supplementary Fig. 3.** Maximum *a posteriori* fits for the "Treatment diversity" model

1107 allowing one parameter to vary between countries. Parameters *c* and *z* can capture the

1108 additional variation in resistance frequency between countries.

1109

1110

Supplementary Fig. 4. Maximum *a posteriori* fits for the "Pathogen diversity" model

allowing one parameter to vary between countries. Parameters c, δ, and z can capture

the additional variation in resistance frequency between countries.

1114

1115

1116 **Supplementary Fig. 5.** Maximum *a posteriori* fits for the "Treatment competition"

1117 model allowing one parameter to vary between countries. Parameters β, *c*, *u*, *k*, and *z*

1118 can capture the additional variation in resistance frequency between countries.

1119

1120

1121 **Supplementary Fig. 6.** Maximum *a posteriori* fits for the "Growth competition" model

1122 allowing one parameter to vary between countries. Parameters *b* and *z* can capture the

1123 additional variation in resistance frequency between countries.

1124

1125

1126 **Supplementary Fig. 7.** Impact of vaccination under the "Treatment diversity" model,

1127 for those parameters able to capture the between-country variation in resistance

1128 frequency. The base model fit (thick grey solid line) is compared with the model fits in

1129 which parameters vary between countries (thin dashed lines).

1130

1131

Supplementary Fig. 8. Impact of vaccination under the "Pathogen diversity" model, for those parameters able to capture the between-country variation in resistance frequency. The base model fit (thick grey solid line) is compared with the model fits in which parameters vary between countries (thin dashed lines).

Supplementary Fig. 9. Impact of vaccination under the "Treatment competition"
model, for those parameters able to capture the between-country variation in resistance
frequency. The base model fit (thick grey solid line) is compared with the model fits in
which parameters vary between countries (thin dashed lines).

1142

1143

1144 **Supplementary Fig. 10.** Impact of vaccination under the "Growth competition" model,

1145 for those parameters able to capture the between-country variation in resistance

1146 frequency. The base model fit (thick grey solid line) is compared with the model fits in

1147 which parameters vary between countries (thin dashed lines).

1148

1149

1150 **Supplementary Fig. 11.** Impact of vaccination under the "Treatment diversity" model,

1151 for those parameters *not* able to capture the between-country variation in resistance

1152 frequency. The base model fit (thick grey solid line) is compared with the model fits in

1153 which parameters vary between countries (thin dashed lines).

1154

1155

1156 **Supplementary Fig. 12.** Impact of vaccination under the "Pathogen diversity" model,

1157 for those parameters *not* able to capture the between-country variation in resistance

1158 frequency. The base model fit (thick grey solid line) is compared with the model fits in

1159 which parameters vary between countries (thin dashed lines).

Supplementary Fig. 13. Impact of vaccination under the "Treatment competition" model, for those parameters *not* able to capture the between-country variation in resistance frequency. The base model fit (thick grey solid line) is compared with the model fits in which parameters vary between countries (thin dashed lines).

Growth competition

1166

1167

1168 **Supplementary Fig. 14.** Impact of vaccination under the "Growth competition" model,

1169 for those parameters *not* able to capture the between-country variation in resistance

1170 frequency. The base model fit (thick grey solid line) is compared with the model fits in

1171 which parameters vary between countries (thin dashed lines).