Title

Post-saccadic face processing is modulated by pre-saccadic preview:

Evidence from fixation-related potentials

Authors

Authors

Antimo Buonocore[1,2*], Olaf Dimigen[3*], David Melcher[4]

1. Werner Reichardt Centre for Integrative Neuroscience, University of Tuebingen,

Tuebingen, Germany.

2. Hertie Institute for Clinical Brain Research, Tuebingen University, Tuebingen, Germany,

72076

3. Department of Psychology, Humboldt-Universitaet zu Berlin, Berlin, Germany

4. Center for Mind/Brain Sciences, University of Trento, Rovereto, Italy

* Equal contribution

Corresponding author:

Antimo Buonocore

Werner Reichardt Centre for Integrative Neuroscience

Otfried-Müller Str. 25

Tübingen, 72076, Germany

Tel: +49 7071 29 88821

antimo.buonocore@cin.uni-tuebingen.de

Abstract

Humans actively sample their environment with saccadic eye movements to bring relevant information into high-acuity foveal vision. Despite being lower in resolution, peripheral information is also available prior to each saccade. How pre-saccadic extrafoveal preview of a visual object influences its post-saccadic processing is still an unanswered question. Here, we investigated this question by simultaneously recording behavior and fixation-related brain potentials while human subjects made saccades to face stimuli. We manipulated the relationship between pre-saccadic "previews" and post-saccadic images to explicitly isolate the influences of the former. Subjects performed a gender discrimination task on a newly foveated face under three preview conditions: phase-scrambled face, incongruent face (different identity from the foveated face), and congruent face (same identity). As expected, reaction times were faster after a congruent-face preview compared to the phase-scrambled and incongruent conditions. Importantly, a face preview (either incongruent or congruent) resulted in a strong reduction of post-saccadic neural responses. Specifically, we analyzed the classic face-selective N170 component at occipito-temporal EEG electrodes, which was still present in our experiments with active looking. We found that this component was attenuated for face preview conditions compared to scrambled conditions. This large and long-lasting decrease in evoked activity is consistent with an active prediction mechanism influencing category-specific neural processing at the start of a new fixation. These findings constrain theories of visual stability and show that the extrafoveal preview methodology can be a useful tool to investigate its underlying mechanisms.

Significance Statement

Neural correlates of object recognition have traditionally been studied by flashing stimuli to the central visual field. This procedure differs in fundamental ways from natural vision, where viewers actively sample the environment with eye movements and also obtain a low-resolution preview of soon-to-be-fixated objects. Here we show that the N170, a classic electrophysiological marker of the structural processing of faces, also occurs during a more natural viewing condition but is massively reduced due to extrafoveal preprocessing (preview benefit). Our results therefore highlight the importance of peripheral vision during trans-saccadic processing in building a coherent and stable representation of the world around us.

Introduction

In natural viewing, visual processing takes place primarily during periods of fixation, which are separated by fast and ballistic eye movements known as saccades. Unlike laboratory experiments, in which stimuli appear suddenly, the image that presents itself in the fovea at the beginning of each fixation is usually the result of a choice to fixate that item, typically based on a low-resolution peripheral preview of that object. Whether such a peripheral preview influences visual processing at the beginning of a new fixation, and how this might fit into different competing theories regarding why visual perception seems stable and continuous across saccades, remains an important question.

One view posits that the integration of pre- and post-saccadic information, at the level of neural mechanisms, involves a form of active prediction (Srinivasan et al., 1982; Rao and Ballard, 1999; Clark, 2013). The peripheral preview and the oculomotor plan for the next saccade might be combined to predict where the eye will land and what visual features will be present (for review, Melcher and Colby, 2008; Melcher, 2011). In the case of reading, a classic finding is the preview benefit effect in behavior (Rayner, 1975): when a word is visible in the periphery prior to the saccade, the subsequent fixation on the word is shorter compared to an invalid preview (Dimigen et al., 2012). When looking at fixation-related brain potentials (fERPs), Dimigen and colleagues have shown that such behavioral benefits are also associated with a reduction of the word-specific neural response, an effect termed the "preview positivity" (Dimigen et al., 2012; Kornrumpf et al., 2016).

Along these lines, several recent studies using fMRI have demonstrated a reduction in BOLD responses when a stimulus is consistent across the saccade (Dunkley et al., 2016; Zimmermann et al., 2016; Fairhall et al., 2017). For more complex images such as faces, there is behavioral evidence for an interaction across the saccade that can influence post-saccadic percepts (Melcher, 2005; Wolfe and Whitney, 2014). Recently, Edwards and colleagues (2017) showed that the decoding of post-saccadic EEG

responses to faces was possible earlier when the preview of the target did not change

during the execution of the saccade, suggesting the use of peripheral information.

An alternative view on active prediction focusses instead on the spatial shift of

attention towards the peripheral stimulus. Prior to saccade execution, attention is directed

towards the saccade target (Hoffman and Subramaniam, 1995; Deubel and Schneider,

1996; Zhao et al., 2012; Buonocore et al., 2017) and this attentional shift has been

implicated in many theories of stable perception (Mathôt and Theeuwes, 2011; Melcher,

2011). A key idea here is that selective attention is immediately present at the beginning of

the new fixation, leading to attentional facilitation of post-saccadic processing (for review,

see Mathôt and Theeuwes, 2011). In contrast to active prediction, which typically results in

a reduction in evoked responses, selective attention tends to amplify neural responses (for

review, Thiele and Bellgrove, 2018). In the case of face stimuli, for example, selective

attention enhances the evoked response to the stimulus (Mohamed et al., 2009;

Sreenivasan et al., 2009; Churches et al., 2010). If overt attention shifts are similar to

covert ones, then the post-saccadic fixation-related ERPs would be expected to be larger

in amplitude when a preview is available, due to the target receiving attentional

enhancement. Testing whether there is an increase in neural activity (due to attention)

versus a reduction (due to prediction) has therefore been suggested to be a marker to

differentiate between these two mechanisms (Kok et al., 2012; Spaak et al., 2016; de

Lange et al., 2018).

The aim of the current study was to investigate whether a peripheral preview of a

face image would influence the post-saccadic processing of that face and, if so, whether it

would lead to an increase (attention) or reduction (prediction) in the neural response. We

tested this hypothesis by measuring the effect of the pre-saccadic preview stimulus (either

an intact or phase-scrambled face) on post-saccadic evoked response.

Materials and Methods

Participants

Seventeen volunteers (12 females) between 18 and 30 years of age ($M = 23.6$)

participated in the study. All were free from neurological and visual impairments. The

experiment was conducted in accordance with the Declaration of Helsinki (2008) and

approved by the University of Trento Research Ethics Committee. All participants provided

informed written consent and received a compensation of €10 per hour.

Apparatus

Stimuli were presented on a 24-inch LED monitor (resolution: 1920×1080 pixels,

subtending 39° × 24.7°) at a vertical refresh of 120 Hz. To reduce head movements,

participants were seated with their head stabilized by a chin and forehead rest. The eyes

were horizontally and vertically aligned with the center of the screen at a viewing distance

of 63 cm. Eye movements were recorded with a video-based eye tracker (EyeLink 1000

with desktop mount; SR Research, Ontario, Canada) at a sampling rate of 1000 Hz

(detection algorithm: pupil and corneal reflex; thresholds for saccade detection: 30 deg/s

velocity and 9500 deg/s^2 acceleration). A five point-calibration and validation of the eye

tracker on a standard rectangular grid was run at the beginning of the experiment and

whenever necessary during the experiment. Programs for stimulus presentation and data

collection were written in MATLAB (MathWorks) using the Psychophysics Toolbox Version

3 (Brainard, 1997; Pelli, 1997) and Eyelink Toolbox extensions (Cornelissen et al., 2002).

Participants' manual responses were recorded on a standard keyboard.

The electroencephalogram (EEG) was recorded from 64 Ag/AgCl electrodes (Brain

Products GmbH, Munich, Germany) placed at standard locations of the International 10-10

system. Signals were recorded with a time constant of 10 s and a high cutoff of 250 Hz,

referenced online against the left mastoid, and digitized at a rate of 1000 Hz. The system

was set up with a parallel port splitter so that trigger pulses were sent simultaneously to

the EyeLink and EEG acquisition computers.

Procedure

Participants were seated in a dimly lit room and then briefly familiarized with the task by

the experimenter. Figure 1 illustrates the trial scheme. Participants started each trial by

pressing the space bar while maintaining their gaze at a central fixation cross (0.5° wide,

shown in white on a black background). One second after this button press, two circular

placeholders (white rings, diameter 4°, line width 1 pixel) appeared to the left and right of

the central fixation cross. Placeholders were centered at eccentricities of ±8° and indicated

the positions of the upcoming preview stimuli. Once the eye tracker detected a stable

fixation for 1000 ms within an area of 2° around the central fixation cross, the preview

display was triggered. Depending on the condition, the preview display consisted either of

two different scrambled faces (*scrambled-face preview* condition) or two different intact

faces (*intact-face preview* condition) that appeared at the previous positions of the

placeholders (see Fig. 1, panel "Preview"). After 500 ms of preview, the fixation cross

changed its color and turned either green or red, thereby cueing the participant to execute

a saccade towards the left or right stimulus, respectively (Fig. 1, panel "Saccade cue").

Participants were instructed to respond as quickly and accurately as possible to the cue

with a single saccade.

During the saccade, once gaze position crossed an invisible vertical boundary

placed a distance of 1° from the fixation cross, a scrambled version of the preview face

(that was always different from those shown as previews in the scrambled-face preview

condition) was transiently presented for just a single display cycle (8.3 ms; see Fig. 1,

panel "Transient"). The purpose of this gaze-contingent display change was to introduce

an intra-saccadic visual transient in all experimental conditions, that is, also in the

congruent-face preview condition in which the same face was presented before and after

the saccade. After the transient was displayed, and still during the saccade, the preview

stimulus always changed into an intact face (Fig. 1, panel "Target"). Participants then

responded with a button press whether the face that they had landed on with their eyes

was male or female. Responses were given with the index fingers of the left and right hand

using two keyboard buttons.

The experimental design comprised three main conditions: *scrambled-face preview*,

incongruent-face preview, and *congruent-face preview* (Fig. 1, panel "Preview"). Each

condition comprised 160 trials, leading to a total of 480 trials. Conditions differed in terms

of the stimulus shown before the saccade (preview stimulus). In the *scrambled-face*

preview condition, the stimuli presented during the preview interval were scrambled faces.

In contrast, in both the *incongruent- and congruent-face preview* conditions, the stimuli

shown as previews were intact faces. After the saccade, participants always looked at a

face as the target stimulus. This means that in the *scrambled-face preview* condition, the

scrambled face shown as a preview changed into a face. In the *incongruent-face* preview

condition, the target face shown after the saccade was different from the preview face

seen before the saccade (in this condition, the face shown at the irrelevant screen location

opposite the cued saccade direction remained the same). Finally, in the *congruent-face*

preview condition, the target stimulus was identical to the face presented at this position

before the saccade. The face seen after the saccade was equiprobably male and female

and the gender of the target face was counterbalanced with the preview condition.

Stimuli

Forty-two grayscale images were selected from the Nottingham face database

(http://pics.stir.ac.uk/zips/nottingham.zip), each showing a frontal view of a face (21

female, 21 male) with a neutral facial expression. To standardize the images and to reduce

differences between the genders, a black mask with a circular aperture was applied to

each face to cover the external facial features (e.g. hair, see Fig. 1). The aperture was

centered on the nose, spanned from the forehead to the chin, and subtended a diameter

of 4° of visual angle at the viewing distance of 63 cm.

For each original face stimulus, we also generated a scrambled counterpart that

was used as the pre-saccadic preview stimulus in the scrambled-face preview condition

(see *Procedure*). For this purpose, we calculated the 2D Fourier transform of each face

image and then added a matrix of random phase angles to the existing phase information

of the image. We then performed an inverse Fourier transform, thereby preserving the

original power spectrum of the image. The same circular aperture as for the intact faces

was also applied to the scrambled images.

Finally, for each face image, we selected a second face stimulus that served as the

saccade target in the condition with an incongruent preview as well as a third scrambled-

face stimulus which was used as a transient during the saccade. Specifically, to control for

low-level differences between the face stimuli shown before and after the saccade, we

randomly selected for each image another face stimulus from the pool of 42 face images,

such that their difference in average image luminance (estimated via their RGB grey

values) was less than 4% (i.e. difference < 11 in 8-bit grey values) and not statistically

significant (as confirmed by a one-way ANOVA). In addition, possible differences in image

luminance between the stimulus shown before and after the saccade (see below) were

also controlled by adding luminance as a predictor in the statistical analysis of the EEG

(see section *Single-subject GLM*).

Behavioral screening & analysis

In an initial analysis step, trials were screened for incorrect oculomotor behavior.

Specifically, we removed all trials in which no saccade was executed towards either

stimulus (0.1% of trials) or an eye blink occurred around the time of saccade execution (-

200 to 600 ms around saccade onset; 1.1%). Furthermore, we removed trials in which the

eyes deviated from the central fixation cross by > 2° during the preview interval (1.9%), the

saccadic reaction time (SRT) was extremely short (< 100 ms; 0.8%) or long (> 700 ms,

7.2%), saccade amplitude was extremely small (< 3°; 2.1%) or large (> 10°; 2.9%), or in

which the saccade went in the wrong direction (6.0%). Finally, we excluded trials in which

the saccade-contingent display change was triggered prematurely by drift movements or

microsaccades during the preview interval (11.5%) or in which the main saccade to the

target was followed by a secondary saccade larger than 3° within 150 ms or less (0.2%).

After applying these behavioral criteria, two participants had to be excluded from the

sample due to excessive trials loss (>60%), reducing the final sample to 15 participants.

Manual RTs and response accuracy in the gender discrimination task were then

submitted to repeated-measures ANOVAs on the three-level factor *Preview*. For the

analysis of the button presses, trials with an extreme manual RT (< 200 or > 1000 ms)

were ignored as outliers. Furthermore, one participant was dropped from the manual RT

analysis due to very slow manual RTs and therefore too few remaining trials.

Electrophysiological data analysis

For the electrophysiological analysis, the EEG was first synchronized with the eye-tracking

channels based on the shared trigger pulses using the EYE-EEG toolbox (Dimigen et al.,

2011). The synchronized EEG was then downsampled to 500 Hz, bandpass-filtered from

0.1 to 40 Hz using EEGLAB's (Delorme and Makeig, 2004) finite response filter

(*pop_eegfiltnew.m*) with default settings, and digitally re-referenced to an average

reference. In the next step, ocular EEG artifacts were removed using an optimized eye-

tracker-guided variant of Infomax ICA in EEGLAB. To optimize the ICA decomposition and

the suppression of the myogenic spike potential peaking at saccade onset (Keren et al.,

2011), the ICA was trained on a copy of the data high pass-filtered at 2 Hz (Winkler et al.,

2015) in which EEG sampling points occurring around saccade onsets (-20 to +10 ms)

were overweighted (see Dimigen, 2018). The resulting unmixing weights computed on this

high-pass filtered and optimized training data were then applied to the original unfiltered

recording, and ocular components were automatically flagged using the eye tracker-

guided procedure by Plöchl et al. (2012) with the saccade/fixation variance ratio threshold

set to 1.1 (Plöchl et al., 2012; Dimigen, 2018).

Based on the trials with a correct oculomotor behavior, we then extracted two sets

of 1000 ms long epochs (-300 to 700 ms) from the artifact-corrected EEG. The first set

was cut around the onset of the preview stimuli on the screen (traditional ERP average).

The second set was cut around the onset of the first fixation on the target face following

the saccade (fERP average). To exclude segments with residual non-ocular artifacts, we

removed all epochs containing peak-to-peak voltage differences > 120 µV in any channel

(2.3% of ERP and 2.8% of fERP epochs). Epochs were then baseline-corrected by

subtracting the mean channel voltages in the 200 ms interval before stimulus/fixation

onset, respectively.

Single-subject GLM (first-level analysis)

Stimulus- and fixation-related potentials were analyzed using a massive univariate model

(Smith and Kutas, 2015a) in which a GLM was fitted on each electrode and time point

separately using the *unfold* toolbox (Ehinger and Dimigen, 2018). Analysis of EEG data

with massive univariate models has advantages in terms of higher sensitivity and unbiased

data-driven analysis (Rousselet et al., 2011; Smith and Kutas, 2015a) and allows to

control the effects of continuous covariates on the waveform. For ERPs, the model only

contained the intercept term and one categorical predictor coding whether the preview

stimuli consisted of two scrambled (0) or two intact faces (1). For the fERP analysis, the

predictors in the regression model were a three-level categorial predictor coding the type

of preview shown before the saccade (scrambled, incongruent, congruent) as well as two

continuous linear covariates: saccade amplitude and the preview-target luminance

difference. Saccade amplitude (in degrees of visual angle) was added in the model

because the size of the incoming saccade has a well-established and strong influence on

the amplitude of the post-saccadic neural response (e.g. Thickbroom et al., 1991;

Dandekar et al., 2011). Including saccade amplitude as a nuisance variable in the model

therefore controlled for the slight difference in incoming saccade amplitude (about 0.3°,

see *Results*) between the preview conditions. In addition, we also found that the fERP was

modulated by the difference in mean luminance between the stimulus shown as preview

and the post-saccadic target. The mean luminance difference between both stimuli was

therefore also included as a continuous covariate.

As a control analysis, we repeated our analysis of the fERP using a GLM-based

linear deconvolution technique (also called continuous-time regression, Dandekar et al.,

2011; Smith and Kutas, 2015b; Ehinger and Dimigen, 2018) that is also implemented in

the *unfold* toolbox. In the current experiment, SRTs were about 30 ms longer for the

scrambled-face preview than for the intact-face preview conditions (see *Results*). This

means that the temporal overlap between the ERP evoked by the onset of the saccade

cue (red/green fixation cross) and the fERP evoked at saccade offset differed

systematically between conditions, potentially biasing the results. GLM-based

deconvolution allows us to control this overlapping activity by modeling the response to

both types of events (cue and fixation onset) in the same statistical model. However, since

the results were virtually identical to those obtained with the simpler univariate model, we

only report the results of the latter here.

Group statistics (second-level analysis)

Second-level statistical analyses were performed using the threshold-free cluster

enhancement method (TFCE, Smith and Nichols, 2009; Mensen and Khatami, 2013), a

variant of a cluster-based permutation tests (Maris and Oostenveld, 2007) which controls

for multiple testing across electrodes and time points without the need to define an

arbitrary cluster-forming threshold. Analyses were run using the Matlab implementation of

TFCE (http://github.com/Mensen/ept_TFCE-matlab) based on 2000 random permutations.

For ERPs, we compared the response following an intact-face vs. scrambled-face preview.

For fERPs, we used the ANOVA variant of the TFCE algorithm, followed up by Bonferroni-

corrected pairwise comparisons between the three preview conditions, again using the

TFCE method. For visualization of the TFCE results in Figures 2 and 4, p-values were

thresholded at $p<0.05$, $p<0.01$, and $p<0.005$.

Results

In the following, we first report the neural response evoked by the onset of the preview

stimuli (ERP to intact vs. scrambled faces). This is followed by an analysis of the behavior

and fERP to the post-saccadic face stimulus.

Preview stimulus onset: evoked response (ERP)

The goal of this analysis was to ensure that our stimuli were effective in eliciting typical

face-related ERP components. Figure 2A shows the scalp-topographic difference maps of

the difference between extrafoveal intact-face previews (i.e. two faces presented bilaterally

at ±8° eccentricity) minus scrambled-face previews (two scrambled faces presented at ±8).

Topographies are shown at three latencies after preview stimulus onset, corresponding to

the peaks of the P1 (124 ms), N1 (226 ms) and P3 (350 ms) components. White dots in

the scalp maps indicate electrodes which showed significant differences between intact-

and scrambled- face previews at the given latency (in a pairwise TFCE-based *t*-test). The

grand-mean waveforms in Figure 2B show the stimulus-ERP elicited by the onset of the

bilateral preview display, averaged across two occipito-temporal electrodes over the left

(PO7) and right hemisphere (PO8).

At the earlier latencies, during the P1 component, there was not yet a clear

difference between the ERP responses for the two types of stimuli (intact- vs. scrambled-

faces) beside a small cluster of activation at right frontocentral sites. However, in the

following N1 time window, a strong bilateral negativity emerged at occipital-temporal

electrode sites that was slightly larger over the right hemisphere, as typical for N170 face

effects (Eimer, 2012). Over frontocentral sites, the posterior N170 effect was accompanied

by a corresponding "vertex positive potential" (VPP); a broad positive potential generally

taken to reflect the positive poles of the bilateral dipole pair giving rise to the occipito-

temporal N170 (Eimer, 2012). These results clearly therefore show how the bilateral

presentation of the face preview (dashed green-pink line) led to a markedly different

evoked response than that of the scrambled-face images (blue line) (Fig. 2B); with faces

eliciting a much more pronounced N170 component over occipital-temporal sites (Halgren

et al., 2000; Hoshiyama et al., 2003; Deffke et al., 2007; Gao and Wilson, 2013). In

contrast, only a weaker effect (i.e. smaller frontocentral cluster) of preview type was

observed during the earlier P1 component (Fig. 2A). With a peak at around 226 ms, the

N170 reached its peak about 50 ms later than typically observed (Bentin et al., 1996). A

likely reason for this delayed N170 peak is that the two face stimuli were presented

bilaterally in the extrafoveal visual field, rather than in the fovea. By looking at the full

matrix of TFCE-corrected *p*-values depicted in Fig. 2C, it is clear how clusters of significant

activation arose at around 160 ms after stimulus onset, both over antero-frontal areas and

occipito-temporal electrodes. Although the difference between the intact-face and

scrambled-face preview condition reached its maximum after 224 ms, this effect remained

topographically stable and statistically significant throughout the entire stimulus-locked

analysis period (i.e. until 600 ms after stimuli onset).

Preview effects: behavioral results

Figure 3 summarizes behavioral performance in the task. A first finding is that saccadic

reaction times (SRTs) were affected by the preview condition: SRTs were about 30 ms

faster in trials with an intact compared to a scrambled-face preview (intact vs. scrambled:

$t(14) = -4.673$; $p < 0.0004$) (Fig. 3A). The same pattern was also reflected in the saccade

amplitudes, which were slightly larger (~0.3°) when the preview was an intact rather than a

scrambled face (Fig. 3B, $t(14) = 8.259$; $p < 0.000001$). This pattern of results indicates that

seeing a possible target stimulus – that is, a face – in the periphery enhanced the

preparation of the oculomotor response toward the target.

For the gender discrimination task following the saccade, response accuracy was

generally high (89% correct) and did not differ between preview conditions ($F(2,26) =$

0.475 $p < 0.627$). However, like SRTs, manual RTs for the button press depended strongly

on the preview condition (main effect; $F(2,26) = 8.535$ $p < 0.001$) with numerically shorter

RTs observed in the two conditions in which a congruent- or an incongruent face was

shown as a preview compared to the scrambled-face condition (Figure 3, right panel).

Bonferroni-corrected post-hoc tests confirmed that congruent face previews produced

significantly shorter RTs than scrambled previews, $t(13) = -3.802$; $p < 0.007$. Importantly,

this effect replicates the classic trans-saccadic preview benefit also observed with other

types of stimuli, in particular words (Rayner, 1975; Dimigen et al., 2012). When the

preview was an incongruent face, there was a only a statistical trend for faster RTs as

compared to the scrambled-preview condition, $t(13) = -2.546$; $p < 0.07$. Manual RTs did

not differ significantly between the congruent and incongruent preview condition.

Taken together, these results replicate a robust trans-saccadic benefit for

previewed human faces compared to a non-informative scrambled-preview condition. Both

the initial oculomotor response towards the peripheral face as well as the subsequent

foveal processing of the facial features (necessary for the gender discrimination task) were

significantly enhanced if the extrafoveal preview provided before the saccade was also a

human face, supporting the hypothesis of preview facilitation for the processing of face

stimuli.

Preview effects: evoked response (fERP)

Main goal of the current study was to compare the fixation-related brain response elicited

by the first direct fixation on the target face as a function of the extrafoveal information

available during the *preceding* fixation: a scrambled face, a different person's face, or the

same face.

Figure 4 summarizes the fERP elicited by the first direct fixation on the target face,

that is, after the end of the critical saccade. Top panels (Figure 4A) show the topographic

difference maps for the three contrasts at the peaks of the P1 (106 ms), N1 (180 ms) and

P3 (350 ms) component. The middle panel shows the corresponding fERP waveforms,

averaged again across occipito-temporal electrodes PO7 and PO8. The bottom panel

(Figure 4C) presents the corresponding statistical comparison (TFCE) between the

congruent- and the scrambled-face preview conditions and between the incongruent- and

the scrambled-face preview conditions (Figure 4D).

The first interesting observation is that when contrasting the activity following a

congruent- as compared to an incongruent-face preview (Fig. 4A, top row), there was no

sign of a significant difference across the entire scalp for any of the main components. In

the second and third rows of Fig. 4A, we contrasted the activity of the incongruent- or

congruent- face, respectively, against the scrambled-face preview. Differently from the

previous comparison, it is now evident that seeing a face rather than a scramble-face

stimulus in the periphery led to a completely different response pattern at the time of the

new fixation, once the target face was foveated. While the fixation-related P1 did not differ

between conditions, the following N1 component was dramatically influenced by the type

of preview visible during the preceding fixation. In particular, we report a strong positivity at

occipito-temporal channels, corresponding to a strong attenuation of the fixation-related

N170 in the conditions in which a congruent or incongruent face was visible before the

saccade. This effect was more pronounced over the right hemisphere, and a

corresponding negative pole over central frontal regions, congruent with the activation

pattern observed for the ERP response time-locked to stimulus onset (see section above).

This is especially clear by looking at the three waveforms for the electrodes

PO7/PO8, whereby the incongruent- (green) and congruent- (pink) face preview showed a

strong reduction in the post-saccadic evoked response at the time of the N170, i.e. a

preview positivity effect, when compared to the scrambled-face preview (blue) (Figure 4B).

Figure 4C visualizes the complete *p*-values matrix for the contrast between the congruent-

face minus scrambled-face preview condition across the entire epoch. This plot confirms

that the preview positivity began at around 160 ms and persisted up to about 300 ms after

fixation onset. This was then followed by a later and weaker cluster of activation between

around 360 to 420 ms, which shared a similar scalp topography as the initial N170 effect.

For completeness, we also report the *p*-values matrix for the contrast between the

incongruent-face minus scrambled-face preview condition (Figure 4D). The TFCE

statistical matrix highlights how landing on a different face from the one that was available

during the preview led to an almost identical pattern of activation over all the electrodes for

the entire epoch as in the congruent preview condition.

Discussion

Neural correlates of face and object recognition have traditionally been studied by flashing stimuli to the central visual field during prolonged visual fixation. In contrast, natural vision typically affords an extrafoveal preview of soon-to-be fixated items before they are brought into the fovea with a saccade. Here we show that the extrafoveal preview of a face stimulus leads to a dramatic reduction in the post-saccadic evoked response to that stimulus, compared to a control condition in which a meaningful preview was withheld by scrambling the spectral phase of the preview stimulus. This reduction occurred even though a face stimulus was always present after the saccade, with only the pre-saccadic preview differing between the conditions. Interestingly, this preview reduction was similar for trials in which the exact same face was present across the saccade (congruent-face condition) and when a different face was present after the saccade (incongruent-face condition). In particular, the N170 component, which is classically linked to the structural processing of faces, was substantially reduced in trials with a face preview than those with a scrambled preview. More generally, the fixation-related evoked response to the post-saccadic face was strongly attenuated, consistent with a "preview benefit" (i.e. reduction) in the evoked response as previously observed for visual words (Dimigen et al., 2012). These findings are consistent with claims that both visual and category-specific information about the saccadic target can influence post-saccadic processing of that stimulus (for review, see: Melcher and Colby, 2008; Melcher and Morrone, 2015).

A critical issue in the context of preview studies is whether the peripheral preview influences early or late ERP components (Edwards et al., 2017). Interestingly, in our data, a significant preview effect arose only after the end of the P1 component, around 160 ms after fixation onset. Current theories suggest that face processing involves several stages of neural processing which differ in terms of their feature-selectivity, neural substrate, and associated ERP components. In particular, the *occipital face area* has been implicated in

processing parts of faces, such as eyes or mouth, and suggested to influence the P100

component (Pitcher et al., 2007; Sadeh et al., 2010), while the *fusiform face area* is

associated with processing facial identity and thought to be reflected in N170 activity,

typically associated with "structural encoding" (Halgren et al., 2000; Hoshiyama et al.,

2003; Deffke et al., 2007; Gao and Wilson, 2013). From our results, the peripheral preview

related modulation is therefore more consistent with processing of facial identity and

structural encoding (Sadeh et al., 2010), rather than individual parts. The lack of any

difference between congruent and incongruent previews is also consistent with processing

at the level of the face representation rather than specific local features. This result is

compatible with the task in which the participants were involved, that was a gender

discrimination, suggesting that participants might have focused more on the global

features of the stimuli presented rather than any specific face part. At the same time, the

preview faces in our study were presented in the periphery as part of a bilateral pair of

stimuli, and so future work is needed to investigate whether earlier components might be

influenced when the competition between two stimuli is removed and only one face

stimulus is presented in isolation in the periphery.

The morphology of the effects observed here also argues strongly against an

interpretation of preview effects as merely the absence of surprise or a change in context,

since such effects are more typically reflected in the later centroparietal P3 component

(Sutton et al., 1965; Duncan-Johnson and Donchin, 1977; Donchin, 1981). Indeed we

observed a persistence of positivity also in the later part of the fERP (as well as in the ERP

during fixation), that in the context of face stimuli might be associated with processing of

dynamic aspects of the face in the Superior Temporal Sulcus (Itier and Taylor, 2004;

Sadeh et al., 2010; Dalrymple et al., 2011). Also, there was a face stimulus present after

the saccade in every trial, meaning that the face target was not a surprise in the traditional

sense.

It is important to note that the overall pattern of results is not consistent with

hypotheses based on a primary role for spatial attention or for priming. Prior to saccade

execution attention is allocated to the saccade target (Hoffman and Subramaniam, 1995;

Deubel and Schneider, 1996; Zhao et al., 2012; Buonocore et al., 2017) and such

allocation might be associated with an enhancement of the P1 and N1 components of the

fixation-related ERP (see also: Eimer, 2000; Mohamed et al., 2009; Sreenivasan et al.,

2009; Churches et al., 2010; Meyberg et al., 2015). Results from a number of paradigms

have in fact shown that stimuli that might be relevant as peripheral previews can lead to an

increased evoked response. For example, repeating a face leads to increased (i.e. more

negative) ERP amplitudes in the time window following the peak of the N170 (N250 early

repetition priming effect; Schweinberger et al., 1995) rather than the decreased amplitude

seen here for trans-saccadic preview. On the other hand, prediction mechanisms would

have led to a reduction in evoked responses. Indeed, our results are consistent with the

prediction hypothesis and the idea that prediction and attention can be dissociated by

looking at the direction of effects, with enhanced evoked activity for attention and

attenuated responses for prediction (Kok et al., 2012; Spaak et al., 2016; de Lange et al.,

2018). Nonetheless, we cannot completely rule out that the relatively stronger response in

the scrambled-face condition might be at least partially modulated by the visual

discrepancy between the foveated face stimulus compared to the scrambled stimulus seen

in the extrafoveal visual field during the preview interval. According to this framework, the

larger N1 for the scrambled-face preview condition would share some features with the

visual mismatch negativity (Stefanics et al., 2014; Kornrumpf et al., 2016). One could then

argue that in the scrambled-face preview condition (i.e. the condition where the prediction

is not matched), the onset of the face stimulus might lead to an enhanced "mismatch"

response compared to conditions where participants were seeing a face before and after

the saccade (i.e. conditions where the prediction is matched). In this regard, the

interpretation of the results would be compatible with the idea of an increase in amplitude

for the "unpredicted" condition.

It is also important to note that the fixation-related ERPs elicited by faces brought to

the fovea by the saccade showed a generally similar N170 component to that traditionally

observed for a sudden face onsets in the fovea (Soto et al., 2018). Given that face

perception in real life is typically in response to a face brought to the fovea from the

periphery, rather than a face appearing out of nowhere, it was critically important to show

the ecological validity of such category-specific components. In future work, it would be

interesting to see whether the fixation-related N170 behaves in a similar way to the classic

ERP component under various manipulations such as inversion (Bentin et al., 1996).

Finally, our results are in line with the pattern of results reported in the decoding

study by Edwards and colleagues (Edwards et al., 2017) but also expand on their work in

several ways. First, we could at least in part address the issue of whether the modulation

of the fERP represents a preview benefit rather than a change-related cost, due to

including a scrambled-face condition that was not part of their design. In their study, there

was only a valid or invalid preview condition. Some reduction in decoding performance

might have resulted, for example, from the added noise in the invalid preview condition

due to a visual mismatch negativity-related signal. Another way in which their results might

reflect an invalid cost (rather than validity benefit) is that the use of a decoding measure

means that information about the invalid preview might still have been available well into

the new fixation, with that information interfering with the decoding of the post-saccadic

stimulus. As described above, there is evidence that visual processing during the first 50-

100 ms of a new fixation is driven at least in part by pre-saccadic information, followed by

a potentially shift to processing the new input at the current foveal position. Future work is

needed to clarify the question of how pre-saccadic object-specific information influences

the earliest stages of visual processing and how this might depend on task requirements

(such as the gender task used here).

More generally, the current study provides a proof of concept for the usefulness of

the fERP paradigm for studying trans-saccadic integration and visual stability. Follow-up

studies can use this technique to investigate the mechanisms underlying trans-saccadic

perception to distinguish between competing theories of how our impression of the visual

world remains stable despite the retinal image changing so dramatically an average of

three times per second.

References

Bentin S, Allison T, Puce A, Perez E, McCarthy G (1996) Electrophysiological Studies of Face Perception in Humans. Journal of cognitive neuroscience 8:551-565.

Brainard DH (1997) The Psychophysics Toolbox. Spatial Vision 10:433-436.

Buonocore A, Fracasso A, Melcher D (2017) Pre-saccadic perception: Separate time courses for enhancement and spatial pooling at the saccade target. PLoS ONE 12:e0178902-0178923.

Churches O, Wheelwright S, Baron-Cohen S, Ring H (2010) The N170 is not modulated by attention in autism spectrum conditions. NeuroReport 21:399-403.

Clark A (2013) Whatever next? Predictive brains, situated agents, and the future of cognitive science. Behavioral and Brain Sciences 36:181-204.

Cornelissen FW, Peters EM, Palmer J (2002) The Eyelink Toolbox: Eye tracking with MATLAB and the Psychophysics Toolbox. Behavior research methods, instruments, & computers : a journal of the Psychonomic Society, Inc 34:613-617.

Dalrymple KA, Oruç I, Duchaine B, Pancaroglu R, Fox CJ, Iaria G, Handy TC, Barton JJS (2011) The anatomic basis of the right face-selective N170 IN acquired prosopagnosia: A combined ERP/fMRI study. Neuropsychologia 49:2553-2563.

Dandekar S, Privitera C, Carney T, Klein SA (2011) Neural saccadic response estimation during natural viewing. Journal of Neurophysiology 107:1776-1790.

de Lange FP, Heilbron M, Kok P (2018) How Do Expectations Shape Perception? Trends in Cognitive Sciences 22:764-779.

Deffke I, Sander T, Heidenreich J, Sommer W, Curio G, Trahms L, Lueschow A (2007) MEG/EEG sources of the 170-ms response to faces are co-localized in the fusiform gyrus. NeuroImage 35:1495-1501.

Delorme A, Makeig S (2004) EEGLAB: an open source toolbox for analysis of single-trial

EEG dynamics including independent component analysis. Journal of Neuroscience

Methods 134:9-21.

Deubel H, Schneider WX (1996) Saccade target selection and object recognition:

Evidence for a common attentional mechanism. Vision Research 36:1827-1837.

Dimigen O, Kliegl R, Sommer W (2012) Trans-saccadic parafoveal preview benefits in

fluent reading: A study with fixation-related brain potentials. NeuroImage 62:381-

393.

Dimigen O, Sommer W, Hohlfeld A, Jacobs AM, Kliegl R (2011) Coregistration of eye

movements and EEG in natural reading: analyses and review. J Exp Psychol Gen

140:552-572.

Donchin E (1981) Surprise!… Surprise? Psychophysiology 18:493-513.

Duncan-Johnson CC, Donchin E (1977) On Quantifying Surprise: The Variation of Event-

Related Potentials With Subjective Probability. Psychophysiology 14:456-467.

Dunkley BT, Baltaretu B, Crawford JD (2016) Trans-saccadic interactions in human

parietal and occipital cortex during the retention and comparison of object

orientation. Cortex 82:263-276.

Edwards G, VanRullen R, Cavanagh P (2017) Decoding trans-saccadic memory. The

Journal of Neuroscience.

Ehinger BV, Dimigen O (2018) Unfold: An integrated toolbox for overlap correction, non-

linear modeling, and regression-based EEG analysis. bioRxiv:360156.

Eimer M (2000) Attentional modulations of event-related brain potentials sensitive to faces.

Cognitive Neuropsychology 17:103-116.

Eimer M (2012) The Face-Sensitive N170 Component of the Event-Related Brain

Potential.

Fairhall SL, Schwarzbach J, Lingnau A, Van Koningsbruggen MG, Melcher D (2017)

Spatiotopic updating across saccades revealed by spatially-specific fMRI

adaptation. NeuroImage 147:339-345.

Gao X, Wilson HR (2013) The neural representation of face space dimensions.

Neuropsychologia 51:1787-1793.

Halgren E, Raij T, Marinkovic K, Jousmäki V, Hari R (2000) Cognitive Response Profile of

the Human Fusiform Face Area as Determined by MEG. Cerebral Cortex 10:69-81.

Hoffman JE, Subramaniam B (1995) The role of visual attention in saccadic eye

movements. Perception & Psychophysics 57:787-795.

Hoshiyama M, Kakigi R, Watanabe S, Miki K, Takeshima Y (2003) Brain responses for the

subconscious recognition of faces. Neuroscience Research 46:435-442.

Itier RJ, Taylor MJ (2004) Source analysis of the N170 to faces and objects. NeuroReport

15:1261-1265.

Kok P, Rahnev D, Jehee JFM, Lau HC, de Lange FP (2012) Attention Reverses the Effect

of Prediction in Silencing Sensory Signals. Cerebral Cortex 22:2197-2206.

Kornrumpf B, Niefind F, Sommer W, Dimigen O (2016) Neural Correlates of Word

Recognition: A Systematic Comparison of Natural Reading and Rapid Serial Visual

Presentation. Journal of Cognitive Neuroscience 28:1374-1391.

Maris E, Oostenveld R (2007) Nonparametric statistical testing of EEG- and MEG-data.

Journal of Neuroscience Methods 164:177-190.

Mathôt S, Theeuwes J (2011) Visual attention and stability. Philosophical Transactions of

the Royal Society B: Biological Sciences 366:516.

Melcher D (2005) Spatiotopic Transfer of Visual-Form Adaptation across Saccadic Eye

Movements. Current Biology 15:1745-1748.

Melcher D (2011) Visual stability. Philosophical Transactions of the Royal Society B:

Biological Sciences 366:468-475.

Melcher D, Colby CL (2008) Trans-saccadic perception. Trends in Cognitive Sciences

12:466-473.

Melcher D, Morrone MC (2015) Nonretinotopic visual processing in the brain. Visual

Neuroscience 32:E017.

Mensen A, Khatami R (2013) Advanced EEG analysis using threshold-free cluster-

enhancement and non-parametric statistics. NeuroImage 67:111-118.

Meyberg S, Werkle-Bergner M, Sommer W, Dimigen O (2015) Microsaccade-related brain

potentials signal the focus of visuospatial attention. NeuroImage 104:79-88.

Mohamed TN, Neumann MF, Schweinberger SR (2009) Perceptual load manipulation

reveals sensitivity of the face-selective N170 to attention. NeuroReport 20:782-787.

Pelli DG (1997) The VideoToolbox software for visual psychophysics: transforming

numbers into movies. Spatial Vision 10:437-442.

Pitcher D, Walsh V, Yovel G, Duchaine B (2007) TMS Evidence for the Involvement of the

Right Occipital Face Area in Early Face Processing. Current Biology 17:1568-1573.

Rao RPN, Ballard DH (1999) Predictive coding in the visual cortex: a functional

interpretation of some extra-classical receptive-field effects. Nature Neuroscience

2:79.

Rayner K (1975) The perceptual span and peripheral cues in reading. Cognitive

Psychology 7:65-81.

Rousselet GA, Gaspar CM, Wieczorek KP, Pernet CR (2011) Modeling Single-Trial ERP

Reveals Modulation of Bottom-Up Face Visual Processing by Top-Down Task

Constraints (in Some Subjects). Frontiers in psychology 2:137-137.

Sadeh B, Podlipsky I, Zhdanov A, Yovel G (2010) Event-related potential and functional

MRI measures of face-selectivity are highly correlated: A simultaneous ERP-fMRI

investigation. Human Brain Mapping 31:1490-1501.

Schweinberger SR, Pfütze E-M, Sommer W (1995) Repetition priming and associative

priming of face recognition: Evidence from event-related potentials. Journal of

Experimental Psychology: Learning, Memory, and Cognition 21:722-736.

Smith NJ, Kutas M (2015a) Regression-based estimation of ERP waveforms: I. The rERP

framework. Psychophysiology 52:157-168.

Smith NJ, Kutas M (2015b) Regression-based estimation of ERP waveforms: II. Nonlinear

effects, overlap correction, and practical considerations. Psychophysiology 52:169-

181.

Smith SM, Nichols TE (2009) Threshold-free cluster enhancement: Addressing problems

of smoothing, threshold dependence and localisation in cluster inference.

NeuroImage 44:83-98.

Soto V, Tyson-Carr J, Kokmotou K, Roberts H, Cook S, Fallon N, Giesbrecht T, Stancak A

(2018) Brain Responses to Emotional Faces in Natural Settings: A Wireless Mobile

EEG Recording Study. Frontiers in Psychology 9.

Spaak E, Fonken Y, Jensen O, de Lange FP (2016) The Neural Mechanisms of Prediction

in Visual Search. Cerebral Cortex 26:4327-4336.

Sreenivasan KK, Goldstein JM, Lustig AG, Rivas LR, Jha AP (2009) Attention to faces

modulates early face processing during low but not high face discriminability.

Attention, Perception, & Psychophysics 71:837-846.

Srinivasan MV, Laughlin SB, Dubs A (1982) Predictive coding: a fresh view of inhibition in

the retina. Proceedings of the Royal Society of London Series B, Biological

sciences 216:427-459.

Stefanics G, Kremláček J, Czigler I (2014) Visual mismatch negativity: a predictive coding

view. Frontiers in Human Neuroscience 8.

Sutton S, Braren M, Zubin J, John ER (1965) Evoked-Potential Correlates of Stimulus

Uncertainty. Science 150:1187.

Thickbroom GW, Knezevicˇ W, Carroll WM, Mastaglia FL (1991) Saccade onset and offset

lambda waves: relation to pattern movement visually evoked potentials. Brain

Research 551:150-156.

Thiele A, Bellgrove MA (2018) Neuromodulation of Attention. Neuron 97:769-785.

Wolfe BA, Whitney D (2014) Facilitating recognition of crowded faces with presaccadic

attention. Frontiers in Human Neuroscience 8.

Zhao M, Gersch TM, Schnitzer BS, Dosher BA, Kowler E (2012) Eye movements and

attention: The role of pre-saccadic shifts of attention in perception, memory and the

control of saccades. Vision Research 74:40-60.

Zimmermann E, Weidner R, Abdollahi RO, Fink GR (2016) Spatiotopic Adaptation in

Visual Areas. Journal of Neuroscience 36:9526-9534.

 Figure captions

Figure 1. Trial scheme. At the beginning of each trial, participants fixated a central fixation

cross 1000 ms. Afterwards, two placeholders appeared in the periphery at ±8 degree to

the left and right of fixation (Placeholders panel). After 1000 ms, two preview stimuli

appeared at the position of the placeholders for 500 ms (Preview panel). These stimuli

could be either scrambled-faces (blue outline) or intact-faces or (dashed green-pink

outline). After the preview interval, the central cross turned either green (left) or red (right),

thereby cueing the participant to execute a saccade towards the left of right placeholder,

respectively (Saccade cue panel). During the saccade, the preview was first changed into

a scrambled image patch for one display cycle (8.3 ms) in order to introduce a peri-

saccadic transient in all three conditions (Transient panel). Afterwards, the stimulus

changed to the target face in all conditions (Target panel). The relationship between the

preview stimulus and the target face yielded three conditions for the behavioral and

fixation-related EEG analysis: a scrambled preview condition (blue outline), an incongruent

preview condition (green outline; different face seen before and after saccade), and a

congruent preview condition (pink outline; same face seen before and after saccade).

Participants were asked to discriminate the gender (male/female) of the face visible after

the saccade with a button press. Note that stimuli are not drawn to scale.

Figure 2. Event-related potentials aligned to the onset of the preview display. (A)

Topographic difference maps between intact-face previews minus scrambled-face

previews for three latencies after stimulus onset that represent the peak latencies of the

P1, N1, and P300, respectively. White dots represent electrodes that show significant

differences between the two preview conditions in the TFCE statistic at this latency. (B)

Grand-mean stimulus-locked ERP, averaged over occipito-temporal electrodes PO7 and

PO8 for intact-face previews (green-pink) and scrambled-face previews (blue). (C) Results

of the TFCE statistic comparing face- and scrambled-face previews at all time points and

channels. For visualization, *p*-values are thresholded at 0.05, 0.01, and 0.005 with

different shades of blue.

Figure 3. Behavioral results. Panels show the average (A) saccadic reaction time, (B)

saccadic amplitude, and (C) manual RT for the scrambled-, incongruent-, and congruent-

face preview condition, respectively. Asterisks denote $p < 0.05$. Error bars denote ±1 SEM.

Figure 4. Fixation-related potentials (fERP). (A) Topographic difference maps for the

difference between the congruent vs. incongruent (top row), incongruent vs. scrambled

(middle row) and congruent vs. scrambled preview condition (bottom row) at three

latencies after fixation onset on the target face. The latencies correspond to the P1, N1,

and P300, respectively. (B) Grand-mean fERP averaged across occipito-temporal

electrodes PO7 and PO8 for the scrambled (blue), incongruent (green) and congruent face

preview condition (blue). Note that the three conditions only differ in terms of the stimulus

seen before the saccade, the target face fixated at time zero of this plot was the same in

all three conditions. (C-D) TFCE results for the pairwise comparison between the

congruent- and scrambled-face preview condition and the incongruent- and scrambled-

face preview condition respectively.

Figure 1.

Figure 2.

724   **Figure 3.**

Figure 4.