1   **Behavioral and metabolic consequences after a single generation of angling**

2   **selection in brown trout**

Jenni M. Prokkola[1,2,*], Nico Alioravainen[1], Lauri Mehtätalo[1], Pekka Hyvärinen[3], Alexandre

Lemopoulos[1,4], Sara Metso[1], Anssi Vainikka[1]

[1] University of Eastern Finland, Department of Environmental and Biological Sciences, P.O.

box 111, FI-80101 Joensuu, Finland

[2] University of Liverpool, Institute of Integrative Biology, Crown Street, L69 7ZB Liverpool,

UK

[3] Natural Resources Institute Finland (Luke), Kainuu Fisheries Research Station,

Manamansalontie 90, FI-88300 Paltamo, Finland

[4] University of Turku, Department of Biology, FI-20014 Turku, Finland

*Corresponding author, email jenni.prokkola@liverpool.ac.uk

Abstract

Human induced selection, including mortality caused by recreational angling, can cause phenotypic changes in wild populations. Brown trout (*Salmo trutta*) is an intensively fished salmonid that is also impacted by unintended hatchery-induced domestication, and thus provides a relevant model to study both angling- and hatchery-induced selection. We produced crosses of fish with high or low vulnerability to angling using two populations of brown trout –one wild and one reared in captivity for several generations– and reared the offspring in common garden conditions. We then assessed minimum and average metabolic rates, boldness and sensitivity to stress in juveniles at the age of 1 year. Our results show that angling selection had population-specific effects on risk taking -related latency and exploration tendency, and that populations differed on average in several measured traits, which could be due to a combination of genetic and non-genetic effects. Our study provides evidence for angling induced selection in juvenile personality and suggests that metabolic rate and stress sensitivity may also be affected. The context-dependent effects of angling selection indicate that easy solutions for fisheries management and conservation purposes to mitigate these changes may not exist.

Introduction

Selective harvesting of natural populations by humans induces strong selection (Fugère and Hendry 2018) and can increase the relative frequencies of maladaptive phenotypes (Allendorf and Hard 2009; Coltman et al. 2003). Empirical studies have shown that responses to human-induced selection can be rapid at both genetic (Cooke et al. 2007; Sutter et al. 2012; Uusi-Heikkilä et al. 2015) and phenotypic levels (Kern et al. 2016; Wong et al. 2012). In empirical and simulation studies of heavily exploited fish stocks, fisheries-induced evolution has decreased the size at maturity and growth rate of fish (Audzijonyte et al. 2013; Biro and Post 2008; Devine et al. 2012; Nussle et al. 2009).

In addition to large-scale fisheries using gillnets, trawls and other commercial gear, recreational and small-scale fisheries can also induce selection on vulnerability to fishing and traits that explain vulnerability (Cooke et al. 2007; Hollins et al. 2018; Redpath et al. 2010; Sutter et al. 2012; Uusi-Heikkilä et al. 2008). Regarding behavioral traits, selection from recreational fishing is expected to particularly affect boldness and exploration tendency (Arlinghaus et al. 2017). Bold and explorative fish are often the most vulnerable to angling, likely due to the required behavioral decisions from the fish (Cooke et al. 2007; Härkönen et al. 2014; Wilson et al. 2015), reviewed in (Lennox et al. 2017), though not in all studies (Louison et al. 2017; Vainikka et al. 2016). Over time, angling selection could increase the frequency of shy, slow-growth phenotypes in the population, which could lead to less efficient resource use and decreased population growth and thus diminished productivity (Andersen et al. 2018; Arlinghaus et al. 2017).

Mechanisms mediating angling-induced selection are poorly understood, but several studies have found a correlation between behaviors affecting energy balance and minimum metabolic

rate (meta-analysis by Mathot et al. 2018), implying that selection acting on personality could

thereby affect metabolism in fish. This relationship may be bidirectional, as metabolic rate is

affected by behavior, but may also be the underlying cause for risk-taking behavior,

depending on food availability (Killen et al. 2011). According to the pace-of-life syndrome

(POLS) theory, boldness should correlate positively with metabolic rate, because a fast

metabolic machinery requires high food intake, which again requires bold behavior (Réale et

al. 2010). In one of the first empirical angling selection studies, standard metabolic rate was

found to be 10% lower in a low vulnerability selection line compared to a high vulnerability

selection line in largemouth bass (*Micropterus salmoides*) (Redpath et al. 2010). This

supports the expectation of a positive correlation between vulnerability to angling and

metabolic rate, however, several studies have found no association between these traits

(Louison et al. 2017; Louison et al. 2018; Väätäinen et al. 2018) and more empirical studies in

common garden conditions are needed to address this question.

Angling selection can be understood also in the light of stress coping styles (Louison et al.

2017). Coping styles can be defined as consistent behavioral differences driven by varying

neurochemical stress responses (Schjolden et al. 2005; Vindas et al. 2017a; Vindas et al.

2017b). Proactive coping style is generally bold, routine-based and relies on a sympathetic

stress response (involving catecholamines), while reactive type is shyer, more flexible in

behavior, and relies on a parasympathetic stress response (involving glucocorticoids)

(Koolhaas et al. 2010; Schjolden et al. 2005), although distinct types have not been identified

in all studies, e.g., (Thomson et al. 2011). Selection by angling may therefore affect the

neurochemical stress response of fish due to underlying correlations with behavior. In this

scenario, a fish that responds to the presence of an angler with a high cortisol response is less

likely caught than a non-stressed fish.

As many other taxa, salmonids can display distinctive behavioral strategies/syndromes and

coping styles (Adriaenssens and Johnsson 2011; Brelin et al. 2008; Huntingford and Adams

2005; Näslund and Johnsson 2016; Vindas et al. 2017b), which may provide resource- and

life stage -dependent survival benefits. Salmonids are also affected by domestication in

hatchery rearing, which impacts their life-history strategies, growth and behavior (Araki et al.

2008; Horreo et al. 2018; Huntingford 2004) and can increase their vulnerability to angling

(Klefoth et al. 2013). Brown trout (*Salmo trutta*), are one of the most targeted game fish

species globally and endangered in parts of its native range in Europe. To understand the

human impact on brown trout populations, empirical research on the consequences of angling

is urgently needed. In this study, we asked whether already one generation of angling

selection could induce observable changes in the behavior, metabolic rate, or cortisol response

of brown trout. We studied fish from both wild and hatchery origin to assess the potential

interplay between fishing- and hatchery-induced selection. We hypothesized that offspring

from angling-vulnerable parents would have 1) higher scores in boldness-related behavior, 2)

higher minimum metabolic rate, and 3) lower stress sensitivity compared to fish from non-

vulnerable parents, and 4) that fish from hatchery stock parents would display more proactive

stress coping styles compared to fish from wild parents.

Material and methods

Angling experiment and fish husbandry

Experiments on brown trout were carried out between 2015 and 2017 at the Natural

Resources Institute Finland (Luke) Kainuu Fisheries Research Station (www.kfrs.fi) under

licence obtained from the national Animal Experiment Board in Finland (licence number

ESAVI/3443/04.10.07/2015). Two strains of brown trout were used. Wild, predominantly

non-migratory, parental fish from River Vaarainjoki were captured by electrofishing

(generally non-selective fishing gear) during spawning time in 2010–2012 and brought to the

research station. The second parental strain used was a hatchery strain (so-called Lake

Oulujärvi hatchery broodstock). The parental fish were taken from two year-classes of the 2nd

generation of the brood stock maintained in the same research station. The founders of the

brood stock came from three hatchery stocks established from two source populations and

reared in nearby hatcheries for 3–4 generations. These stocks originated from predominantly

migratory (adfluvial) populations in the region (further details in Lemopoulos et al. (2019)).

Despite originating from the same River Varisjoki watershed, the used populations showed

moderate genetic divergence based on fixation index (F_{ST}-value) of 0.11 (Lemopoulos et al.

2019). The wild population had been exposed to angling more recently than the hatchery

population, although fishing pressure had been weaker than the fishing pressure on the

migratory strain prior to hatchery rearing (P. Hyvärinen, unpublished observation).

During the whole study, fish were fed with commercial fish pellets (Raisio Oyj). In 2015,

hatchery-origin and wild-origin adult fish were exposed to experimental fly fishing and

divided into captured (high vulnerability, HV) and uncaptured (low vulnerability, LV) groups.

Fish were fished in two size-assortative pools for each population during June and July with

fly fishing gear adjusted by the size of the fish in the pools. The wild fish were fished in semi-

natural 50-m^2 ponds with a gravel-bottom outer riffle sections and *ca*. 1 m deep, concrete

inner pool sections (53 and 91 visually size-sorted fish in two ponds). The hatchery fish were

fished in 75-m^2 concrete ponds with no structures (64 larger and 167 smaller fish from two

different cohorts in two ponds). Angling was performed by experienced fly fishers (mainly

A.V.) using unnaturally coloured woolly bugger -type fly patterns tied to barbless hooks.

During angling sessions, an angler fished a pond until a fish took the fly or five minutes

passed, after which angling was continued at earliest one hour later. If a fish was captured,

angling was continued immediately after processing, which included anaesthesia with

benzocaine (40 mg L^{-1}), identification of passive integrated transponder (Oregon RFID) code

or tagging when a pre-existing tag was missing, and measuring total length (to 1 mm) and

weight (to 2 g). Fish that were missing PIT-tags were tagged under the skin next to the dorsal

fin using 12 mm tags at this point. After processing, the fish were transferred to similar ponds

(hatchery fish to a 50 m^2 otherwise similar concrete pond) as used for each population during

angling. After angling trials were finished, on 25 June 2015, all remaining wild fish that were

not captured were collected by dip-netting after draining the experimental angling ponds,

anaesthetized, measured and weighed (mean body lengths of fish uncaptured and captured by

angling: in large fish 457 and 475 mm, respectively, and in small fish 344 and 354,

respectively). Uncaptured wild fish were then combined in the same ponds as the fish

captured by angling. The captured hatchery strain fish were subjected to a second round of

angling ~2 weeks later, where in total eight fish were captured and prioritized for breeding the

HV-line, but this was not done on wild fish due to their limited availability. Angling trials

finished on 8 July 2015, and also hatchery fish were transferred back to their original ponds.

Because of the warm water at the time of finishing the second round of angling, the

uncaptured hatchery fish were not measured to avoid handling-induced stress and mortality.

One deep-hooked small hatchery fish was found dead 5 days and one large hatchery fish 41

days after capture, but otherwise no mortality occurred between angling trials and the

breeding.

The offspring used in this study were obtained from fish bred in four groups (i.e. high- and

low-vulnerability [HV and LV, respectively] within each population) in the autumn of 2015.

A replicated, fully factorial 3 × 3 breeding design was used to create the F_1-generation; males

were crossed with females in all combinations in one matrix, and the matrices replicated three

times for each group, details in Electronic Supplemental Material (ESM1, available online).

In the autumn of 2016, the one-summer-old fish were tagged with individual 12-mm PIT-tags

in the abdominal cavity under anaesthesia (benzocaine). After tagging, the selection lines

were mixed together in two 3.2 m^2 fibreglass rearing tanks.

Photoperiod acclimations

In mid-March 2017, after being reared under constant light, 100 fish were divided into two

different photoperiod groups in 0.4-m^2 green, plastic, flow-through tanks. The tanks were

covered with green nets. The first group continued to be reared under constant light (at water

surface approximately 9 lux, N= 10/group, 40 fish in tank), and the second group received a

12h:12h light-dark (L:D) acclimation (at water surface approximately 12 lux during light

period, N = 15/group divided equally in two tanks, details in ESM1, available online). Fish

were fed using automatic belt feeders (~0.3% fish mass per day) on 5–6 days per week during

approx. 4 h between 8:00 and 20:00 to avoid the entrainment of endogenous rhythms by

feeding. After a minimum two-week acclimation, the metabolic rate measurements were

started.

Measurement of O_2 consumption

The O_2 consumption ($\dot{M}O_2$) was measured as a proxy of metabolic rate (Nelson 2016) using

intermittent flow-through respirometry (Svendsen et al. 2016) with 15–17-min cycles. The

fish were caught by dip-netting under a dim red light into 10-L buckets, identified with a PIT-

reader and transferred to the flow-through measurement chambers immersed in a water bath,

which was also immersed in a flow-through buffer tank. Measurements were started

immediately and continued for approximately 23h, corresponding to 90–96 measurement

cycles for all individuals. After measurements, fish were anesthetized with benzocaine,

measured for total length (to 1 mm) and weighed (to 0.1 g), after which they were transferred

to new 0.4 m^2 tanks similar to those used prior to measurements, with the same photoperiods

as before the measurements. Respirometer chamber oxygen levels were then measured empty

for one cycle to quantify bacterial respiration rates. No measurable respiration was detected

without fish. The slope of the decrease in oxygen level during each 3.5-minute measurement

period was calculated using linear regression in AV Bio-Statistics v. 5.2 (by A.V., available at

http://www.kotikone.fi/ansvain/). Because the $\dot{M}O_2$ of fish was extremely low due to cold

water temperature, we accepted all measurement periods with regression coefficients $R^2 > 0.2$

in the calculation (in total 28 slopes were excluded across all measurements). This was

justified as visual inspection of the data revealed clear negative trends and excluding slopes

with low R^2 would have biased $\dot{M}O_2$ estimates strongly upwards. Further details of the

method are given in ESM1, available online.

The minimum oxygen consumption ($\dot{M}O_{2,min}$) was calculated from the average of the four

least negative slopes after discarding the first, the last and the least negative slope. Values

from three individuals were discarded as outliers (>3x SD difference to the mean). In

addition, we calculated the average consumption across all measurements, excluding the first

and last slope, for each fish ($\dot{M}O_{2,ave}$) because the stress of being confined in the measurement

chamber is reflected in oxygen consumption (Morgan and Iwama 1996; Murray et al. 2017).

The coefficient of the relationship of $\log_{10}(\dot{M}O_{2,min})$ and $\log_{10}$(body mass in kg) was used to

calculate mass-specific $\dot{M}O_{2,min}$ for visualization, after (Killen et al. 2011).

Behavioral trial setup

Quantification of boldness in animals should involve an element of risk-taking. In

experimental settings different measures, such as latency to explore a novel environment, are

often used as proxies for boldness (Conrad et al. 2011; Johnsson and Näslund 2018). Here, we

quantified the boldness of fish using different behaviors expressed in the presence of predator

cues in a novel environment.

The fish were allowed to recover from respirometry for at least four days before behavioral

trials to minimize potential effects of handling stress on behavior. They were not fed for 24-h

prior to behavioral trials. The trials were conducted in custom-made mazes (Fig. 1) (size 400

mm wide x 1500 mm long, water depth 100 mm in the open area). During the trials,

temperature in the maintenance tanks and test arenas was on average $4.5 \pm$ SD $1.3°C$. Water

flow rate during the trials was adjusted to $\sim$8 L min^{-1} ($\sim$7.6–8.8 L min^{-1}). This allowed for at

minimum 1.26 times the arena volume of water to flow between consecutive trials, which was

considered sufficient to minimize potential carry-over effects of chemical cues between trials.

The arena was lit by LED lights (CRI90 LED chain in waterproof silicon tube, 3000-3300K,

4.8W m^2) situated along one long edge of the arena (>70 lux across the arena depending on

distance from light source). Half-way across the arena was a brick gate situated next to one

side, allowing entry from the other side. Behind the brick, natural pebbles ($\sim$3–5 cm in

diameter) were scattered unevenly on the floor, and one large stone was provided for shelter.

A second large stone was placed in the center of the arena in front of the start box. Four

similar arenas were used in the experiment, but they differed in the visual appearance of the

natural stones and two of the arenas were mirror images of the other two with respect to the

location of the gate.

Upstream from the flow-through test arena was a section divided by a metal grid (5 mm mesh

size) where a hatchery-reared burbot (*Lota lota*) (length ~30-40 cm) was placed to introduce

olfactory cues of a natural predator of juvenile brown trout. Burbot are nocturnal bottom-

dwelling predators that are likely difficult for prey to detect visually, but their odor induces

antipredator responses in prey species (Ylönen et al. 2007). Burbot were regularly fed with

pieces of various cyprinids and vendace (*Coregonus albula*) during rearing, and only with

fresh pieces of brown trout for two days prior to and during the trials. Burbot were moved to

the test arenas at least one day before the trials. The burbot were fed with trout pieces in

separate tanks and changed in each arena every 10–15 trials (2–3 days).

Before each trial, individual brown trout were haphazardly removed from their rearing tanks

using a dip-net under red light and placed into black 10-L buckets filled with ~8L of water

from the flow-through system. Fish were identified by PIT tags and left undisturbed for 10

min before being transferred into the start box located downstream from the test arena by

pouring. During each trial, the trout was acclimatized in the start box for 3 min, after which

the door of the box was opened by pulling a string from behind a curtain, and fish movements

recorded from above using two CCTV infrared cameras (two arenas simultaneously filmed

using the same camera) for 10 min (of which first 9 min 45 s was included in the behavior

analysis). The behavioral trial was repeated three times between 8:00 and 11:00 for each focal

fish, with an average time of 4.3 days (range 1–8 days) between consecutive trials. One trial

from four fish was omitted from analysis due to error in data collection. The order in which

batches of four fish were captured on the same day from the same tank for the four arenas was

recorded (batch from hereon, levels 1–5, four individuals from batch 6/7 combined to batch

5).

Testing behavioral responses to burbot

To confirm that burbot odor was perceived risky in the personality assays, we tested for the response of brown trout to burbot in separate controlled tests using individuals from wild HV and wild LV groups (N=10 in each). These fish were acclimated to similar tanks as the personality-tested fish at 12h:12h L:D photoperiod for one week before trials started. One individual died after the first trial. The behavior of each individual was tested on six different days in the presence and absence of predator (3 trials in each condition in haphazard order). 3–4 different arenas were used for each fish on different days to reduce fish habituation to the arenas. These trials were conducted between 14:40 and 17:00. Control arenas were emptied and thoroughly rinsed with pressurized tap water and water flow maintained for >2h before the trials to avoid carry-over effects from burbot odor in earlier experiments. The water used in the flow-through system originates from lake Kivesjärvi, where burbot is a common species; thus, traces of burbot odor may have been present in all trials.

Analysis of video recordings

Behavioral data were collected from videos using manual tracking with AV Bio-Statistics 5.2 timing software. The observer was blind to the identity of fish in all recordings. Analyses were conducted in haphazard order, and each trial was analyzed once. In total four people analyzed the videos. Four behaviors were recorded from the arena trials: 1) *latency* as the time from the start of the experiment until the whole body of fish emerged from the start box, (after Boulton et al. 2014; Moran et al. 2016; Vainikka et al. 2016); 2) time until fish passed the gate to the upstream section of the arena (arrow in Fig. 1), but this was not analyzed because of many fish not entering this section; instead we recorded 3) *exploration tendency* as a binary variable indicating whether the whole body of the fish passed the gate within the arena; and 4) *activity* of fish as the proportion of time spent actively swimming after emerging

from the start box. We used the proportion of time rather than absolute time active to reduce

the dependence of activity from latency. Activity was thus calculated by dividing the total

time when fish did not move when outside the start box by the total time spent outside the

start box and subtracting the value from 1. Stillness was characterized as the fish not moving

forward, backward or sideways for longer than ~2 s. Notably, activity by our definition refers

to short-term activity in a risky, novel environment, not in a familiar environment as it is

classically defined (Conrad et al. 2011), and it was recorded only from the trials in which the

fish emerged from the start box.

Cortisol response to confinement stress

We measured the plasma cortisol levels from of a subset of the fish after exposure to a

standardized confinement stress. During the tests, the fish were transferred to individual dark

brown 10-L plastic buckets with 1.5 L water for 30 min (except for one fish in each Wild HV

and LV and Hatchery HV when the time was 36 min by mistake). The water was aerated

using air stones and pump (Sera Air 550R and Sera AS30 air stone) during the test. The

buckets were placed in a flow-through buffer tank at a temperature matching the acclimation

tanks (temperatures increased during the days of the measurement, 26–29 June 2017, from

13.4 to 16.1°C), and left undisturbed in the dark for the duration of the confinement. Fish

were then removed from buckets by dip-netting, anaesthetized using benzocaine solution,

measured (to 1 mm) and weighed (to 0.1 g). Blood samples were collected within 2–5 min

from the start of anesthesia. The sampling order of fish from the same tank during the same

day was recorded. Blood was collected using 23 G heparinized needles and syringes and kept

on ice temporarily until centrifuged at 4000 x g for 10 min. Plasma was collected in

Eppendorf tubes and frozen at –20°C until analysis. Control samples for establishing baseline

plasma cortisol concentrations were collected after terminal anesthesia as described above,

omitting the confinement stress treatment. Plasma cortisol concentration was determined

using enzyme-linked immunosorbent assay (ELISA) (Enzo cortisol assay) as described in

ESM1, available online.

Sex determination from DNA samples

To consider potential sex differences in the studied traits, we identified the sex of fish using

PCR amplification of the sexually dimorphic *sdY* locus, which identifies the correct sex in

brown trout with nearly 100% accuracy (Quéméré et al. 2014); details in ESM1, available

online.

Statistical analyses

The number of individuals included in each analysis is shown in Table 1. We built univariate

models for each response variable (metabolic and behavioral variables and cortisol level) to

assess the differences between breeding group and acclimation conditions (Table 2). All

analyses were conducted in R v.3.3.2 (R Core Team, 2016). Linear (LMM) and generalized

mixed-effects models (GLMM) were fitted using package *lme4* (Bates et al. 2015) with

lmerTest (Kuznetsova et al. 2017) and the frailty models using package *coxme* (Therneau,

2018). The data were visualized using ggplot2 (Wickham 2009) and patchwork

(https://github.com/thomasp85/patchwork). Statistical significance was determined as $\alpha =$

0.05 in all models. Predicted means within groups were estimated for behavior traits with

package *ggeffects* (Lüdecke 2018). The effect of sex was analyzed in separate models,

including the fixed effect of sex as well as the effects from original models, except

photoperiod or its interactions due to limited sample size with known sex. All linear models

were checked for homoscedasticity and normality of residuals.

Log_{10}-transformed $\dot{M}O_{2,min}$ or $\dot{M}O_{2,ave}$ were analyzed using an LMM with function *lmer*. The

main effects of population, selection line, photoperiod and log_{10}-body mass (in kg) were

separately tested using linear hypothesis testing (function *lht* in package *car*) using restricted

models, where each respective main effect and its interactions were defined zero and

compared to the full model using F-tests.

The difference in cortisol level of control fish and fish exposed to confinement stress was first

tested using a one-tailed t-test. The post-confinement stress cortisol level was then analyzed

using a linear model using function *lm*.

Behavioral traits were analyzed using an LMM (*activity*), a frailty model (i.e. mixed effect

Cox proportional hazards models for time-to event data (Collett 2015)) (*latency*) and a

GLMM (Bernoulli-distributed *exploration tendency*). Trial repeats were encoded as -1, 0, and

1 in data from angling selection experiment as 1–6 from burbot vs. control experiment. In 8

trials, the fish jumped out of the start box prior to the trial and their behavior was analyzed for

9 min 45 s min after the jump. Correlations between metabolic traits and activity were

calculated from model residuals and best linear unbiased predictions (BLUPs), respectively,

to assess potential underlying associations between the traits across all individuals.

Correlations were not calculated for time-to-event data (*latency*) or binary data (*exploration*

tendency). For further details see ESM1, available online, and Data accessibility.

Results

$\dot{M}O_{2,min}$, $\dot{M}O_{2,ave}$ and stress-sensitivity

The LMM indicated significantly higher $\dot{M}O_{2,min}$ in the offspring of wild fish than of hatchery

fish, and a moderate interaction effect between photoperiod and population, wild population

having higher values than hatchery population in the 12:12 L:D photoperiod. Interaction was

also found for population and angling selection line, hatchery LV fish tending to have higher

oxygen uptake than HV fish, while selection lines did not differ in the wild population (Fig.

2A; Table 3). $\dot{M}O_{2,ave}$ was higher in wild than in hatchery population, with a modest

interaction effect of angling selection in the two populations (non-significant, $P = 0.085$),

observed as higher $\dot{M}O_{2,ave}$ in hatchery LV compared to hatchery HV, but no effect of angling

selection in the wild population (Fig. 2B). Sex did not have a significant effect on either

$\dot{M}O_{2,min}$ (Type III test, $F_{1,58} = 0.901$, $P = 0.346$) or $\dot{M}O_{2,ave}$ ($F_{1,58.577} = 0.1823$, $P = 0.671$).

Plasma cortisol increased ~seven-fold in individuals subjected to confinement stress (mean =

140.62 ng mL^{-1}, SD = 41.00) compared to non-stressed fish (mean = 19.22 ng mL^{-1}, SD =

20.65), (t-test, $t = 11.125$, $df = 29.523$, $P < 0.001$). Angling selection or population did not

significantly affect the level of post-stress plasma cortisol, although it showed a similar

tendency as observed in $\dot{M}O_{2,ave}$ (Table 3, Fig. S3, available online).

Behavior in angling selection lines

Fish emerged from the start box during the recorded time in ~84% of the trials. There was a

slightly non-significant interaction effect ($P = 0.054$) of population background and angling

selection on latency (Table 4). This was observed as an elevated probability to emerge in fish

from LV background compared to HV background in the hatchery population, but not in the

wild population (Fig. 3A).

Fish were less active after acclimation in constant light compared to the 12:12 L:D

photoperiod, but activity did not differ between populations or angling selection lines (Fig.

3B; Table 4). Angling selection had contrasting effects on exploration tendency in each

population: in the hatchery population, a higher proportion of fish from LV selection line

were explorative than from HV selection line, while there was an opposite tendency in the

wild population (Fig. 3C; Table 4). In addition, exploration tendency increased with repeats

of the behavioral trial. Sex did not have a significant effect on any behavior trait (female vs

male, Activity: $F_{1,39.612} = 1.217$, $P = 0.277$; Latency: $e^{coef} = 1.03$, $z = 0.29$, $P = 0.770$;

Exploration tendency: $z = –0.514$, $P = 0.607$). There was no correlation between the BLUPs

of activity and residual $\dot{M}O_{2,min}$ (*Pearson r* = 0.02) or $\dot{M}O_{2,ave}$ (*Pearson r* = –0.04).

Behavioral responses to predator presence

The fish tended to be less active ($P = 0.072$) in the presence of burbot than under control

conditions (Table 5). The variance of activity between individuals appeared higher in the

presence of burbot, but this was not significant in Levene's test of homogeneity of variance

($F_{1,93} = 0.214$, $P = 0.645$). Activity decreased slightly with increasing behavior trial repeats.

Latency was not affected by predator cues (non-significant increase in probability to emerge

by 9%), but it increased with increasing behavior trial repeats and between-individual

variation in latency was high (~10% higher variance in burbot vs control data compared to

data from angling selection lines). The exploration tendency of fish was not affected by

predator cues.

Discussion

Population-specific effects of angling selection on boldness

Captured and non-captured parent brown trout produced offspring that differed in boldness-

related traits during their second summer. Surprisingly, boldness, measured as latency to

explore a novel arena, was lower in the HV fish than in the LV fish in the hatchery

population. This was a contradictory finding as angling is expected to select against boldness

(Arlinghaus et al. 2017). The wild population showed a weaker difference between selection

lines, but its direction was more in line with the theory, with HV fish being bolder than LV

fish. Activity during the behavioral trials was not affected by angling selection. In contrast,

activity was affected by photoperiod, which suggests a connection between energy

demand/expenditure and photoperiod through activity.

The unexpected result of angling-vulnerability related low boldness in hatchery population

may be explained by the hatchery rearing environment through several indirect effects. First,

the parent fish that were vulnerable to angling may have had the least strict preference for the

standard feeds, as they accepted the colourful, unnatural fly patterns. Second, the vulnerable

fish may have had the lowest status in the dominance hierarchy within the ponds, and

therefore been the hungriest and likeliest to attack lures. Third, the hatchery-history itself, i.e.

inadvertent domestication, might have selected for the most proactive individuals. This, in

turn, could have shifted the vulnerability in this population towards more reactive fish that

may have more flexibility in their behavior.

In contrast to the hatchery population, angling trials on wild parent fish were more

representative of real angling situations in the field. The wild fish had natural invertebrate

food available in their ponds, and the structured ponds offered more hiding places. The wild

fish had clearly lower catchability than the hatchery fish, and the wild fish could only be

captured when approaching the undisturbed pond from a distance. Very few wild fish were

captured in one angling session (maximum 4) compared to the hatchery fish (maximum 11).

The captured and non-captured parent fish did not show evident size-differences, indicating

that the effects of angling were most likely mediated by size-independent traits.

Stress coping styles and angling selection

A possible consequence of angling selection is a change in the frequencies of not only bold

and timid individuals, but also stress-tolerant and stress-sensitive individuals, due to the

association of angling vulnerability, behavior and stress coping styles (Vindas et al. 2017a).

Furthermore, the POLS theory suggests that stress coping styles, along with metabolic rate

and other traits, are connected to life-history traits, which vary along a slow-fast continuum

(e.g., (Dammhahn et al. 2018; Reale et al. 2010). Stress coping styles and POLS concepts

therefore provide frameworks for testing the consequences of angling induced selection.

In largemouth bass, cortisol response to a standard stressor was negatively associated with

capture probability, as expected (Louison et al. 2017). Koeck et al. (2018) found a weak

negative effect (~0.5% change in risk) of high cortisol response on vulnerability to angling in

a domestic strain of rainbow trout (*Oncorhynchus mykiss*), but there was no similar

relationship in a wild strain of brown trout. In our study, stress sensitivity, expressed as a

cortisol response to confinement, was not affected by angling-selection, although these tests

suffered from low statistical power. However, a higher response in hatchery LV fish

compared to HV was more visible through $\dot{M}O_{2,ave}$, which is notable given that confinement

in the respirometer can induce a stress response in fish (Murray et al. 2017) and increase their

oxygen uptake. Thus, the trends between HV and LV in the hatchery population observed in

the cortisol response and $\dot{M}O_{2,ave}$ suggest potential for angling selection for increased stress

sensitivity, which might become more visible after multiple generations of selection. These

results suggest that the individuals of LV selection line within the hatchery population

showed a more reactive stress coping style than HV line. Differences in coping styles could

also partly explain why the behavior interpreted as boldness showed a pattern contradicting

our expectations; if the LV fish were more reactive compared to the HV fish, their behavior in

the personality trial may have indicated a higher stress response to the experiment and

heightened escape behavior (Laskowski et al. 2016). The stress test and $\dot{M}O_{2,ave}$ results are in

agreement with the hypothesis that the wild population displayed a more reactive coping style

compared to the hatchery population.

Whether $\dot{M}O_{2,min}$ is connected to the coping styles/POLS's or angling selection remains

inconclusive based on our results. A lack of association between metabolic rate and

personality has been reported previously in other species, such as the Trinidadian guppy

(*Poecilia reticulata*) (White et al. 2016). We found a trend of higher $\dot{M}O_{2,min}$ in the hatchery

LV line compared to HV line, and an opposite trend in wild fish, under 12:12 L:D

photoperiod. The contrast between the populations in the response to angling selection is

therefore in line with the behavioral results in latency and exploration tendency, but also with

activity in the response to photoperiod, although there was no correlation between activity and

$\dot{M}O_{2,min}$ after accounting for group differences. Again, as in boldness, the result in hatchery

population was opposite to the prevailing theory, and it is possible that stronger effects from

angling could be observed over several generations of selection in natural conditions. Growth

rate is unlikely to explain the differences in $\dot{M}O_{2,min}$ between groups, as the body mass of fish

at the end of the experiment did not differ between groups (Table 1). Despite not directly

addressing questions on trait covariances, as physiological traits were measured only once

(Mathot and Frankenhuis 2018; Niemelä and Dingemanse 2018), our results add to the

literature to promote the understanding of evolution in traits due to angling induced selection.

From an angling selection perspective, some of the most interesting traits to include in further

experiments would be neurochemical stress responses and their links to the bold-shy

behavioral axis, and a role for metabolic rate remains to be understood better in connection to

these traits.

Genetic and parental effects in population divergence and angling selection

Populations frequently differ in e.g., metabolic rate and behavioral syndromes (Dingemanse et

al. 2007; Lahti et al. 2002; Polverino et al. 2018), driven by environmental differences,

natural selection, founder effects, and genetic drift. The differences we found between

populations can therefore be explained by several factors, including the level of

domestication, as the hatchery stock had been reared in captivity for several generations. They

also differed in their life-histories, with the wild population being clearly resident and the

hatchery population migratory (A. Lemopoulos, unpublished data). In addition, although we

reared offspring under common garden conditions and maximized genetic diversity within

each group, it is possible that differences in the early rearing environments of wild and

hatchery parents could have had contrasting effects on offspring through parental or

epigenetic effects (Crews et al. 2012; Reddon 2012). We studied individuals in their second

summer, and parental effects usually affect early life-stages the most; for instance, maternal

effects on metabolic traits have been shown to be negligible from 90 days post hatching in

coral reef fish (Munday et al. 2017), although maternal stress affects many life-stages in

three-spined stickleback (*Gasterosteus aculeatus*) (Bell et al. 2016; Metzger and Schulte

2016). Additionally, parental effects may have also contributed to differences between the

selection lines *via* stress resulting from angling (briefly increased cortisol level after angling

shown, e.g., in Wilson et al. (2011)). It is nevertheless likely that for both population and

angling selection line differences, genetic inheritance may explain our results at least partly,

as both angling vulnerability and personality traits can be heritable in the studied populations

(Ågren et al. 2019) or in other species (Dingemanse et al. 2009; Philipp et al. 2009).

Potential effects of photoperiod on energy balance

We incorporated environmental variation in our study as two different photoperiods. The

results demonstrate, on one hand, that metabolic rate and swimming activity are sensitive to

photoperiod, and on the other hand, that the other behavioral traits lack this sensitivity.

Constant light is not encountered by brown trout during the winter months; hence the 24-hour

light regime could be considered unnatural and potentially stressful for the fish. Constant light

can disrupt entrainment of endogenous rhythms by inhibiting the synthesis of melatonin and

by directly affecting photosensitive proteins (Falcón et al. 2010; Peirson et al. 2009). Based

on our results, constant light had an inhibiting effect on fish swimming activity, and also

decreased $\dot{M}O_{2,min}$ in the wild population, indicating that energy metabolism in brown trout

can be affected by (an unnatural) photoperiod. In general, non-tropical species are expected to

be particularly sensitive to photoperiod disturbances due to the role of day length in

anticipating seasonal changes in environmental conditions (Borniger et al. 2017).

Innate vs learned antipredatory responses

Our goal was to study risk-taking behavior/boldness of offspring by subjecting fish to the

olfactory cues of a natural predator that had fed on conspecifics. Wild brown trout typically

increase the use of refuges under predation threat, while hatchery brown trout do not (Álvarez

and Nicieza 2003). None of the individuals in the behavior trials in this study had been

exposed to predators before the trials apart from potential traces of piscivore odors in the

rearing water. The scarcity of responses to the presence of predator odor, measured in the

offspring of wild fish, suggests only weak innate responses. Nevertheless, the tendency for

lower activity in the presence of burbot than in control conditions resembles previously

shown antipredator responses in fish (Álvarez and Nicieza 2003; Kopack et al. 2015).

Conclusions

Our results demonstrate the potential for rapid fishing-induced evolution in a very popular

target species. The effects of angling selection are somewhat contradictory between wild and

hatchery populations of fish, which leads to new questions on the mechanisms behind the

observed differences. For practical fisheries management, the results with the wild fish are

more representative and generalizable and showed little evidence of selection in physiological

or behavioral traits. Overall, our study supports earlier findings according to which angling

may be a potentially significant driver of evolution in behavioral and physiological traits in

harvested populations.

Author contributions

A.V. and P.H. produced the selection lines, J.M.P, N.A. and A.V. designed the

experiment, J.M.P., N.A., S.M. and A.L. collected the data, J.M.P. and L.M. analysed the

data, J.M.P. wrote the initial draft of the manuscript. All authors contributed to preparing

the manuscript.

Conflict of Interest

The authors declare that they have no conflict of interest.

Acknowledgements

We thank the staff of Kainuu Fisheries Research Station for their help in catching, breeding

and rearing fish, Dr. Hannu Huuskonen for advice in setting up the respirometry, and Dr.

Chris Elvidge for comments on the manuscript. J.M.P., A.V. and A.L. were supported by the

Academy of Finland grant for A.V. (nr. 286261). J.M.P. was also supported by Oskar

Öflund's foundation and by the Finnish Cultural Foundation. All applicable institutional

and/or national guidelines for the care and use of animals were followed.

Data accessibility

All data and R codes for the models in this manuscript are available in Github

(https://github.com/jprokkola/Strutta_repo). Videos of behavior trials will be made publicly

available in Figshare (accession) upon acceptance for publication.

Electronic Supplementary Material

ESM1. Pdf-file including supplemental figures and methods.

Literature Cited

Adriaenssens B, Johnsson JI. 2011. Shy trout grow faster: exploring links between personality and fitness-related traits in the wild. Behav Ecol 22:135-143. doi: 10.1093/beheco/arq185

Ågren A, Vainikka A, Janhunen M, Hyvärinen P, Piironen J, Kortet R. 2019. Experimental crossbreeding reveals strain-specific variation in mortality, growth and personality in the brown trout (*Salmo trutt*a). Sci Rep 9. doi: 10.1038/s41598-018-35794-6

Allendorf FW, Hard JJ. 2009. Human-induced evolution caused by unnatural selection through harvest of wild animals. Proc Natl Acad Sci USA 106:9987-9994. doi: 10.1073/pnas.0901069106

Álvarez D, Nicieza AG. 2003. Predator avoidance behaviour in wild and hatchery-reared brown trout: the role of experience and domestication. J Fish Biol 63:1565-1577. doi: 10.1111/j.1095-8649.2003.00267.x

Andersen KH, Marty L, Arlinghaus R. 2018. Evolution of boldness and life history in response to selective harvesting. Can J Fish Aquat Sci 75:271-281. doi: 10.1139/cjfas-2016-0350

Araki H, Berejikian BA, Ford MJ, Blouin MS. 2008. Fitness of hatchery-reared salmonids in the wild. Evol Appl 1:342-355. doi: 10.1111/j.1752-4571.2008.00026.x

Arlinghaus R, Laskowski KL, Alós J, Klefoth T, Monk CT, Nakayama S, Schröder A. 2017. Passive gear-induced timidity syndrome in wild fish populations and its potential ecological and managerial implications. Fish Fish 18:360-373. doi: 10.1111/faf.12176

Audzijonyte A, Kuparinen A, Fulton EA. 2013. How fast is fisheries-induced evolution? Quantitative analysis of modelling and empirical studies. Evol Appl 6:585-595. doi: 10.1111/eva.12044

Bates D, Maechler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using

lme4. J Stat Softw, 67:1-48. doi:10.18637/jss.v067.i01

Bell AM, Laura KEM, Stein LR. 2016. Effects of mothers' and fathers' experience with

predation risk on the behavioral development of their offspring in threespined

sticklebacks. Curr Opin Behav Sci 7:28-32. doi: 10.1016/j.cobeha.2015.10.011

Biro PA, Post JR. 2008. Rapid depletion of genotypes with fast growth and bold personality

traits from harvested fish populations. Proc Natl Acad Sci USA 105:2919-2922. doi:

10.1073/pnas.0708159105

Borniger JC, Cisse YM, Nelson RJ, Martin LB. 2017. Seasonal Variation in Stress Responses.

Pp 411-419 in Fink, G., ed. Stress: Neuroendocrinology and Neurobiology,

Amsterdam : Elsevier/AP. doi: 10.1016/b978-0-12-802175-0.00041-3

Boulton K, Grimmer AJ, Rosenthal GG, Walling CA, Wilson AJ. 2014. How stable are

personalities? A multivariate view of behavioural variation over long and short

timescales in the sheepshead swordtail, *Xiphophorus birchmanni*. Behav Ecol

Sociobiol 68:791-803. doi: 10.1007/s00265-014-1692-0

Brelin D, Petersson E, Dannewitz J, Dahl J, Winberg S. 2008. Frequency distribution of

coping strategies in four populations of brown trout (*Salmo trutta*). Horm Behav

53:546-556. doi: 10.1016/j.yhbeh.2007.12.011

Collett D. 2015. Modelling Survival Data in Medical Research, Third Edition. Chapman &

Hall/CRC Texts in Statistical Science.

Coltman DW, O'Donoghue P, Jorgenson JT, Hogg JT, Strobeck C, Festa-Bianchet M. 2003.

Undesirable evolutionary consequences of trophy hunting. Nature 426:655-658. doi:

10.1038/nature02177

Conrad JL, Weinersmith KL, Brodin T, Saltz JB, Sih A. 2011. Behavioural syndromes in

fishes: a review with implications for ecology and fisheries management. J Fish Biol

78:395-435. doi: 10.1111/j.1095-8649.2010.02874.x

Cooke SJ, Suski CD, Ostrand KG, Wahl DH, Philipp DP. 2007. Physiological and behavioral

consequences of long-term artificial selection for vulnerability to recreational angling

in a teleost fish. Physiol Biochem Zool 80:480-490. doi: 10.1086/520618

Crews D, Gillette R, Scarpino SV, Manikkam M, Savenkova MI, Skinner MK. 2012.

Epigenetic transgenerational inheritance of altered stress responses. Proc Natl Acad

Sci USA 109:9143-9148. doi: 10.1073/pnas.1118514109

Dammhahn M, Dingemanse NJ, Niemelä PT, Réale D. 2018. Pace-of-life syndromes: a

framework for the adaptive integration of behaviour, physiology and life history.

Behav Ecol Sociobiol 72:62. doi: 10.1007/s00265-018-2473-y

Devine JA, Wright PJ, Pardoe HE, Heino M. 2012. Comparing rates of contemporary

evolution in life-history traits for exploited fish stocks. Can J Fish Aquat Sci 69:1105-

1120. doi: 10.1139/f2012-047

Dingemanse NJ, Van der Plas F, Wright J, Réale D, Schrama M, Roff DA, Van der Zee E,

Barber I. 2009. Individual experience and evolutionary history of predation affect

expression of heritable variation in fish personality and morphology. Proc Biol Sci

276:1285-1293. doi: 10.1098/rspb.2008.1555

Dingemanse NJ, Wright J, Kazem AJN, Thomas DK, Hickling R, Dawnay N. 2007.

Behavioural syndromes differ predictably between 12 populations of three-spined

stickleback. J Anim Ecol 76:1128-1138. doi: 10.1111/j.1365-2656.2007.01284.x

Falcón J, Migaud H, Muñoz-Cueto JA, Carrillo M. 2010. Current knowledge on the

melatonin system in teleost fish. Gen Comp Endocrinol 165:469-482. doi:

10.1016/j.ygcen.2009.04.026

Fugère V, Hendry AP. 2018. Human influences on the strength of phenotypic selection. Proc

Natl Acad Sci USA 115:10070-10075. doi: 10.1073/pnas.1806013115

Härkönen L, Hyvärinen P, Paappanen J, Vainikka A. 2014. Explorative behavior increases

vulnerability to angling in hatchery-reared brown trout. *Salmo trutta*). Can J Fish

Aquat Sci 71:1900-1909. doi: 10.1139/cjfas-2014-0221

Hollins J, Thambithurai D, Koeck B, Crespel A, Bailey DM, Cooke SJ, Lindström J, Parsons

KJ, Killen SS. 2018. A physiological perspective on fisheries-induced evolution. Evol

Appl 11:561-576. doi: 10.1111/eva.12597

Horreo JL, Valiente AG, Ardura A, Blanco A, Garcia-Gonzalez C, Garcia-Vazquez E. 2018.

Nature versus nurture? Consequences of short captivity in early stages. Ecol Evol

8:521-529. doi: 10.1002/ece3.3555

Huntingford F, Adams C. 2005. Behavioural syndromes in farmed fish: implications for

production and welfare. Behaviour 142:1207-1221.

Huntingford FA. 2004. Implications of domestication and rearing conditions for the behaviour

of cultivated fishes. J Fish Biol 65:122-142. doi: 10.1111/j.1095-8649.2004.00562.x

Johnsson JI, Näslund J. 2018. Studying behavioural variation in salmonids from an ecological

perspective: observations questions methodological considerations. Rev Fish Biol Fish

28:795-823. doi: 10.1007/s11160-018-9532-3

Kern EMA, Robinson D, Gass E, Godwin J, Langerhans RB. 2016. Correlated evolution of

personality, morphology and performance. Anim Behav 117:79-86. doi:

10.1016/j.anbehav.2016.04.007

Killen SS, Marras S, McKenzie DJ. 2011. Fuel, fasting, fear: routine metabolic rate and food

deprivation exert synergistic effects on risk-taking in individual juvenile European sea

bass. J Anim Ecol 80:1024-1033. doi: 10.1111/j.1365-2656.2011.01844.x

Klefoth T, Pieterek T, Arlinghaus R. 2013. Impacts of domestication on angling vulnerability

of common carp, *Cyprinus carpio*: the role of learning, foraging behaviour and food

preferences. Fisheries Management and Ecology 20:174-186. doi: 10.1111/j.1365-

2400.2012.00865.x

Koolhaas JM, de Boer SF, Coppens CM, Buwalda B. 2010. Neuroendocrinology of coping

styles: Towards understanding the biology of individual variation. Front

Neuroendocrinol 31:307-321. doi: 10.1016/j.yfrne.2010.04.001

Kopack CJ, Broder ED, Lepak JM, Fetherman ER, Angeloni LM. 2015. Behavioral responses

of a highly domesticated, predator naive rainbow trout to chemical cues of predation.

Fish Res 169:1-7. doi: 10.1016/j.fishres.2015.04.005

Kuznetsova A, Brockhoff PB, Christensen RHB. 2017). lmerTest Package: Tests in Linear

Mixed Effects Models. J Stat Softw, 82:1–26. doi: 10.18637/jss.v082.i13.

Lahti K, Huuskonen H, Laurila A, Piironen J. 2002. Metabolic rate and aggressiveness

between Brown Trout populations. Funct Ecol 16:167-174. doi: 10.1046/j.1365-

2435.2002.00618.x

Laskowski KL, Monk CT, Polverino G, Alos J, Nakayama S, Staaks G, Mehner T,

Arlinghaus R. 2016. Behaviour in a standardized assay, but not metabolic or growth

rate, predicts behavioural variation in an adult aquatic top predator *Esox lucius* in the

wild. J Fish Biol 88:1544-1563. doi: 10.1111/jfb.12933

Lemopoulos A, Prokkola JM, Uusi-Heikkilä S, Vasemägi A, Huusko A, Hyvärinen P,

Koljonen ML, Koskiniemi J, Vainikka A. 2019. Comparing RADseq and

microsatellites for estimating genetic diversity and relatedness Implications for brown

trout conservation. Ecol Evol 9:2106-2120. doi: 10.1002/ece3.4905

Lennox RJ, Alos J, Arlinghaus R, Horodysky A, Klefoth T, Monk CT, Cooke SJ. 2017. What

688   makes fish vulnerable to capture by hooks? A conceptual framework and a review of

689   key determinants. Fish Fish 18:986-1010. doi: 10.1111/faf.12219

690   Louison MJ, Adhikari S, Stein JA, Suski CD. 2017. Hormonal responsiveness to stress is

691   negatively associated with vulnerability to angling capture in fish. J Exp Biol

692   220:2529-2535. doi: 10.1242/jeb.150730

693   Louison MJ, Stein JA, Suski CD. 2018. Metabolic phenotype is not associated with

694   vulnerability angling in bluegill sunfish (*Lepomis macrochirus*). Can J Zool 96:1264-

695   1271. doi: 10.1139/cjz-2017-0363

696   Lüdecke D. 2018. ggeffects: Tidy Data Frames of Marginal Effects from Regression Models.

697   Journal of Open Source Software 3: 772. doi: 10.21105/joss.00772

698   Mathot KJ, Dingemanse NJ, Nakagawa S. 2019. The covariance between metabolic rate and

699   behaviour varies across behaviours and thermal types: meta-analytic insights. Biol

700   Rev 4:1056-1074. doi:10.1111/brv.12491

701   Mathot KJ, Frankenhuis WE. 2018. Models of pace-of-life syndromes (POLS): a systematic

702   review. Behav Ecol Sociobiol 72. doi: 10.1007/s00265-018-2459-9

703   Metzger DCH, Schulte PM. 2016. Maternal stress has divergent effects on gene expression

704   patterns in the brains of male and female threespine stickleback. Proc Biol Sci 283.

705   doi: 10.1098/rspb.2016.1734

706   Moran NP, Mossop KD, Thompson RM, Wong BBM. 2016. Boldness in extreme

707   environments: temperament divergence in a desert-dwelling fish. Anim Behav

708   122:125-133. doi: 10.1016/j.anbehav.2016.09.024

709   Morgan JD, Iwama GK. 1996. Cortisol-induced changes in oxygen consumption and ionic

710   regulation in coastal cutthroat trout (*Oncorhynchus clarki clarki*) parr. Fish Physiol

711   Biochem 15:385-394. doi: 10.1007/bf01875581

Munday PL, Donelson JM, Domingos JA. 2017. Potential for adaptation to climate change in

a coral reef fish. Global Chang Biol 23:307-317. doi: 10.1111/gcb.13419

Murray L, Rennie MD, Svendsen JC, Enders EC. 2017. Respirometry increases cortisol levels

in rainbow trout *Oncorhynchus mykiss*: implications for measurements of metabolic

rate. J Fish Biol 90:2206-2213. doi: 10.1111/jfb.13292

Näslund J, Johnsson JI. 2016. State-dependent behavior and alternative behavioral strategies

in brown trout (*Salmo trutta* L.) fry. Behav Ecol Sociobiol 70:2111-2125. doi:

10.1007/s00265-016-2215-y

Nelson JA. 2016. Oxygen consumption rate v. rate of energy utilization of fishes: a

comparison and brief history of the two measurements. J Fish Biol 88:10-25. doi:

10.1111/jfb.12824

Niemelä PT, Dingemanse NJ. 2018. Meta-analysis reveals weak associations between

intrinsic state and personality. Proc Biol Sci 285. doi: 10.1098/rspb.2017.2823

Nussle S, Bornand CN, Wedekind C. 2009. Fishery-induced selection on an Alpine whitefish:

quantifying genetic and environmental effects on individual growth rate. Evol Appl

2:200-208. doi: 10.1111/j.1752-4571.2008.00054.x

Peirson SN, Halford S, Foster RG. 2009. The evolution of irradiance detection: melanopsin

and the non-visual opsins. Philos Trans R Soc Lond B Biol Sci 364:2849-2865. doi:

10.1098/rstb.2009.0050

Philipp DP, Cooke SJ, Claussen JE, Koppelman JB, Suski CD, Burkett DP. 2009. Selection

for vulnerability to angling in largemouth bass. Trans Am Fish Soc 138:189-199. doi:

10.1577/t06-243.1

Polverino G, Santostefano F, Diaz-Gil C, Mehner T. 2018. Ecological conditions drive pace-

of-life syndromes by shaping relationships between life history, physiology and

behaviour in two populations of Eastern mosquitofish. Sci Rep 8. doi:

10.1038/s41598-018-33047-0

Quéméré E, Perrier C, Besnard AL, Evanno G, Baglinière JL, Guiguen Y, Launey S. 2014.

An improved PCR-based method for faster sex determination in brown trout (*Salmo*

trutta) and Atlantic salmon (*Salmo salar*). Conserv Genet Resour 6:825-827. doi:

10.1007/s12686-014-0259-8

Réale D, Garant D, Humphries MM, Bergeron P, Careau V, Montiglio PO. 2010. Personality

and the emergence of the pace-of-life syndrome concept at the population level. Philos

Trans R Soc Lond B Biol Sci 365:4051-4063. doi: 10.1098/rstb.2010.0208

Reddon AR. 2012. Parental effects on animal personality. Behav Ecol 23:242-245. doi:

10.1093/beheco/arr210

Redpath TD, Cooke SJ, Suski CD, Arlinghaus R, Couture P, Wahl DH, Philipp DP. 2010.

The metabolic and biochemical basis of vulnerability to recreational angling after

three generations of angling-induced selection in a teleost fish. Can J Fish Aquat Sci

67:1983-1992. doi: 10.1139/f10-120

Schjolden J, Backström T, Pulman KGT, Pottinger TG, Winberg S. 2005. Divergence in

behavioural responses to stress in two strains of rainbow trout (*Oncorhynchus mykiss*)

with contrasting stress responsiveness. Horm Behav 48:537-544. doi:

10.1016/j.yhbeh.2005.04.008

Sutter DAH, Suski CD, Philipp DP, Klefoth T, Wahl DH, Kersten P, Cooke SJ, Arlinghaus R.

2012. Recreational fishing selectively captures individuals with the highest fitness

potential. Proc Natl Acad Sci USA 109:20960-20965. doi: 10.1073/pnas.1212536109

Svendsen MBS, Bushnell PG, Steffensen JF. 2016. Design and setup of intermittent-flow

respirometry system for aquatic organisms. J Fish Biol 88:26-50. doi:

10.1111/jfb.12797

Therneau TM. 2018). coxme: Mixed Effects Cox Models. R package version 2.2-7.

https://CRAN.R-project.org/package=coxme

Thomson JS, Watts PC, Pottinger TG, Sneddon LU. 2011. Physiological and genetic

correlates of boldness: Characterising the mechanisms of behavioural variation in

rainbow trout, *Oncorhynchus mykiss*. Horm Behav 59:67-74. doi:

10.1016/j.yhbeh.2010.10.010

Uusi-Heikkilä S, Whiteley AR, Kuparinen A, Matsumura S, Venturelli PA, Wolter C, Slate J,

Primmer CR, Meinelt T, Killen SS, Bierbach D, Polverino G, Ludwig A, Arlinghaus

R. 2015. The evolutionary legacy of size-selective harvesting extends from genes to

populations. Evol Appl 8:597-620. doi: 10.1111/eva.12268

Uusi-Heikkilä S, Wolter C, Klefoth T, Arlinghaus R. 2008. A behavioral perspective on

fishing-induced evolution. Trends Ecol Evol 23:419-421. doi:

10.1016/j.tree.2008.04.006

Väätäinen R, Huuskonen H, Hyvärinen P, Kekäläinen J, Kortet R, Arnedo MT, Vainikka A.

2018. Do metabolic traits, vulnerability to angling, or capture method explain

boldness variation in Eurasian perch? Physiol Biochem Zool 91:1115-1128. doi:

10.1086/700434

Vainikka A, Tammela I, Hyvärinen P. 2016. Does boldness explain vulnerability to angling in

Eurasian perch *Perca fluviatilis*? Curr Zool 62:109-115. doi: 10.1093/cz/zow003

Vindas MA, Gorissen M, Höglund E, Flik G, Tronci V, Damsgard B, Thörnqvist PO, Nilsen

TO, Winberg S, Øverli Ø, Ebbesson LOE. 2017a. How do individuals cope with

stress? Behavioural, physiological and neuronal differences between proactive and

reactive coping styles in fish. J Exp Biol 220:1524-1532. doi: 10.1242/jeb.153213

Vindas MA, Magnhagen C, Brännäs E, Øverli Ø, Winberg S, Nilsson J, Backström T. 2017b.

Brain cortisol receptor expression differs in Arctic charr displaying opposite coping

styles. Physiol Behav 177:161-168. doi: 10.1016/j.physbeh.2017.04.024

White SJ, Kells TJ, Wilson AJ. 2016. Metabolism, personality and pace of life in the

Trinidadian guppy, *Poecilia reticulata*. Behaviour 153:1517-1543. doi:

10.1163/1568539x-00003375

Wickham H. 2009. ggplot2: Elegant Graphics for Data Analysis. Ggplot2: Elegant Graphics

for Data Analysis:1-212. doi: 10.1007/978-0-387-98141-3

Wilson ADM, Binder TR, McGrath KP, Cooke SJ, Godin JGJ. 2011. Capture technique and

fish personality: angling targets timid bluegill sunfish, *Lepomis macrochirus*. Can J

Fish Aquat Sci 68:749-757. doi: 10.1139/f2011-019

Wilson ADM, Brownscombe JW, Sullivan B, Jain-Schlaepfer S, Cooke SJ. 2015. Does

Angling Technique Selectively Target Fishes Based on Their Behavioural Type?

PLoS One 10. doi: 10.1371/journal.pone.0135848

Wong RY, Perrin F, Oxendine SE, Kezios ZD, Sawyer S, Zhou LR, Dereje S, Godwin J.

2012. Comparing behavioral responses across multiple assays of stress and anxiety in

zebrafish (*Danio rerio*). Behaviour 149:1205-1240. doi: 10.1163/1568539x-00003018

Ylönen H, Kortet R, Myntti J, Vainikka A. 2007. Predator odor recognition and antipredatory

response in fish: does the prey know the predator diel rhythm? Acta Oecol 31:1-7. doi:

10.1016/j.actao.2005.05.007

Tables

Table 1. The number of individuals in each group in each analysis, and fish total body length and mass (mean ± SD) at the end of the experiment.

Photoperiod	Group	N (metabolic rate)	N (latency)	N (activity)	N (stress response)	N (body size)	N males / females (unknown)	Body length / mm	Body mass / g
	Hatchery						9 / 5 (1)		
	HV	15	15	15	7	14		117 ± 10	17.1 ± 4.3
12:12	Hatchery LV	7	14	14	6	14	6 / 2 (7)	115 ± 13	16.5 ± 6.7
	Wild HV	11	14	14	4	14	6 / 6 (3)	117 ± 7	17.4 ± 3.2
	Wild LV	14	15	15	9	14	7 / 7 (1)	115 ± 9	16.6 ± 4.3
	Hatchery						6 / 3 (1)		
	HV	10	10	10	0	9		119 ± 9	17.7 ± 4.7
24	Hatchery LV	7	10	10	0	6	0 / 4 (6)	124 ± 9	19.6 ± 3.9
	Wild HV	8	10	10	0	6	2 / 3 (5)	116 ± 9	17.8 ± 7.1
	Wild LV	6	10	10	0	5	2 / 1 (7)	119 ± 9	17.8 ± 3.0

808 Table 2. The main statistical models used in this study. Abbreviations explained below the

809 table.

Study section	Response variable	Model
I. Angling selection experiment	$Log_{10}(\dot{M}O_{2,min})$ or $Log_{10}(\dot{M}O_{2,ave})$	$y_{ij} = \beta_0 + \beta_1 PHO_{ij} + \beta_2 POP_{ij} + \beta_3 SEL_{ij} + \beta_4 POP_{ij} \times SEL_{ij} + \beta_5 POP_{ij} \times PHO_{ij} + \beta_6 POP_{ij} \times logBM_{ij} + \beta_7 logBM_{ij} + \beta_8 WT_{ij} + p_l + e_{ij}$
	Stress sensitivity (post-stress plasma cortisol)	$y_i = \beta_0 + \beta_1 POP_i + \beta_2 SEL_i + \beta_3 POP_i \times SEL_i + \beta_4 ORD_i + \beta_5 BL_i + \beta_6 WT_i + e_i$
	Activity (proportion of time spent swimming during the trial)	$y_{ijk} = \beta_0 + \beta_1 PHO_{ijk} + \beta_2 POP_{ijk} + \beta_3 SEL_{ijk} + \beta_4 POP_{ijk} \times SEL_{ijk} + \beta_5 POP_{ijk} \times PHO_{ijk} + \beta_6 REP_{ijk} + b_i + c_j + d_k + e_{ijk}$
	Latency to emerge from the box	$\lambda(t) = \lambda_0(t) e^{\beta_1 PHO_{ijk} + \beta_2 POP_{ijk} + \beta_3 SEL_{ijk} + \beta_4 POP_{ijk} \times SEL_{ijk} + \beta_5 POP_{ijk} \times PHO_{ijk} + \beta_6 REP_{ijk} + b_i + c_j + d_k}$
	Exploration tendency (1 = explorative, 0 = unexplorative)	$y_{ij} \sim Bernoulli(p_{ijk})$ $logit(p_{ijk}) = \beta_0 + \beta_1 PHO_{ijk} + \beta_2 POP_{ijk} + \beta_3 SEL_{ijk} + \beta_4 POP_{ijk} \times SEL_{ijk} + \beta_5 POP_{ijk} \times PHO_{ijk} + \beta_6 REP_{ijk} + b_i + c_j + d_k$
II. Behaviour responses to burbot olfactory cues	Activity	$y_{ijk} = \beta_0 + \beta_1 SEL_{ijk} + \beta_2 TRE_{ijk} + \beta_3 REP_{ijk} + \beta_4 BL_{ijk} + b_i^{(1)} CON + b_i^{(2)} BUR + c_j + d_k + e_{ijk}$
	Latency to emerge from the box	$\lambda(t) = \lambda_0(t) e^{\beta_1 SEL_{ijk} + \beta_2 GR_{ijk} + \beta_3 REP_{ijk} + \beta_4 BL_{ijk} + b_i + c_j + d_k}$
	Exploration tendency	$y_{ijk} \sim Bernoulli(p_{ijk})$ $logit(p_{ijk}) = \beta_0 + \beta_1 SEL_{ijk} + \beta_2 GR_{ijk} + \beta_3 REP_{ijk} + b_i + c_j + d_k$

β_0 Intercept, *PHO* Photoperiod, *POP* Population, *SEL* Selection, *logBM* Log_{10} body mass in kg, *WT* Water temperature in °C, *ORD* Capture order from the same tank, *BL* Body length in mm – mean (118.8182 mm for I, 122.4464 mm for II), *REP* Trial repeat, *TRE* Treatment, p_l the random effect for chamber l, b_i random effect for fish i, c_j the random effect for arena j, d_k the random effect for batch k, *e* Residual, λ_0 baseline hazard, *t* time, *CON / BUR* binary explanatory variables for burbot and control treatments.

Table 3. Results from models for $\dot{M}O_{2,min}$ and $\dot{M}O_{2,ave}$ and post-stress plasma cortisol. The

zero levels for contrasts were: photoperiod 12:12, population hatchery, and selection line HV.

For the LMMs, F and P- values for the interactions and temperature effect were obtained from

Type III sums of squares and Satterthwaite approximation for degrees of freedom. For the

other fixed effects, linear hypothesis tests using F-test on restricted models with each main

effect and its interactions set to zero were used – residual degrees of freedom are given for

these tests. For cortisol, Type III F-test shown with population and selection line fixed effects

estimated using linear hypothesis tests. Significant ($P < 0.05$) effects shown in bold. For

intercepts, t-test values are shown.

	Fixed effects	Estimate ± SE	Num df	Res / Den df	F	P
$Log_{10}(\dot{M}O_{2,min})$ (LMM)	Intercept	1.82 ± 0.29	1	70	6.32 (t)	<0.001
	Photoperiod	**0.0024 ± 0.02**	**2**	**67.48**	**4.03**	**0.022**
	Population	**-0.84 ± 0.38**	**4**	**52.71**	**3.01**	**0.026**
	Selection	0.054 ± 0.02	2	68.63	2.50	0.090
	Temperature	-0.0009 ± 0.02	1	70	0.002	0.968
	Log₁₀ body mass	**1.15 ± 0.15**	**2**	**68.86**	**35.22**	**<0.001**
	Pop × selection	-0.068 ± 0.03	1	70	3.86	0.053
	Pop × photoperiod	**-0.07 ± 0.03**	**1**	**70**	**4.51**	**0.037**
	Pop × log₁₀ body mass	**-0.51 ± 0.21**	**1**	**70**	**5.65**	**0.020**
	Random effects	**Variance (SD²)**				
	Chamber	0				
	Residual	0.07^2				

	Fixed effects	Estimate ± SE	Num df	Res / Den df	F	P
$Log_{10}(\dot{M}O_{2,ave})$ (LMM)	Intercept	1.18 ±0.44	1	72.50	2.66 (t)	0.010
	Photoperiod	-0.027 ± 0.04	2	70.178	1.52	0.226
	Population	**-0.72 ± 0.54**	**4**	**70.271**	**5.38**	**<0.001**
	Selection	**0.09 ± 0.04**	**2**	**70.763**	**3.17**	**0.048**
	Temperature	-0.03 ± 0.04	1	72.742	0.68	0.412
	Log₁₀ body mass	0.38 ± 0.22	2	71.883	1.60	0.209
	Pop × selection	-0.09 ± 0.05	1	70.660	3.05	0.085
	Pop × photoperiod	-0.03 ± 0.05	1	70.373	0.36	0.550
	Pop × log₁₀ body mass	-0.49 ± 0.30	1	71.343	2.67	0.107
	Random effects	**Variance (SD²)**				
	Chamber	0.029^2				
	Residual	0.109^2				

	Fixed effects	Estimate ± SE	Sum sq.	Df	Test statistic	P
Post-stress cortisol (LM)	Intercept	331.91 ± 225.90	2963.0	1	2.16 (t)	0.162
	Population	47.84 ± 30.43	3392.6	2	2.47	0.142
	Selection	19.35 ± 25.88	767.1	2	0.56	0.466
	Temperature	-15.22 ± 15.38	1343.9	1	0.98	0.338
	Body length	0.07 ± 0.84	8.3	1	0.006	0.939
	Sampling order	-0.75 ± 3.93	50.4	1	0.04	0.851
	Pop × selection	-19.81 ± 37.59	381.1	1	0.28	0.606
	Residual		20587.6	15		

821
822

Table 4. Results of models for behaviour traits in brown trout from hatchery and wild

populations and two angling selection lines (HV and LV). The zero levels for contrasts in all

models were: photoperiod 12:12, population hatchery, and selection line HV. For model

equations, see Table 2. For Activity, F and P-values for the interactions and trial repeat were

obtained from Type III test, and for the other main effects from linear hypothesis tests using

restricted models with each main effect and its interactions set to zero. Fixed effects with $P <$

0.05 shown in bold. For intercepts, t- or z-test values shown.

	Fixed effects	Estimate ± SE	Num Df	Res/Den Df	F	P
Activity (LMM)	**Intercept**	**0.30 ± 0.04**	**1**	**33.18**	**7.78 (t)**	**<0.001**
	Photoperiod	**-0.75 ± 0.04**	**3**	**84.03**	**6.53**	**0.001**
	Population	0.08 ± 0.06	3	42.50	1.40	0.242
	Selection	-0.01 ± 0.04	2	74.67	0.10	0.903
	Pop × selection	0.025 ± 0.06	1	197.41	1.28	0.259
	Photoperiod × pop	-0.070 ± 0.06	1	81.43	1.37	0.245
	Trial repeat	-0.017 ± 0.02	1	44.24	0.20	0.655
	Random effects	**Variance (SD^2)**				
	ID	0.054^2				
	Batch	0.036^2				
	Arena	0.016^2				

	Fixed effects	Coef	e^{coef}	SE	z	P
Latency (frailty model)	Photoperiod	0.13	1.14	0.26	0.49	0.630
	Population	-0.083	0.92	0.30	-0.27	0.780
	Selection	**0.57**	**1.76**	**0.26**	**2.21**	**0.027**
	Pop × selection	-0.71	0.49	0.37	-1.92	0.054
	Photoperiod × pop	-0.060	0.94	0.38	-0.16	0.870
	Trial repeat	**0.23**	**1.26**	**0.082**	**2.83**	**0.005**
	Random effects	**Variance (SD^2)**				
	ID	0.625^2				
	Batch	0.079^2				
	Arena	0.097^2				

	Fixed effects	Estimate ± SE	Wald χ^2	Df	P	
Exploration tendency (GLMM)	Intercept	0.55 ± 0.63	0.88 (z)	1	0.380	
	Photoperiod	-0.63 ± 0.63	1.80	1	0.179	
	Population	1.07 ± 0.72	0.058	1	0.810	
	Selection	1.54 ± 0.64	0.43	1	0.511	
	Pop × selection	**-2.46 ± 0.92**	**7.18**	**1**	**0.007**	
	Photoperiod × pop	0.07 ± 0.88	0.006	1	0.936	
	Trial repeat	**0.58 ± 0.21**	**7.79**	**1**	**0.005**	
	Random effects	**Variance (SD^2)**				
	ID	1.398^2				
	Batch	0.270^2				
	Arena	0.762^2				

Table 5. Results of models for activity, latency and exploration tendency in the presence of

predatory olfactory cues and control conditions in brown trout. For activity, the t-test was

used with Satterthwaite approximations to degrees of freedom. The model was fit with

restricted maximum likelihood. For latency, proportional hazard estimates (± standard error)

are shown with hazard ratios (e^{coef}). For latency and exploration tendency, Wald Chisquare

test was used to determine significance of fixed effects. The zero levels for contrasts in all

models were: treatment control and selection line HV. Significant effects ($P < 0.05$) shown in

bold.

	Fixed effects	Estimate ± SE	Den Df	t	P
	Intercept	**0.435 ± 0.060**	**23.25**	**7.198**	**<0.001**
	Selection line	0.012 ± 0.065	9.59	0.18	0.861
	Treatment	-0.081 ± 0.042	15.85	-1.928	0.072
	Trial repeat	**-0.029 ± 0.010**	**71.62**	**-2.97**	**0.004**
Activity (LMM)	Body length	-0.00009 ± 0.004	8.63	-0.02	0.984
	Random effects	**Variance (SD²)**			
	ID (burbot)	0.122^2			
	ID (control)	0.077^2			
	Batch	0.000			
	Arena	0.000			
	Residual	1.400^2			
	Fixed effects	Coef ± SE	e^{coef}	z	P
	Selection line	0.336 ± 0.372	1.399	0.900	0.370
	Treatment	0.089 ± 0.220	1.093	0.400	0.690
Latency (frailty model)	Trial repeat	0.113 ± 0.065	1.120	1.759	0.080
	Body length	-0.006 ± 0.026	0.994	-0.230	0.820
	Random effects	**Variance (SD²)**			
	ID	0.661^2			
	Batch	0.210^2			
	Arena	0.080^2			
	Fixed effects	Estimate ± SE	z	P	
	Intercept	1.449 ± 0.801	1.808	0.0706	
Exploration tendency (GLMM)	Selection line	0.659 ± 0.724	0.910	0.363	
	Treatment	-0.63 ± 0.487	-1.294	0.196	
	Trial repeat	-0.066 ± 0.142	-0.465	0.642	
	Random effects	**Variance (SD²)**			

ID	1.140^2
Batch	0.000
Arena	$(1.847 \times 10^{-5})^2$

Figure Legends

Fig.1. 3D-illustration of the arena used in personality trials without the left side wall. Water

flow direction is left-right. Burbot was placed in the area indicated by blue colour, upstream

from the net (inaccessible to the brown trout). The grey box indicates the start box, where the

fish was placed before the start of a trial. Latency was measured as time to emerge from the

box. Activity was measured as swimming activity outside the start box arena after emergence.

Exploration tendency was measured as the whole body of fish passing the gate indicated by

an arrow.

Fig. 2. Means and 75% confidence intervals for A) mass-specific $\dot{M}O_{2,min}$, using mass-scaling

exponent 0.928; and B) mass-specific $\dot{M}O_{2,ave}$ using raw body mass (scaling exponent was

0.19 and was not used). For statistical significance, see Table 3. Wild and hatchery

populations shown with two angling selection lines produced from individuals with high

(HV) or low (LV) vulnerability to angling. 12:12 = L:D rhythm, 24 = continuous light. N in

each group shown in Table 1. Legend shown in A.

Fig. 3. Behavioural differences between two angling vulnerability selection lines (HV – high

vulnerability, LV – low vulnerability) within the hatchery and wild populations. A) Curves

showing the proportion of individuals emerged from the start box, drawn with Kaplan-Meier

estimator. Photoperiods are combined within each breeding group, confidence intervals

omitted for clarity. Higher proportion indicates higher boldness. B) Predicted activity from

LMM with 75% confidence intervals for predicted values. Significantly lower activity was

observed in 24 (constant light) compared to 12:12 (light-dark rhythm) (Table 4). C) Predicted

exploration tendency from GLMM with 75% confidence intervals for predicted values.

Angling selection had opposing effects on exploration tendency in the two populations (Table

4). In B and C, predictions were made for the first trial repeat. For N in each group, see Table

1.

Fig. 1.
873

875   Fig. 2.
876

877
878

 Fig. 3.

880