Behavioural and metabolic consequences after a single generation of angling selection in brown trout

Jenni M. Prokkola[1,2,*], Nico Alioravainen[1], Lauri Mehtätalo[1], Pekka Hyvärinen[3], Alexandre Lemopoulos[1,4], Sara Metso[1], Anssi Vainikka[1]

[1] University of Eastern Finland, Department of Environmental and Biological Sciences, P.O. box 111, FI-80101 Joensuu, Finland

[2] University of Liverpool, Institute of Integrative Biology, Crown Street, L69 7ZB Liverpool, UK

[3] Natural Resources Institute Finland (Luke), Kainuu Fisheries Research Station, Manamansalontie 90, FI-88300 Paltamo, Finland

[4] University of Turku, Department of Biology, FI-20014 Turku, Finland

*Corresponding author, email jenni.prokkola@liverpool.ac.uk, tel. +358 40 747 8387

Author contributions

A.V. and P.H. conceived and produced the selection lines, J.M.P, N.A. and A.V. designed the experiment, J.M.P., N.A., S.M. and A.L. collected the data, J.M.P. and L.M. analysed the data, J.M.P. wrote the initial draft of the manuscript. All authors contributed to preparing the manuscript.

Abstract

Human induced selection, including mortality caused by recreational angling, can cause phenotypic changes in wild populations. Brown trout (*Salmo trutta*) is an intensively fished salmonid that is also impacted by unintended hatchery-induced domestication, and thus provides a relevant model to study both fishing- and hatchery-induced selection. We produced crosses of fish with high or low vulnerability to angling using two populations of brown trout – one wild and one reared in captivity for several generations – and reared the offspring in common garden conditions. We then assessed minimum and average metabolic rates, boldness and sensitivity to stress in juveniles at the age of 1 year. Our results show that angling selection had population-specific effects on boldness-related latency and exploration tendency, and that populations differed on average in several measured traits, which could be due to a combination of genetic and non-genetic effects. Our study provides evidence for angling induced selection in juvenile personality and suggests that metabolic rate and stress sensitivity may also be affected. The context-dependent effects of angling selection indicate that easy solutions for fisheries management and conservation purposes to mitigate these changes may not exist.

Keywords

Fisheries, metabolism, animal personality, transgenerational effects

Introduction

Selective harvesting of natural populations by humans induces strong selection (Fugère and

Hendry 2018) and can increase the relative frequencies of maladaptive phenotypes (Allendorf

and Hard 2009; Coltman et al. 2003). Empirical studies have shown that responses to human-

induced selection can be rapid at both genetic (Cooke et al. 2007; Sutter et al. 2012; Uusi-

Heikkilä et al. 2015) and phenotypic levels (Kern et al. 2016; Wong et al. 2012). In empirical and

simulation studies of heavily exploited fish stocks, fisheries-induced evolution has decreased the

size at maturity and growth rate of fish (Audzijonyte et al. 2013; Biro and Post 2008; Devine et

al. 2012; Nussle et al. 2009).

In addition to large-scale fisheries using gillnets, trawls and other commercial gear, recreational

and small-scale fisheries can also induce selection on vulnerability to fishing and traits that

explain vulnerability (Cooke et al. 2007; Hollins et al. 2018; Redpath et al. 2010; Sutter et al.

2012; Uusi-Heikkilä et al. 2008). Regarding behavioural traits, selection from recreational

fishing is expected to particularly affect boldness and exploration tendency (Arlinghaus et al.

2017). Bold and explorative fish are often the most vulnerable to angling, likely due to the

required behavioural decisions from the fish (Cooke et al. 2007; Härkönen et al. 2014; Wilson et

al. 2015), reviewed in (Lennox et al. 2017), though not in all studies (Louison et al. 2017;

Vainikka et al. 2016). Over time, angling selection could increase the frequency of shy, slow-

growth phenotypes in the population, which could lead to less efficient resource use and

decreased population growth and thus diminished productivity (Andersen et al. 2018;

Arlinghaus et al. 2017).

Mechanisms mediating angling-induced selection are poorly understood, but several studies

have found a correlation between behaviours affecting energy balance and minimum metabolic

rate (meta-analysis by Mathot et al. 2018), implying that selection acting on personality could

thereby affect metabolism in fish. This relationship may be bidirectional, as metabolic rate is

affected by behaviour, but may also be the underlying cause for risk-taking behaviour,

depending on food availability (Killen et al. 2011). According to the pace-of-life syndrome

(POLS) theory, boldness should correlate positively with metabolic rate, because a fast

metabolic machinery requires high food intake, which again requires bold behaviour (Réale et

al. 2010). In one of the first empirical angling selection studies, standard metabolic rate was

found to be 10% lower in a low vulnerability selection line compared to a high vulnerability

selection line in largemouth bass (*Micropterus salmoides*) (Redpath et al. 2010). This supports

the expectation of a positive correlation between vulnerability to angling and metabolic rate,

however, several studies have found no association between these traits (Louison et al. 2017;

Louison et al. 2018; Väätäinen et al. 2018) and more empirical studies in common garden

conditions are needed to address this question.

Angling selection can be understood also in the light of stress coping styles (Louison et al. 2017).

Coping styles can be defined as consistent behavioural differences driven by varying

neurochemical stress responses (Schjolden et al. 2005; Vindas et al. 2017a; Vindas et al. 2017b).

Proactive coping style is generally bold, routine-based and relies on a sympathetic stress

response (involving catecholamines), while reactive type is shyer, more flexible in behaviour,

and relies on a parasympathetic stress response (involving glucocorticoids) (Koolhaas et al.

2010; Schjolden et al. 2005), although distinct types have not been identified in all studies, e.g.,

(Thomson et al. 2011). Selection by angling may therefore affect the neurochemical stress

response of fish due to underlying correlations with behaviour. In this scenario, a fish that

responds to the presence of an angler with a high cortisol response is less likely caught than a

non-stressed fish.

As many other taxa, salmonids can display distinctive behavioural strategies/syndromes and

coping styles (Adriaenssens and Johnsson 2011; Brelin et al. 2008; Huntingford and Adams

2005; Näslund and Johnsson 2016; Vindas et al. 2017b), which may provide resource- and life

stage -dependent survival benefits. Salmonids are also affected by domestication in hatchery

rearing, which impacts their life-history strategies, growth and behaviour (Araki et al. 2008;

Horreo et al. 2018; Huntingford 2004) and can increase their vulnerability to angling (Klefoth et

al. 2013). Brown trout (*Salmo trutta*), are one of the most targeted game fish species globally

and endangered in parts of its native range in Europe. To understand the human impact on

brown trout populations, empirical research on the consequences of angling is urgently needed.

In this study, we asked whether already one generation of angling selection could induce

observable changes in the behaviour, metabolic rate, or cortisol response of brown trout. We

studied fish from both wild and hatchery origin to assess the potential interplay between

fishing- and hatchery-induced selection. We hypothesized that offspring from angling-

vulnerable parents would have 1) higher scores in boldness-related behaviour, 2) higher

minimum metabolic rate, and 3) lower stress sensitivity compared to fish from non-vulnerable

parents, and 4) that fish from hatchery stock parents would display more proactive stress

coping styles compared to fish from wild parents.

Material and methods

Angling experiment and fish husbandry

Experiments on brown trout were carried out between 2015 and 2017 at the Natural Resources

Institute Finland (Luke) Kainuu Fisheries Research Station (www.kfrs.fi) under licence obtained

from the national Animal Experiment Board in Finland (licence number

ESAVI/3443/04.10.07/2015). Two strains of brown trout were used. Wild, predominantly non-

migratory, parental fish from River Vaarainjoki were captured by electrofishing (generally non-

selective fishing gear) during spawning time in 2010–2012 and brought to the research station.

The second parental strain used was a hatchery strain (so-called Lake Oulujärvi hatchery

broodstock). The parental fish were taken from two year-classes of the 2nd generation of the

brood stock maintained in the same research station. The founders of the brood stock came from

three hatchery stocks established from two source populations and reared in nearby hatcheries

for 3–4 generations. These stocks originated from predominantly migratory (adfluvial)

populations in the region (further details in Lemopoulos et al. (2019)). Despite originating from

the same River Varisjoki watershed, the used populations showed moderate genetic divergence

based on fixation index (F_{ST}-value) of 0.11 (Lemopoulos et al. 2019). The wild population had

been exposed to angling more recently than the hatchery population, although fishing pressure

had been weaker than the fishing pressure on the migratory strain prior to hatchery rearing (P.

Hyvärinen, unpublished observation).

During the whole study, fish were fed with commercial fish pellets (Raisio Oyj). In 2015,

hatchery-origin and wild-origin adult fish were exposed to experimental fly fishing and divided

into captured (high vulnerability, HV) and uncaptured (low vulnerability, LV) groups. Fish were

fished in two size-assortative pools for each population during June and July with fly fishing gear

adjusted by the size of the fish in the pools. The wild fish were fished in semi-natural 50-m²

ponds with a gravel-bottom outer riffle sections and *ca.* 1 m deep, concrete inner pool sections

(53 and 91 visually size-sorted fish in two ponds). The hatchery fish were fished in 75-m²

concrete ponds with no structures (64 larger and 167 smaller fish from two different cohorts in

two ponds). Angling was performed by experienced fly fishers (mainly A.V.) using unnaturally

coloured woolly bugger -type fly patterns tied to barbless hooks. During angling sessions, an

angler fished a pond until a fish took the fly or five minutes passed, after which angling was

continued at earliest one hour later. If a fish was captured, angling was continued immediately

after processing, which included anaesthesia with benzocaine (40 mg L^{-1}), identification of

passive integrated transponder (Oregon RFID) code or tagging when a pre-existing tag was

missing, and measuring total length (to 1 mm) and weight (to 2 g). Fish that were missing PIT-

tags were tagged under the skin next to the dorsal fin using 12 mm tags at this point. After

processing, the fish were transferred to similar ponds (hatchery fish to a 50 m² otherwise

similar concrete pond) as used for each population during angling. After angling trials were

finished, on 25 June 2015, all remaining wild fish that were not captured were collected by dip-

netting after draining the experimental angling ponds, anaesthetized, measured and weighed

(mean body lengths of fish uncaptured and captured by angling: in large fish 457 and 475 mm,

respectively, and in small fish 344 and 354, respectively). Uncaptured wild fish were then

combined in the same ponds as the fish captured by angling. The captured hatchery strain fish

were subjected to a second round of angling ~2 weeks later, where in total eight fish were

captured and prioritized for breeding the HV-line, but this was not done on wild fish due to their

limited availability. Angling trials finished on 8 July 2015, and also hatchery fish were

transferred back to their original ponds. Because of the warm water at the time of finishing the

second round of angling, the uncaptured hatchery fish were not measured to avoid handling-

induced stress and mortality. One deep-hooked small hatchery fish was found dead 5 days and

one large hatchery fish 41 days after capture, but otherwise no mortality occurred between

angling trials and the breeding.

The offspring used in this study were obtained from fish bred in four groups (i.e. high- and low-

vulnerability [HV and LV, respectively] within each population) in the autumn of 2015. A

replicated, fully factorial 3 × 3 breeding design was used to create the F_1-generation; males were

crossed with females in all combinations in one matrix, and the matrices replicated three times

for each group, details in Electronic Supplemental Material (ESM1). In the autumn of 2016, the

one-summer-old fish were tagged with individual 12-mm PIT-tags in the abdominal cavity under

anaesthesia (benzocaine). After tagging, the selection lines were mixed together in two 3.2 m^2

fibreglass rearing tanks.

Photoperiod acclimations

In mid-March 2017, after being reared under constant light, 100 fish were divided into two

different photoperiod groups in 0.4-m^2 green, plastic, flow-through tanks. The tanks were

covered with green nets. The first group continued to be reared under constant light (at water

surface approximately 9 lux, N= 10/group, 40 fish in tank), and the second group received a

12h:12h light-dark (L:D) acclimation (at water surface approximately 12 lux during light period,

N = 15/group divided equally in two tanks, details in ESM1). Fish were fed using automatic belt

feeders (~0.3% fish mass per day) on 5–6 days per week during approx. 4 h between 8:00 and

20:00 to avoid the entrainment of endogenous rhythms by feeding. After a minimum two-week

acclimation, the metabolic rate measurements were started.

Measurement of O_2 consumption

The O_2 consumption ($\dot{M}O_2$) was measured as a proxy of metabolic rate (Nelson 2016) using

intermittent flow-through respirometry (Svendsen et al. 2016) with 15–17-min cycles. The fish

were caught by dip-netting under a dim red light into 10-L buckets, identified with a PIT-reader

and transferred to the flow-through measurement chambers immersed in a water bath, which

was also immersed in a flow-through buffer tank. Measurements were started immediately and

continued for approximately 23h, corresponding to 90–96 measurement cycles for all

individuals. After measurements, fish were anesthetized with benzocaine, measured for total

length (to 1 mm) and weighed (to 0.1 g), after which they were transferred to new 0.4 m² tanks

similar to those used prior to measurements, with the same photoperiods as before the

measurements. Respirometer chamber oxygen levels were then measured empty for one cycle

to quantify bacterial respiration rates. No measurable respiration was detected without fish. The

slope of the decrease in oxygen level during each 3.5-minute measurement period was

calculated using linear regression in AV Bio-Statistics v. 5.2 (by A.V., available at

http://www.kotikone.fi/ansvain/). Because the $\dot{M}O_2$ of fish was extremely low due to cold water

temperature, we accepted all measurement periods with regression coefficients $R^2 > 0.2$ in the

calculation (in total 28 slopes were excluded across all measurements). This was justified as

visual inspection of the data revealed clear negative trends and excluding slopes with low R^2

would have biased $\dot{M}O_2$ estimates strongly upwards. Further details of the method are given in

ESM1.

The minimum oxygen consumption ($\dot{M}O_{2,min}$) was calculated from the average of the four least

negative slopes after discarding the first, the last and the least negative slope. Values from three

individuals were discarded as outliers (>3x SD difference to the mean). In addition, we

calculated the average consumption across all measurements, excluding the first and last slope,

for each fish ($\dot{M}O_{2,ave}$) because the stress of being confined in the measurement chamber is

reflected in oxygen consumption (Morgan and Iwama 1996; Murray et al. 2017). The coefficient

of the relationship of $\log_{10}(\dot{M}O_{2,min})$ and $\log_{10}$(body mass in kg) was used to calculate mass-

specific $\dot{M}O_{2,min}$ for visualization, after (Killen et al. 2011).

Behavioural trial setup

Quantification of boldness in animals should involve an element of risk-taking. In experimental

settings different measures, such as latency to explore a novel environment, are often used as

proxies for boldness (Conrad et al. 2011; Johnsson and Näslund 2018). Here, we quantified the

boldness of fish using different behaviours expressed in the presence of predator cues in a novel

environment.

The fish were allowed to recover from the $\dot{M}O_{2,min}$ measurement for at least four days before

behavioural trials to minimize potential effects of handling stress on behaviour. They were not

fed for 24-h prior to behavioural trials. The trials were conducted in custom-made mazes (Fig. 1)

(size 400 mm wide x 1500 mm long, water depth 100 mm in the open area). During the trials,

temperature in the maintenance tanks and test arenas was on average 4.5 ± SD 1.3°C. Water

flow rate during the trials was adjusted to ~8 L min^{-1} (~7.6–8.8 L min^{-1}). This allowed for at

minimum 1.26 times the arena volume of water to flow between consecutive trials, which was

considered sufficient to minimize potential carry-over effects of chemical cues between trials.

The arena was lit by LED lights (CRI90 LED chain in waterproof silicon tube, 3000-3300K, 4.8W

m^2) situated along one long edge of the arena (>70 lux across the arena depending on distance

from light source). Half-way across the arena was a brick gate situated next to one side, allowing

entry from the other side. Behind the brick, natural pebbles (~3–5 cm in diameter) were

scattered unevenly on the floor, and one large stone was provided for shelter. A second large

stone was placed in the centre of the arena in front of the start box. Four similar arenas were

used in the experiment, but they differed in the visual appearance of the natural stones and two

of the arenas were mirror images of the other two with respect to the location of the gate.

Upstream from the flow-through test arena was a section divided by a metal grid (5 mm mesh

size) where a hatchery-reared burbot (*Lota lota*) (length ~30-40 cm) was placed to introduce

olfactory cues of a natural predator of juvenile brown trout. Burbot are nocturnal bottom-

dwelling predators that are likely difficult for prey to detect visually, but their odour induces

antipredator responses in prey species (Ylönen et al. 2007). Burbot were regularly fed with

pieces of various cyprinids and vendace (*Coregonus albula*) during rearing, and only with fresh

pieces of brown trout for two days prior to and during the trials. Burbot were moved to the test

arenas at least one day before the trials. The burbot were fed with trout pieces in separate tanks

and changed in each arena every 10–15 trials (2–3 days).

Before each trial, individual brown trout were haphazardly removed from their rearing tanks

using a dip-net under red light and placed into black 10-L buckets filled with ~8L of water from

the flow-through system. Fish were identified by PIT tags and left undisturbed for 10 min before

being transferred into the start box located downstream from the test arena by pouring. During

each trial, the trout was acclimatized in the start box for 3 min, after which the door of the box

was opened by pulling a string from behind a curtain, and fish movements recorded from above

using two CCTV infrared cameras (two arenas simultaneously filmed using the same camera) for

10 min (of which first 9 min 45 s was included in the behaviour analysis). The behavioural trial

was repeated three times between 8:00 and 11:00 for each focal fish, with an average time of 4.3

days (range 1–8 days) between consecutive trials. One trial from four fish was omitted from

analysis due to error in data collection. The order in which batches of four fish were captured on

the same day from the same tank for the four arenas was recorded (batch from hereon, levels 1–

5, four individuals from batch 6/7 combined to batch 5).

Testing behavioural responses to burbot

To confirm that burbot odour was perceived risky in the personality assays, we tested for the

response of brown trout to burbot in separate controlled tests using individuals from wild HV

and wild LV groups (N=10 in each). These fish were acclimated to similar tanks as the

personality-tested fish at 12h:12h L:D photoperiod for one week before trials started. One

individual died after the first trial. The behaviour of each individual was tested on six different

days in the presence and absence of predator (3 trials in each condition in haphazard order). 3–

4 different arenas were used for each fish on different days to reduce fish habituation to the

arenas. These trials were conducted between 14:40 and 17:00. Control arenas were emptied and

thoroughly rinsed with pressurised tap water and water flow maintained for >2h before the

trials to avoid carry-over effects from burbot odour in earlier experiments. The water used in

the flow-through system originates from lake Kivesjärvi, where burbot is a common species;

thus, traces of burbot odour may have been present in all trials.

Analysis of video recordings

Behavioural data were collected from videos using manual tracking with AV Bio-Statistics 5.2

timing software. The observer was blind to the identity of fish in all recordings. Analyses were

conducted in haphazard order, and each trial was analysed once. In total four people analysed

the videos. Four behaviours were recorded from the arena trials: 1) *latency* as the time from the

start of the experiment until the whole body of fish emerged from the start box, (after Boulton et

al. 2014; Moran et al. 2016; Vainikka et al. 2016); 2) time until fish passed the gate to the

upstream section of the arena (arrow in Fig. 1), but this was not analysed because of many fish

not entering this section; instead we recorded 3) *exploration tendency* as a binary variable

indicating whether the whole body of the fish passed the gate within the arena; and 4) *activity* of

fish as the proportion of time spent actively swimming after emerging from the start box. We

used the proportion of time rather than absolute time active to reduce the dependence of

activity from latency. Activity was thus calculated by dividing the total time when fish did not

move when outside the start box by the total time spent outside the start box and subtracting

the value from 1. Stillness was characterized as the fish not moving forward, backward or

sideways for longer than ~2 s. Notably, activity by our definition refers to short-term activity in

a risky, novel environment, not in a familiar environment as it is classically defined (Conrad et al.

2011), and it was recorded only from the trials in which the fish emerged from the start box.

Cortisol response to confinement stress

We measured the plasma cortisol levels from of a subset of the fish after exposure to a

standardized confinement stress. During the tests, the fish were transferred to individual dark

brown 10-L plastic buckets with 1.5 L water for 30 min (except for one fish in each Wild HV and

LV and Hatchery HV when the time was 36 min by mistake). The water was aerated using air

stones and pump (Sera Air 550R and Sera AS30 air stone) during the test. The buckets were

placed in a flow-through buffer tank at a temperature matching the acclimation tanks

(temperatures increased during the days of the measurement, 26–29 June 2017, from 13.4 to

16.1°C), and left undisturbed in the dark for the duration of the confinement. Fish were then

removed from buckets by dip-netting, anaesthetized using benzocaine solution, measured (to 1

mm) and weighed (to 0.1 g). Blood samples were collected within 2–5 min from the start of

anaesthesia. The sampling order of fish from the same tank during the same day was recorded.

Blood was collected using 23 G heparinized needles and syringes and kept on ice temporarily

until centrifuged at 4000 x g for 10 min. Plasma was collected in Eppendorf tubes and frozen at –

20°C until analysis. Control samples for establishing baseline plasma cortisol concentrations

were collected after terminal anaesthesia as described above, omitting the confinement stress

treatment. Plasma cortisol concentration was determined using enzyme-linked immunosorbent

assay (ELISA) (Enzo cortisol assay) as described in ESM1.

Sex determination from DNA samples

To consider potential sex differences in the studied traits, we identified the sex of fish using PCR

amplification of the sexually dimorphic *sdY* locus, which identifies the correct sex in brown trout

with nearly 100% accuracy (Quéméré et al. 2014); details in ESM1.

Statistical analyses

The number of individuals included in each analysis is shown in Table 1. We built univariate

models for each response variable (metabolic and behavioural variables and cortisol level) to

assess the differences between breeding group and acclimation conditions (Table 2). All

analyses were conducted in R v.3.3.2 (R Core Team, 2016). Linear (LMM) and generalized

mixed-effects models (GLMM) were fitted using package *lme4* (Bates et al. 2015) with *lmerTest*

(Kuznetsova et al. 2017) and the frailty models using package *coxme* (Therneau, 2018). The data

were visualized using ggplot2 (Wickham 2009) and patchwork

(https://github.com/thomasp85/patchwork). Statistical significance was determined as $\alpha =$

0.05 in all models. Predicted means within groups were estimated for behaviour traits with

package *ggeffects* (Lüdecke 2018). The effect of sex was analysed in separate models, including

the fixed effect of sex as well as the effects from original models, except photoperiod or its

interactions due to limited sample size with known sex. All linear models were checked for

homoscedasticity and normality of residuals.

Log$_{10}$-transformed $\dot{M}O_{2,min}$ or $\dot{M}O_{2,ave}$ were analysed using an LMM with function *lmer*. The main

effects of population, selection line, photoperiod and log$_{10}$-body mass (in kg) were separately

tested using linear hypothesis testing (function *lht* in package *car*) using restricted models,

where each respective main effect and its interactions were defined zero and compared to the

full model using F-tests.

The difference in cortisol level of control fish and fish exposed to confinement stress was first

tested using a one-tailed t-test. The post-confinement stress cortisol level was then analysed

using a linear model using function *lm*.

Behavioural traits were analysed using an LMM (*activity*), a frailty model (i.e. mixed effect Cox

proportional hazards models for time-to event data (Collett 2015)) (*latency*) and a GLMM

(Bernoulli-distributed *exploration tendency*). Trial repeats were encoded as -1, 0, and 1 in data

from angling selection experiment as 1–6 from burbot vs. control experiment. In 8 trials, the fish

jumped out of the start box prior to the trial and their behaviour was analysed for 9 min 45 s

min after the jump. Correlations between metabolic traits and activity were calculated from

model residuals and best linear unbiased predictions (BLUPs), respectively, to assess potential

underlying associations between the traits across all individuals. Correlations were not

calculated for time-to-event data (*latency*) or binary data (*exploration tendency*). For further

details see ESM1 and Data accessibility.

Results

$\dot{M}O_{2,min}$, $\dot{M}O_{2,ave}$ and stress-sensitivity

The LMM indicated significantly higher $\dot{M}O_{2,min}$ in the offspring of wild fish than of hatchery fish,

and a moderate interaction effect between photoperiod and population, wild population having

higher values than hatchery population in the 12:12 L:D photoperiod. Interaction was also found

for population and angling selection line, hatchery LV fish tending to have higher oxygen uptake

than HV fish, while selection lines did not differ in the wild population (Fig. 2A; Table 3). $\dot{M}O_{2,ave}$

was higher in wild than in hatchery population, with a modest interaction effect of angling

selection in the two populations (non-significant, $P = 0.085$), observed as higher $\dot{M}O_{2,ave}$ in

hatchery LV compared to hatchery HV, but no effect of angling selection in the wild population

(Fig. 2B). Sex did not have a significant effect on either $\dot{M}O_{2,min}$ (Type III test, $F_{1,58} = 0.901$, $P =$

0.346) or $\dot{M}O_{2,ave}$ ($F_{1,58.577} = 0.1823$, $P = 0.671$).

Plasma cortisol increased ~seven-fold in individuals subjected to confinement stress (mean =

140.62 ng mL^{-1}, SD = 41.00) compared to non-stressed fish (mean = 19.22 ng mL^{-1}, SD = 20.65),

(t-test, t = 11.125, df = 29.523, P < 0.001). Angling selection or population did not significantly

affect the level of post-stress plasma cortisol, although it showed a similar tendency as observed

in $\dot{M}O_{2,ave}$ (Table 3, Fig. S3).

Behaviour in angling selection lines

Fish emerged from the start box during the recorded time in ~84% of the trials. There was a

slightly non-significant interaction effect (P = 0.054) of population background and angling

selection on latency (Table 4). This was observed as an elevated probability to emerge in fish

from LV background compared to HV background in the hatchery population, but not in the wild

population (Fig. 3A).

Fish were less active after acclimation in constant light compared to the 12:12 L:D photoperiod,

but activity did not differ between populations or angling selection lines (Fig. 3B; Table 4).

Angling selection had contrasting effects on exploration tendency in each population: in the

hatchery population, a higher proportion of fish from LV selection line were explorative than

from HV selection line, while there was an opposite tendency in the wild population (Fig. 3C;

Table 4). In addition, exploration tendency increased with repeats of the behavioural trial. Sex

did not have a significant effect on any behaviour trait (female vs male, Activity: $F_{1,39.612}$ = 1.217,

P = 0.277; Latency: e^{coef} = 1.03, z = 0.29, P = 0.770; Exploration tendency: z = −0.514, P = 0.607).

There was no correlation between the BLUPs of activity and residual $\dot{M}O_{2,min}$ (*Pearson r* = 0.02)

or $\dot{M}O_{2,ave}$ (*Pearson r* = −0.04).

Behavioural responses to predator presence

The fish tended to be less active (P = 0.072) in the presence of burbot than under control

conditions (Table 5). The variance of activity between individuals appeared higher in the

presence of burbot, but this was not significant in Levene's test of homogeneity of variance ($F_{1,93}$

= 0.214, P = 0.645). Activity decreased slightly with increasing behaviour trial repeats. Latency

was not affected by predator cues (non-significant increase in probability to emerge by 9%), but

it increased with increasing behaviour trial repeats and between-individual variation in latency

was high (~10% higher variance in burbot vs control data compared to data from angling

selection lines). The exploration tendency of fish was not affected by predator cues.

Discussion

Population-specific effects of angling selection on boldness

Captured and non-captured parent brown trout produced offspring that differed in boldness-

related traits during their second summer. Surprisingly, boldness, measured as latency to

explore a novel arena, was lower in the HV fish than in the LV fish in the hatchery population.

This was a contradictory finding as angling is expected to select against boldness (Arlinghaus et

al. 2017). The wild population showed a weaker difference between selection lines, but its

direction was more in line with the theory, with HV fish being bolder than LV fish. Activity

during the behavioural trials was not affected by angling selection. In contrast, activity was

affected by photoperiod, which suggests a connection between energy demand/expenditure and

photoperiod through activity.

The unexpected result of angling-vulnerability related low boldness in hatchery population may

be explained by the hatchery rearing environment through several indirect effects. First, the

parent fish that were vulnerable to angling may have had the least strict preference for the

standard feeds, as they accepted the colourful, unnatural fly patterns. Second, the vulnerable fish

may have had the lowest status in the dominance hierarchy within the ponds, and therefore

been the hungriest and likeliest to attack lures. Third, the hatchery-history itself, i.e. inadvertent

domestication, might have selected for the most proactive individuals. This, in turn, could have

shifted the vulnerability in this population towards more reactive fish that may have more

flexibility in their behaviour.

In contrast to the hatchery population, angling trials on wild parent fish were more

representative of real angling situations in the field. The wild fish had natural invertebrate food

available in their ponds, and the structured ponds offered more hiding places. The wild fish had

clearly lower catchability than the hatchery fish, and the wild fish could only be captured when

approaching the undisturbed pond from a distance. Very few wild fish were captured in one

angling session (maximum 4) compared to the hatchery fish (maximum 11). The captured and

non-captured parent fish did not show evident size-differences, indicating that the effects of

angling were most likely mediated by size-independent traits.

Stress coping styles and angling selection

A possible consequence of angling selection is a change in the frequencies of not only bold and

timid individuals, but also stress-tolerant and stress-sensitive individuals, due to the association

of angling vulnerability, behaviour and stress coping styles (Vindas et al. 2017a). Furthermore,

the POLS theory suggests that stress coping styles, along with metabolic rate and other traits,

are connected to life-history traits, which vary along a slow-fast continuum (e.g., (Dammhahn et

al. 2018; Reale et al. 2010). Stress coping styles and POLS concepts therefore provide

frameworks for testing the consequences of angling induced selection.

In largemouth bass, cortisol response to a standard stressor was negatively associated with

capture probability, as expected (Louison et al. 2017). Koeck et al. (2018) found a weak negative

effect (~0.5% change in risk) of high cortisol response on vulnerability to angling in a domestic

strain of rainbow trout (*Oncorhynchus mykiss*), but there was no similar relationship in a wild

strain of brown trout. In our study, stress sensitivity, expressed as a cortisol response to

confinement, was not affected by angling-selection, although these tests suffered from low

statistical power. However, a higher response in hatchery LV fish compared to HV was more

visible through $\dot{M}O_{2,ave}$, which is notable given that confinement in the respirometer can induce a

stress response in fish (Murray et al. 2017) and increase their oxygen uptake. Thus, the trends

between HV and LV in the hatchery population observed in the cortisol response and $\dot{M}O_{2,ave}$

suggest potential for angling selection for increased stress sensitivity, which might become more

visible after multiple generations of selection. These results suggest that the individuals of LV

selection line within the hatchery population showed a more reactive stress coping style than

HV line. Differences in coping styles could also partly explain why the behaviour interpreted as

boldness showed a pattern contradicting our expectations; if the LV fish were more reactive

compared to the HV fish, their behaviour in the personality trial may have indicated a higher

stress response to the experiment and heightened escape behaviour (Laskowski et al. 2016).

The stress test and $\dot{M}O_{2,ave}$ results are in agreement with the hypothesis that the wild population

displayed a more reactive coping style compared to the hatchery population.

Whether $\dot{M}O_{2,min}$ is connected to the coping styles/POLS's or angling selection remains

inconclusive based on our results. A lack of association between metabolic rate and personality

has been reported previously in other species, such as the Trinidadian guppy (*Poecilia*

reticulata) (White et al. 2016). We found a trend of higher $\dot{M}O_{2,min}$ in the hatchery LV line

compared to HV line, and an opposite trend in wild fish, under 12:12 L:D photoperiod. The

contrast between the populations in the response to angling selection is therefore in line with

the behavioural results in latency and exploration tendency, but also with activity in the

response to photoperiod, although there was no correlation between activity and $\dot{M}O_{2,min}$ after

accounting for group differences. Again, as in boldness, the result in hatchery population was

opposite to the prevailing theory, and it is possible that stronger effects from angling could be

observed over several generations of selection in natural conditions. Growth rate is unlikely to

explain the differences in $\dot{M}O_{2,min}$ between groups, as the body mass of fish at the end of the

experiment did not differ between groups (Table 1). Despite not directly addressing questions

on trait covariances, as physiological traits were measured only once (Mathot and Frankenhuis

2018; Niemelä and Dingemanse 2018), our results add to the literature to promote the

understanding of evolution in traits due to angling induced selection. From an angling selection

perspective, some of the most interesting traits to include in further experiments would be

neurochemical stress responses and their links to the bold-shy behavioural axis, and a role for

metabolic rate remains to be understood better in connection to these traits.

Genetic and parental effects in population divergence and angling selection

Populations frequently differ in e.g., metabolic rate and behavioural syndromes (Dingemanse et

al. 2007; Lahti et al. 2002; Polverino et al. 2018), driven by environmental differences, natural

selection, founder effects, and genetic drift. The differences we found between populations can

therefore be explained by several factors, including the level of domestication, as the hatchery

stock had been reared in captivity for several generations. They also differed in their life-

histories, with the wild population being clearly resident and the hatchery population migratory

(Lemopoulos et al. unpublished data). In addition, although we reared offspring under common

garden conditions and maximized genetic diversity within each group, it is possible that

differences in the early rearing environments of wild and hatchery parents could have had

contrasting effects on offspring through parental or epigenetic effects (Crews et al. 2012;

Reddon 2012). We studied individuals in their second summer, and parental effects usually

affect early life-stages the most; for instance, maternal effects on metabolic traits have been

shown to be negligible from 90 days post hatching in coral reef fish (Munday et al. 2017),

although maternal stress affects many life-stages in three-spined stickleback (*Gasterosteus*

aculeatus) (Bell et al. 2016; Metzger and Schulte 2016). Additionally, parental effects may have

also contributed to differences between the selection lines *via* stress resulting from angling

(briefly increased cortisol level after angling shown, e.g., in Wilson et al. (2011)). It is

nevertheless likely that for both population and angling selection line differences, genetic

inheritance may explain our results at least partly, as both angling vulnerability and personality

traits can be heritable in the studied populations (Ågren et al. 2019) or in other species

(Dingemanse et al. 2009; Philipp et al. 2009).

Potential effects of photoperiod on energy balance

We incorporated environmental variation in our study as two different photoperiods. The

results demonstrate, on one hand, that metabolic rate and swimming activity are sensitive to

photoperiod, and on the other hand, that the other behavioural traits lack this sensitivity.

Constant light is not encountered by brown trout during the winter months, hence the 24-hour

light regime could be considered unnatural and potentially stressful for the fish. Constant light

can disrupt entrainment of endogenous rhythms by inhibiting the synthesis of melatonin and by

directly affecting photosensitive proteins (Falcón et al. 2010; Peirson et al. 2009). Based on our

results, constant light had an inhibiting effect on fish swimming activity, and also decreased

$\dot{M}O_{2,min}$ in the wild population, indicating that energy metabolism in brown trout can be affected

by (an unnatural) photoperiod. In general, non-tropical species are expected to be particularly

sensitive to photoperiod disturbances due to the role of day length in anticipating seasonal

changes in environmental conditions (Borniger et al. 2017).

Innate vs learned antipredatory responses

Our goal was to study risk-taking behaviour/boldness of offspring by subjecting fish to the

olfactory cues of a natural predator that had fed on conspecifics. Wild brown trout typically

increase the use of refuges under predation threat, while hatchery brown trout do not (Álvarez

and Nicieza 2003). None of the individuals in the behaviour trials in this study had been exposed

to predators before the trials apart from potential traces of piscivore odours in the rearing

water. The scarcity of responses to the presence of predator odour, measured in the offspring of

wild fish, suggests only weak innate responses. Nevertheless, the tendency for lower activity in

the presence of burbot than in control conditions resembles previously shown antipredator

responses in fish (Álvarez and Nicieza 2003; Kopack et al. 2015).

Conclusions

Our results demonstrate the potential for rapid fishing-induced evolution in a very popular

target species. The effects of angling selection are somewhat contradictory between wild and

hatchery populations of fish, which leads to new questions on the mechanisms behind the

observed differences. For practical fisheries management, the results with the wild fish are more

representative and generalizable and showed little evidence of selection in physiological or

behavioural traits. Overall, our study supports earlier findings according to which angling may

be a potentially significant driver of evolution in behavioural and physiological traits in

harvested populations.

Conflict of Interest

The authors declare that they have no conflict of interest.

Acknowledgements

We thank the staff of Kainuu Fisheries Research Station for their help in catching, breeding and

rearing fish, Dr. Hannu Huuskonen for advice in setting up the respirometry, and Dr. Chris

Elvidge for comments on the manuscript. J.M.P., A.V. and A.L. were supported by the Academy of

Finland grant for A.V. (nr. 286261). J.M.P. was also supported by Oskar Öflund's foundation and

by the Finnish Cultural Foundation. All applicable institutional and/or national guidelines for the

care and use of animals were followed.

Data accessibility

All data and R codes for the models in this manuscript are available in Github

(https://github.com/jprokkola/Strutta_repo). Videos of personality trials will be made publicly

available in Figshare (accession) upon acceptance for publication.

Electronic Supplementary Material

ESM1. Pdf-file including supplemental figures and methods.

References

Adriaenssens B, Johnsson JI (2011) Shy trout grow faster: exploring links between personality

550   and fitness-related traits in the wild. Behavioral Ecology 22:135-143. doi:

551   10.1093/beheco/arq185

552   Ågren A, Vainikka A, Janhunen M, Hyvärinen P, Piironen J, Kortet R (2019) Experimental

553   crossbreeding reveals strain-specific variation in mortality, growth and personality in

554   the brown trout (*Salmo trutt*a). Scientific Reports 9. doi: 10.1038/s41598-018-35794-6

555   Allendorf FW, Hard JJ (2009) Human-induced evolution caused by unnatural selection through

556   harvest of wild animals. Proceedings of the National Academy of Sciences of the United

557   States of America 106:9987-9994. doi: 10.1073/pnas.0901069106

558   Álvarez D, Nicieza AG (2003) Predator avoidance behaviour in wild and hatchery-reared brown

559   trout: the role of experience and domestication. Journal of Fish Biology 63:1565-1577.

560   doi: 10.1111/j.1095-8649.2003.00267.x

561   Andersen KH, Marty L, Arlinghaus R (2018) Evolution of boldness and life history in response to

562   selective harvesting. Canadian Journal of Fisheries and Aquatic Sciences 75:271-281. doi:

563   10.1139/cjfas-2016-0350

564   Araki H, Berejikian BA, Ford MJ, Blouin MS (2008) Fitness of hatchery-reared salmonids in the

565   wild. Evolutionary Applications 1:342-355. doi: 10.1111/j.1752-4571.2008.00026.x

Arlinghaus R, Laskowski KL, Alós J, Klefoth T, Monk CT, Nakayama S, Schröder A (2017) Passive

gear-induced timidity syndrome in wild fish populations and its potential ecological and

managerial implications. Fish and Fisheries 18:360-373. doi: 10.1111/faf.12176

Audzijonyte A, Kuparinen A, Fulton EA (2013) How fast is fisheries-induced evolution?

Quantitative analysis of modelling and empirical studies. Evolutionary Applications

6:585-595. doi: 10.1111/eva.12044

Bates D, Maechler M, Bolker B, Walker S (2015). Fitting Linear Mixed-Effects Models Using lme4.

Journal of Statistical Software, 67:1-48. doi:10.18637/jss.v067.i01

Bell AM, Laura KEM, Stein LR (2016) Effects of mothers' and fathers' experience with predation

risk on the behavioral development of their offspring in threespined sticklebacks.

Current Opinion in Behavioral Sciences 7:28-32. doi: 10.1016/j.cobeha.2015.10.011

Biro PA, Post JR (2008) Rapid depletion of genotypes with fast growth and bold personality

traits from harvested fish populations. Proceedings of the National Academy of Sciences

of the United States of America 105:2919-2922. doi: 10.1073/pnas.0708159105

Borniger JC, Cisse YM, Nelson RJ, Martin LB (2017) Seasonal Variation in Stress Responses.

Stress: Neuroendocrinology and Neurobiology 2:411-419. doi: 10.1016/b978-0-12-

802175-0.00041-3

Boulton K, Grimmer AJ, Rosenthal GG, Walling CA, Wilson AJ (2014) How stable are

personalities? A multivariate view of behavioural variation over long and short

timescales in the sheepshead swordtail, *Xiphophorus birchmanni*. Behavioral Ecology and

Sociobiology 68:791-803. doi: 10.1007/s00265-014-1692-0

Brelin D, Petersson E, Dannewitz J, Dahl J, Winberg S (2008) Frequency distribution of coping

strategies in four populations of brown trout (*Salmo trutta*). Hormones and Behavior

53:546-556. doi: 10.1016/j.yhbeh.2007.12.011

Collett D (2015) Modelling Survival Data in Medical Research, Third Edition. Chapman &

Hall/CRC Texts in Statistical Science.

Coltman DW, O'Donoghue P, Jorgenson JT, Hogg JT, Strobeck C, Festa-Bianchet M (2003)

Undesirable evolutionary consequences of trophy hunting. Nature 426:655-658. doi:

10.1038/nature02177

Conrad JL, Weinersmith KL, Brodin T, Saltz JB, Sih A (2011) Behavioural syndromes in fishes: a

review with implications for ecology and fisheries management. Journal of Fish Biology

78:395-435. doi: 10.1111/j.1095-8649.2010.02874.x

Cooke SJ, Suski CD, Ostrand KG, Wahl DH, Philipp DP (2007) Physiological and behavioral

consequences of long-term artificial selection for vulnerability to recreational angling in

a teleost fish. Physiological and Biochemical Zoology 80:480-490. doi: 10.1086/520618

Crews D, Gillette R, Scarpino SV, Manikkam M, Savenkova MI, Skinner MK (2012) Epigenetic

transgenerational inheritance of altered stress responses. Proceedings of the National

Academy of Sciences of the United States of America 109:9143-9148. doi:

10.1073/pnas.1118514109

Dammhahn M, Dingemanse NJ, Niemelä PT, Réale D (2018) Pace-of-life syndromes: a framework

for the adaptive integration of behaviour, physiology and life history. Behavioral Ecology

and Sociobiology 72:62. doi: 10.1007/s00265-018-2473-y

Devine JA, Wright PJ, Pardoe HE, Heino M (2012) Comparing rates of contemporary evolution in

life-history traits for exploited fish stocks. Canadian Journal of Fisheries and Aquatic

Sciences 69:1105-1120. doi: 10.1139/f2012-047

Dingemanse NJ, Van der Plas F, Wright J, Réale D, Schrama M, Roff DA, Van der Zee E, Barber I

(2009) Individual experience and evolutionary history of predation affect expression of

heritable variation in fish personality and morphology. Proceedings of the Royal Society

B-Biological Sciences 276:1285-1293. doi: 10.1098/rspb.2008.1555

Dingemanse NJ, Wright J, Kazem AJN, Thomas DK, Hickling R, Dawnay N (2007) Behavioural

syndromes differ predictably between 12 populations of three-spined stickleback.

Journal of Animal Ecology 76:1128-1138. doi: 10.1111/j.1365-2656.2007.01284.x

Falcón J, Migaud H, Muñoz-Cueto JA, Carrillo M (2010) Current knowledge on the melatonin

system in teleost fish. General and Comparative Endocrinology 165:469-482. doi:

10.1016/j.ygcen.2009.04.026

Fugère V, Hendry AP (2018) Human influences on the strength of phenotypic selection.

Proceedings of the National Academy of Sciences of the United States of America

115:10070-10075. doi: 10.1073/pnas.1806013115

Härkönen L, Hyvärinen P, Paappanen J, Vainikka A (2014) Explorative behavior increases

vulnerability to angling in hatchery-reared brown trout (*Salmo trutta*). Canadian Journal

of Fisheries and Aquatic Sciences 71:1900-1909. doi: 10.1139/cjfas-2014-0221

Hollins J, Thambithurai D, Koeck B, Crespel A, Bailey DM, Cooke SJ, Lindström J, Parsons KJ,

Killen SS (2018) A physiological perspective on fisheries-induced evolution.

Evolutionary Applications 11:561-576. doi: 10.1111/eva.12597

Horreo JL, Valiente AG, Ardura A, Blanco A, Garcia-Gonzalez C, Garcia-Vazquez E (2018) Nature

versus nurture? Consequences of short captivity in early stages. Ecology and Evolution

8:521-529. doi: 10.1002/ece3.3555

Huntingford F, Adams C (2005) Behavioural syndromes in farmed fish: implications for

production and welfare, pp 1207-1221

Huntingford FA (2004) Implications of domestication and rearing conditions for the behaviour

of cultivated fishes. Journal of Fish Biology 65:122-142. doi: 10.1111/j.1095-

8649.2004.00562.x

Johnsson JI, Näslund J (2018) Studying behavioural variation in salmonids from an ecological

perspective: observations questions methodological considerations. Reviews in Fish

Biology and Fisheries 28:795-823. doi: 10.1007/s11160-018-9532-3

Kern EMA, Robinson D, Gass E, Godwin J, Langerhans RB (2016) Correlated evolution of

personality, morphology and performance. Animal Behaviour 117:79-86. doi:

10.1016/j.anbehav.2016.04.007

Killen SS, Marras S, McKenzie DJ (2011) Fuel, fasting, fear: routine metabolic rate and food

deprivation exert synergistic effects on risk-taking in individual juvenile European sea

bass. Journal of Animal Ecology 80:1024-1033. doi: 10.1111/j.1365-2656.2011.01844.x

Klefoth T, Pieterek T, Arlinghaus R (2013) Impacts of domestication on angling vulnerability of

common carp, *Cyprinus carpio*: the role of learning, foraging behaviour and food

preferences. Fisheries Management and Ecology 20:174-186. doi: 10.1111/j.1365-

2400.2012.00865.x

Koolhaas JM, de Boer SF, Coppens CM, Buwalda B (2010) Neuroendocrinology of coping styles:

Towards understanding the biology of individual variation. Frontiers in

Neuroendocrinology 31:307-321. doi: 10.1016/j.yfrne.2010.04.001

Kopack CJ, Broder ED, Lepak JM, Fetherman ER, Angeloni LM (2015) Behavioral responses of a

highly domesticated, predator naive rainbow trout to chemical cues of predation.

Fisheries Research 169:1-7. doi: 10.1016/j.fishres.2015.04.005

Kuznetsova A, Brockhoff PB, Christensen RHB (2017). lmerTest Package: Tests in Linear Mixed

Effects Models. Journal of Statistical Software, 82:1–26. doi: 10.18637/jss.v082.i13.

Lahti K, Huuskonen H, Laurila A, Piironen J (2002) Metabolic rate and aggressiveness between

Brown Trout populations. Functional Ecology 16:167-174. doi: 10.1046/j.1365-

2435.2002.00618.x

Laskowski KL, Monk CT, Polverino G, Alos J, Nakayama S, Staaks G, Mehner T, Arlinghaus R

(2016) Behaviour in a standardized assay, but not metabolic or growth rate, predicts

behavioural variation in an adult aquatic top predator *Esox lucius* in the wild. Journal of

Fish Biology 88:1544-1563. doi: 10.1111/jfb.12933

Lemopoulos A, Prokkola JM, Uusi-Heikkilä S, Vasemägi A, Huusko A, Hyvärinen P, Koljonen ML,

Koskiniemi J, Vainikka A (2019) Comparing RADseq and microsatellites for estimating

genetic diversity and relatedness Implications for brown trout conservation. Ecology and

Evolution 9:2106-2120. doi: 10.1002/ece3.4905

Lennox RJ, Alos J, Arlinghaus R, Horodysky A, Klefoth T, Monk CT, Cooke SJ (2017) What makes

fish vulnerable to capture by hooks? A conceptual framework and a review of key

determinants. Fish and Fisheries 18:986-1010. doi: 10.1111/faf.12219

Louison MJ, Adhikari S, Stein JA, Suski CD (2017) Hormonal responsiveness to stress is

negatively associated with vulnerability to angling capture in fish. Journal of

Experimental Biology 220:2529-2535. doi: 10.1242/jeb.150730

Louison MJ, Stein JA, Suski CD (2018) Metabolic phenotype is not associated with vulnerability

angling in bluegill sunfish (*Lepomis macrochirus*). Canadian Journal of Zoology 96:1264-

1271. doi: 10.1139/cjz-2017-0363

Lüdecke D (2018). ggeffects: Tidy Data Frames of Marginal Effects from Regression Models.

Journal of Open Source Software 3: 772. doi: 10.21105/joss.00772

Mathot KJ, Dingemanse NJ, Nakagawa S (2019) The covariance between metabolic rate and

behaviour varies across behaviours and thermal types: meta-analytic insights. Biological

Reviews. In press. doi:10.1111/brv.12491

Mathot KJ, Frankenhuis WE (2018) Models of pace-of-life syndromes (POLS): a systematic

review. Behavioral Ecology and Sociobiology 72. doi: 10.1007/s00265-018-2459-9

Metzger DCH, Schulte PM (2016) Maternal stress has divergent effects on gene expression

patterns in the brains of male and female threespine stickleback. Proceedings of the

Royal Society B-Biological Sciences 283. doi: 10.1098/rspb.2016.1734

Moran NP, Mossop KD, Thompson RM, Wong BBM (2016) Boldness in extreme environments:

temperament divergence in a desert-dwelling fish. Animal Behaviour 122:125-133. doi:

10.1016/j.anbehav.2016.09.024

Morgan JD, Iwama GK (1996) Cortisol-induced changes in oxygen consumption and ionic

regulation in coastal cutthroat trout (*Oncorhynchus clarki clarki*) parr. Fish Physiology

and Biochemistry 15:385-394. doi: 10.1007/bf01875581

Munday PL, Donelson JM, Domingos JA (2017) Potential for adaptation to climate change in a

coral reef fish. Global Change Biology 23:307-317. doi: 10.1111/gcb.13419

Murray L, Rennie MD, Svendsen JC, Enders EC (2017) Respirometry increases cortisol levels in

rainbow trout *Oncorhynchus mykiss*: implications for measurements of metabolic rate.

Journal of Fish Biology 90:2206-2213. doi: 10.1111/jfb.13292

Näslund J, Johnsson JI (2016) State-dependent behavior and alternative behavioral strategies in

brown trout (*Salmo trutta* L.) fry. Behavioral Ecology and Sociobiology 70:2111-2125.

doi: 10.1007/s00265-016-2215-y

Nelson JA (2016) Oxygen consumption rate v. rate of energy utilization of fishes: a comparison

and brief history of the two measurements. Journal of Fish Biology 88:10-25. doi:

10.1111/jfb.12824

Niemelä PT, Dingemanse NJ (2018) Meta-analysis reveals weak associations between intrinsic

state and personality. Proceedings of the Royal Society B-Biological Sciences 285. doi:

10.1098/rspb.2017.2823

Nussle S, Bornand CN, Wedekind C (2009) Fishery-induced selection on an Alpine whitefish:

quantifying genetic and environmental effects on individual growth rate. Evolutionary

Applications 2:200-208. doi: 10.1111/j.1752-4571.2008.00054.x

Peirson SN, Halford S, Foster RG (2009) The evolution of irradiance detection: melanopsin and

the non-visual opsins. Philosophical Transactions of the Royal Society B-Biological

Sciences 364:2849-2865. doi: 10.1098/rstb.2009.0050

Philipp DP, Cooke SJ, Claussen JE, Koppelman JB, Suski CD, Burkett DP (2009) Selection for

vulnerability to angling in largemouth bass. Transactions of the American Fisheries

Society 138:189-199. doi: 10.1577/t06-243.1

Polverino G, Santostefano F, Diaz-Gil C, Mehner T (2018) Ecological conditions drive pace-of-life

syndromes by shaping relationships between life history, physiology and behaviour in

two populations of Eastern mosquitofish. Scientific Reports 8. doi: 10.1038/s41598-018-

33047-0

Quéméré E, Perrier C, Besnard AL, Evanno G, Baglinière JL, Guiguen Y, Launey S (2014) An

improved PCR-based method for faster sex determination in brown trout (*Salmo trutta*)

and Atlantic salmon (*Salmo salar*). Conservation Genetics Resources 6:825-827. doi:

10.1007/s12686-014-0259-8

Réale D, Garant D, Humphries MM, Bergeron P, Careau V, Montiglio PO (2010) Personality and

the emergence of the pace-of-life syndrome concept at the population level.

Philosophical Transactions of the Royal Society B-Biological Sciences 365:4051-4063.

doi: 10.1098/rstb.2010.0208

Reddon AR (2012) Parental effects on animal personality. Behavioral Ecology 23:242-245. doi:

10.1093/beheco/arr210

Redpath TD, Cooke SJ, Suski CD, Arlinghaus R, Couture P, Wahl DH, Philipp DP (2010) The

metabolic and biochemical basis of vulnerability to recreational angling after three

generations of angling-induced selection in a teleost fish. Canadian Journal of Fisheries

and Aquatic Sciences 67:1983-1992. doi: 10.1139/f10-120

Schjolden J, Backström T, Pulman KGT, Pottinger TG, Winberg S (2005) Divergence in

behavioural responses to stress in two strains of rainbow trout (*Oncorhynchus mykiss*)

with contrasting stress responsiveness. Hormones and Behavior 48:537-544. doi:

10.1016/j.yhbeh.2005.04.008

Sutter DAH, Suski CD, Philipp DP, Klefoth T, Wahl DH, Kersten P, Cooke SJ, Arlinghaus R (2012)

Recreational fishing selectively captures individuals with the highest fitness potential.

Proceedings of the National Academy of Sciences of the United States of America

109:20960-20965. doi: 10.1073/pnas.1212536109

Svendsen MBS, Bushnell PG, Steffensen JF (2016) Design and setup of intermittent-flow

respirometry system for aquatic organisms. Journal of Fish Biology 88:26-50. doi:

10.1111/jfb.12797

Therneau TM (2018). coxme: Mixed Effects Cox Models. R package version 2.2-7.

https://CRAN.R-project.org/package=coxme

Thomson JS, Watts PC, Pottinger TG, Sneddon LU (2011) Physiological and genetic correlates of

boldness: Characterising the mechanisms of behavioural variation in rainbow trout,

Oncorhynchus mykiss. Hormones and Behavior 59:67-74. doi:

10.1016/j.yhbeh.2010.10.010

Uusi-Heikkilä S, Whiteley AR, Kuparinen A, Matsumura S, Venturelli PA, Wolter C, Slate J,

Primmer CR, Meinelt T, Killen SS, Bierbach D, Polverino G, Ludwig A, Arlinghaus R

(2015) The evolutionary legacy of size-selective harvesting extends from genes to

populations. Evolutionary Applications 8:597-620. doi: 10.1111/eva.12268

Uusi-Heikkilä S, Wolter C, Klefoth T, Arlinghaus R (2008) A behavioral perspective on fishing-

induced evolution. Trends in Ecology & Evolution 23:419-421. doi:

10.1016/j.tree.2008.04.006

Väätäinen R, Huuskonen H, Hyvärinen P, Kekäläinen J, Kortet R, Arnedo MT, Vainikka A (2018)

Do metabolic traits, vulnerability to angling, or capture method explain boldness

variation in Eurasian perch? Physiological and Biochemical Zoology 91:1115-1128. doi:

10.1086/700434

Vainikka A, Tammela I, Hyvärinen P (2016) Does boldness explain vulnerability to angling in

Eurasian perch *Perca fluviatilis*? Current Zoology 62:109-115. doi: 10.1093/cz/zow003

Vindas MA, Gorissen M, Höglund E, Flik G, Tronci V, Damsgard B, Thörnqvist PO, Nilsen TO,

Winberg S, Øverli Ø, Ebbesson LOE (2017a) How do individuals cope with stress?

Behavioural, physiological and neuronal differences between proactive and reactive

coping styles in fish. Journal of Experimental Biology 220:1524-1532. doi:

10.1242/jeb.153213

Vindas MA, Magnhagen C, Brännäs E, Øverli Ø, Winberg S, Nilsson J, Backström T (2017b) Brain

cortisol receptor expression differs in Arctic charr displaying opposite coping styles.

Physiology & Behavior 177:161-168. doi: 10.1016/j.physbeh.2017.04.024

White SJ, Kells TJ, Wilson AJ (2016) Metabolism, personality and pace of life in the Trinidadian

guppy, *Poecilia reticulata*. Behaviour 153:1517-1543. doi: 10.1163/1568539x-00003375

Wickham H (2009) ggplot2: Elegant Graphics for Data Analysis. Ggplot2: Elegant Graphics for

Data Analysis:1-212. doi: 10.1007/978-0-387-98141-3

Wilson ADM, Binder TR, McGrath KP, Cooke SJ, Godin JGJ (2011) Capture technique and fish

personality: angling targets timid bluegill sunfish, *Lepomis macrochirus*. Canadian

Journal of Fisheries and Aquatic Sciences 68:749-757. doi: 10.1139/f2011-019

Wilson ADM, Brownscombe JW, Sullivan B, Jain-Schlaepfer S, Cooke SJ (2015) Does Angling

Technique Selectively Target Fishes Based on Their Behavioural Type? PLoS One 10. doi:

10.1371/journal.pone.0135848

Wong RY, Perrin F, Oxendine SE, Kezios ZD, Sawyer S, Zhou LR, Dereje S, Godwin J (2012)

Comparing behavioral responses across multiple assays of stress and anxiety in

zebrafish (*Danio rerio*). Behaviour 149:1205-1240. doi: 10.1163/1568539x-00003018

Ylönen H, Kortet R, Myntti J, Vainikka A (2007) Predator odor recognition and antipredatory

response in fish: does the prey know the predator diel rhythm? Acta Oecologica-

International Journal of Ecology 31:1-7. doi: 10.1016/j.actao.2005.05.007

Tables

Table 1. The number of individuals in each group in each analysis, and fish total body length and mass (mean ± SD) at the end of the experiment.

Photoperiod	Group	N (metabolic rate)	N (latency)	N (activity)	N (stress response)	N (body size)	N males / females (unknown)	Body length / mm	Body mass / g
12:12	Hatchery HV	15	15	15	7	14	9 / 5 (1)	117 ± 10	17.1 ± 4.3
	Hatchery LV	7	14	14	6	14	6 / 2 (7)	115 ± 13	16.5 ± 6.7
	Wild HV	11	14	14	4	14	6 / 6 (3)	117 ± 7	17.4 ± 3.2
	Wild LV	14	15	15	9	14	7 / 7 (1)	115 ± 9	16.6 ± 4.3
24	Hatchery HV	10	10	10	0	9	6 / 3 (1)	119 ± 9	17.7 ± 4.7
	Hatchery LV	7	10	10	0	6	0 / 4 (6)	124 ± 9	19.6 ± 3.9
	Wild HV	8	10	10	0	6	2 / 3 (5)	116 ± 9	17.8 ± 7.1
	Wild LV	6	10	10	0	5	2 / 1 (7)	119 ± 9	17.8 ± 3.0

794   Table 2. The main statistical models used in this study. Abbreviations explained below the table.

Study section	Response variable	Model
I. Angling selection experiment	$Log_{10}(\dot{M}O_{2,min})$ or $Log_{10}(\dot{M}O_{2,ave})$	$y_{ij} = \beta_0 + \beta_1 PHO_{ij} + \beta_2 POP_{ij} + \beta_3 SEL_{ij} + \beta_4 POP_{ij} \times SEL_{ij} + \beta_5 POP_{ij} \times PHO_{ij} + \beta_6 POP_{ij} \times logBM_{ij} + \beta_7 logBM_{ij} + \beta_8 WT_{ij} + p_l + e_{ij}$
	Stress sensitivity (post-stress plasma cortisol)	$y_i = \beta_0 + \beta_1 POP_i + \beta_2 SEL_i + \beta_3 POP_i \times SEL_i + \beta_4 ORD_i + \beta_5 BL_i + \beta_6 WT_i + e_i$
	Activity (proportion of time spent swimming during the trial)	$y_{ijk} = \beta_0 + \beta_1 PHO_{ijk} + \beta_2 POP_{ijk} + \beta_3 SEL_{ijk} + \beta_4 POP_{ijk} \times SEL_{ijk} + \beta_5 POP_{ijk} \times PHO_{ijk} + \beta_6 REP_{ijk} + b_i + c_j + d_k + e_{ijk}$
	Latency to emerge from the box	$\lambda(t) = \lambda_0(t) e^{\beta_1 PHO_{ijk} + \beta_2 POP_{ijk} + \beta_3 SEL_{ijk} + \beta_4 POP_{ijk} \times SEL_{ijk} + \beta_5 POP_{ijk} \times PHO_{ijk} + \beta_6 REP_{ijk} + b_i + c_j + d_k}$
	Exploration tendency (1 = explorative, 0 = unexplorative)	$y_{ij} \sim Bernoulli(p_{ijk})$ $logit(p_{ijk}) = \beta_0 + \beta_1 PHO_{ijk} + \beta_2 POP_{ijk} + \beta_3 SEL_{ijk} + \beta_4 POP_{ijk} \times SEL_{ijk} + \beta_5 POP_{ijk} \times PHO_{ijk} + \beta_6 REP_{ijk} + b_i + c_j + d_k$
II. Behaviour responses to burbot olfactory cues	Activity	$y_{ijk} = \beta_0 + \beta_1 SEL_{ijk} + \beta_2 TRE_{ijk} + \beta_3 REP_{ijk} + \beta_4 BL_{ijk} + b_i^{(1)} CON + b_i^{(2)} BUR + c_j + d_k + e_{ijk}$
	Latency to emerge from the box	$\lambda(t) = \lambda_0(t) e^{\beta_1 SEL_{ijk} + \beta_2 GR_{ijk} + \beta_3 REP_{ijk} + \beta_4 BL_{ijk} + b_i + c_j + d_k}$
	Exploration tendency	$y_{ijk} \sim Bernoulli(p_{ijk})$ $logit(p_{ijk}) = \beta_0 + \beta_1 SEL_{ijk} + \beta_2 GR_{ijk} + \beta_3 REP_{ijk} + b_i + c_j + d_k$

β_0 Intercept, *PHO* Photoperiod, *POP* Population, *SEL* Selection, *logBM* Log$_{10}$ body mass in kg, *WT* Water temperature in °C, *ORD* Capture order from the same tank, *BL* Body length in mm – mean (118.8182 mm for I, 122.4464 mm for II), *REP* Trial repeat, *TRE* Treatment, p_l the random effect for chamber l, b_i random effect for fish i, c_j the random effect for arena j, d_k the random effect for batch k, *e* Residual, λ_0 baseline hazard, *t* time, *CON / BUR* binary explanatory variables for burbot and control treatments.

Table 3. Results from models for $\dot{M}O_{2,min}$ and $\dot{M}O_{2,ave}$ and post-stress plasma cortisol. The zero

levels for contrasts were: photoperiod 12:12, population hatchery, and selection line HV. For the

LMMs, F and P- values for the interactions and temperature effect were obtained from Type III

sums of squares and Satterthwaite approximation for degrees of freedom. For the other fixed

effects, linear hypothesis tests using F-test on restricted models with each main effect and its

interactions set to zero were used – residual degrees of freedom are given for these tests. For

cortisol, Type III F-test shown with population and selection line fixed effects estimated using

linear hypothesis tests. Significant ($P < 0.05$) effects shown in bold. For intercepts, t-test values

are shown.

	Fixed effects	Estimate ± SE	Num df	Res / Den df	F	P
$Log_{10}(\dot{M}O_{2,min})$ (LMM)	Intercept	1.82 ± 0.29	1	70	6.32 (t)	<0.001
	Photoperiod	**0.0024 ± 0.02**	**2**	**67.48**	**4.03**	**0.022**
	Population	**-0.84 ± 0.38**	**4**	**52.71**	**3.01**	**0.026**
	Selection	0.054 ± 0.02	2	68.63	2.50	0.090
	Temperature	-0.0009 ± 0.02	1	70	0.002	0.968
	Log$_{10}$ body mass	**1.15 ± 0.15**	**2**	**68.86**	**35.22**	**<0.001**
	Pop × selection	-0.068 ± 0.03	1	70	3.86	0.053
	Pop × photoperiod	**-0.07 ± 0.03**	**1**	**70**	**4.51**	**0.037**
	Pop × log$_{10}$ body mass	**-0.51 ± 0.21**	**1**	**70**	**5.65**	**0.020**
	Random effects	**Variance (SD^2)**				
	Chamber	0				
	Residual	0.07^2				
	Fixed effects	**Estimate ± SE**	**Num df**	**Res / Den df**	**F**	**P**
$Log_{10}(\dot{M}O_{2,ave})$ (LMM)	Intercept	1.18 ±0.44	1	72.50	2.66 (t)	0.010
	Photoperiod	-0.027 ± 0.04	2	70.178	1.52	0.226
	Population	**-0.72 ± 0.54**	**4**	**70.271**	**5.38**	**<0.001**
	Selection	**0.09 ± 0.04**	**2**	**70.763**	**3.17**	**0.048**
	Temperature	-0.03 ± 0.04	1	72.742	0.68	0.412
	Log$_{10}$ body mass	0.38 ± 0.22	2	71.883	1.60	0.209
	Pop × selection	-0.09 ± 0.05	1	70.660	3.05	0.085
	Pop × photoperiod	-0.03 ± 0.05	1	70.373	0.36	0.550
	Pop × log$_{10}$ body mass	-0.49 ± 0.30	1	71.343	2.67	0.107
	Random effects	**Variance (SD^2)**				
	Chamber	0.029^2				
	Residual	0.109^2				
	Fixed effects	**Estimate ± SE**	**Sum sq.**	**Df**	**Test statistic**	**P**
Post-stress cortisol (LM)	Intercept	331.91 ± 225.90	2963.0	1	2.16 (t)	0.162
	Population	47.84 ± 30.43	3392.6	2	2.47	0.142
	Selection	19.35 ± 25.88	767.1	2	0.56	0.466
	Temperature	-15.22 ± 15.38	1343.9	1	0.98	0.338
	Body length	0.07 ± 0.84	8.3	1	0.006	0.939
	Sampling order	-0.75 ± 3.93	50.4	1	0.04	0.851
	Pop × selection	-19.81 ± 37.59	381.1	1	0.28	0.606
	Residual		20587.6	15		

806
807

Table 4. Results of models for behaviour traits in brown trout from hatchery and wild

populations and two angling selection lines (HV and LV). The zero levels for contrasts in all

models were: photoperiod 12:12, population hatchery, and selection line HV. For model

equations, see Table 2. For Activity, F and P-values for the interactions and trial repeat were

obtained from Type III test, and for the other main effects from linear hypothesis tests using

restricted models with each main effect and its interactions set to zero. Fixed effects with $P <$

0.05 shown in bold. For intercepts, t- or z-test values shown.

	Fixed effects	Estimate ± SE	Num Df	Res/Den Df	F	P
	Intercept	**0.30 ± 0.04**	**1**	**33.18**	**7.78 (t)**	**<0.001**
	Photoperiod	**-0.75 ± 0.04**	**3**	**84.03**	**6.53**	**0.001**
	Population	0.08 ± 0.06	3	42.50	1.40	0.242
Activity	Selection	-0.01 ± 0.04	2	74.67	0.10	0.903
(LMM)	Pop × selection	0.025 ± 0.06	1	197.41	1.28	0.259
	Photoperiod × pop	-0.070 ± 0.06	1	81.43	1.37	0.245
	Trial repeat	-0.017 ± 0.02	1	44.24	0.20	0.655
	Random effects	**Variance (SD^2)**				
	ID	0.054^2				
	Batch	0.036^2				
	Arena	0.016^2				

	Fixed effects	Coef	e^{coef}	SE	z	P
	Photoperiod	0.13	1.14	0.26	0.49	0.630
	Population	-0.083	0.92	0.30	-0.27	0.780
Latency	**Selection**	**0.57**	**1.76**	**0.26**	**2.21**	**0.027**
(frailty	Pop × selection	-0.71	0.49	0.37	-1.92	0.054
model)	Photoperiod × pop	-0.060	0.94	0.38	-0.16	0.870
	Trial repeat	**0.23**	**1.26**	**0.082**	**2.83**	**0.005**
	Random effects	**Variance (SD^2)**				
	ID	0.625^2				
	Batch	0.079^2				
	Arena	0.097^2				

	Fixed effects	Estimate ± SE	Wald χ^2	Df	P	
	Intercept	0.55 ± 0.63	0.88 (z)	1	0.380	
	Photoperiod	-0.63 ± 0.63	1.80	1	0.179	
	Population	1.07 ± 0.72	0.058	1	0.810	
Exploration	Selection	1.54 ± 0.64	0.43	1	0.511	
tendency	**Pop × selection**	**-2.46 ± 0.92**	**7.18**	**1**	**0.007**	
(GLMM)	Photoperiod × pop	0.07 ± 0.88	0.006	1	0.936	
	Trial repeat	**0.58 ± 0.21**	**7.79**	**1**	**0.005**	
	Random effects	**Variance (SD^2)**				
	ID	1.398^2				
	Batch	0.270^2				
	Arena	0.762^2				

Table 5. Results of models for activity, latency and exploration tendency in the presence of

predatory olfactory cues and control conditions in brown trout. For activity, the t-test was used

with Satterthwaite approximations to degrees of freedom. The model was fit with restricted

maximum likelihood. For latency, proportional hazard estimates (± standard error) are shown

with hazard ratios (e^{coef}). For latency and exploration tendency, Wald Chisquare test was used to

determine significance of fixed effects. The zero levels for contrasts in all models were:

treatment control and selection line HV. Significant effects ($P < 0.05$) shown in bold.

	Fixed effects	Estimate ± SE	Den Df	t	P
Activity (LMM)	**Intercept**	**0.435 ± 0.060**	**23.25**	**7.198**	**<0.001**
	Selection line	0.012 ± 0.065	9.59	0.18	0.861
	Treatment	-0.081 ± 0.042	15.85	-1.928	0.072
	Trial repeat	**-0.029 ± 0.010**	**71.62**	**-2.97**	**0.004**
	Body length	-0.00009 ± 0.004	8.63	-0.02	0.984
	Random effects	**Variance (SD²)**			
	ID (burbot)	0.122²			
	ID (control)	0.077²			
	Batch	0.000			
	Arena	0.000			
	Residual	1.400²			

	Fixed effects	Coef ± SE	e^{coef}	z	P
Latency (frailty model)	Selection line	0.336 ± 0.372	1.399	0.900	0.370
	Treatment	0.089 ± 0.220	1.093	0.400	0.690
	Trial repeat	0.113 ± 0.065	1.120	1.759	0.080
	Body length	-0.006 ± 0.026	0.994	-0.230	0.820
	Random effects	**Variance (SD²)**			
	ID	0.661²			
	Batch	0.210²			
	Arena	0.080²			

	Fixed effects	Estimate ± SE	z	P
Exploration tendency (GLMM)	Intercept	1.449 ± 0.801	1.808	0.0706
	Selection line	0.659 ± 0.724	0.910	0.363
	Treatment	-0.63 ± 0.487	-1.294	0.196
	Trial repeat	-0.066 ± 0.142	-0.465	0.642
	Random effects	**Variance (SD²)**		
	ID	1.140²		
	Batch	0.000		
	Arena	$(1.847^{-5})^2$		

Figure captions

Fig.1. 3D-illustration of the arena used in personality trials without the left side wall. Water flow

direction is left-right. Burbot was placed in the area indicated by blue colour, upstream from the

net (inaccessible to the brown trout). The grey box indicates the start box, where the fish was

placed before the start of a trial. Latency was measured as time to emerge from the box. Activity

was measured as swimming activity outside the start box arena after emergence. Exploration

tendency was measured as the whole body of fish passing the gate indicated by an arrow.

Fig. 2. Means and 75% confidence intervals for A) mass-specific $\dot{M}O_{2,min}$, using mass-scaling

exponent 0.928; and B) mass-specific $\dot{M}O_{2,ave}$ using raw body mass (scaling exponent was 0.19

and was not used). For statistical significance, see Table 3. Wild and hatchery populations shown

with two angling selection lines produced from individuals with high (HV) or low (LV)

vulnerability to angling. 12:12 = L:D rhythm, 24 = continuous light. N in each group shown in

Table 1. Legend shown in A.

Fig. 3. Behavioural differences between two angling vulnerability selection lines (HV – high

vulnerability, LV – low vulnerability) within the hatchery and wild populations. A) Curves

showing the proportion of individuals emerged from the start box, drawn with Kaplan-Meier

estimator. Photoperiods are combined within each breeding group, confidence intervals omitted

for clarity. Higher proportion indicates higher boldness. B) Predicted activity from LMM with

75% confidence intervals for predicted values. Significantly lower activity was observed in 24

(constant light) compared to 12:12 (light-dark rhythm) (Table 4). C) Predicted exploration

tendency from GLMM with 75% confidence intervals for predicted values. Angling selection had

opposing effects on exploration tendency in the two populations (Table 4). In B and C,

predictions were made for the first trial repeat. For N in each group, see Table 1.

Fig. 1.
854

856 Fig. 2.
857

858
859

860   Fig. 3.