The cryptic impacts of invasion: functional homogenization of tropical ant communities by invasive fire ants

Mark K. L. Wong[1],*, Benoit Guénard[2] and Owen T. Lewis[1]

[1]Department of Zoology, University of Oxford, South Parks Road, Oxford OX1 3PS, UK

[2]School of Biological Sciences, The University of Hong Kong, Kadoorie Biological Sciences Building, Pok Fu Lam Road, Hong Kong SAR, China

*Author for correspondence (E-mail: mark.wong@zoo.ox.ac.uk; Tel.: +44 1865 271234)

Abstract

1. Invasive insects represent major threats to ecosystems, with known impacts on the taxonomic diversity of invaded communities. However, their effects on the functional dimension of biodiversity, measured as the diversity and distribution of traits, are poorly known. Such measures can determine the resilience of a community and of the ecosystem processes it modulates.

2. The fire ant *Solenopsis invicta* is a widespread and highly problematic invasive species. Its impacts on the taxonomic diversity of native ant communities have been widely studied, but not its impacts on their functional diversity. Comparing invaded and uninvaded plots in tropical grasslands of Hong Kong, we investigated how the presence of *S. invicta* affects the diversity and distribution of ant species and traits

within and across communities, the functional identities of communities, and

functionally unique species.

3. Using a trait-probability density framework, we estimated the functional diversity of

individual species, including the trait variation from intraspecific polymorphisms. We

then scaled up these values to calculate functional diversity at the community level.

4. Invasion had only limited effects on species richness and functional richness, which

were 13% and 8.5% lower in invaded communities respectively. In contrast, invasion

had pronounced effects on taxonomic and functional composition due to turnover in

species and trait values. Furthermore, invaded communities were functionally more

homogeneous, displaying 23% less turnover and 56% more redundancy than

uninvaded communities, as well as greater clustering and lower divergence in trait

values. Invaded communities had fewer functionally-unique individuals and were

characterized by ant species with narrower heads and bodies and shorter mandibles.

5. *Synthesis and applications*. Our results suggest that the full ramifications of invasions

are likely to be underestimated if based solely on taxonomic measures of diversity or

indices describing trait variety. Invasions by *S. invicta* led to significantly altered

functional identity and increased functional homogenization of native ant

communities, but only marginally lower taxonomic and functional richness.

Investigating the diversity and distributions of traits at species, community and

landscape levels can reveal the 'cryptic impacts' of invasions which, despite causing

little taxonomic change, may substantially modify the structure and functioning of

ecological communities.

Keywords

Beta diversity, functional diversity, functional trait, functional uniqueness, insect,

intraspecific, invasive species, turnover

Introduction

Invasions by alien species reduce native species richness, drive cascading impacts on

ecosystems, cause economic damage and undermine human health (Pyšek & Richardson,

2010). As with many ecological phenomena, invasions have long been studied by

summarising ecological communities using metrics based on species' taxonomic identities:

abundance, richness and diversity. Changes in these metrics, however, may provide only

limited insight into the specific mechanisms (such as novel niches and enemy escape)

underlying the causes of invasions (MacDougall, Gilbert & Levine, 2009). Furthermore, the

effects of biodiversity on ecosystem processes often depend on the functions performed by

the different species, rather than on species numbers and identities *per se* (Gagic et al., 2015).

Summarising biodiversity in terms of the traits that directly impact organisms' ecological

interactions (Functional Diversity) (McGill et al., 2006) may advance understanding of the

causes and consequences of invasions.

Traits are the specific phenotypic properties of organisms which modulate their ecological

interactions. The interactions of key 'functional traits' influence organism fitness and may

also contribute to ecosystem functions (Wong, Guénard & Lewis, 2018). Thus, studying the

diversity of functional traits in an ecological community can simultaneously reveal how

biodiversity and ecosystem processes are impacted by disturbances such as invasive species.

For instance, trait-based studies of plant and vertebrate communities undergoing invasion by

alien species revealed declines in functional richness (the variety of trait values in individual

communities) and a tendency for functional homogenization (i.e., an increased similarity in

trait values between communities) (Villéger, Grenouillet & Brosse, 2014; Castro-Díez et al.,

2016). Crucially, invasion-driven changes in functional structure were linked to altered

ecosystem functions (Castro-Díez et al., 2016).

One general limitation of earlier trait-based studies, including those on invasions, has been

the exclusive use of mean trait values for species to estimate community functional diversity.

This approach underestimates intraspecific trait variation, which can strongly influence

community dynamics and ecosystem processes (Des Roches et al., 2018). Intraspecific trait

variation has also been implicated in the success of some invasive species (González-Suárez,

Bacher, & Jeschke, 2015). Recently-developed statistical tools such as the Trait Probability

Density (TPD) framework can incorporate intraspecific trait variation into estimates of

functional diversity (Carmona et al., 2016), but few studies have explored this within the

context of invasions. Additionally, although trait-based studies are advancing understanding

of invasions by plants and vertebrates, there is a shortage of similar work on invasive insect

species (Wong et al., 2018), despite their ubiquity and widespread impacts on biodiversity

and ecosystem services (Bradshaw et al., 2016).

Many ants, for instance, are contenders for the world's most harmful invasive species (Lowe

et al., 2000). Numerous studies have documented the consequences of ant invasions for

native ant communities, which include declines in species richness, taxonomic

homogenization, and phylogenetic clustering (reviewed in Lessard et al., 2009). These studies

focus on taxonomic and phylogenetic patterns, and rarely consider trait values directly. Few

such studies document the phenotypes that were lost or gained at the community level, and

none directly assessed the impacts of invasions on the functional diversity of individual

communities (functional alpha diversity) or patterns across multiple communities (functional

beta diversity). Furthermore, no trait-based studies on ants, and few for insects in general

include intraspecific trait variation in estimates of community functional diversity (Wong et

al., 2018). Intraspecific trait variation is expected to be high in polymorphic species and it

may influence how such species respond to or effect ecological change. Some ant species

display marked variation in the body size and morphology of their worker caste (worker

polymorphism), a feature which may contribute to colony fitness and ecological success

(Tschinkel, 1988; Wilson, 2003). It has been observed that polymorphic ant species can

surpass monomorphic ones in their abilities to collect resources varying in size, and to access

environments varying in rugosity (Farji-Brener, Barrantes & Ruggiero, 2004); that is,

polymorphic species may access a wider variety of niches than monomorphic species. Thus, a

basic yet apparently untested assumption is that polymorphic species have higher functional

(trait) richness than monomorphic species.

The Red Imported Fire Ant, *Solenopsis invicta* Buren, 1972, is native to South America but

its expanding global range already encompasses five continents (Guénard et al., 2017). The

species has been an extensive ecological problem throughout the United States for decades

due to its strong impacts on native biodiversity at multiple trophic levels (including native

ants) and the subsequent cascading effects on ecosystems (Porter & Savignano, 1990;

Vinson, 1997). *Solenopsis invicta* are dietary generalists and mature colonies consist of a

polymorphic worker caste (Tschinkel, 2006), a factor which, alongside their strong

interspecific aggression, may contribute to the success of invasive populations (Tschinkel,

1988; 2006). *Solenopsis invicta* was first reported in Taiwan in 2003, and in continental

China in 2005 (Ascunce et al., 2011). In spite of their devastating impacts elsewhere, studies

on the ecology of *S. invicta* invasions in Asia have been limited. Preliminary observations

from agrosystems in China suggest *S. invicta* invasions may be associated with declines in

the species richness of arthropod communities (Wang et al., 2018).

Here, we investigate the impacts of the invasion of *S. invicta* on taxonomic and functional

diversity within (alpha diversity) and between (beta diversity) native ant communities in

Hong Kong. This is the first comprehensive trait-based study of an ant invasion's impact on

the functional facet of biodiversity, as well as the first to incorporate polymorphisms in

calculations of functional diversity at the species level.

At the scale of the individual community, we examined how invasion by *S. invicta* affected

(i) species and functional richness, (ii) abundance-weighted indices of multidimensional

functional diversity, and (iii) functional identity, the dominant value of a trait in the

community. We predicted lower species and functional richness as well as altered functional

identities in communities invaded by *S. invicta* (Porter & Savignano, 1990; Castro-Díez et al.,

2016). At the multi-community (landscape) scale, we investigated how *S. invicta* invasion

affected taxonomic and functional beta diversity (the dissimilarities in species and traits

between communities). We predicted that invasion by *S. invicta* would lead to taxonomic

homogenization (Lessard et al., 2009), which would be associated with functional

homogenization, as observed for other alien taxa (Villéger et al., 2014). Using measures of

species-level functional diversity, we also identified functionally unique species and

compared their relative uniqueness to uninvaded and invaded communities. Lastly, we

measured and compared the functional richness of different species, with the prediction that

the functional richness of polymorphic species such as *S. invicta* would exceed that of

monomorphic species.

Materials and methods

Study area and sampling design

Our study sites are two (<2 km apart) wetland reserves in northern Hong Kong: Lok Ma Chau (22.512°N, 114.063°E) and Mai Po (22.485°N, 114.036°E). Both reserves encompass abandoned fish farms that have since been conserved for >35 years as habitats for migratory birds. Each contains a network of bunds (width ≤5 m) which separate individual ponds (Fig. S1). The habitat is relatively homogeneous and comprises exposed grasslands with native tree species interspersed throughout. Ant communities in this landscape are comprised mostly of native species but pilot surveys from 2015 to 2017 revealed that colonies of *S. invicta* are present at high densities at multiple locations. We marked these locations, and in 2018 selected a total of 61 plots, each a 4 × 4 m quadrat, to reflect two ant community types: communities with *S. invicta* absent (uninvaded; 37 plots), and those with *S. invicta* present (invaded; 24 plots). A minimum distance of 20 m between individual plots facilitated independent observations since most ant species in the region forage no further than 5 m from their nests (Eguchi, Bui & Yamane, 2004) and *S. invicta* forage within 4 m of their nests (Weeks, Wilson & Vinson, 2004). Given the homogeneity of the landscape we assumed that any community differences observed between uninvaded and invaded plots would primarily be a consequence of invasion by *S. invicta*; environmental data collected at fine spatial resolutions were used to test this assumption (see below). Sampling was conducted from April to September 2018. At each plot, six pitfall traps (55 mm in diameter) were installed to sample the ant community over 48 hours. All specimens were sorted into morphospecies and subsequently most were identified to species using taxonomic keys.

Environmental data

We used local GIS models (Morgan & Guénard, 2019) to obtain high-resolution data (30 ×

30 m rasters) for three environmental variables corresponding to each plot: Normalized

Difference Vegetation Index (NDVI), mean annual temperature, and mean annual

precipitation.

Assembling the individual-level trait dataset

Here we aimed to obtain values of functional diversity that incorporated intraspecific trait

variation, including the variation arising from worker polymorphisms. We assembled an

individual-level trait dataset comprising data for seven morphological traits that regulate ant

physiology and behaviour and that are hypothesized to impact performance and fitness (Table

1). Using mounted specimens from the pitfall traps and a Leica M165c stereo microscope

paired with Leica Application Suite software, we recorded high-resolution images and

performed trait measurements on at least 10 individual workers of every species (N=319).

For dimorphic species of *Camponotus* and *Pheidole* where workers comprise two distinct

sub-castes (minors and majors), we included trait data for individuals of both sub-castes

based on the relative proportions (i.e., ratio of minors to majors) observed in natural colonies

(Passera, 1984; Wilson, 2003). The invader *S. invicta* has a polymorphic worker caste, and

Tschinkel (1988) showed that this polymorphism is mainly expressed in the morphological

variation displayed by the 'majors' (head width >0.7 mm), which are present only in mature

colonies where they comprise 35% of the worker population; 'minors' of head width <0.7

mm comprise the remaining 65% in mature colonies and juvenile colonies only consist of

minors. We observed that all invaded plots contained majors (head width >0.7 mm),

suggesting they were mature colonies; thus, our trait data for *S. invicta* (n=20) included both

minors (65% of individuals) and majors (35% of individuals).

Table 1. The seven traits measured on each individual, and each trait's hypothesized links to

the performance and fitness of ants.

Trait	Measurement	Hypothesized link to performance and fitness
size	Weber's length: diagonal length of mesosoma	Modulates vital and physiological rates, determines physical constraints and exposure to predators, influences resource type and acquisition efficiency (Silva & Brandão, 2010).
head width	Width of head including eyes	Determines the size of gaps through which an individual can pass (Schofield, Bishop & Parr, 2016) and the volume of muscles powering the mandibles during foraging (Richter et al., 2019).
eye width	Width of left eye	Determines ability in navigation, foraging, predator and prey detection, and indicative of activity times (Silva & Brandão, 2010).
mandible length	Length of left mandible	Responds to selection on diet type and specialization (Silva & Brandão, 2010).
scape length	Length of scape of left antenna	Responds to selection on navigation and sensory abilities (Silva & Brandão, 2010).
pronotum width	Width of pronotum	Determines volume of muscles for head-support and load-bearing (Keller et al., 2014).
leg length	Combined length of femur and tibia of left hind leg	Determines mobility; leg length influences running speed, which affects success in foraging or escape from predators (Grevé et al., 2019).

Compressing trait variation

We divided the measurements of six traits (*head width*, *eye width, mandible length*, *scape*

length, *pronotum width* and *leg length*) by *size* to reduce correlation with body size. We then

log-transformed the values of all seven traits to reduce the influence of extreme values, and

standardized trait values to have mean of zero and unit variance. Next, we used Principal

Components Analysis (PCA) to synthesize the major axes of variation in multidimensional

trait space and to reduce the number of dimensions used to calculate functional diversity

indices. We performed the PCA using the mean trait values of each species and subsequently

predicted the values of the PCA components for all individuals in the dataset. We used

species means instead of individual trait values in the PCA because using the latter could bias

the analysis if some species had disproportionately large numbers of individuals in the

dataset. We retained the first two components of the PCA, which had eigenvalues greater

than unity, and which captured 76.9% of the total variance in traits. We then predicted the

values of these two components for every individual in the trait dataset and used these new

'traits' to calculate functional diversity indices.

Functional diversity from species to communities

All functional diversity indices were calculated using the Trait Probability Density

framework which incorporates intraspecific variation, the multidimensional nature of traits,

species abundances, and probabilistic trait distributions (see Carmona et al., 2016). First, we

used multidimensional probability density functions to calculate trait probability distributions

(which reflect the probabilities of observing different trait values) at the level of individual

species (TPDsp). Next, we scaled up TPDsp to local community levels (TPDcom) by

summing the TPDsp of all species in each local ant community, weighted by their relative

abundances – which we estimated as frequencies of occurrence. Finally, five different indices

for functional diversity were calculated using each community's TPDcom. The indices were

Functional Richness (FRic), the volume of functional space occupied by the community;

Functional Evenness (FEve), the regularity of the distribution of abundance in functional

space; Functional Divergence (FDiv), a measure of how abundances tend to be on the outer

margins of the functional space while controlling for functional richness; Rao, the

abundance-weighted dispersion of individuals (or species) in functional space; and

Functional Redundancy (FRed), the degree to which trait values are represented by multiple

species in the community (Carmona et al., 2016). We used this multi-index approach to

measure functional diversity because no one index can encapsulate the independent

components of functional diversity (Mouchet et al., 2010). In addition to calculating the

observed values of the functional diversity indices, we calculated Standardized Effect Size

(SES) values for all indices so as to estimate community-level functional diversity that had

been corrected for potential effects of species richness (Swenson, 2014). SES values were

calculated by comparing the observed values to values generated from 999 constrained null

models randomizing the community data matrix using the "Independent Swap" algorithm.

The formula for calculating SES is:

$SES = (Mean_{observed} - Mean_{null})/s.d._{null}$

Functional identity

242   To estimate functional identity, we calculated the community-weighted mean (CWM) for

243   each trait in every local ant community. A CWM reflects the dominant value of a given trait

244   in a given community (Swenson, 2014). We calculated CWMs using mean trait values of

245   species weighted according to their relative abundances in the different communities. Size-

246   correction was applied to all traits except *size* (see above).

Taxonomic and functional beta diversity

We calculated six pairwise measures of taxonomic and functional beta diversity for all

possible pairs of local ant communities. We used matrices of species' abundances to calculate

the Taxonomic Dissimilarity between each pair, and further decomposed this into Taxonomic

Turnover (dissimilarities arising from the replacement of species between communities), and

Taxonomic Nestedness (dissimilarities in the relative abundances of species that occurred in

both communities). We calculated the Functional Dissimilarity between paired communities

using their TPDcom, and further decomposed this into Functional Turnover (dissimilarities in

the trait values between communities), and Functional Nestedness (dissimilarities in the

relative abundances of trait values shared between communities). In addition to observed

values, we calculated SES values (from comparisons with 999 constrained null models using

the "Independent Swap" algorithm) for all components of functional beta diversity. Although

the "Independent Swap" algorithm may not be optimal for generating null models of beta

diversity patterns shaped by dispersal limitation (Swenson, 2014), this is unlikely to be a

problem for the present study because all species disperse by flying alates that can travel

distances exceeding the scale of the study landscape (the maximum distance between any two

plots was 4 km).

Species' functional richness and functional uniqueness

We calculated functional richness and functional uniqueness values for all species.

Functional richness was calculated based on each species' trait probability distribution

(TPDsp) (Carmona et al., 2016). Species' functional uniqueness values were calculated

relative to individual local communities, based on the degree to which a species' functional

space (TPDsp) did not overlap with a local community's functional space (TPDcom)

(Carmona et al., 2016). We calculated each species' 'relative uniqueness' with respect to the

different uninvaded and invaded communities, as well as its 'objective uniqueness' in the

species pool (using a hypothetical community containing all species at equal abundance).

Statistical analysis

Taxonomic alpha diversity, functional diversity and CWMs

We used separate linear mixed-effects models to assess whether the values of alpha

taxonomic diversity, functional diversity indices (including observed and SES values) and

CWMs differed significantly between uninvaded and invaded local ant communities, while

including a random effect of environmental variation that was captured in the first component

of a PCA for the three environmental variables, which had eigenvalues greater than unity.

Taxonomic and functional beta diversity

We used non-metric multidimensional scaling (NMDS) to scrutinize the relationships

between and among invaded and uninvaded local ant communities in multidimensional space

(Fig. S2). We used PERMANOVA (9,999 permutations) to quantify dissimilarity, turnover

and nestedness between the observed taxonomic and functional compositions of uninvaded

and invaded communities. We used permutation tests for multivariate dispersions to assess

whether the levels of taxonomic and functional beta diversity (in three components) observed

among uninvaded communities differed from those observed among invaded communities.

We also used nonparametric Mann-Whitney U tests to compare SES values of the three

functional beta diversity components between uninvaded and invaded communities.

Functional uniqueness of individual species

We calculated each species' average relative uniqueness to uninvaded and invaded

communities and regressed these against its objective uniqueness in a linear model.

Software

We used the following packages in R software version 3.3.3 (R Core Team, 2017): *TPD*

(Carmona, 2018) for calculating trait probability distributions, functional diversity indices,

functional dissimilarity and functional uniqueness measures, *FD* (Laliberté, Legendre &

Shipley, 2014) for calculating CWMs, *betapart* (Baselga et al., 2018) for beta diversity

analyses, *lme4* (Bates et al., 2015) for linear mixed-effects models, *MASS* (Venables &

Ripley, 2002) for NMDS, and *ggplot2* (Wickham, 2009) for the production of graphics.

Results

Community composition and species richness

A total of 29 ant species (including *S. invicta*) were collected from 366 pitfall traps in 37

uninvaded plots and 24 invaded plots (Table S1). The species composition across invaded

and uninvaded communities was similar overall, with 27 of the 28 native species occurring in

both community types, and only one species not found in invaded communities. On average,

the species richness of invaded communities was marginally and non-significantly lower (by

13%) than that of uninvaded communities (Table 2; Fig. 1).

Functional diversity and CWMs

Uninvaded and invaded communities had similar levels of FRic and FEve for both observed

and SES values. However, in linear mixed-effects models, the observed FDiv and Rao of

invaded communities were significantly lower than those of uninvaded communities by 11%

and 7% respectively, and the FRed of invaded communities was significantly higher than that

of uninvaded communities by 56% (Table 2; Fig. 1); similar relationships were observed for

SES values. CWMs for the traits *size*, *scape length*, *eye width* and *leg length* did not differ

significantly between uninvaded and invaded communities. By contrast, the CWMs for *head*

width, *pronotum width* and *mandible length* were significantly smaller (by 4–7%) in the

invaded communities (Table 2; Fig. 2).

Taxonomic and functional beta diversity

Uninvaded and invaded communities were significantly dissimilar in both taxonomic and

functional composition, and these dissimilarities were driven by turnover in species as well as

trait values (Table 3; Fig. 3). The observed levels of total taxonomic and functional

dissimilarities among both uninvaded and invaded communities were comparable (Table 4),

but SES values revealed that total functional dissimilarity was lower among invaded

communities when corrected for species richness (Mann-Whitney U test: $P < 0.001$) (Fig. 3).

Invaded communities had significantly lower levels of functional turnover (by 23%) and

higher functional nestedness (by 20%) in observed values; these relationships were

maintained in SES values (Mann-Whitney U tests: $P < 0.001$) (Table 4). Likewise, invaded

communities were significantly more taxonomically nested than uninvaded communities (by

42%, Table 4; Fig. 3). That is, in comparison to uninvaded communities, relatively greater

proportions of the total taxonomic and functional dissimilarities among invaded communities

were driven by losses of species than by replacements of species, and by changes in the

abundances of trait values than by changes in the trait values themselves, respectively.

340   Species' functional richness and functional uniqueness

Functional richness varied over four-fold among species (Min. = 0.86, Max. = 3.61) (Fig. 4).

The four most functionally-rich species were two dimorphic species of *Camponotus*, another

dimorphic species, *Pheidole nodus*, followed by the polymorphic invader *S. invicta*. In

separate linear regressions, species' relative uniqueness to both uninvaded and invaded

communities increased with their objective uniqueness (Fig. 5). However, there was

relatively more overlap between the functional spaces of less unique species and the

functional spaces of invaded communities (Intercept$_{Invaded}$ = -0.15; Intercept$_{Uninvaded}$ = 0.45).

Furthermore, the relative uniqueness of species to invaded communities increased more

steeply with an increase in objective uniqueness (Slope$_{Invaded}$ = 1.17; Slope$_{Uninvaded}$ = 0.52), such that

very unique species were more unique to invaded communities than to uninvaded

communities (Fig. 5).

Table 2. Summary statistics for response variables in separate linear mixed-effects models

with community type (uninvaded vs. invaded) as fixed effects and environmental variation as

a random effect.

Index	Uninvaded (mean ± SE)	Invaded (mean ± SE)	χ^2	df	*P*
Species Richness	8.97 ± 0.42	7.80 ± 0.67	3.06	1	0.08
FRic	14.1 ± 0.61	12.9 ± 0.97	1.59	1	0.21
FRic.SES	-0.008 ± 0.15	0.22 ± 0.24	0.86	1	0.35
FEve	0.60 ± 0.007	0.59 ± 0.01	0.24	1	0.62
FEve.SES	0.14 ± 0.16	-0.09 ± 0.26	0.83	1	0.36
FDiv	0.73 ± 0.01	0.65 ± 0.02	17.8	1	<0.001***
FDiv.SES	0.61 ± 0.15	-0.46 ± 0.24	20.7	1	<0.001***
Rao	0.82 ± 0.01	0.76 ± 0.02	7.97	1	<0.01**
Rao.SES	0.44 ± 0.13	-0.42 ± 0.2	18.0	1	<0.001***
FRed	0.34 ± 0.05	0.53 ± 0.07	13.8	1	<0.01**
FRed.SES	-0.48 ± 0.13	0.55 ± 0.21	23.9	1	<0.001***
CWM*head*	0.76 ± 0.005	0.72 ± 0.007	23.7	1	<0.001***
CWM*pron*	0.52 ± 0.003	0.48 ± 0.004	85.8	1	<0.001***
CWM*mand*	0.42 ± 0.003	0.40 ± 0.005	10.1	1	<0.01**

Figure 1. Boxplots showing species richness and observed values of five functional diversity

indices in 37 uninvaded and 24 invaded local communities. Dots show values of individual

communities, thick bars show means, boxes show inter-quartile range and vertical lines

extend to maximum and minimum values (excluding outliers). Asterisks indicate statistical

significance (*** $P<0.001$, ** $P<0.01$, * $P<0.05$, NS, not significant).

Figure 2. Boxplots displaying community-weighted mean values for size-corrected *head*

width, *pronotum width* and *mandible length* in 37 uninvaded and 24 invaded communities.

Dots show values of individual local communities, thick bars show mean values, box edges

371 show standard deviations, and vertical lines extend towards minimum and maximum values.

372 Asterisks indicate statistical significance (*** $P<0.001$, ** $P<0.01$).

375 Table 3. Results of PERMANOVA tests for dissimilarities between uninvaded and invaded

376 communities in their observed taxonomic and functional compositions.

Beta diversity	Component	F	R^2	P
Taxonomic	Total Dissimilarity	22.4	0.28	<0.001***
	Turnover	31.9	0.35	<0.001***
	Nestedness	13.6	0.31	1.0
Functional	Total Dissimilarity	33.1	0.36	<0.001***
	Turnover	32.7	0.36	<0.001***
	Nestedness	14.7	0.33	1.0

379 Table 4. Permutation tests for multivariate dispersions, with calculations based on the

380 average distances to centroids of uninvaded and invaded communities for different

381 components of taxonomic and functional beta diversity. These tests compare the levels of

382 beta diversity observed among uninvaded communities to those observed among invaded

383 communities.

Beta diversity	Component	Uninvaded	Invaded	F	P
Taxonomic	Total Dissimilarity	0.35	0.36	0.10	0.77
	Turnover	0.29	0.28	0.20	0.66
	Nestedness	0.07	0.10	5.89	0.02*
Functional	Total Dissimilarity	0.32	0.31	0.01	0.91
	Turnover	0.30	0.23	5.01	0.03*
	Nestedness	0.41	0.49	7.37	<0.01**

Figure 3. Observed levels of taxonomic beta diversity and functional beta diversity corrected

for species-richness (SES values) in three measures of dissimilarity (Total, Turnover, and

Nestedness). Boxplots show values among uninvaded communities, among invaded

communities, and between uninvaded and invaded communities. Asterisks indicate statistical

significance (*** $P<0.001$, * $P<0.05$, NS, not significant).

Figure 4. Functional richness of ant species with three different degrees of polymorphism.

Bars show values for 22 monomorphic species (light grey), six dimorphic species (dark grey),

and the polymorphic species *S. invicta* (black). Full species names are listed in Table S1, and

images of monomorphic, dimorphic and polymorphic species are shown in Fig. S4.

Figure 5. Species' relative functional uniqueness to local communities that were uninvaded

(grey dotted line; circles) and invaded (black solid line; triangles) plotted against their

objective functional uniqueness in the species pool.

Discussion

Most concerns surrounding biological invasions cite declines in the species richness of

invaded communities. However, invasions can also impact the structure and function of

ecological communities in ways that are less detectable with taxonomic measures of

diversity. Here we investigated the impacts of an invasion on the diversity and distribution of

species and traits in individual communities (alpha diversity) and across multiple

communities (beta diversity). We found that invasion by *S. invicta*, one of "the world's worst

invasive species" (Lowe et al., 2000), led to the functional homogenization and altered

functional identity of ant communities in Hong Kong despite effecting marginal changes in

species and functional richness. Additionally, we observed higher functional richness in

polymorphic species than monomorphic species. We discuss the possible ecological drivers

and ecosystem implications of our findings below.

Polymorphic species are more functionally rich

Polymorphisms are among the most conspicuous and pervasive sources of intraspecific trait

variation (Ford, 1957). It is conceivable that polymorphic species, with greater variation in

morphology, would display higher functional richness. We generally found this to be the case

among the 29 ant species studied. Dimorphic species of the genera *Camponotus* and *Pheidole*

and the polymorphic *S. invicta* displayed 2–322% more functional richness than

monomorphic species (Fig. 4). However, the functional richness of three dimorphic species

was higher than that of the polymorphic *S. invicta*. This likely resulted from the pronounced

allometry of the discrete major subcastes of dimorphic species, which generated extreme trait

values; polymorphic *S. invicta* workers, by contrast, display a continuous variation with size

(Figs. 4 and S4). While it is unlikely that the morphological traits measured were

representative of all species' multidimensional niches, our findings demonstrate that, at least

across multiple aspects of morphology linked to foraging, mobility and physiology (Table 1),

polymorphic species occupy more functional space and may have access to a greater variety

of niches than monomorphic species, as previously suggested (Farji-Brener et al., 2004).

Future work investigating whether functional richness predicts niche variety or specialisation

across polymorphic and monomorphic species could further our understanding of how

species and ecosystem processes respond to ecological change.

Lower species and functional richness in invaded communities

Disturbances drive local extinctions by decreasing the abundances of particular species with

vulnerable trait combinations; this process can initially occur without significantly modifying

community composition or affecting total species and functional richness (Mouillot et al.,

2013). Invasions by *S. invicta* in Hong Kong generally follow these patterns. Two thirds of

the native species had lower abundances in the invaded communities (Fig. S3), but the same

communities displayed only marginally lower species richness (by 13%) and functional

richness (by 8.5%) (Table 2 and Fig. 1). These results differ from previously reported

extensive declines (by 69%) in the species richness of ant communities invaded by *S. invicta*

in North America (Porter & Savignano, 1990). The current absence of an extensive decline in

species richness may be associated with the relatively younger *S. invicta* invasion in Hong

Kong (Ascunce et al., 2011), and potentially the ecological differences (and thus responses)

between the tropical grassland ant communities studied here, which comprised many

disturbance specialists, and the ant communities of temperate forests in previous studies

(Porter & Savignano, 1990). Aside from the marginal differences in species and functional

richness, multiple indices sensitive to changes in the abundance and distribution of trait

values differed significantly between uninvaded and invaded communities (Table 2: CWMs,

FDiv, Rao, and FRed). Although each index measures a unique aspect of functional identity

and diversity (Mouchet et al., 2010), the results collectively indicate that invasions by *S.*

invicta exert a non-random selection on native communities through the trait values of

individuals; we discuss these patterns in the two sections that follow.

Altered functional identity of invaded communities

The CWMs for size-corrected *head width*, *pronotum width* and *mandible length* of ants in

invaded communities decreased significantly by 4–7% (Fig. 1), in line with our hypothesis

that invasion would alter communities' functional identities. The results suggest that invasion

selects for individuals with narrower heads and pronotums and shorter mandibles. One

hypothesis is that the observed patterns relate to mobility. It has been shown that the width of

an ant's head and pronotum determine the size of gaps through which it can pass (Schofield

et al., 2016). Narrower heads and pronotums of ants in invaded communities could thus

reflect demands for moving through tighter spaces to avoid the behaviourally dominant *S.*

invicta during foraging (Tschinkel, 2006) or to reach resources in less accessible locations.

For instance, studies on other ant invasions reported that native hypogaeic species,

specialised to forage and move through soil, increased in relative abundance by 50% (Human

& Gordon, 1997). A second hypothesis relates to diet. In ants, long mandibles of many

predatory species are specialized adaptations for prey capture (Silva & Brandão, 2010), and

larger heads and pronotums afford more space for the musculature powering snapping,

gripping and load-bearing abilities (Keller et al., 2014; Richter et al., 2019). Hence, the

shorter mandibles and narrower heads and pronotums of ants in invaded communities may

reflect a less specialized or more herbivorous diet. Such ants may use more liquid foods (e.g.,

honeydew from hemipterans) that can be ingested relatively quickly, and which require less

manipulation (capture, ripping, transport) than solid foods. The presence of the behaviourally

dominant *S. invicta* may select for ants that 'grab and go' over those which remain at food

sources for longer periods.

Functional clustering and redundancy of invaded communities

In invaded communities, values of FDiv and Rao decreased significantly by 11% and 7 %

respectively, while FRed increased significantly by 56% (Fig. 2). These patterns arose due to

the presence of more individuals with similar trait values in invaded communities.

Specifically, FDiv reflects the degree to which the distribution of species' abundances in

functional space maximizes total community variation in trait values (and niches) (Mouchet

et al., 2010). Hence, the results suggest that the most abundant species in uninvaded

communities are more differentiated in their niches (higher FDiv), while those in invaded

communities have more similar niches (lower FDiv). As observed for FDiv, the lower Rao in

invaded communities is indicative of niche clustering (species' abundances are relatively

more clustered in functional space) (Mouchet et al., 2010). FRed reflects the degree to which

specific trait values are represented by multiple species in the community (Carmona et al.,

2016). In uninvaded communities (lower FRed) fewer species share the same trait values,

suggesting less niche overlap. By contrast, more species share the same trait values in

invaded communities (higher FRed), suggesting more niche overlap. Therefore, as observed

for the CWMs, the results for FDiv, Rao and FRed show a general pattern of species'

abundances converging towards particular trait values in invaded communities, which is one

signature of selection (Vellend, 2016).

Functional homogenization across invaded communities

Our analysis of functional alpha diversity suggests that *S. invicta* invasions are associated

with a selection for specific trait values in individual communities. Because such selection

has repeated over separate communities invaded by *S. invicta*, functional beta diversity

patterns across multiple communities show a trend towards functional homogenization. This

is evident from the significantly lower functional dissimilarity among communities where *S.*

invicta is present (Fig. 3). Contrary to our hypothesis, however, functional turnover did not

track taxonomic turnover; changes in the species found in different invaded communities

were not matched proportionately by changes in those communities' trait values. The invaded

communities actually retained similar levels of taxonomic turnover to uninvaded

communities (Fig. 3). However, the former displayed significantly less functional turnover in

observed structure (by 23%; Table 4), as well as in SES values of functional beta diversity

corrected for the effects of species richness (Fig. 3). In previous analyses using computer

simulations, such patterns of low functional turnover amid higher taxonomic turnover were

predicted to emerge most frequently when there are high levels of functional redundancy in

individual communities (Baiser & Lockwood, 2011). Given that communities invaded by *S.*

invicta displayed 56% more functional redundancy than uninvaded communities (Fig. 2), our

observations in an invasion context provide empirical support for the theoretical predictions

(Baiser & Lockwood, 2011).

Functionally ordinary winners and functionally unique losers across invaded

communities

Invasions and other disturbances can result in losses of functionally unique species before

functionally redundant species (Flynn et al., 2009). Thus, examining the abundance and

distribution of functionally unique species may promote the advanced detection of invasion

impacts. Previous studies used trait patterns of aggregated communities to define functionally

unique groups or species, then analysed their abundances within each community (e.g.,

Coetzee & Chown, 2016). A species' functional uniqueness, however, is a relative property,

dependent on the value and abundance of other traits present in the particular community.

Thus, also measuring species' functional uniqueness as *relative to specific communities* may

improve the understanding of changes in functional space and how shifting species

abundances contribute to these changes. Here we first assigned each species an objective

value of functional uniqueness in the species pool using a community containing all species

at equal abundance. We then validated this measure by showing that objectively unique

species were on the whole more unique than others across different uninvaded and invaded

communities (Fig. 5: positive linear relationships for both lines). Next, we found that

objectively non-unique (functionally ordinary) species constituted more of the functional

spaces of invaded communities than uninvaded communities (Fig. 5: lower intercept of the

'invaded' line). We further found that objectively very-unique species constituted less of the

functional spaces of invaded communities than uninvaded communities (Fig. 5: steeper slope

of 'invaded' line). Together, these findings suggest that the *S. invicta* invasion has led to

invaded communities becoming more comprised of a subset of species (winners) sharing trait

values which are common in the species pool, and less comprised of other species (losers)

with trait values that are rare in the species pool. These patterns mirror the decline of

functionally unique species before functionally redundant species observed in other

disturbances (Flynn et al., 2009).

Implications for ecosystem function

The consequences of an invasion will extend to the ecosystem if the affected taxa are also

key modulators of ecosystem functions. Ants are such organisms, and the effects of *S. invicta*

invasion on various ant-modulated ecosystem functions such as predation, nutrient cycling

and bioturbation is a pertinent question to tackle in future research. These effects will hinge

on the particular relationships between ant diversity and ant-modulated ecosystem functions

in the tropical grassland communities of Hong Kong. For instance, if ecosystem functions

mainly respond to the functional identities of the ant communities (i.e., selection effects),

they may be impacted significantly by the altered CWMs of invaded communities.

Alternatively, functional homogenization and the decline of functionally unique species in

invaded communities could impact ecosystem functions driven by functional

complementarity. Ecosystem functions may also respond to both functional identity and

complementarity across different spatial and temporal scales (Isbell et al., 2018). Whatever

the case, the present findings show *S. invicta* invasions in Hong Kong to have significant

impacts on the functional structure of ant communities, which may also affect ecosystem-

wide changes.

Implications for understanding and managing invasions

Using data on the traits of individuals we have shown that an alien invasive species alters – in

a selective, non-random manner – the functional properties of native communities that may

influence ecosystem processes. Crucially, our findings further suggest that such impacts may

unfold in the absence of similar changes in both taxonomic and functional richness. Thus,

assessments exclusively using taxonomic measures of diversity, or indices that only describe

trait variety, may fail to detect various consequences of invasions for the structure and

function of ecological communities. These other ecologically significant consequences of

invasion (e.g., functional clustering and homogenization) can be uncovered by investigating

patterns in the diversity and distribution of traits at the species, community, and landscape

levels. Future work using such comprehensive functional approaches may not only improve

the management of invasive species, but also help to identify other alien species driving

'cryptic impacts' on ecosystem-relevant community structure amid modest taxonomic

change.

Authors' Contributions

582 M.K.L.W., B.G. and O.T.L. designed the study. M.K.L.W. conducted fieldwork, analysed the

583 data and wrote the first draft of the manuscript. All authors contributed substantially to

584 manuscript revisions.

Acknowledgements

587 We are grateful to Brett Morgan for providing access to environmental data, Mac Pierce,

588 Roger Lee and Roy Cheung for field assistance, Carlos Carmona for help with *TPD*, and staff

589 of AEC and WWF Hong Kong for logistical support. This work was supported by a National

590 Geographic Grant (60-16) and a University of Oxford Clarendon Scholarship to M.K.L.W.

References

Ascunce, M. S., Yang, C. C., Oakey, J., Calcaterra, L., Wu, W. J., Shih, C. J., ... &

Shoemaker, D. (2011). Global invasion history of the fire ant Solenopsis invicta. Science,

331, 1066-1068.

Baiser, B., & Lockwood, J. L. (2011). The relationship between functional and taxonomic

homogenization. Global Ecology and Biogeography, 20, 134-144.

Baselga, A., Orme, D., Villeger, S., De Bortoli, J., & Leprieur, F. (2018). betapart:

partitioning beta diversity into turnover and nestedness components. R package version 1.5.1.

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models

using lme4. Journal of Statistical Software, 67, 1-48.

Bradshaw, C. J., Leroy, B., Bellard, C., Roiz, D., Albert, C., Fournier, A., ... & Courchamp,

F. (2016). Massive yet grossly underestimated global costs of invasive insects. Nature

Communications, 7, 12986.

Carmona, C. P., de Bello, F., Mason, N. W., & Lepš, J. (2016). Traits without borders:

integrating functional diversity across scales. Trends in Ecology and Evolution, 31, 382-394.

Carmona, C. P. (2018). TPD: methods for measuring functional diversity based on trait

probability density. R package version 1.0.0.

Castro-Díez, P., Pauchard, A., Traveset, A., & Vilà, M. (2016). Linking the impacts of plant

invasion on community functional structure and ecosystem properties. Journal of Vegetation

Science, 27, 1233-1242.

Coetzee, B. W., & Chown, S. L. (2016). Land-use change promotes avian diversity at the

expense of species with unique traits. Ecology and Evolution, 6(21), 7610-7622.

Des Roches, S., Post, D. M., Turley, N. E., Bailey, J. K., Hendry, A. P., Kinnison, M. T., ... &

Palkovacs, E. P. (2018). The ecological importance of intraspecific variation. Nature Ecology

and Evolution, 2, 57-64.

Eguchi, K., Bui, T. V., & Yamane, S. (2004). A preliminary study on foraging distance and

nesting sites of ants in Indo-Chinese lowland vegetation. Sociobiology, 43, 445-457.

Farji-Brener, A. G., Barrantes, G., & Ruggiero, A. (2004). Environmental rugosity, body size

and access to food: a test of the size-grain hypothesis in tropical litter ants. Oikos, 104, 165-

171.

Flynn, D. F., Gogol-Prokurat, M., Nogeire, T., Molinari, N., Richers, B. T., Lin, B. B., ... &

DeClerck, F. (2009). Loss of functional diversity under land use intensification across

multiple taxa. Ecology Letters, 12, 22-33.

Ford, E. B. (1957). Polymorphism in plants, animals and man. Nature, 180, 1315-1319.

Gagic, V., Bartomeus, I., Jonsson, T., Taylor, A., Winqvist, C., Fischer, C., ... & Tscharntke,

T. (2015). Functional identity and diversity of animals predict ecosystem functioning better

than species-based indices. Proceedings of the Royal Society B: Biological Sciences, 282,

20142620.

González-Suárez, M., Bacher, S., & Jeschke, J. M. (2015). Intraspecific trait variation is

correlated with establishment success of alien mammals. The American Naturalist, 185, 737-

746.

Grevé, M. E., Bláha, S., Teuber, J., Rothmaier, M., & Feldhaar, H. (2019). The effect of

ground surface rugosity on ant running speed is species-specific rather than size dependent.

Insectes Sociaux. doi.org/10.1007/s00040-019-00694-z

Guénard, B., Weiser, M., Gomez, K., Narula, N., & Economo, E.P. (2017). The Global Ant

Biodiversity Informatics (GABI) database: a synthesis of ant species geographic

distributions. Myrmecological News, 24, 83-89.

Human, K. G., & Gordon, D. M. (1997). Effects of Argentine ants on invertebrate

biodiversity in northern California. Conservation Biology, 11, 1242-1248.

Isbell, F., Cowles, J., Dee, L. E., Loreau, M., Reich, P. B., Gonzalez, A., ... & Schmid, B.

(2018). Quantifying effects of biodiversity on ecosystem functioning across times and places.

Ecology Letters, 21, 763-778.

Keller, R. A., Peeters, C., & Beldade, P. (2014). Evolution of thorax architecture in ant castes

highlights trade-off between flight and ground behaviors. Elife, 3, e01539.

Laliberté, E., Legendre, P., & Shipley, B. (2014). FD: measuring functional diversity from

multiple traits, and other tools for functional ecology. R package version 12.0.

Lessard, J. P., Fordyce, J. A., Gotelli, N. J., & Sanders, N. J. (2009). Invasive ants alter the

phylogenetic structure of ant communities. Ecology, 90, 2664-2669.

Lowe, S., Browne, M., Boudjelas, S., & De Poorter, M. (2000). 100 of the world's worst

invasive alien species: a selection from the global invasive species database. Auckland:

Invasive Species Specialist Group.

MacDougall, A. S., Gilbert, B., & Levine, J. M. (2009). Plant invasions and the niche.

Journal of Ecology, 97, 609-615.

McGill, B. J., Enquist, B. J., Weiher, E., & Westoby, M. (2006). Rebuilding community

ecology from functional traits. Trends in Ecology and Evolution, 21, 178-185.

Morgan, B., & Guénard, B. (2019). Local models reveal greater spatial variation than global

grids in an urban mosaic: Hong Kong climate, vegetation, and topography rasters. Earth

System Science Data Discussion. doi.org/10.5194/essd-2018-132 (preprint).

Mouchet, M. A., Villéger, S., Mason, N. W., & Mouillot, D. (2010). Functional diversity

measures: an overview of their redundancy and their ability to discriminate community

assembly rules. Functional Ecology, 24, 867-876.

Mouillot, D., Graham, N. A., Villéger, S., Mason, N. W., & Bellwood, D. R. (2013). A

functional approach reveals community responses to disturbances. Trends in Ecology and

Evolution, 28, 167-177.

Passera, L. (1984). L'organisation sociale des Fourmis. Toulouse: Privat.

Porter, S. D., & Savignano, D. A. (1990). Invasion of polygyne fire ants decimates native

ants and disrupts arthropod community. Ecology, 71, 2095-2106.

Pyšek, P., & Richardson, D. M. (2010). Invasive species, environmental change and

management, and health. Annual Review of Environment and Resources, 35, 25-55.

R Core Team. (2017). R: a language and environment for statistical computing. R.

Richter, A., Keller, R. A., Rosumek, F. B., Economo, E. P., Garcia, F. H., & Beutel, R. G.

(2019). The cephalic anatomy of workers of the ant species *Wasmannia affinis* (Formicidae,

Hymenoptera, Insecta) and its evolutionary implications. Arthropod Structure and

Development, 49, 26-49.

Schofield, S. F., Bishop, T. R., & Parr, C. L. (2016). Morphological characteristics of ant

assemblages (Hymenoptera: Formicidae) differ among contrasting biomes. Myrmecological

News, 23, 129-137.

Silva, R. R., & Brandão, C. R. F. (2010). Morphological patterns and community

organization in leaf-litter ant assemblages. Ecological Monographs, 80, 107-124.

Swenson, N. G. (2014). Functional and phylogenetic ecology in R. New York, NY: Springer.

Tschinkel, W. R. (1988). Colony growth and the ontogeny of worker polymorphism in the

fire ant, *Solenopsis invicta*. Behavioral Ecology and Sociobiology, 22, 103-115.

Tschinkel, W. R. (2006). The fire ants. Cambrdige, MA: Harvard University Press.

Vellend, M. (2016). The theory of ecological communities. New Jersey, NJ: Princeton

University Press.

Venables, W. N., & Ripley, B. D. (2002). Modern applied statistics with S. New York, NY:

Springer.

Villéger, S., Grenouillet, G., & Brosse, S. (2014). Functional homogenization exceeds

taxonomic homogenization among European fish assemblages. Global Ecology and

Biogeography, 23, 1450-1460.

Vinson, S. B. (1997). Invasion of the red imported fire ant (Hymenoptera: Formicidae):

spread, biology, and impact. American Entomologist, 43(1), 23-39.

Wang, L., Xu, Y., Zeng, L., & Lu, Y. (2018). A review of the impact of the red imported fire

ant *Solenopsis invicta* Buren on biodiversity in South China. Journal of Integrative

Agriculture, 17: 60345-60347

Weeks, R. D., Wilson, L. T., & Vinson, S. B. (2004). Resource partitioning among colonies

of polygyne red imported fire ants (Hymenoptera: Formicidae). Environmental Entomology,

33, 1602-1608.

Wickham, H. (2009). ggplot2: elegant graphics for data analysis. New York, NY: Springer-

Verlag.

Wilson, E. O. (2003). *Pheidole* in the New World: a dominant, hyperdiverse ant genus.

Cambridge, MA: Harvard University Press.

Wong, M. K. L., Guénard, B., & Lewis, O. T. (2018). Trait-based ecology of terrestrial

arthropods. Biological Reviews. doi.org/10.1111/brv.12488