Reference plasmid pHXB2_D is an HIV-1 molecular clone that exhibits identical LTRs and a single integration site indicative of an HIV provirus

Alejandro R. Gener[1,2,3,4]§, Brian T. Foley[5], Deborah P. Hyink[2], Paul E. Klotman[2]

[1]Integrative Molecular and Biomedical Sciences Program, Baylor College of Medicine, Houston, Texas, USA
[2]Margaret M. and Albert B. Alkek Department of Medicine, Nephrology, Baylor College of Medicine, Houston, Texas, USA
[3]Department of Genetics, MD Anderson Cancer Center, Houston, Texas, USA
[4]School of Medicine, Universidad Central del Caribe, Bayamón, Puerto Rico, USA
[5]Theoretical Biology and Biophysics Group T-6, Los Alamos National Laboratory, Los Alamos, New Mexico, USA

§Corresponding author: Alejandro R. Gener

One Baylor Plaza
Mail Stop 710
Houston, Texas, 77030, USA
9045715562
gener@bcm.edu ; itspronouncedhenner@gmail.com

Keywords: HIV-1, reagent verification, nanopore DNA sequencing, provirus, plasmid, sequence variability, resequencing

Abstract

Objective: To compare long-read nanopore DNA sequencing (DNA-seq) with short-read sequencing-by-synthesis for sequencing a full-length (e.g., non-deletion, nor reporter) HIV-1 model provirus in plasmid pHXB2_D.

Design: We sequenced pHXB2_D with long- and short-read DNA-seq, evaluating sample variability with resequencing (sequencing and mapping to reference HXB2) and *de novo* viral genome assembly.

Methods: We prepared pHXB2_D for long-read nanopore DNA-seq, varying DNA polymerases Taq (Sigma-Aldrich) and Long Amplicon (LA) Taq (Takara). Nanopore basecallers were compared. After aligning reads to the reference HXB2 to evaluate sample coverage, we looked for variants. We next assembled reads into contigs, followed by finishing and polishing. We hired an external core to sequence-verify pHXB2_D with single-end 150 base-long Illumina reads, after masking sample identity.

Results: We achieved full-coverage (100%) of HXB2 HIV-1 from 5' to 3' long terminal repeats (LTRs), with median per-base coverage of over 9000x in one experiment on a single MinION flow cell. The longest HIV-spanning read to-date was generated, at a length of 11,487 bases, which included full-length HIV-1 and plasmid backbone with flanking host sequences supporting a single HXB2 integration event. We discovered 20 single nucleotide variants in pHXB2_D compared to reference, verified by short-read DNA sequencing.

Conclusions: Nanopore sequencing performed as-expected, even covering full-length HIV. The discovery of variants in a reference plasmid demonstrates the need for sequence verification moving forward, in line with calls from funding agencies for reagent verification. These results

23   illustrate the utility of long-read DNA-seq to advance the study of HIV at single integration site

24   resolution.

Introduction

27   Much of what we know about human acquired immunodeficiency syndrome (AIDS)

28 came after isolating the causative agent – the human immunodeficiency virus type 1 (HIV-1) –

29 and describing the viral genome information content. The HIV-1 isolate HXB2 (also known as

30 HTLV-III and HIV-1$_{LAI}$ or LAV/BRU [1], [2]) was the first full-length replication-competent

31 HIV genome sequenced [3]. Derivative clones commonly called "HXB2" are still used for *in*

32 *vitro* infection assays, including RNA (almost always cDNA [4]) sequencing (**Figure 1A** and

33 **Supplemental Table 1**). Despite the availability of the HXB2 HIV-1 reference sequence [3], no

34 sequence is available for any complete and readily available HXB2 clone.

35   HIV clones were originally made by choosing non-cutter restriction enzymes to digest

36 intact proviral sequences upstream and downstream of unknown integration sites from infected

37 host cells, followed by ligation into an *E. coli* cloning vector (plasmid) (**Figure 1B**), allowing for

38 low-error (but not error-free) propagation [5]. These clones became available before tractable

39 sequencing methods permitted routine sequence verification. As such, it was uncommon to

40 sequence them. While funding agencies now require investigators to include in their proposals

41 plans to validate their key reagents, these funders tend to leave the process up to investigators

42 and may not always follow up on whether a given reagent is ever actually validated (or

43 revalidated between changes of hand). Investigators do not regularly validate their clones, in part

44 because there is no universally accepted standard. Instead, a common practice is to assume a

45 given clone, often kindly gifted from a colleague, is as reported. As such, we often do not truly

46 know what we have been working with for 35+ years.

47   Making sense of the information from HIV sequencing experiments is complicated by

48 many factors, including the cycling that all orterviruses [6] undergo between two major states (as

infectious virion RNA and integrated proviral DNA **Figure 1B**), repetitive viral sequences like

long terminal repeats (LTRs), non-integrated forms [7], rarity of integration events *in vivo*

(reviewed in [8]), and alternative splicing of viral mRNAs [9]. Short-read DNA sequencing

(<150 base pairs (bp) in most reported experiments, but up to 500 bp for either Illumina

sequencing-by-synthesis or <1,000 bp for chain termination sequencing) provides some

information, but analyses require high coverage and/or extensive effort (non-exhaustive

examples [10], [11]). These factors limit the ability to assign variants to specific loci within each

provirus, as well as at the proviral integration site(s) (reviewed in [12]). Despite progress (HIV

DNA) [13], (HIV RNA) [14], [15], researchers have yet to observe the genome of HIV-1 as

complete provirus (integrated DNA) in a single read, hindering locus-specific studies. To this

end, current long-read DNA sequencing clearly surpasses the limitations of read length of

leading next-generation/short-read sequencing platforms. Here we used the MinION sequencer

to sequence HIV-1 plasmid pHXB2_D in a pilot study focusing on coverage acquisition (as

opposed to full-length sequencing), with the goal of evaluating the technology for future

applications.

Methods

This work did not include human or animal subjects. Nanopore libraries for this work were prepared in their entirety by ARG in a Biosafety Level 2 laboratory on main campus at Baylor College of Medicine. Nanopore sequencing was completed between April and May of 2018 as two of several control experiments included in the Student Genomics pilot run (**Supplemental Information**). Short-read sequencing was completed in April 2019.

HIV-1 plasmid

A plasmid, "pHXB2_D" (alternate names pHXB2, pHXB-2D), believed to contain the HIV-1 reference strain HXB2 [16] was acquired from the NIH AIDS Reagent and Reference Program (ARP) via BioServe. pHXB2_D is believed to be a molecular clone (likely a restriction product of HXB2 proviral DNA inserted into an unknown cloning plasmid backbone) from one of the earliest clinical "HXB2" HIV-1 isolates. At the time of this work, it was unknown whether this plasmid was ever sequence-verified before or after the reference sequence for HXB2 was deposited.

HIV-1 reference sequence

The reference sequence of HXB2 is from the National Center for Biotechnology Information (NCBI), Genbank accession number K03455.1. It runs from the beginning of the 5' LTR to the end of the 3' LTR, and is 9,719 bp.

Long-read DNA sequencing

A plasmid containing HXB2 was sequence-verified with long-read nanopore sequencing on a MinION Mk1B (Oxford Nanopore Technologies, Oxford, UK). Unless otherwise noted, reagents (and software) were purchased (or acquired) from Oxford Nanopore. Briefly, stock

plasmid was diluted to 5 ng final amount in ultrapure water (as two samples) and processed with

Rapid PCR Barcoding kit SQK-RPB004 along with 10 other barcoded samples (not discussed

further in this manuscript) following ONT protocol RPB_9059_V1_REVA_08MAR2018

(**Figure 1C**), a public description of which is here: https://store.nanoporetech.com/us/sample-

prep/rapid-pcr-barcoding-kit.html. Two DNA polymerases were evaluated (barcode 10 used

high-fidelity LA (for "long amplicon") Taq (Takara); barcode 11 Taq (Sigma-Aldrich). Libraries

were loaded onto a MinION flow cell version R9.4.1 and a 48-hour sequencing run was

completed with MinKNOW (version 1.10.11). Residual reads from subsequent runs were pooled

for final analyses.

Raw data was basecalled (converted from FAST5 to FASTQ format) with Albacore

version 2.3.4 (older basecaller), Guppy version 2.3.1 (current official at time of work), and

FlipFlop (Guppy development config). Mapping to reference was done with Minimap2 [17] and

BWA-MEM [18], implemented in Galaxy (usegalaxy.org) [19]. Alignments (.bam and .bai files)

were visualized in the Integrative Genomics Viewer [20] unless otherwise noted. For *de novo*

assembly, demultiplexed basecalled reads were fed into Canu version 1.8 [21]. Genome size was

estimated to be 16 Kb from agarose gel of undigested, but naturally degraded linearized

pHXB2_D (data not shown). SnapGene version 4.3.4 was used to manually annotate contigs

from Canu. Blastn (NCBI) was used to identify unknown regions of pHXB2_D. Polishing was

performed on ONT-only assemblies with Medaka (https://github.com/nanoporetech/medaka), in

Galaxy. Medaka models: r941_min_fast_g303, r941_min_high_g303, r941_min_high_g330.

Inference batch size (-b) = 100. The final pHXB2_D assembly and other full-length HIV clones

from the ARP were aligned to the most recent human reference genome (hg38) with Minimap2

in Galaxy with the following parameters: Long assembly to reference mapping (-k19 -w19 -A1 -

B19 -O39,81 -E3,1 -s200 -z200 --min-occ-floor=100).

Statistics

Two-tailed Mann-Whitney U tests were used to compare distributions in long-read data.

P-values are reported over brackets delineating relevant comparisons. Calculations and graphing

were done with GraphPad Prism for macOS version 8.0.2.

Short-read DNA sequencing

pHXB2_D was provided as 35 ul at 63 ng/ul to the Center for Computational &

Integrative Biology DNA Core at Massachusetts General Hospital, an external DNA sequencing

core specializing in high-throughput next generation (short-read) plasmid sequencing and

assembly. Neither HXB2 reference sequence nor pHXB2_D draft assemblies (from this work)

were provided to core staff at the time of sequencing so that testing would remain masked. While

the core's exact library prep is proprietary, multiplexed library prep and 150 single-end Illumina

(ILMN) sequencing were most likely performed on a MiSeq with platform-specific reagents (V2

chemistry, per their website) and barcoding. Data was returned as FASTQ. FASTQC [22] was

used in Galaxy for in-house data quality control, and read lengths were all 142 bp per this tool.

Mapping as above.

Sequence comparisons

We used MAFFT v7.475 [23], [24] to compare the LTR sequences of pHXB2_D and

HXB2. For cladistics, we used BLAST at HIV-DB

(https://www.hiv.lanl.gov/content/sequence/BASIC_BLAST/basic_blast.html) to find other

HXB2-like genomes. The top 50 BLAST hits included many sequences pNL43 clones. pNL4-3

is an artificial recombinant of the NY5 clone with the HXB2 clone. We then made a multi-

sequence alignment with the final pHXB2_D assembly, the top BAST hits, and the HIV-1 M

group subtype reference set using GeneCutter

(https://www.hiv.lanl.gov/content/sequence/GENE_CUTTER/cutter.html), and built the

maximum likelihood tree using IQ-tree

(https://www.hiv.lanl.gov/content/sequence/IQTREE/iqtree.html).

Results

Viewing mapped data in IGV, the long reads (median read length >2000 bp, **Figures 1E**) from both ONT experiments clearly covered each LTR (**Figure 1F, Supplemental Figures 1**, **2**), while shorter reads collapsed into one of either LTR (**Figures 1F, Supplemental Figures 2D,2E**). This was also seen when long reads were shorter than LTRs (<600 bp). Mappers BWA-MEM and Minimap2 were chosen based on their ability to handle long and short reads. Other mappers were not evaluated. BWA-MEM mapped more ambiguously, piling partially mapped reads between each LTR; Minimap2 mapped with higher fidelity to reference without splitting reads. Coverage as sequencing depth was higher and more even from the higher-fidelity LA Taq library (**Supplemental Figure 1**).

We counted 20 single nucleotide variants (SNVs) in this reference clone of HXB2 (**Table 1, Supplemental Table 3, Supplemental Figure 2E**). These mismatches were seen in all Canu assemblies (**Supplemental Figure 3A,3B**), verified in IGV and/or SnapGene, and were orthogonally verified by short-read sequencing performed by the external core given masked samples (**Supplemental Figure 2E**). These mismatches represent a ~0.21% divergence from reference HXB2 K03455.1 (20/9719), which was assumed to have perfect identity (0% divergence). Transitions were more common (14/20) (**Table 1**), coinciding with a previous report of increased transitions over transversions in infection models, because transversions are more likely to be deleterious to viral replication (i.e.: to cause protein-coding changes) [25]. Indeed, almost half (9/20) of the observed SNVs occurred in protein-coding regions, even though 92% of HXB2 is coding (791/9719). Of those 9 SNVs in protein-coding regions, 4 caused non-synonymous mutations. One of those occurs in a region overlapping both gag and pol regions, however only pol exhibited a non-synonymous change from valine to isoleucine in p6, at

position 2259 relative to HXB2. Other non-synonymous variants occurred at 4609 (in p31

integrase, arginine to lysine), 7823 (in ASP antisense protein, glycine to arginine), and 9253 (in

nef, isoleucine to valine). 11/20 SNVs were in LTRs (see **Supplemental Figure 2** for counting

based on mapping); 8/20 of these would have been missed with mapping-only variant calling or

consensus. The longest HIV-mapping read (**Figure 2**) phased 16/20 SNVs (failed at sites

2,8,10,12, **Table 1**).

We assembled the previously undefined plasmid pHXB2_D (**Supplemental Figure**

3A,3B). Canu's final output was a set of contiguous DNA sequences (contigs) as FASTA files. A

consequence of assembling plasmid sequences with this tool was partial redundancy at contig

ends (**Supplemental Figure 3C**). Manual end-trimming of contigs was performed in SnapGene

based on an estimated length of 16 kilobases. Top blastn hits from barcode 10/LA Taq pHXB2

basecalled with FlipFlop were as follows: for the main backbone (with origin of replication and

antibiotic selection cassette for cloning), shuttle vector pTB101-CM DNA, complete sequence

(based on pBR322), from 4352-8340; for the upstream element (relative to 5' LTR), Homo

sapiens chromosome 3 clone RP11-83E7 map 3p, complete sequence from 58,052 to 59,165; for

the downstream element, cloning vector pNHG-CapNM from 10,204 to 11,666. Other identified

elements included Enterobacteria phage SP6 (the SP6 promoter, per SnapGene's "Detect

common features"), complete sequence from 39,683 to 39,966. Identities of query to HXB2 and

hits were all approximately 99%. The MGH CCIB DNA Core's proprietary *de novo* UltraCycler

v1.0 assembler (Brian Seed and Huajun Wang, unpublished) was able to assemble both 5' and 3'

LTRs with short-read data only but may have collapsed SNVs into an artificial single consensus.

Long-read mapping and assembly (and polished assemblies) supported a single HIV-1 HXB2_D

haplotype (**Supplemental Figure 3,5**); a final LTR-phased and annotated assembly leveraging

short and long reads is provided as pHXB2_D Genbank:MW079479 (embargoed until

publication). Importantly, for pHXB2_D, each LTR was identical, which is distinct from the

current HXB2 (K03455.1) (**Figure 3**). To determine whether pHXB2_D was an isolated provirus

(as opposed to a cDNA clone), the pHXB2_D assembly was aligned to the current human

reference hg38, returning a single complete insertion site on 3p24.3 (**Figure 4, Supplemental**

Table 2). We sought to put our pHXB2_D assembly into context of other HXB2-like references

available (**Figure 5**). pHXB2_D (red) clusters closely with HXB2 reference (K03455) and

related clone sequences (green). pNL4-3 clones in blue. The LTR-masked HIV-spanning

segment of pHXB2_D is most homologous to B.FR.1983.DM461230 and B.FR.1983.CS793683,

which are identical except for areas in nef and a GFP insertion (verified by blastn). This finding

suggests they were from the same stock. HIV-1 M group subtype reference set (HIV Sequence

Database) was added to put HXB2s and pNL4-3 clones into perspective. HXB2 (believed to be a

complete isolate) and NL4-3 (synthetic clone based on two early isolates) are examples of HIV

type 1 (HIV-1), group M, subgroup B.

As previously reported [26], per-read variability in ONT data was higher near

homopolymers (runs of the same base) (**Supplemental Figure 4A**). For the datasets generated in

the present study, homopolymers were counted and classified as continuous (unbroken run of a

given nucleobase) vs. discontinuous (broken run of a given nucleobase) (**Supplemental Figure**

4B,4D,4F,4H). A/T (2 hydrogen bonds; 2H) and G/C (3 hydrogen bonds; 3H) were evaluated.

Because runs longer than 4 or 5 were rare in these datasets, it was impossible to evaluate longer

homopolymers. A simple calculation $\text{Abs}(\Delta)=\text{Abs}(\#\text{homopolymers}_{reference}$ -

$\#\text{homopolymers}_{assembly})$ helped to evaluate the performance of basecallers, such that better

basecallers had smaller $\text{Abs}(\Delta)$ (**Supplemental Figure 4C,4E,4G,4I,4K**). At the level of

consensus (made from sequences mapped to reference HXB2), homopolymers contributed few,

if any, obvious errors. A special case of homopolymer, dimer runs, was noted to cause persistent

errors regardless of ONT basecaller (**Supplemental Figure 4J,4K**). While dips occurred at

certain points near homopolymers, the consensus did not change much at the sequencing depth

used in this study for either barcoded pHXB2_D samples (**Supplemental Figures 1,2,3**).

Another interpretation is that homopolymers tend to seem truncated with ONT, with more reads

in support of shorter homopolymers. Canu assemblies showed basecaller-dependent variability

(**Supplemental Table 3**). That said, newer basecallers tended to produce fewer and smaller per-

read truncations. Assemblies without polishing did not correct all homopolymer truncations

(**Supplemental Figure 3A**). Polishing assemblies tended to correct these toward the final

pHXB2_D assembly (**Supplemental Figure 3B,5**). Data from polished ONT-only assemblies

and short-read sequencing do not support the truncations (gaps relative to reference) suggested

by unpolished ONT-only assemblies, representing a known current limitation of ONT. These are

not the same as the 20 SNVs supported by BOTH long- and short-read sequencing performed in

this study. The ratio of per-read deletions to per-read insertions (DEL/INS) was much higher for

SNVs occurring at homopolymers and near the same base, and this difference was maintained

between all basecallers used (**Supplemental Figure 4L**). These changes created more

problematic (longer) homopolymers.

Discussion

This work represents the first instance of complete and unambiguous sequencing of HIV-

1 provirus as plasmid and contributed to the identification of single nucleotide variants which

may not have been easily determined using other sequencing modalities, illustrating the

importance of validating molecular reagents in their entirety, and with complementary

approaches. Nanopore sequencing surpassed the read length limitations of traditional sequencing

modalities used for HIV such as Sanger sequencing and sequencing-by-synthesis by at least two

orders of magnitude. Other long-read DNA sequencing technologies such as PacBio's zero-mode

waveguide DNA sequencing were not evaluated in this work, but in principle would be

interchangeable for nanopore sequencing. Paired-end sequencing (as either DNA-seq or RNA-

seq) was not evaluated in this work, but has shown promise phasing LTRs in our hands [27]–

[29].

First complete pass over all HIV information in reference plasmid pHXB2_D

HIV provirus is believed to occur naturally as one or a few copies of reverse-transcribed

DNA forms integrated into the host nuclear genome. Depending on where integration occurs,

local GC or AT content might cause problems for detecting integrants with PCR. HIV also has

conserved transitions from areas of higher GC content (~60%) to content approximating average

human GC content (~40%). To limit PCR sequencing bias and to accommodate for the potential

heterogeneity of HIV sequences, we fractionated whole sample directly (as opposed to PCR-

barcoding select amplicons) with tagmentation provided in the Rapid PCR-Barcoding kit (ONT).

Tagmentation uses transposon-mediated cleavage and ligation of barcode adapters for later PCR

amplification. A consequence of this fractionation was a distribution of reads (**Figure 1E**)

shorter than longer reads reported elsewhere for ONT experiments [30]. Based on this

distribution and the level of coverage, it was expected that HIV might be covered from end to

end, but this would have been exceptional. That said, an example is presented here (**Figure 2**).

The provirus status of pHXB2_D is supported by recovery of both upstream and downstream

homology arms which map to a single human integration site.

Long reads enable LTR discrimination and HIV haplotype definition

We created 6 assemblies for pHXB2_D from ONT-only data (**Supplemental Figure 3**),

each with a common set of 20 SNVs (11 in LTRs), and 1 final assembly (a single HIV-1

HXB2_D haplotype) leveraging long- and short-read data. The external core's *de novo* assembly

pipeline identified the same 20 SNVs, and variants in the LTRs were supported by ONT

unambiguously. That the core's assembler was able to phase LTR variants in these samples may

have been because the samples had high amounts of the same upstream and downstream

sequences because of coming from one plasmid. The core's assembler thus may have had

additional sequencing information at the edges of HXB2, helping it to map deeper into each

LTR. This approach would likely fail in samples with multiple integrations (as in various animal

models of HIV disease [31]), which have unknown upstream and downstream sequences, or in

samples from natural human infection, which is well known to exhibit multiple pseudo-random

integration sites between cells [32], [33], but with mostly single integration events per cell [8].

Inverse PCR (iPCR) is an alternative method [34] with its own issues (e.g., PCR biases, HIV

concatemers, host repeats). While current PCR reagents have extended the range of what can be

seen with iPCR, current approaches are likewise limited by long DNA extraction methods,

sample amount, and remain to be optimized. If coverage is sufficient ($\geq$10 reads in non-

homopolymers and non-dimer runs), long-read sequencing can provide linked variant

information to individual integration sites. Identical 5' and '3 LTRs (**Figure 3**) in the context of

a single integration event (**Figure 4**) support this integrant being a *bona fide* provirus [35]. Other

proviruses also had identical LTR pairs (**Supplemental Table 2**). Technical limitations such as

PCR errors before earlier sequencing may explain the variability in the HXB2 reference LTRs.

These were sequenced at a time before paired-end 150 or long-read DNA-seq were available to

phase LTRs, raising the possibility that these LTRs were incorrectly annotated by depositors

assuming identity and copy-and-pasting the sequence of one LTR for both without being able to

unambiguously resolve each LTR.

Mutations in a reference HIV-1 plasmid illustrate the need for reagent verification

Up until 2020, HIV had been the most studied human pathogen, but reagents are not

routinely re(verified). The pHXB2_D sequenced was allegedly a reference plasmid, with

unknown divergence between the published reference HXB2. Three independent experiments

(two long-read with PCR-barcoded libraries made with regular and long-amplicon Taq master

mixes, one short-read) yielded at least 20 single nucleotide variants in pHXB2_D which differed

from the HXB2 reference sequence (**Table 1, Supplemental Figure 2**), which were also

concordant across the three basecallers used (**Supplemental Table 3**), and are therefore not PCR

errors. By leveraging long reads with the MinION, we were able to find mutations in highly

repetitive LTRs relative to HXB2 Genbank:K03455.1 which are often assumed (but until now

never proven) to be identical (**Table 1**, **Figure 1**, **Supplemental Figures 1, 2E**), as well as

mutations in protein-coding regions (**Table 1**). We were also able to confirm that the backbone

of this plasmid is from pSP62 [16], a pBR322 derivative with the SP6 promoter [36], aiding in

the continued use of this important reagent, and illustrating the need of full-length reagent

validation moving forward. We suggest that all clinical reagents (e.g., vectors) be sequence-

verified at the level of single-molecule sequencing as standard quality control to protect against

sample heterogeneity.

Improvement in ONT basecallers over time

Albacore, Guppy, and FlipFlop basecallers were compared. Each produced reads of

similar length distributions (relative to polymerase used), while Guppy and FlipFlop produced

improved and best performance relative to quality score distributions (**Figure 1D**). Interestingly,

while read length distributions were affected by fidelity of polymerases evaluated in this work,

mean quality distributions were not. This is important because of the differences in cost between

higher fidelity Taq and classic Taq enzymes. That said, higher fidelity LA Taq produced much

higher coverage compared to Taq (**Supplemental Figure 1**). In consideration of library prep,

choice of enzyme used should be based on the desired read-length distribution and coverage.

Regarding read mapping, the increase in mean quality score between these basecallers improved

overall mapping, in part by facilitating demultiplexing, resulting in approximately ~10%

increases number of reads in barcoded libraries before mapping (shift in reads from unclassified

to a given barcode). FlipFlop tended to handle homopolymers better than previous basecallers

(**Supplemental Figures 4,5**). Homopolymers in HXB2 tended to exhibit apparent deletions near

5' ends of homopolymers (upstream due to technical artifact from mapping), but because

consensus is near conserved (example, at least 80% of base in called read set is identical to

reference), and because short-read data lacks INDELS at these sites, it is unlikely that any of

these homopolymer deletions are real in these experiments. Dimer runs – stretches of repeating

2-mers (pronounced "two-mers") – proved challenging regardless of basecaller. Mapping as

above may be used to aid in manually calling these when they occur. Albacore is currently

deprecated, and current versions of Guppy now incorporate a version of FlipFlop called Guppy

High-ACcuracy (HAC). Guppy HAC and subsequence versions were not evaluated in this work.

Polishing is becoming standard practice for processing assemblies from ONT data because it

redresses most homopolymer errors propagated into long-read-only assemblies. The best

manually finished and polished contig had 1 error out of 16,722 bases, illustrating the utility of

ONT hardware when paired with burgeoning software.

Conclusions

HIV informatics, the study of HIV sequence information, has been limited by the common assumption that sequence fidelity exists between reference genomes available in sequence databases and similarly named HIV clones. Modern DNA sequencing methods, such as long- and short-read sequencing, are available to redress this issue. Long-read sequencing fills in gaps left behind by short-read interrogation of HIV-1. Current limitations of the approaches used in the present work to study HIV are 1.) the cost of long-read sequencing, regardless of platform, compared to the cheaper short reads from sequencing-by-synthesis, 2.) long DNA extraction methods in diseased tissue (Gener, unpublished), and 3.) the lower per-base accuracy (low-mid 90's with ONT vs. 98-99% with ILMN or newer PacBio HiFi), including difficulty near homopolymers and dimer runs (**Supplemental Figure 4**). A nontrivial but redressable limitation is availability of personnel trained to prepare sequencing libraries, to run sequencing, and to analyze results. As the price of long-read sequencing decreases, hardware and software used in basecalling and library protocols improve, and with the advent of more user-friendly tools, the cost of obtaining usable data from long reads will become negligible compared to the ability to answer historically intractable questions. This work raises the possibility of being able to detect at least some recombination events, in a reference-free manner requiring only the comparison of LTRs from the same integrants. While pHXB2_D and other HIV reference clones had identical LTR pairs, this remains to be tested in other clones, since other clones were generated with shorter sequencing methods. Future work will include optimizing DNA extraction protocols with the goal of capturing higher-coverage fuller glimpses of each HIV proviral integration site in *in vivo* HIV models and patient samples. This work has broad implications for all cells infected by both integrating and non-integrating viruses, and for the characterization of targeted regions in

348   the genome which may be recalcitrant to previous sequencing methods. Long-read sequencing is

349   an important emerging tool defining the post-scaffold genomic era, allowing for the

350   characterization of anatomical landmarks of hosts and pathogens at the genomic scale.

Disclaimer

Erratum: Preprint version 1 of this work [37] incorrectly cited the Integrated Genome Browser for work that was completed with the Integrative Genomics Viewer. Apologies for the mistake.

Funding

This work was funded in part by institutional support from Baylor College of Medicine; the Human Genome Sequencing Center at Baylor College of Medicine; private funding by Bob Ostendorf, CEO of East Coast Oils, Inc., Jacksonville, Florida; ARG's own private funding, including Student Genomics (manuscripts in prep). Compute resources from the Computational and Integrative Biomedical Research Center at BCM ("sphere" cluster managed by Dr. Steven Ludtke) and the Department of Molecular and Human Genetics at BCM ("taco" cluster managed by Mr. Tanner Beck and Dr. Charles Lin) greatly facilitated the completion of this work. ARG has also received the PFLAG of Jacksonville scholarship for multiple years.

Competing interests

ARG received travel bursaries from Oxford Nanopore Technologies (ONT). The present work was completed independently of ONT. Other authors declare no conflicts of interest.

Authors' contributions

ARG conceived of this project, performed experiments, analyzed results, and drafted the manuscript. BTF, DPH and PK discussed data and edited the manuscript. ARG and PK provided funding.

Acknowledgements

As part of a summer bioinformatics internship in the Paul E. Klotman Laboratory at

Baylor College of Medicine, Akash Naik supervised by ARG performed *in silico* mapping

analyses/experiments, generated and/or aided in the synthesis of **Supplemental Figure 3**, and

assisted in writing relevant portions, discussing, and editing this manuscript. During a second

summer internship with American Physician Scientists Association Virtual Summer Research

Program, the following students were supervised by ARG helped to create **Figure 1A** and

Supplemental Table 1: Yini Liang, Kirk Niekamp, Maliha Jeba, Delmarie M. Rivera

Rodríguez. Orthogonal sequence verification was performed as a service by staff at the Center

for Computational & Integrative Biology DNA Core at Massachusetts General Hospital, Boston,

MA, USA.

We would like to thank the staff at the DNA Core for their exceptional services,

including expert analyses and rapid turnaround time. We would like to thank Drs. Steven

Richards, Qingchang Meng and the staff of the Human Genome Sequencing Center Research

(HGSC) and Development (R&D) team for their earlier support in nanopore adoption. We would

like to thank the team at Oxford Nanopore Technologies for their timely improvements and

continued R&D. I would also like to thank Ms. Taneasha Monique Washington (current) and

former members of the Paul E. Klotman lab, Dr. Gokul C. Das and Alexander Batista. I would

like to thank Dr. Alana Canupp and the late Dr. Jim Maruniak for their early interest in my

scientific development, and for the passion that they show in everything that they do.

Available additional files

Albacore basecalled barcode 10

Guppy basecalled barcode 10

FlipFlop basecalled barcode 10

Albacore basecalled barcode 11

Guppy basecalled barcode 11

FlipFlop basecalled barcode 11

Minimap2 and BWA-MEM alignments (.bam and .bai)

Clipboards from points of interest (verified SNVs; n=20)

.dna files of contigs (n=6)

MGH data (raw + contig)

Supplemental Tables

Supplemental Figures

References

[1]   F. Barré-Sinoussi *et al.*, "Isolation of a T-lymphotropic retrovirus from a patient at risk for acquired immune deficiency syndrome (AIDS).," *Science*, vol. 220, no. 4599, pp. 868–871, May 1983, doi: 10.1126/science.6189183.

[2]   S. Wain-Hobson *et al.*, "LAV revisited: origins of the early HIV-1 isolates from Institut Pasteur.," *Science*, vol. 252, no. 5008, pp. 961–965, May 1991, doi: 10.1126/science.2035026.

[3]   L. Ratner *et al.*, "Complete nucleotide sequence of the AIDS virus, HTLV-III.," *Nature*, vol. 313, no. 6000, pp. 277–284, Jan. 1985, doi: 10.1038/313277a0.

[4]   A. R. Gener and J. T. Kimata, "Full-coverage native RNA sequencing of HIV-1 viruses," *bioRxiv*, p. 845610, Jan. 2019, doi: 10.1101/845610.

[5]   G. M. Shaw, B. H. Hahn, S. K. Arya, J. E. Groopman, R. C. Gallo, and F. Wong-Staal, "Molecular characterization of human T-cell leukemia (lymphotropic) virus type III in the acquired immune deficiency syndrome.," *Science*, vol. 226, no. 4679, pp. 1165–1171, Dec. 1984, doi: 10.1126/science.6095449.

[6]   M. Krupovic *et al.*, "Ortervirales: New Virus Order Unifying Five Families of Reverse-Transcribing Viruses," *J. Virol.*, vol. 92, no. 12, pp. e00515-18, May 2018, doi: 10.1128/JVI.00515-18.

[7]   E. H. Graf *et al.*, "Elite suppressors harbor low levels of integrated HIV DNA and high levels of 2-LTR circular HIV DNA compared to HIV+ patients on and off HAART," *PLoS Pathog.*, vol. 7, no. 2, 2011, doi: 10.1371/journal.ppat.1001300.

[8] Y. Ito *et al.*, "Number of infection events per cell during HIV-1 cell-free infection," *Sci.*

Rep., vol. 7, no. 1, p. 6559, 2017, doi: 10.1038/s41598-017-03954-9.

[9] I. Cuesta, A. Mari, A. Ocampo, C. Miralles, S. Pérez-castro, and M. M. Thomson,

"Sequence Analysis of In Vivo -Expressed HIV-1 Spliced RNAs Reveals the Usage of

New and Unusual Splice Sites by Viruses of Different Subtypes," pp. 1–24, 2016, doi:

10.1371/journal.pone.0158525.

[10] C. Wymant *et al.*, "Easy and accurate reconstruction of whole HIV genomes from short-

read sequence data with shiver," *Virus Evol.*, vol. 4, no. 1, pp. 1–13, 2018, doi:

10.1093/ve/vey007.

[11] K. M. Bruner *et al.*, "A quantitative approach for measuring the reservoir of latent HIV-1

proviruses," *Nature*, vol. 566, no. 7742, pp. 120–125, 2019, doi: 10.1038/s41586-019-

0898-8.

[12] M. R. Pinzone and U. O'Doherty, "Measuring integrated HIV DNA ex vivo and in vitro

provides insights about how reservoirs are formed and maintained," *Retrovirology*, vol.

15, no. 1, pp. 1–12, 2018, doi: 10.1186/s12977-018-0396-3.

[13] K. B. Einkauf *et al.*, "Intact HIV-1 proviruses accumulate at distinct chromosomal

positions during prolonged antiretroviral therapy Find the latest version : Intact HIV-1

proviruses accumulate at distinct chromosomal positions during prolonged antiretroviral

therapy," vol. 129, no. 3, pp. 988–998, 2019.

[14] D. Bonsall *et al.*, "THAA0101 - HIV genotyping and phylogenetics in the HPTN 071

(PopART) study: Validation of a high-throughput sequencing assay for viral load

quantification, genotyping, resistance testing and high-resolution transmission

networking," in *22nd International AIDS Conference (AIDS2018)*, 2018, p. Oral Abstract.

[15] A. N. Banin *et al.*, "Development of a Versatile, Near Full Genome Amplification and

Sequencing Approach for a Broad Variety of HIV-1 Group M Variants," *Viruses*, vol. 11,

no. 4, p. 317, Apr. 2019, doi: 10.3390/v11040317.

[16] A. G. Fisher, E. Collalti, L. Ratner, R. C. Gallo, and F. Wong-Staal, "A molecular clone of

HTLV-III with biological activity," *Nature*, vol. 316, no. 6025, pp. 262–265, 1985, doi:

10.1038/316262a0.

[17] H. Li, "Minimap2: pairwise alignment for nucleotide sequences," *Bioinformatics*, vol. 34,

no. 18, pp. 3094–3100, May 2018, doi: 10.1093/bioinformatics/bty191.

[18] H. Li and R. Durbin, "Fast and accurate long-read alignment with Burrows – Wheeler

transform," vol. 26, no. 5, pp. 589–595, 2010, doi: 10.1093/bioinformatics/btp698.

[19] E. Afgan *et al.*, "The Galaxy platform for accessible, reproducible and collaborative

biomedical analyses: 2016 update," *Nucleic Acids Res.*, vol. 44, no. W1, pp. W3–W10,

2016, doi: 10.1093/nar/gkw343.

[20] J. T. Robinson *et al.*, "Integrative genomics viewer," *Nat Biotechnol*, vol. 29, no. 1, pp.

24–26, 2011, doi: 10.1038/nbt0111-24.

[21] B. P. Walenz, S. Koren, N. H. Bergman, A. M. Phillippy, J. R. Miller, and K. Berlin,

"Canu: scalable and accurate long-read assembly via adaptive k -mer weighting and repeat

separation," *Genome Res.*, vol. 27, no. 5, pp. 722–736, 2017, doi: 10.1101/gr.215087.116.

[22] S. Andrews, "FastQC A Quality Control tool for High Throughput Sequence Data."

470 [23] K. Katoh and D. M. Standley, "MAFFT multiple sequence alignment software version 7:

471    Improvements in performance and usability," *Mol. Biol. Evol.*, vol. 30, no. 4, pp. 772–780,

472    2013, doi: 10.1093/molbev/mst010.

473 [24] K. Katoh, J. Rozewicki, and K. D. Yamada, "MAFFT online service: multiple sequence

474    alignment, interactive sequence choice and visualization," *Brief. Bioinform.*, vol. 20, no. 4,

475    pp. 1160–1166, Sep. 2017, doi: 10.1093/bib/bbx108.

476 [25] D. M. Lyons and A. S. Lauring, "Evidence for the Selective Basis of Transition-to-

477    Transversion Substitution Bias in Two RNA Viruses," *Mol. Biol. Evol.*, vol. 34, no. 12,

478    pp. 3205–3215, 2017, doi: 10.1093/molbev/msx251.

479 [26] N. J. Loman, J. Quick, and J. T. Simpson, "A complete bacterial genome assembled de

480    novo using only nanopore sequencing data," *Nat. Methods*, vol. 12, no. 8, pp. 733–735,

481    2015, doi: 10.1038/nmeth.3444.

482 [27] A. Gener *et al.*, "PEA0011 - Insights from HIV-1 transgene insertions in the murine

483    model of HIV-associated nephropathy," in *23rd International AIDS Conference*

484    *(AIDS2020)*, 2020, vol. ePoster.

485 [28] A. R. Gener *et al.*, "P39 - Insights from comprehensive transcript models of HIV-1," in

486    *Genome Informatics 2020*, 2020, p. ePoster.

487 [29] A. R. Gener, T. Washington, D. Hyink, and P. Klotman, "3264 - The Multiple HIV-1

488    Transgenes in the Murine Model of HIV-Associated Nephropathy Fail to Segregate as

489    Expected," in *American Society of Human Genetics Annual Meeting*, 2020, p. ePoster.

490 [30] A. Payne, N. Holmes, V. Rakyan, and M. Loose, "Whale watching with BulkVis: A

491 graphical viewer for Oxford Nanopore bulk fast5 files," *bioRxiv*, p. 312256, Jan. 2018,

492 doi: 10.1101/312256.

493 [31] P. Rosenstiel, A. Gharavi, V. D'Agati, and P. Klotman, "Transgenic and infectious animal

494   models of HIV-associated nephropathy.," *J. Am. Soc. Nephrol.*, vol. 20, no. 11, pp. 2296–

495   304, 2009, doi: 10.1681/ASN.2008121230.

496 [32] M. Kvaratskhelia, A. Sharma, R. C. Larue, E. Serrao, and A. Engelman, "Molecular

497   mechanisms of retroviral integration site selection.," *Nucleic Acids Res.*, vol. 42, no. 16,

498   pp. gku769-, 2014, doi: 10.1093/nar/gku769.

499 [33] B. Marini *et al.*, "Nuclear architecture dictates HIV-1 integration site selection," *Nature*,

500   vol. 521, pp. 227–233, 2015, doi: 10.1038/nature14226.

501 [34] H. Ochman, A. S. Gerber, and D. L. Hartl, "Genetic applications of an inverse polymerase

502   chain reaction," *Genetics*, vol. 120, no. 3, pp. 621–623, 1988.

503 [35] W.-S. Hu and S. H. Hughes, "HIV-1 Reverse Transcription," *Cold Spring Harb. Perspect.*

504   *Med.* , vol. 2, no. 10, Oct. 2012, doi: 10.1101/cshperspect.a006882.

505 [36] M. R. Green, T. Maniatis, and D. A. Melton, "Human beta-globin pre-mRNA synthesized

506   in vitro is accurately spliced in Xenopus oocyte nuclei.," *Cell*, vol. 32, no. 3, pp. 681–

507   694, Mar. 1983, doi: 10.1016/0092-8674(83)90054-5.

508 [37] A. R. Gener, "Full-coverage sequencing of HIV-1 provirus from a reference plasmid,"

509   *bioRxiv*, p. 611848, Jan. 2019, doi: 10.1101/611848.

510 [38] A. Adachi *et al.*, "Production of acquired immunodeficiency syndrome-associated

511   retrovirus in human and nonhuman cells transfected with an infectious molecular clone.,"

J. Virol., vol. 59, no. 2, pp. 284–91, 1986.

[39]   B. Lucic *et al.*, "Spatially clustered loci with multiple enhancers are frequent targets of

HIV-1," *bioRxiv*, 2018.

[40]   W. J. Kent *et al.*, "The Human Genome Browser at UCSC," *Genome Res.* , vol. 12, no. 6,

pp. 996–1006, Jun. 2002, doi: 10.1101/gr.229102.

[41]   Y. Peng, H. C. M. Leung, S. M. Yiu, and F. Y. L. Chin, "IDBA – A Practical Iterative de

Bruijn Graph De Novo Assembler BT - Research in Computational Molecular Biology,"

2010, pp. 426–440.

[42]   P. J. A. Cock, B. A. Grüning, K. Paszkiewicz, and L. Pritchard, "Galaxy tools and

workflows for sequence analysis with applications in molecular plant pathology," *PeerJ*,

vol. 1, p. e167, 2013, doi: 10.7717/peerj.167.

[43]   C. B, W. T, and S. S, "Genome sequence assembly using trace signals and additional

sequence information.," *Comput. Sci. Biol. Proc. Ger. Conf. Bioinforma.*, vol. 99, pp. 45–

56.

[44]   A. Bankevich *et al.*, "SPAdes: A New Genome Assembly Algorithm and Its Applications

to Single-Cell Sequencing," *J. Comput. Biol.*, vol. 19, no. 5, pp. 455–477, Apr. 2012, doi:

10.1089/cmb.2012.0021.

[45]   G. Cuccuru *et al.*, "Orione, a web-based framework for NGS analysis in microbiology,"

Bioinformatics, vol. 30, no. 13, pp. 1928–1929, Jul. 2014, doi:

10.1093/bioinformatics/btu135.

[46]   R. L. Warren, G. G. Sutton, S. J. M. Jones, and R. A. Holt, "Assembling millions of short

DNA sequences using SSAKE," *Bioinformatics*, vol. 23, no. 4, pp. 500–501, Feb. 2007,

doi: 10.1093/bioinformatics/btl629.

Tables

Table 1: Summary of pHXB2 sample divergence from reference HXB2.

Site	Position	Change	Substitution Class	Change	Mutation Class (Syn/Non /Stop)	Homopolymer-adjacent?	Same as neighbor?	LANL Feature	Subfeature	Frame
1	24	C>A	transversion	NA	NA	yes	yes	5'LTR	U3	NA
2	108	A>G	transition	NA	NA	yes	yes	5'LTR	U3	NA
3	164	G>T	transversion	NA	NA	yes	no	5'LTR	U3	NA
4	168	T>G	transversion	NA	NA	yes	yes	5'LTR	U3	NA
5	176	A>G	transition	NA	NA	yes	yes	5'LTR	U3	NA
6	182	C>T	transition	NA	NA	yes	no	5'LTR	U3	NA
7	227	A>G	transition	NA	NA	yes	yes	5'LTR	U3	NA
8	291	A>G	transition	NA	NA	no	no	5'LTR	U3	NA
9	333	C>T	transition	NA	NA	no	no	5'LTR	U3	NA
10	654	C>T	transition	NA	NA	no	no	None	None	NA
11	1659	aaG>aaA	transition	None	Syn	yes	yes	gag	p24, p55	gag frame 1
12	2259	gag:agG>agA pol:Gtc>Atc	transition	gag:Arg>Arg pol:Val>Ile	Syn/Non	no	no	gagpol	p6	gag frame 1 pol frame 3
13	2927	aaG>aaA	transition	None	Syn	yes	yes	pol	p51 RT	pol frame 3
14	3812	ccC>ccT	transition	None	Syn	yes	yes	pol	p51 RT	pol frame 3
15	4574	acT>acA	transversion	None	Syn	no	no	pol	p31 IN	pol frame 3
16	4596	Ggt>Agt	transition	None	Syn	yes	no	pol	p31 IN	pol frame 3

17	4609	aGg>aAg	transition	Arg>Lys	Non	yes	yes	pol	p31 IN	pol frame 3
18	7823	gcC>gcG Ggc>Cgc	transversion	ASP:Gly>Arg	Syn/Non	no	no	gp41	RRE, also ASP	gp41 frame 3, ASP -2
19	9253	Ata>Gta	transition	Ile>val	Non	no	yes	nef/3'LTR	also U3	nef frame 1
20	9418	C>T	transition	NA	NA	no	no	3'LTR	U3	NA

Coverage numbers vary by input (albacore, guppy, FlipFlop basecalled FASTQ) and mapping

method (Minimap2 vs. BWA-MEM). This information is provided as Supplemental Digital

Content. Base-1 (first base is numbered 1, 2nd 2, etc.), relative to HXB2, Genbank:K03455.1.

Changed base represented as upper-case. Annotated as codon if in protein-coding region. No

deletions or insertions were predicted from manual inspection or supported by short-read

sequencing. Abbreviations, ASP: antisense protein, RRE: rev-response element, NA: not

applicable. Syn: synonymous mutation. Non: non-synonymous mutation. Stop: stop codon/non-

sense mutation. LTR: long terminal repeat. RT: reverse transcriptase. IN: integrase. LANL: Los

Alamos National Laboratory HIV Sequence Database. Data from three separate sequencing

experiments on the same plasmid sample support these 20 sites. Note site 1-8 variants in 5'LTR

have been previously reported (LANL), albethey ambiguously. These may also be incorrectly

annotated as variants in nef.

Figures

Figure 1: HIV information in pHXB2_D is recovered by long-read sequencing and

mapping.

Figure 1A: HXB2 is still a commonly used resource. It is the reference HIV-1 genome, derived

from one of the earliest clinical isolates. While older HIV samples are occasionally rediscovered,

they are not made routinely available to researchers. All public HIV-1 RNA-seq datasets were

obtained from the NCBI SRA with the following search phrase: "HIV-1" AND "RNA-seq".

Metadata from these 2527 runs (number current as of 7/21/2020) were used to make a pie chart

summary.

Figure 1B: HIV information comes from three main sources: proviruses (HIV sandwiched

between two assumedly identical full-length long terminal repeats (LTRs)), unspliced HIV

mRNAs (also known as viral genomes) starting from the transcription start site and ending in the

3' LTR [4], and engineered proviruses recovered in their entirety or stitched together from

multiple isolates like NL4-3 [38].

Figure 1C: ONT library prep pipeline. Tagmentation cleaves double-stranded DNA, ligating

barcoded PCR adapters (magenta). PCR-adapted DNA may be amplified. After amplification

and cleanup, ONT sequencing adapters (green) are ligated. Barcoded samples may be pooled and

sequenced.

Figure 1D: Newer basecallers increase read mean quality. Median (big dash) and quartiles (little

dash). Effect of enzyme version was not statistically significant.

Figure 1E: Read stats with different callers/aligners. Median (big dash) and quartiles (little dash).

Read lengths increase with higher fidelity Taq.

Figure 1F: Sequencing coverage with long- vs. short-read single-end 150 bp (trimmed to 142 bp)

DNA sequencing. Long-read sequencing covers ambiguously mappable areas missed by short-

read in HXB2 reference Genbank:K03455.1 (**Supplemental Figure 2D,2E**), but at the expense

of accuracy near homopolymers longer than about 4 nucleobases (**Supplemental Figure 4**).

Short-read mapping fails at repetitive elements longer than their read lengths (**Supplemental**

Figure 2D,2E). Long read Minimap2 settings: map-ont -k15. Short read Minimap2 settings:

Short reads without splicing (-k21 -w11 --sr -F800 -A2 -B8 -O12,32 -E2,1 -r50 -p.5 -N20 -

f1000,5000 -n2 -m20 -s40 -g200 -2K50m --heap-sort=yes --secondary=no) (sr).

Figure 2: Longest read containing complete full-length HIV-1 reference HXB2

The 5th longest read in the barcode 10 set (read ID 6fbf0205-5195-460e-8e28-930db50e5d79)

contained full-length HIV-1. Query (full read) blastn against HIV (taxid:11676) returned 92.95%

identity to HIV-1, complete genome (Genbank:AF033819.3). Limiting query to HXB2 (red)

blastn against Nucleotide collection nr/nt returned 100% coverage and 93.02% identity to HIV-1

HXB2. This read was 11,487 bases long, with mean quality score 11.984396. Basecalled using

Guppy 2.3.1 with FlipFlop config.

Figure 3: pHXB2_D has identical LTRs, resolving likely errors in HXB2 (K03455.1)

```
598   CLUSTAL format alignment by MAFFT (v7.475)
599
600
601   K03455.1_5'LTR   tggaagggctaattcactcccaacgaagacaagatatccttgatctgtggatctaccaca
602   pHXB2_D_5'LTR    tggaagggctaattcactcccaaagaagacaagatatccttgatctgtggatctaccaca
603   pHXB2_D_3'LTR    tggaagggctaattcactcccaaagaagacaagatatccttgatctgtggatctaccaca
604   K03455.1_3'LTR   tggaagggctaattcactcccaaagaagacaagatatccttgatctgtggatctaccaca
605                    ********************** **************************************
606
607   K03455.1_5'LTR   cacaaggctacttccctgattagcagaactacacaccagggccagggatcagatatccac
608   pHXB2_D_5'LTR    cacaaggctacttccctgattagcagaactacacaccagggccaggggtcagatatccac
609   pHXB2_D_3'LTR    cacaaggctacttccctgattagcagaactacacaccagggccaggggtcagatatccac
610   K03455.1_3'LTR   cacaaggctacttccctgattagcagaactacacaccagggccaggggtcagatatccac
611                    *******************************************.************
612
613   K03455.1_5'LTR   tgacctttggatggtgctacaagctagtaccagttgagccagagaagttagaagaagcca
614   pHXB2_D_5'LTR    tgacctttggatggtgctacaagctagtaccagttgagccagataaggtagaagaggcca
615   pHXB2_D_3'LTR    tgacctttggatggtgctacaagctagtaccagttgagccagataaggtagaagaggcca
616   K03455.1_3'LTR   tgacctttggatggtgctacaagctagtaccagttgagccagataagatagaagaggcca
617                    ********************************************* *** *******.****
618
619   K03455.1_5'LTR   acaaaggagagaacaccagcttgttacaccctgtgagcctgcatggaatggatgacccgg
620   pHXB2_D_5'LTR    ataaaggagagaacaccagcttgttacaccctgtgagcctgcatgggatggatgacccgg
621   pHXB2_D_3'LTR    ataaaggagagaacaccagcttgttacaccctgtgagcctgcatgggatggatgacccgg
622   K03455.1_3'LTR   ataaaggagagaacaccagcttgttacaccctgtgagcctgcatgggatggatgacccgg
623                    *.*********************************************.*************
624
625   K03455.1_5'LTR   agagagaagtgttagagtggaggtttgacagccgcctagcatttcatcacatggcccgag
626   pHXB2_D_5'LTR    agagagaagtgttagagtggaggtttgacagccgcctagcatttcatcacgtggcccgag
627   pHXB2_D_3'LTR    agagagaagtgttagagtggaggtttgacagccgcctagcatttcatcacgtggcccgag
628   K03455.1_3'LTR   agagagaagtgttagagtggaggtttgacagccgcctagcatttcatcacgtggcccgag
629                    **************************************************.*********
630
631   K03455.1_5'LTR   agctgcatccggagtacttcaagaactgctgacatcgagcttgctacaagggactttccg
632   pHXB2_D_5'LTR    agctgcatccggagtacttcaagaactgctgatatcgagcttgctacaagggactttccg
633   pHXB2_D_3'LTR    agctgcatccggagtacttcaagaactgctgatatcgagcttgctacaagggactttccg
634   K03455.1_3'LTR   agctgcatccggagtacttcaagaactgctgacatcgagcttgctacaagggactttccg
635                    ********************************.***************************
636
637   K03455.1_5'LTR   ctggggactttccagggaggcgtggcctgggcgggactggggagtggcgagccctcagat
638   pHXB2_D_5'LTR    ctggggactttccagggaggcgtggcctgggcgggactggggagtggcgagccctcagat
```

```
639    pHXB2_D_3'LTR    ctggggactttccagggaggcgtggcctgggcgggactggggagtggcgagccctcagat
640    K03455.1_3'LTR   ctggggactttccagggaggcgtggcctgggcgggactggggagtggcgagccctcagat
641                     ************************************************************

643    K03455.1_5'LTR   cctgcatataagcagctgcttttttgcctgtactgggtctctctggttagaccagatctga
644    pHXB2_D_5'LTR    cctgcatataagcagctgcttttttgcctgtactgggtctctctggttagaccagatctga
645    pHXB2_D_3'LTR    cctgcatataagcagctgcttttttgcctgtactgggtctctctggttagaccagatctga
646    K03455.1_3'LTR   cctgcatataagcagctgcttttttgcctgtactgggtctctctggttagaccagatctga
647                     ************************************************************

649    K03455.1_5'LTR   gcctgggagctctctggctaactagggaacccactgcttaagcctcaataaagcttgcct
650    pHXB2_D_5'LTR    gcctgggagctctctggctaactagggaacccactgcttaagcctcaataaagcttgcct
651    pHXB2_D_3'LTR    gcctgggagctctctggctaactagggaacccactgcttaagcctcaataaagcttgcct
652    K03455.1_3'LTR   gcctgggagctctctggctaactagggaacccactgcttaagcctcaataaagcttgcct
653                     ************************************************************

655    K03455.1_5'LTR   tgagtgcttcaagtagtgtgtgcccgtctgttgtgtgactctggtaactagagatccctc
656    pHXB2_D_5'LTR    tgagtgcttcaagtagtgtgtgcccgtctgttgtgtgactctggtaactagagatccctc
657    pHXB2_D_3'LTR    tgagtgcttcaagtagtgtgtgcccgtctgttgtgtgactctggtaactagagatccctc
658    K03455.1_3'LTR   tgagtgcttcaagtagtgtgtgcccgtctgttgtgtgactctggtaactagagatccctc
659                     ************************************************************

661    K03455.1_5'LTR   agacccttttagtcagtgtggaaaatctctagca
662    pHXB2_D_5'LTR    agacccttttagtcagtgtggaaaatctctagca
663    pHXB2_D_3'LTR    agacccttttagtcagtgtggaaaatctctagca
664    K03455.1_3'LTR   agacccttttagtcagtgtggaaaatctctagca
665                     **********************************
```

668 **Figure 4: HXB2 integration site**

pHXB2_D's, and therefore HXB2's, integration site is unambiguously singular (falls outside of

annotated repeat), and in the same orientation (minus strand relative to hg38) as target gene

TBC1D5. Alignment quality is 60 for both homology arms (**Supplemental Table 2**). Features

captured by homology arms in pHXB2_D and other clones verified as proviruses in the present

study are consistent with HIV-1 integration behavior [39]. Visualized in UCSC Genome Browser

[40].

Figure 5: pHXB2_D provenance and top 50 neighbors

Supplemental Information

Data exploration with long- and short-read mapping

To assemble pHXB2_D, we used tried the following short read assemblers on short-read data from the external core: IDBA [41], MIRA [42], [43], SPAdes [44], and SSAKE [45], [46]. These were chosen as a convenience because they were already stably implemented in Galaxy (specifically usegalaxy.eu). Of these, SSAKE produced discontinuous assemblies with default parameters. The discontinuous contigs did however map to the core's assembly (not shown).

Enabling STEM outreach

This work was performed as two control experiments with identically prepared libraries for a STEM outreach initiative, Student Genomics (Gener, et al., manuscript in prep). Given the constraints of the Student Genomics pilot, a rapid sequencing kit with tagmentation (explained below) with PCR barcoding was used to pool samples for ONT sequencing, with the consequence of fragmenting plasmid DNA more than what would have been ideal for capturing full-length HIV. That said, these controls could have been just as easily replaced by any samples/experiments benefiting from long-read sequencing at moderate-to-high coverage.

Supplemental Tables

Supplemental Table 1: HXB2 is still a common HIV clone.

See **Supplemental Digital Content.**

See also **Figure 1A**.

Supplemental Table 2: HIV provirus clones

See **Supplemental Digital Content.**

Of the HIV clones available through ARP, the table represents the only validated proviruses with

both upstream and downstream homology arms mapping to the same integration sites. pNL4-3 is

included as a known chimera with two integration half-sites. Other clones were made with

cDNA cloning, usually TA cloning (per ARP entries). Note: Reference hg38. Aligner: minimap2

with "Long Assembly" mapping settings. All homology arms had Alignment quality = 60.

Upstream = host plus strand; independent of integration orientation. Coordinates reported from

UCSC. ARP = NIH AIDS Reagent and Reference Program. IS = integration site.

Supplemental Table 3: Variation in assemblies at the feature level.

	Mismatches						Gaps (INDEL)					
	Taq			LA Taq			Taq			LA Taq		
	Albacore	Guppy	FlipFlop	Albacore	Guppy	FlipFlop	Albacore	Guppy	FlipFlop	Albacore	Guppy	FlipFlop
5' LTR	NA	9	9	9	9	9	NA	NA	0	2	0	0
gag	2	2	2	2	2	2	12	10	9	9	8	8
5' LTR+ψ	10	10	NA	NA	NA	NA	5	2	NA	NA	NA	NA
pol	7	6	6	6	6	6	26	22	9	18	11	10
vif	0	0	0	0	0	0	3	3	2	3	1	1
vpr	0	0	0	0	0	0	1	1	1	0	0	0
tat	2	1	1	1	1	1	10	6	3	7	4	5
rev	2	1	1	1	1	1	10	7	4	7	4	5
vpu	1	0	0	0	0	0	0	0	0	0	0	0
gp160	2	1	1	1	1	1	11	7	4	8	4	5
nef	1	1	1	1	1	1	3	2	2	3	2	1
3' LTR	2	2	NA	NA	NA	NA	2	0	NA	NA	NA	NA
nef+3' LTR	NA	2	2	2	2	2	NA	2	2	5	2	1
HXB2	22	20	20	20	20	20	61	46	28	47	27	25
Downstream bridge	0	0	0	0	0	0	4	5	1	2	1	1
pBR322-related	0	0	0	0	0	0	19	19	13	18	19	15
Upstream bridge	2	2	3	2	2	2	8	6	5	7	7	3

718   Assembled with Canu. NA denotes features which may not have matched exactly, but which

719   were collapsed with adjacent features to facilitate counting. Variants called manually by

720   mapping assemblies over HXB2 features with SnapGene.

723 **Supplemental Figures**

725   **Supplemental Figure 1A: Unbiased nanopore DNA sequencing coverage over HXB2**

726   **depends on DNA polymerase and mapper. ONT basecaller = Albacore (worst).**

727

728

Supplemental Figure 1B: Unbiased nanopore DNA sequencing coverage over HXB2

depends on DNA polymerase and mapper. ONT basecaller = Guppy.

Supplemental Figure 1C: Unbiased nanopore DNA sequencing coverage over HXB2

depends on DNA polymerase and mapper. ONT basecaller = FlipFlop (best).

Top two Coverage and Alignment panels from barcoded library 10 (bar10 = LA Taq). Bottom

two from Barcode 11 (bar11 = Taq). Minimap2 and BWA-MEM were used to map reads

basecalled with Albacore (worst), Guppy, or FlipFlop (best) to HXB2. Color-coding: Red below

genome scale marks 20 SNVs across the HIV segment of pHXB2_D. Purple is an insertion in a

given read relative to reference. White is either a deletion in a given read or space between two

aligned reads. Gray in alignment field means base same as reference, and in coverage field

means major allele is at least 95% the same as reference. Per-read "insertions" and "deletions"

do not necessarily represent true insertions or deletions actually present in the sample, because

each read is likely an imperfect independent observation. Automated assembly followed by

manual consensus building converts these overlapping reads into approximations of the ground

truth. "Unbiased" refers to not amplifying a given region (e.g., pol) before ligating ONT

sequencing adapters. In the present approach, the tagmentation process randomly cuts DNA,

creating ~2000 bp pieces. Tagmented DNA is then amplified based on tagmentation adapters.

Supplemental Figure 2A: HIV single nucleotide variants (SNVs) in pHXB2_D. ONT

basecaller = Albacore (worst).

755 **Supplemental Figure 2B: HIV single nucleotide variants (SNVs) in pHXB2_D. ONT**

756 **basecaller = Guppy.**

757

758

759 **Supplemental Figure 2C: HIV single nucleotide variants (SNVs) in pHXB2_D. ONT**

760 **basecaller = FlipFlop (best).**

763   **Supplemental Figure 2D: HIV single nucleotide variants (SNVs) in pHXB2_D, long vs.**

764   **short reads (HIV genome)**

767   Supplemental Figure 2E: HIV single nucleotide variants (SNVs) in pHXB2_D, long vs.

768   short reads (20 SNV-focused)

771 Supplemental Figure 2A: HIV single nucleotide variants (SNVs) in pHXB2_D. ONT basecaller

772 = Albacore (worst). Gray indicates per-base consensus accuracy ≥ 80%. These alignments are

773 the noisiest (less gray and most divergent from reference) between Supplemental Figures 2A,

774 2B, and 2C.

775 Supplemental Figure 2B: HIV single nucleotide variants (SNVs) in pHXB2_D. ONT basecaller

776 = Guppy.

777 Supplemental Figure 2C: HIV single nucleotide variants (SNVs) in pHXB2_D. ONT basecaller

778 = FlipFlop (best). These alignments are the least noisy (most gray and like reference) between

779 Supplemental Figures 2A, 2B, and 2C.

780 Supplemental Figure 2D: HIV single nucleotide variants (SNVs) in pHXB2_D, long vs. short

781 reads (HIV genome). Long reads outperform short reads at HIV-1 LTRs. ONT basecaller =

782 FlipFlop. Short read as single-end 150, clipped to 142, provided by external core. Mappers =

783 Minimap2 (better), BWA-MEM.

784 Supplemental Figure 2E: HIV single nucleotide variants (SNVs) in pHXB2_D, long vs. short

785 reads (SNV-focused).

Supplemental Figure 3A: Assembling pHXB2_D from long reads only, varying basecaller

and polymerase.

Each pane (n=6) summarizes the results of contig curation. Divergence from reference decreases

with newer basecallers, and with long amplicon DNA polymerase (Sigma-Aldrich Taq vs. LA

Taq by Takara). Errors in assembly occurred at homopolymers (most often deletions not visible

at this resolution; see **Supplemental Figure 5**), dimer or trimer runs. bar10 = LA Taq library.

bar11 = Taq library. pHXB2_D Genbank:MW079479. Best contigs presented, manually curated

to match pHXB2_D coordinates. Note LTRs (beginning and terminal 634 bp of red bar) are

resolved in almost all assemblies. See **Supplemental Table 3** for differences between assemblies

and the reference (left red). Plasmid backbone (right) differences are not reported.

Supplemental Figure 3B: ONT errors corrected by polishing ONT-only assemblies.

Assemblies polished with Medaka (ONT). Top: pHXB2_D genome. Bottom: HIV-only segment.

The best polished assembly had one error in the entire plasmid (1 error out of 16,722 bases), with

a corresponding consensus accuracy of 99.99402%. This happened to be in HIV segment (HIV-1

between position 1 and 9719; 1 error out of 9719 bases), with corresponding accuracy of

99.989711%. Note the conserved 52 bp gap in the backbone of pHXB2_D was redundant

sequence included in the short-read assembly from the core. It was not supported by long-read

data, and therefor was validated as a technical artifact from the core's pipeline. Reference: short-

read assembly. LTRs (beginning and terminal 634 bp of red bar) are resolved in polished

assemblies.

Supplemental Figure 3C: Mappability of long reads over contigs during assembly quality

control

Coverage depends on context. Abrupt changes in coverage from terminal regions of HXB2

(**Figure 1F, Supplemental Figures 1 and 2)** were artifacts from supplying mappers with an

HIV reference without a plasmid backbone. Long reads from barcode 10 (LA Taq) mapped with

minimap2 [17] and "reference" contig from assembly (Canu v1.8) with basecalled data

(FlipFlop). Stripes in this figure are not SNVs. They represent technical variability at

homopolymers. Assemblies were manually curated to start with 5' LTR in the sense orientation,

leaving the plasmid backbone on the left. Because there were not real insertions in the HIV

segment, the HIV coordinates are the same as HXB2 (both 9719 bp long). Compare with 52 bp

technical artifact from the core's short read assembly in **Supplemental Figure 3A, 3B**.

Supplemental Figure 4: Homopolymers and dimer runs are ONT artifacts in unpolished

assemblies.

Supplemental Figure 4A: A set of homopolymer tracks from HXB2 plasmid. Alignments with

BWA-MEM shown from FlipFlop (top) and Albacore (bottom) basecalled reads. Mapping is

pre-assembly.

Supplemental Figure 4B: Continuous 2H counts in unpolished assemblies. 2H = A or T

homodimers.

Supplemental Figure 4C: Continuous 2H Absolute Difference.

Supplemental Figure 4D: Discontinuous 2H counts in unpolished assemblies.

Supplemental Figure 4E: Discontinuous 2H Absolute Difference.

Supplemental Figure 4F: Continuous 3H counts in unpolished assemblies. 3H = C or G

homodimers.

Supplemental Figure 4G: Continuous 3H Absolute Difference.

Supplemental Figure 4H: Discontinuous 3H counts in unpolished assemblies.

Supplemental Figure 4I: Discontinuous 3H Absolute Difference.

Supplemental Figure 4J: Dimer run counts in unpolished assemblies.

Supplemental Figure 4K: Dimer run Absolute Difference. Dimer runs as pairs are the most

problematic, with runs as triplets being resolvable by ONT.

Supplemental Figure 4L: The ratio of deletions to insertions is higher at mismatches both

adjacent to homopolymers and similar to neighbor bases. Box plot shows median ("x" is mean)

and quartile ranges. Y-axis is ratio. HPA: homopolymer-adjacent. ==: same as neighbor base. Δ:

different than neighbor base. Higher coverage (above ~10) usually makes up for current error

profile. Above true for Albacore, Guppy, and FlipFlop.

Supplemental Figure 5: Assembly partially resolved homopolymers, which are improved

by polishing

Top: Six ONT-only assemblies. Bottom: polished ONT-only assemblies, varying Medaka

models. Deletions at 5' of G homopolymers were not corrected, regardless of basecaller or Taq

isoform. Note that polishing was not performed. IGV window is Linear:4,781-4,820. Bottom:

polishing canu assemblies with medaka abrogated most ONT artifacts. Best medaka setting

tested: r941_min_high_g330.

Word count:

Abstract: **245**/250

96 brackets – 46 references = 50 citations

Main text: 3935-57 (subheadings) – 50 citations = **3828**

Tables: 1

Figures: 5