HIV-1 controllers exhibit an enhanced antiviral state characterized by overexpression of p21 and MCPIP1

Suwellen S. D. de Azevedo[1][*], Marcelo Ribeiro-Alves[2], Fernanda H. Côrtes[1], Edson Delatorre[3],
Brenda Hoagland[2], Beatriz Grinsztejn[2], Valdilea G. Veloso[2], Mariza G. Morgado[1], Thiago
Moreno L. Souza[4][,][5], and Gonzalo Bello[1]

[1] Laboratório de AIDS & Imunologia Molecular. Instituto Oswaldo Cruz – IOC, FIOCRUZ. Rio de
Janeiro, Brazil.

[2] Laboratório de Pesquisa Clínica em DST-AIDS. Instituto Nacional de Infectologia Evandro Chagas
- INI, FIOCRUZ, Rio de Janeiro, Brazil.

[3] Laboratório de Genética Molecular de Microrganismos. Instituto Oswaldo Cruz – IOC, FIOCRUZ.
Rio de Janeiro, Brazil.

[4] National Institute for Science and Technology on Innovation on Diseases of Neglected Populations
(INCT/IDPN), Center for Technological Development in Health – CDTS, FIOCRUZ, Rio de Janeiro,
Brazil.

[5] Laboratório de Imunofarmacologia. Instituto Oswaldo Cruz – IOC, FIOCRUZ. Rio de Janeiro,
Brazil.

*Correspondence:

Suwellen S.D. de Azevedo

suwellen@ioc.fiocruz.br/suwellendias@gmail.com

Keywords: HIV-1 controllers, restriction factors, p21, MCPIP1, immune activation,

Words count abstract: 323

Words count text: 2.669

Figures: 3

Table: 1

Abstract

Several cellular proteins have been identified as antiviral restriction factors (RF), one of the arms of
the innate immune response. Among the RF targeting the human immunodeficiency virus type 1
(HIV-1), the cyclin-dependent kinase inhibitor p21 and the monocyte chemotactic protein-induced
protein 1 (MCPIP1) are recognized as multifunctional modulators of cell proliferation, immune
homeostasis and innate response to viral infections. Recently, MCPIP1 overexpression was
associated with an increase of both p21 transcripts and proteins levels in a model of renal carcinoma
cells. Here, we explored the potential associations between MCPIP1 and p21 expression, as well as
with cellular activation in individuals who naturally control HIV-1 replication (HIC), sustaining
undetectable (elite controllers – EC) or low (viremic controllers – VC) viral loads. We found a
selective upregulation of p21 and MCPIP1 mRNA levels in PBMC from HIC compared with both
ART-suppressed and HIV–negative control groups ($P \leq 0.02$) and a strong positive correlation ($r \geq$
0.57; $P \leq 0.014$) between expressions of both RF independently of the VL, treatment condition and
HIV status. The mRNA levels of p21, but not of MCPIP1, were positively correlated with activated
CD4$^+$ T cells levels in HIC and EC ($r \geq 0.53$; $P \leq 0.017$). In relation to the monocyte activation, the
mRNA levels of both p21 ($r = 0.74$; $P = 0.005$) and MCPIP1 ($r = 0.58$; $P = 0.040$) were positively
correlated with plasmatic levels of sCD14 only in EC. Multivariate analysis confirmed the
association among p21, MCPIP1 mRNA levels, and frequency of activated CD4$^+$ T cells. These data
show for the first time the simultaneous overexpression of p21 and MCPIP1 host RF in the setting of
natural suppression of HIV-1 replication *in vivo*. The positive correlation between p21 and MCPIP1
transcripts supports a common regulatory pathway connecting these RF and a possible synergistic
effect on HIV-1 replication control. Pharmacological manipulation of these innate RF may open
novel therapeutic perspectives to prevent HIV-1 replication and disease progression.

1   Introduction

Among the individuals infected by the human immunodeficiency virus type 1 (HIV-1), a rare group
called HIV controllers (HIC) suppress viral replication in absence of antiretroviral therapy,
maintaining RNA viral loads (VL) undetectable (< 50-80 copies/mL; elite controllers, EC) or at low
levels (50-2,000 copies/mL; viremic controllers, VC). Natural control of HIV-1 replication is
probably a multifactorial feature that involves different combinations of host and/or viral factors (1).
Several host restriction factors (RF), which constitute components of the innate immune response,
have been shown to limit HIV-1 replication *in vitro* (2). Importantly, a few RF are elevated in
peripheral blood mononuclear cells (PBMC) of HIC compared to antiretroviral (ART)-suppressed
and/or HIV-uninfected individuals (3–10), suggesting a role of host cell RF in the natural control of
HIV-1 replication *in vivo*.

The cyclin-dependent kinase (CDK) inhibitor p21 is a multifunctional modulator of a broad range of
biological processes like cell proliferation, that also indirectly limit the HIV-1 replication by blocking
the biosynthesis of dNTPs required to viral reverse transcription and by inhibiting the CDK9 activity
required by HIV-1 for viral mRNA transcription (6). The protein p21 has been shown to inhibit HIV-
1 replication *in vitro* in various cellular systems (6,11–15) and is further expressed at high levels *ex*
vivo in CD4[+] T cells from HICs (4,6,9,16). Another study revealed that p21 mRNA levels correlated
with CD4[+] T cell activation in EC, but not in other HIV-infected groups, suggesting that the
inducibility of this RF to immune activation is a singular characteristic of EC and may contribute to
this special phenotype (3).

The monocyte chemotactic protein-induced protein 1 (MCPIP1) is a protein with a critical role in the
regulation of the inflammatory response and immune homeostasis that also block HIV-1 replication
by promoting the viral mRNA degradation through its RNase activity (17,18). MCPIP1 was pointed
as an important RF of HIV-1 replication *in vitro* in quiescent CD4[+] T cells (17), but its role in
controlling HIV-1 infection *in vivo* is undefined. Interestingly, a recent study revealed that MCPIP1
overexpression in renal carcinoma cells (Caki-1 cells) reduces the cellular growth by increasing the
levels of p21 transcripts, along with other proteins involved in cell cycle progression/arrest (19).
According to this study, MCPIP1 expression triggers activation of p21 by down-modulation of
damage-specific DNA binding protein 1 (DDB1) and upregulation of chromatin licensing and DNA
replication factor 1 (CDT1) (19).

This evidence prompted us to ask whether the expression of MCPIP1 and p21 could be positively
correlated in the setting of natural control of HIV-1 infection. To test this hypothesis, we quantified
the *in vivo* expression of these and others antiviral cell host RF mRNAs in PBMC from HIC, ART-
suppressed and HIV-uninfected individuals. We further explored the potential relationship between
antiviral host RF expression and levels of systemic cellular activation in HIC.

2   Methods

2.1   Study Subjects

We analyzed a cohort of 21 HIC subjects followed-up at the Instituto Nacional de Infectologia
Evandro Chagas (INI) in Rio de Janeiro, Brazil. All HIC maintained RNA VL of < 2,000 copies/ml
without antiretroviral therapy for at least five years and were subdivided in two sub-groups: EC (*n* =
13) when most (≥ 70%) plasma VL determinations were below the detection limit (< 40-80 copies/ml

depending on the VL method used along patient follow-up), and VC ($n = 8$) when most ($\geq 70\%$) VL
determinations were between 41-2,000 copies/ml. Virological and immunological characteristics of
these subjects were described in detail in previous studies (20,21). Two groups of ART-suppressed
subjects (ART, $n = 8$) and healthy HIV-1-uninfected subjects (NEG, $n = 10$) were used as controls.

2.2   mRNA gene-expression analysis

Total RNA was extracted from 1 x 10^7 PBMC using RNeasy mini kit (Qiagen, Hilden, North Rhine-
Westphalia, Germany) in which buffer RLT was supplemented with β-mercaptoethanol and displaced
on-column DNase treatment using a Qiagen RNase-Free DNase Set (Qiagen, Hilden, North Rhine-
Westphalia, Germany) according to manufacturer's instruction. Total RNA yield and quality were
determined using NanoDrop® 8000 spectrophotometer and an Agilent® 2100 Bioanalyzer. Only
samples with an RNA integrity number (RIN) greater than 8.0 were used. Purified RNA (1 µg) was
reverse-transcribed to cDNA using RT^2 First Strand Kit (Qiagen, Hilden, North Rhine-Westphalia,
Germany). The cDNA was mixed with RT^2 SYBR Green/ROX qPCR Master Mix (Qiagen, Hilden,
North Rhine-Westphalia, Germany) and the mixture was added into customized RT^{2}RNA PCR Array
(Qiagen, Hilden, North Rhine-Westphalia, Germany) to measure the mRNA expression of cellular
RF (MCPIP1, p21, SAMHD1, APOBEC3G, Tetherin, IFITM1, Mx1, Mx2, Schlafen 11
[SLFN11]and IFITM3) and three housekeeping genes (GAPDH, β-actin, and RNase-P), according to
manufacturer's instructions. Values of the crossing point at the maximum of the second derivative of
the four-parameters fitted sigmoid curve second derivative, Cp, was determined for each sample. The
efficiency of each amplification reaction was calculated as the ratio between the fluorescence of the
cycle of quantification and fluorescence of the cycle immediately preceding that. Genes used in the
normalization among samples were selected by the geNorm method (22). Data were expressed as
fold-changes in mRNA abundance calculated as the normalized gene expression in any test sample
divided by the mean normalized gene expression in the control HIV-negative group.

2.3   CD4+ T cell and monocyte activation analyses

We used data of CD4+ T cell and monocyte activation obtained in a previous study conducted by our
group including these patients (21), in which plasma levels of soluble CD14 (sCD14) were
determined by ELISA-sCD14 Quantikine assay (R&D Systems Minneapolis, MN) according to the
manufacturer's protocol and surface expression of comnined HLA-DR and CD38 on CD4+ T cells
was analyzed by flow cytometry.

2.4   Data analyses

The comparisons of mean log-fold changes in mRNA abundance were performed by either t-tests or
one-way ANOVA nonparametric permutation tests (B = 1,000 permutations), followed by pair-wise
comparisons with Holm-Bonferroni adjustment (23), for two or more groups respectively. Spearman
coefficient was used for correlation analyses. A first-order log-Normal multiple regression analysis
was fitted to model p21 gene expression as a function of MCPIP1 gene expression, CD4+ T cell
activation (HLA-DR+CD38+), and HIC groups (EC and VC). The threshold for statistical
significance was set to $P < 0.05$. Data were analyzed with R software (version 3.5.2) (24).

3   Results

Twenty-nine HIV-1 positive (21 HIC and 8 ART-supressed), and 10 HIV-negative individuals were
included in this cross-sectional study. Most HIV-positive (59%) and HIV-negative (60%) individuals
were females and all individuals displayed CD4+ T cells counts above 500 cells/µl (Table 1).

Analyses of host RF expression revealed a significant upregulation of p21 and MCPIP1 in HIC
(Figure 1). The p21 mRNA was up-regulated in PBMCs from HIC compared to cells from both
ART-suppressed (1.63-fold increase; P = 0.003) and HIV-negative (1.55-fold increase; P = 0.003)
subjects (Figure 1A). Significant overexpression of the MCPIP1 mRNA in PBMC from HIC
compared to ART-suppressed (1.68-fold increase; P = 0.003) and HIV-negative (1.37-fold increase;
P = 0.02) individuals was also observed (Figure 1B). Expression of p21 mRNA was significantly (P
< 0.025) elevated in PBMCs from both EC and VC compared to control groups (Figure 1C).
Expression of MCPIP1 mRNA was also elevated in EC and VC subsets compared with ART-
suppressed and HIV-negative groups, although the difference with HIV-negative group did not reach
statistical significance (Figure 1D). With respect to EC and VC, p21 and MCPIP1 mRNA expression
levels were equally significant. Of note, the expression of other antiretroviral RF shows no statistical
differences between the HIC and control groups, with exception of IFITM1 that was significantly
elevated (1.15-fold increase; P = 0.03) in HIC in comparison to the HIV-negative group
(Supplementary Figure S1). Moreover, we found a significant positive correlation between p21 and
MCPIP1 mRNA expression (r ≥ 0.57; P ≤ 0.014) independently of the VL, treatment condition and
HIV status (Figure 2). With respect to other RF, only a modest, but significant, negative correlation
between p21/MCPIP1with APOBEC3G mRNA in HIC was observed (Supplementary Figure S2).

To explore the potential relationship of p21 or MCPIP1 expression with immune activation and
inflammation, we measured the frequency of CD4+HLA-DR+CD38+ T cells and plasma sCD14 levels
(monocyte activation) in our cohort. Frequencies of activated CD4+ T cell populations in VC and
ART-suppressed subjects were similar; nevertheless, higher than in EC (P < 0.0001) and HIV-
negative (P = 0.0002) individuals (Figure 3A). The median concentration of sCD14 in plasma was
not significantly different across the groups (Figure 3B). The mRNA levels of p21 were positively
associated with activated CD4+ T cells levels in HIC (r = 0.53; P = 0.016) and EC (r = 0.68; P =
0.017) (Figure 3C). No significant correlations between mRNA levels of MCPIP1 and CD4+ T cell
activation were observed for HIC or EC subsets (Figure 3D). Levels of sCD14 were positively
correlated with both p21 (r = 0.74; P = 0.005) and MCPIP1 (r = 0.58; P = 0.04) mRNA levels only in
the EC subset (Figure 3E and F). No significant correlations between mRNA levels of p21/MCPIP1
and CD4+ T cell activation or sCD14 levels were observed when ART-suppressed and HIV-negative
individuals were included (Supplementary Figures S3). Multivariate analysis showed that the
upregulation of MCPIP1 was positively associated with the increase of p21 expression in HIC (1.44-
fold increase; P = 0.0035) (Supplementary Figure S4A). The frequency of activated CD4+ T cells
also was positively associated with the increase of p21 expression in both EC and VC (1.48-fold
increase; P = 0.0116), although this increase of the p21 expression was down-regulated by the
increase of activated CD4+ T cells in VC when compared to EC (1.30-fold decrease by an increase of
1% CD4+HLA-DR+CD38+ T cells; P = 0.0284) (Supplementary Figure S4B). Overall, the model was
highly significant (P = 0.003) and could explain as much as 70% (R^2 = 0.492) of p21 expression.

4   Discussion

In this study, we observed that p21 and MCPIP1 mRNA expression were significantly increased in
PBMC of HIC compared to cells of HIV-negative and -positive/ART-suppressed individuals. The
elevated expression of p21 in PBMC of HIC had already been previously described (3,4,6,9,16).
Originally, this is the first study to show overexpression of MCPIP1 alongside with p21 in these
individuals. The mRNA levels of p21 and MCPIP1 were positively correlated in HIC as well as in
HIV-positive and –negative individuals. This supports a coordinated expression of these cellular
genes in different settings, consistent with what has been shown for a renal carcinoma cell line (19).

Other host RF, such as SAMHD1, SLFN11, and Mx1, previously found to be overexpressed in CD4$^+$
T cells from HIC (3,7,8), were not up-regulated in PBMC from our HIC cohort, which may reflect
differences in the cellular populations analyzed (8). A recent study (10), however, described
upregulation of tetherin and SLFN11 in PBMC from HIC, supporting that the contrasting findings
across studies might also reflect inter-cohort diversity (1). Indeed, we detected in our HIC cohort a
few individuals that displayed mRNA levels of SAMHD1 and/or SLFN11 well above the normal
range (Supplementary Figure S1). These observations suggest that there might not be a unique host
RF expression signature common to all HIC, but that different combinations of host RF could be
associated with natural control of HIV-1 replication in distinct individuals. Thus, the particular set of
increased host RF may vary across different HIC cohorts.

In contrast to p21 and MCPIP1, the other antiretroviral RF analyzed here are interferon (IFN)-
stimulated genes (ISGs) (25). In the chronic phase of HIV−1 infection in viremic untreated patients,
most ISGs (mainly those responding to IFN type I) are upregulated in CD4$^+$ T cells (26–28) and their
expression is positively correlated with the percentage of activated T cells and negatively correlated
with CD4$^+$ T cell counts (26–30). We found no significant upregulation of ISGs in our HIC cohort,
likewise other studies (3,27,28,31), except for IFITM1 that is also induced by type II IFN (32),
indicating that another pathway may have stimulated its expression. These results suggest that
residual or low-level viremia might not be enough to induce a generalized upregulation of ISGs
during chronic infection (28). In addition, MCPIP1 negatively regulates the NF-κB cascade (33,34)
and its overexpression in HIC may also contribute to limit the chronic overexpression of ISGs in
HIC.

Our results confirm previous observations that levels of p21 transcripts are positively correlated with
CD4$^+$ T cell activation in EC and HIC groups (3) and further suggest a positive correlation of p21
and MCPIP1 mRNA with sCD14, a marker of monocyte activation, in EC. Increased expression of
p21/MCPIP1 associated with T cell and/or monocyte activation seems to be a unique characteristic of
HIC/EC, because similar correlations were not observed for HIV-infected/ART-suppressed or HIV-
negative subjects. Furthermore, previous studies have shown that viremic progressors displayed
reduced levels of p21 even though cellular activation was much higher than in HIC or HIV-
uninfected individuals (6). These results suggest that p21/MCPIP1 overexpression may be a
distinctive homoeostatic innate response of HIC to limit the deleterious effects of chronic immune
activation driven by HIV-1 infection.

Selective upregulation of p21 and MCPIP1 in CD4$^+$ T, macrophages and/or dendritic cells may
directly limit HIV-1 replication by 1) reducing the reverse transcription and chromosomal integration
of HIV-1 in quiescent cells and thus limiting the size of the latent proviral reservoir (11–13,35–37);
2) restricting HIV-1 LTR transcription (33,34,38,39); and, 3) degrading viral mRNA and miRNA
(17,40–42). Upregulation of p21 and MCPIP1 may also indirectly limit HIV-1 replication and further
prevent CD4$^+$ T cells loss by reducing chronic IFN-I signaling, generalized inflammation and over-
activation of the immune system (43–52), without affecting the activation of antiviral cellular
responses. Although the enhanced antiviral and anti-inflammatory state may not be enough to fully
restrict HIV-1 replication (53), it could act in concert with other innate and adaptive immune
mechanisms to control HIV replication in HIC.

The enhanced expression of a few select host RF, including p21, was strongly associated with
reduced CD4$^+$ T cell-associated HIV RNA during ART, indicating that this host RF may contribute
to the control of viral expression and ongoing replication during ART (54). Another study
demonstrates that atorvastatin, a lipid-lowering medication, exert a broad spectrum of anti-

inflammatory functions and further reduced HIV infection in both rested and activated CD4[+] T cells
in vitro via p21 upregulation (14). Interestingly, atorvastatin was found to up-regulates p21 through a
p53 independent pathway, which is consistent with a potential role of MCPIP1 in that antiviral
mechanism. These observations suggest that pharmacological manipulation of p21 and MCPIP1 may
open novel therapeutic perspectives to prevent HIV-1 replication and to attenuate HIV-associated
inflammation and immune activation during ART.

An important limitation of our study is the impossibility of assigning which cell(s) population(s) has
increased expression of p21 and MCPIP1 in HIC. The expression profile of many RF and ISGs may
be different between CD4[+] T cells and monocytes (8), suggesting that the individualization of these
cell types might better decipher the mechanisms of RF regulation in the setting of natural control of
HIV-1 infection. Another potential limitation is that only RF mRNA levels were analyzed. Previous
studies showed that p21 mRNA levels mirror p21 protein levels in CD4[+] T cells from HIC (6) and
that MCPIP1 mRNA levels reflect MCPIP1 protein levels in HCV-infected hepatoma cells (55).
Although these evidences indicate a close match between transcripts and protein expression levels,
measuring the levels/activity of p21 and MCPIP1 proteins in cells from HIC should also help to
elucidate the relevance of these RF for HIV control.

In summary, our data confirm the high levels of p21 mRNA expression and shows for the first-time
the concurrent overexpression of MCPIP1 mRNA in HIC. Moreover, we found a positive correlation
between p21 and MCPIP1 transcripts in HIC, indicating a possible synergistic effect of both innate
host RF on natural suppression of HIV-1 replication *in vivo*. Further studies are needed to better
understand the role of p21 and MCPIP1 in the natural control of HIV-1 replication and disease
progression in HIC. These findings may also have important implications for the development of new
immune-based therapeutic strategies for a functional cure of HIV-1 infection.

5   Figure legends

Figure 1. p21 and MCPIP1 mRNA levels are up-regulated in PBMC from HIC. Boxplots
represent the interquartile and sample median (central solid black line) of the relative changes (fold-
change values relative to the mean of HIV-1-uninfected (NEG) subjects) of p21 and MCPIP1
expression comparing NEG and ART-suppressed subjects (ART) with HIV controllers (HIC) (A and
B) or elite (EC) and viremic (VC) controllers (C and D). P-values < 0.05 were considered statistically
significant.

Figure 2: **p21 and MCPIP1 mRNA levels in PBMC from HIC are positively correlated.** The p21
and MCPIP1 normalized expression correlations were calculated considering all groups (A), HIV-
infected (B), HIC (C), and EC (D). The points' colors indicate the patient group, accordingly to the
legend. Correlation coefficients (Spearman's ρ) are shown in the upper right corner of each graph. P-
values < 0.05 were considered statistically significant.

Figure 3. p21 transcripts are positively correlated with CD4$^+$ T cell and monocyte activation
while MCPIP1 transcripts are positively correlated only with monocyte activation in EC. The
mean of activated CD4$^+$ T cells (HLA-DR$^+$CD38$^+$) counts (A) and soluble CD14 (sCD14) in plasma
(B) were compared for each group. The color of each dot represents the group as indicated in the
legend at right. P-values < 0.05 were considered statistically significant. The correlations were made
evaluating the relationship between activated CD4$^+$ T cells (C and D) or sCD14 levels (E and F) with
the normalized expression of p21 and MCPIP1 for EC and HIC groups. The points' colors present in
each graph indicate the groups present according to the legend. Correlations coefficient (Spearman's
ρ) are shown in the upper left corner of each graph.

6   Ethics Statement

This study was carried out in accordance with the recommendations of the ethical committee of
Instituto Nacional de Infectologia Evandro Chagas (INI-Fiocruz) that approved the study protocol
(CAAE 1717.0.000.009-07). All subjects gave written informed consent in accordance with the
Declaration of Helsinki.

7   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

8   Author Contributions

GB and TMLS conceived and designed the study and supervised the experiments. SSDA conducted
experiments and analyzed the data together with MR-A and GB. FH performed the CD4$^+$ T cell and
monocyte activation assays. ED collaborated with mRNA gene-expression analysis. BH, BG, and
VGV conducted patient recruitment and follow-up. FH, ED and MGM provided intellectual input for
results interpretations. SSDA, GB and MR-A wrote the first draft and all authors assisted with the
writing and approved the final manuscript.

9   Acknowledgments

The authors thank the patients, who participated in the study, as well as all the technical staff
involved in the clinical follow-up of these patients. We also thank Ms Marilia Alves Figueira de

Melo for the excellent technical support in RNA quantification and integrity analyses and the
Plataforma de PCR em Tempo Real – RJ (RPT09A) – FIOCRUZ and Plataforma de Sequenciamento
de Ácidos Nucléicos de Nova Geração – RJ (RPT01J) – FIOCRUZ.

10 Funding

This work was supported by the Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro –
FAPERJ (grant number E-26/110.123/2014) and the Conselho Nacional de Desenvolvimento
Científico e Tecnológico – CNPq (Grant Number 401220/2016-8). SSDA was supported by funding
from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Fundação de
Amparo à Pesquisa do Estado do Rio de Janeiro – FAPERJ. ED was financed by a Postdoctoral
fellowship from the "Programa Nacional de Pós-Doutorado (PNPD)" by the Coordenação de
Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

11 References

1. Walker BD, Yu XG. Unravelling the mechanisms of durable control of HIV-1. *Nat Rev*
Immunol (2013) **13**:487–98. doi:10.1038/nri3478

2. Colomer-Lluch M, Ruiz A, Moris A, Prado JG. Restriction Factors: From Intrinsic Viral
Restriction to Shaping Cellular Immunity Against HIV-1. *Front Immunol* (2018) **9**:2876.
doi:10.3389/fimmu.2018.02876

3. Abdel-Mohsen M, Raposo RAS, Deng X, Li M, Liegler T, Sinclair E, Salama MS, Ghanem
HEA, Hoh R, Wong JK, et al. Expression profile of host restriction factors in HIV-1 elite
controllers. *Retrovirology* (2013) **10**:1–13. doi:10.1186/1742-4690-10-106

4. Madlala P, Van de Velde P, Van Remoortel B, Vets S, Van Wijngaerden E, Van Laethem K,
Gijsbers R, Schrijvers R, Debyser Z. Analysis of ex vivo HIV-1 infection in a controller-
discordant couple. *J virus Erad* (2018) **4**:170–173.

5. Krishnan S, Wilson EMP, Sheikh V, Rupert A, Mendoza D, Yang J, Lempicki R, Migueles
SA, Sereti I. Evidence for innate immune system activation in HIV type 1-infected elite
controllers. *J Infect Dis* (2014) **209**:931–939. doi:10.1093/infdis/jit581

6. Chen H, Li C, Huang J, Cung T, Seiss K, Beamon J, Carrington MF, Porter LC, Burke PS,
Yang Y, et al. CD4+ T cells from elite controllers resist HIV-1 infection by selective
upregulation of p21. *J Clin Invest* (2011) **121**:1549–1560. doi:10.1172/JCI44539

7. Riveira-Muñoz E, Ruiz A, Pauls E, Permanyer M, Badia R, Mothe B, Crespo M, Clotet B,
Brander C, Ballana E, et al. Increased expression of SAMHD1 in a subset of HIV-1 elite
controllers. *J Antimicrob Chemother* (2014) **69**:3057–3060. doi:10.1093/jac/dku276

8. Canoui E, Noël N, Lécuroux C, Boufassa F, Sáez-Cirión A, Bourgeois C, Lambotte O, ANRS
CO21 CODEX Study Group. Strong ifitm1 Expression in CD4 T Cells in HIV Controllers Is
Correlated With Immune Activation. *J Acquir Immune Defic Syndr* (2017) **74**:e56–e59.
doi:10.1097/QAI.0000000000001166

9. Sáez-Cirión A, Hamimi C, Bergamaschi A, David A, Versmisse P, Mélard A, Boufassa F,
Barré-Sinoussi F, Lambotte O, Rouzioux C, et al. Restriction of HIV-1 replication in

macrophages and CD4+ T cells from HIV controllers. *Blood* (2011) **118**:955–964.
doi:10.1182/blood-2010-12-327106

10. Van Hecke C, Trypsteen W, Malatinkova E, De Spiegelaere W, Vervisch K, Rutsaert S,
Kinloch-de Loes S, Sips M, Vandekerckhove L. Early treated HIV-1 positive individuals
demonstrate similar restriction factor expression profile as long-term non-progressors.
EBioMedicine (2019)1–12. doi:10.1016/j.ebiom.2019.02.006

11. Zhang J, Scadden DT, Crumpacker CS. Primitive hematopoietic cells resist HIV-1 infection
via p21Waf1/Cip1/Sdi1. *J Clin Invest* (2007) **117**:473–481. doi:10.1172/JCI28971

12. Bergamaschi A, David A, Le Rouzic E, Nisole S, Barre-Sinoussi F, Pancino G. The CDK
Inhibitor p21Cip1/WAF1 Is Induced by Fc R Activation and Restricts the Replication of
Human Immunodeficiency Virus Type 1 and Related Primate Lentiviruses in Human
Macrophages. *J Virol* (2009) **83**:12253–12265. doi:10.1128/JVI.01395-09

13. Valle-Casuso JC, Allouch A, Lenzi G, Studdard L, Barré-sinoussi F, Müller-trutwin M, Kim
B. p21 Restricts HIV-1 in Monocyte-Derived Dendritic Cells through the Reduction of
Deoxynucleoside Triphosphate Biosynthesis and Regulation of SAMHD1 Antiviral Activity. *J*
Virol (2017) **91**:1–18.

14. Elahi S, Weiss RH, Merani S. Atorvastatin restricts HIV replication in CD4+ T cells by
upregulation of p21. *Aids* (2016) **30**:171–183. doi:10.1097/QAD.0000000000000917

15. Elahi S, Niki T, Hirashima M, Horton H. Galectin-9 binding to Tim-3 renders activated human
CD4+ T cells less susceptible to HIV-1 infection. *Blood* (2012) **119**:4192–4204.
doi:10.1182/blood-2011-11-389585

16. Moosa Y, Tanko RF, Ramsuran V, Singh R, Madzivhandila M, Yende-Zuma N, Abrahams
MR, Selhorst P, Gounder K, Moore PL, et al. Case report: Mechanisms of HIV elite control in
two African women. *BMC Infect Dis* (2018) **18**:1–7. doi:10.1186/s12879-018-2961-8

17. Liu S, Qiu C, Miao R, Zhou J, Lee A, Liu B, Lester SN, Fu W, Zhu L, Zhang L, et al. MCPIP1
restricts HIV infection and is rapidly degraded in activated CD4+ T cells. *Proc Natl Acad Sci*
U S A (2013) **110**:19083–8. doi:10.1073/pnas.1316208110

18. Fu M, Blackshear PJ. RNA-binding proteins in immune regulation: A focus on CCCH zinc
finger proteins. *Nat Rev Immunol* (2017) **17**:130–143. doi:10.1038/nri.2016.129

19. Lichawska-Cieslar A, Pietrzycka R, Ligeza J, Kulecka M, Paziewska A, Kalita A, Dolicka
DD, Wilamowski M, Miekus K, Ostrowski J, et al. RNA sequencing reveals widespread
transcriptome changes in a renal carcinoma cell line. *Oncotarget* (2018) **9**:8597–8613.
doi:10.18632/oncotarget.24269

20. Azevedo SSD, Caetano DG, Côrtes FH, Teixeira SLM, Santos Silva K, Hoagland B,
Grinsztejn B, Veloso VG, Morgado MG, Bello G. Highly divergent patterns of genetic
diversity and evolution in proviral quasispecies from HIV controllers. *Retrovirology* (2017)
14:1–13. doi:10.1186/s12977-017-0354-5

21. Côrtes FH, de Paula HHS, Bello G, Ribeiro-Alves M, de Azevedo SSD, Caetano DG, Teixeira

SLM, Hoagland B, Grinsztejn B, Veloso VG, et al. Plasmatic levels of IL-18, IP-10, and activated CD8+T cells are potential biomarkers to identify HIV-1 elite controllers with a true functional cure profile. *Front Immunol* (2018) **9**:1–11. doi:10.3389/fimmu.2018.01576

22. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* (2002) **3**:RESEARCH0034. doi:10.1186/gb-2002-3-7-research0034

23. Basso D, Pesarin F, Salmaso L, Solari A. "Nonparametric One-Way ANOVA", in *Permutation Tests for Stochastic Ordering and ANOVA* (New York, NY: Springer), 105–132. doi:10.1007/978-0-387-85956-9_5

24. R Core Team. *R: A language and environment for statistical computing*. , org. R Foundation for Statistical Computing Vienna, Austria. (2018).

25. Doyle T, Goujon C, Malim MH. HIV-1 and interferons: who's interfering with whom? *Nat Rev Microbiol* (2015) **13**:403–413. doi:10.1038/nrmicro3449

26. Sedaghat AR, German J, Teslovich TM, Cofrancesco J, Jie CC, Talbot CC, Siliciano RF. Chronic CD4+ T-Cell Activation and Depletion in Human Immunodeficiency Virus Type 1 Infection: Type I Interferon-Mediated Disruption of T-Cell Dynamics. *J Virol* (2008) **82**:1870–1883. doi:10.1128/JVI.02228-07

27. Hyrcza MD, Kovacs C, Loutfy M, Halpenny R, Heisler L, Yang S, Wilkins O, Ostrowski M, Der SD. Distinct Transcriptional Profiles in Ex Vivo CD4+ and CD8+ T Cells Are Established Early in Human Immunodeficiency Virus Type 1 Infection and Are Characterized by a Chronic Interferon Response as Well as Extensive Transcriptional Changes in CD8+ T Cells. *J Virol* (2007) **81**:3477–3486. doi:10.1128/JVI.01552-06

28. Rotger M, Dang KK, Fellay J, Heinzen EL, Feng S, Descombes P, Shianna K V., Ge D, Günthard HF, Goldstein DB, et al. Genome-Wide mRNA Expression Correlates of Viral Control in CD4+ T-Cells from HIV-1-Infected Individuals. *PLoS Pathog* (2010) **6**:e1000781. doi:10.1371/journal.ppat.1000781

29. Hardy GAD, Sieg S, Rodriguez B, Anthony D, Asaad R, Jiang W, Mudd J, Schacker T, Funderburg NT, Pilch-Cooper HA, et al. Interferon-α Is the Primary Plasma Type-I IFN in HIV-1 Infection and Correlates with Immune Activation and Disease Markers. *PLoS One* (2013) **8**:e56527. doi:10.1371/journal.pone.0056527

30. Fernandez S, Tanaskovic S, Helbig K, Rajasuriar R, Kramski M, Murray JM, Beard M, Purcell D, Lewin SR, Price P, et al. CD4+ T-Cell Deficiency in HIV Patients Responding to Antiretroviral Therapy Is Associated With Increased Expression of Interferon-Stimulated Genes in CD4+ T Cells. *J Infect Dis* (2011) **204**:1927–1935. doi:10.1093/infdis/jir659

31. Vigneault F, Woods M, Buzon MJ, Li C, Pereyra F, Crosby SD, Rychert J, Church G, Martinez-Picado J, Rosenberg ES, et al. Transcriptional profiling of CD4 T cells identifies distinct subgroups of HIV-1 elite controllers. *J Virol* (2011) **85**:3015–9. doi:10.1128/JVI.01846-10

405   32.   Diamond MS, Farzan M. The broad-spectrum antiviral functions of IFIT and IFITM proteins.
406   *Nat Rev Immunol* (2013) **13**:46–57. doi:10.1038/nri3344

407   33.   Skalniak L, Mizgalska D, Zarebski A, Wyrzykowska P, Koj A, Jura J. Regulatory feedback
408   loop between NF-κB and MCP-1-induced protein 1 RNase. *FEBS J* (2009) **276**:5892–5905.
409   doi:10.1111/j.1742-4658.2009.07273.x

410   34.   Liang J, Saad Y, Lei T, Wang J, Qi D, Yang Q, Kolattukudy PE, Fu M. MCP-induced protein
411   1 deubiquitinates TRAF proteins and negatively regulates JNK and NF-κB signaling. *J Exp
412   Med* (2010) **207**:2959–2973. doi:10.1084/jem.20092641

413   35.   Pauls E, Ruiz A, Riveira-Muñoz E, Permanyer M, Badia R, Clotet B, Keppler OT, Ballana E,
414   Este JA. p21 regulates the HIV-1 restriction factor SAMHD1. *Proc Natl Acad Sci U S A*
415   (2014) **111**:E1322-4. doi:10.1073/pnas.1322059111

416   36.   Allouch A, David A, Amie SM, Lahouassa H, Chartier L, Margottin-Goguet F, Barre-Sinoussi
417   F, Kim B, Saez-Cirion A, Pancino G. p21-mediated RNR2 repression restricts HIV-1
418   replication in macrophages by inhibiting dNTP biosynthesis pathway. *Proc Natl Acad Sci*
419   (2013) **110**:E3997–E4006. doi:10.1073/pnas.1306719110

420   37.   Leng J, Ho H-P, Buzon MJ, Pereyra F, Walker BD, Yu XG, Chang EJ, Lichterfeld M. A Cell-
421   Intrinsic Inhibitor of HIV-1 Reverse Transcription in CD4+ T Cells from Elite Controllers.
422   *Cell Host Microbe* (2014) **15**:717–728. doi:10.1016/j.chom.2014.05.011

423   38.   Wang D, de la Fuente C, Deng L, Wang L, Zilberman I, Eadie C, Healey M, Stein D, Denny
424   T, Harrison LE, et al. Inhibition of Human Immunodeficiency Virus Type 1 Transcription by
425   Chemical Cyclin-Dependent Kinase Inhibitors. *J Virol* (2001) **75**:7266–7279.
426   doi:10.1128/JVI.75.16.7266-7279.2001

427   39.   Kumari N, Iordanskiy S, Kovalskyy D, Breuer D, Niu X, Lin X, Xu M, Gavrilenko K,
428   Kashanchi F, Dhawan S, et al. Phenyl-1-Pyridin-2yl-Ethanone-Based Iron Chelators Increase
429   IκB-α Expression, Modulate CDK2 and CDK9 Activities, and Inhibit HIV-1 Transcription.
430   *Antimicrob Agents Chemother* (2014) **58**:6558–6571. doi:10.1128/AAC.02918-14

431   40.   Lin R-J, Chien H-L, Lin S-Y, Chang B-L, Yu H-P, Tang W-C, Lin Y-L. MCPIP1 ribonuclease
432   exhibits broad-spectrum antiviral effects through viral RNA binding and degradation. *Nucleic
433   Acids Res* (2013) **41**:3314–3326. doi:10.1093/nar/gkt019

434   41.   Suzuki HI, Arase M, Matsuyama H, Choi YL, Ueno T, Mano H, Sugimoto K, Miyazono K.
435   MCPIP1 ribonuclease antagonizes dicer and terminates microRNA biogenesis through
436   precursor microRNA degradation. *Mol Cell* (2011) **44**:424–436.
437   doi:10.1016/j.molcel.2011.09.012

438   42.   Happel C, Ramalingam D, Ziegelbauer JM. Virus-Mediated Alterations in miRNA Factors and
439   Degradation of Viral miRNAs by MCPIP1. *PLoS Biol* (2016) **14**:1–23.
440   doi:10.1371/journal.pbio.2000998

441   43.   Santiago-Raber M-L, Lawson BR, Dummer W, Barnhouse M, Koundouris S, Wilson CB,
442   Kono DH, Theofilopoulos AN. Role of Cyclin Kinase Inhibitor p21 in Systemic
443   Autoimmunity. *J Immunol* (2001) **167**:4067–4074. doi:10.4049/jimmunol.167.7.4067

44.   Khanna AK, Plummer M, Nilakantan V, Pieper GM. Recombinant p21 Protein Inhibits
Lymphocyte Proliferation and Transcription Factors. *J Immunol* (2005) **174**:7610–7617.
doi:10.4049/jimmunol.174.12.7610

45.   Arias CF, Ballesteros-Tato A, Garcia MI, Martin-Caballero J, Flores JM, Martinez-A C,
Balomenos D. p21CIP1/WAF1 Controls Proliferation of Activated/Memory T Cells and
Affects Homeostasis and Memory T Cell Responses. *J Immunol* (2007) **178**:2296–2306.
doi:10.4049/jimmunol.178.4.2296

46.   Lloberas J, Celada A. P21waf1/CIP1, a CDK inhibitor and a negative feedback system that
controls   macrophage   activation.   *Eur   J   Immunol*   (2009)   **39**:691–694.
doi:10.1002/eji.200939262

47.   Scatizzi JC, Mavers M, Hutcheson J, Young B, Shi B, Pope RM, Ruderman EM, Samways
DSK, Corbett JA, Egan TM, et al. The CDK domain of p21 is a suppressor of IL-1beta-
mediated inflammation in activated macrophages. *Eur J Immunol* (2009) **39**:820–5.
doi:10.1002/eji.200838683

48.   Trakala M, Arias CF, García MI, Moreno-Ortiz MC, Tsilingiri K, Fernández PJ, Mellado M,
Díaz-Meco MT, Moscat J, Serrano M, et al. Regulation of macrophage activation and septic
shock   susceptibility   via   p21   (WAF1/CIP1).   *Eur   J   Immunol*   (2009)   **39**:810–819.
doi:10.1002/eji.200838676

49.   Liang J, Song W, Tromp G, Kolattukudy PE, Fu M. Genome-wide survey and expression
profiling of CCCH-zinc finger family reveals a functional module in macrophage activation.
PLoS One (2008) **3**: doi:10.1371/journal.pone.0002880

50.   Matsushita K, Takeuchi O, Standley DM, Kumagai Y, Kawagoe T, Miyake T, Satoh T, Kato
H, Tsujimura T, Nakamura H, et al. Zc3h12a is an RNase essential for controlling immune
responses   by   regulating   mRNA   decay.   *Nature*   (2009)   **458**:1185–1190.
doi:10.1038/nature07924

51.   Li Y, Huang X, Huang S, He H, Lei T, Saaoud F, Yu X-Q, Melnick A, Kumar A, Papasian CJ,
et al. Central role of myeloid MCPIP1 in protecting against LPS-induced inflammation and
lung injury. *Signal Transduct Target Ther* (2017) **2**:17066. doi:10.1038/sigtrans.2017.66

52.   Uehata T, Takeuchi O. Regnase-1 Is an Endoribonuclease Essential for the Maintenance of
Immune Homeostasis. *J Interf Cytokine Res* (2017) **37**:220–229. doi:10.1089/jir.2017.0001

53.   De Pablo A, Bogoi R, Bejarano I, Toro C, Valencia E, Moreno V, Martín-Carbonero L,
Gómez-Hernando C, Rodés B. Short communication: p21/CDKN1A expression shows broad
interindividual diversity in a subset of HIV-1 elite controllers. *AIDS Res Hum Retroviruses*
(2016) **32**:1–5. doi:10.1089/aid.2015.0137

54.   Abdel-Mohsen M, Wang C, Strain MC, Lada SM, Deng X, Cockerham LR, Pilcher CD, Hecht
FM, Liegler T, Richman DD, et al. Select host restriction factors are associated with HIV
persistence   during   antiretroviral   therapy.   *AIDS*   (2015)   **29**:411–20.
doi:10.1097/QAD.0000000000000572

55.   Lin R-J, Chu J-S, Chien H-L, Tseng C-H, Ko P-C, Mei Y-Y, Tang W-C, Kao Y-T, Cheng H-

Y, Liang Y-C, et al. MCPIP1 suppresses hepatitis C virus replication and negatively regulates
virus-induced proinflammatory cytokine responses. *J Immunol* (2014) **193**:4159–68.
doi:10.4049/jimmunol.1400337

490 **Table 1.** Main clinical and epidemiologic characteristics of individuals of this study.

Characteristics	HIC (n = 21)		ART-suppressed (n = 8)	HIV-1 negative (n = 10)
	EC (n = 13)	VC (n = 8)		
Sex, no. (%)				
Female	10 (77)	3 (38)	4 (50)	6 (60)
Male	3 (23)	5 (62)	4 (50)	4 (40)
Age (years)*	45 (39-60)	43.5 (39-47)	47 (38-53)	47 (36-51)
Study point				
Time since HIV-1 diagnosis (years)	9 (5.5-15)	12.5 (7-16)	NA	-
CD4+ T cell (cells/µl)	1027 (834-1255)	664 (563-1228)	889 (678-1097)	1043 (784-1581)
Plasma HIV RNA (copies/ml)	<50	641 (327-915)	<40	-
CD4/CD8 ratio	1.33 (1.24-1.61)	0.91 (0.67-1.23)	1.06 (0.73-1.5)	1.69 (1.62-2.07)

498 * Age at study point; Interquartile ranges are shown in parenthesis. HIC, HIV controllers; ART, antiretroviral therapy;
499 EC, elite controllers; VC, viremic controllers. NA, not available.