

On the efficacy of restoration in stream networks: comments, critiques, and prospective
recommendations

David Murray-Stoker^{1,2}

¹Odum School of Ecology, University of Georgia, Athens, GA 30602, U.S.A.

²Present Address: Department of Ecology and Evolutionary Biology, University of Toronto,
Toronto, ON M5S 3B2, Canada

ORCID: <https://orcid.org/0000-0002-4774-6948>

E-mail: dstoker92@gmail.com

Abstract

Swan and Brown (2017) recently addressed the effects of restoration on stream communities under the meta-community framework. Using a combination of headwater and mainstem streams, Swan and Brown (2017) evaluated how position within a stream network affected the outcome of restoration on invertebrate communities. Ostensibly, their hypotheses were partially supported as restoration had stronger effects in headwater streams: invertebrate taxonomic richness was increased and temporal variability decreased in restored reaches; however, these results were not consistent upon closer scrutiny for both the original paper (Swan and Brown 2017) and the later erratum (Swan and Brown 2018). Here, I provide a secondary analysis of the data, with hypotheses and interpretations in the context of stream, metacommunity, and restoration ecology. I did not find any effects of restoration on local diversity, spatial dissimilarity, or temporal variability, let alone differential effects of restoration between headwaters and mainstems; these results are contrary Swan and Brown (2017, 2018), who reported that restoration increased taxonomic richness, increased spatial dissimilarity, and decreased temporal variability in restored headwater streams. I demonstrate further that the statistical tests conducted by Swan and Brown (2017, 2018) were invalid and, therefore, recommend the use of the results presented here. More broadly, I suggest that river and stream restoration will likely have greater success if a regional approach is taken to designing and implementing restoration projects.

Keywords: biodiversity, community ecology, freshwater ecology, metacommunity theory, open science, restoration ecology

Introduction

In a recent study, Swan and Brown (2017) evaluated how restoration affected community diversity in streams through the use of metacommunity theory. Under this framework, local effects are associated with species' niches while regional effects are more associated with dispersal (Leibold et al. 2004). In the context of stream networks, headwaters are isolated patches more likely to be impacted by niches and environmental characteristics and mainstems are well-connected more likely to be affected by dispersal (Heino et al. 2003, Leibold et al. 2004, Grant et al. 2007, Altermatt 2013, Heino 2013). Restoration of stream habitats was therefore expected to have a greater impact on communities in headwaters relative to mainstems (Swan and Brown 2017).

Although Swan and Brown (2017, 2018) noted that restoration techniques can vary in intrusiveness on stream ecosystems, they did not account for this in their experimental design and statistical analyses. Restored streams in their study received various combinations of bank stabilization, in-channel manipulation, and riparian reforestation (i.e. tree planting) treatments, and these treatments were not applied in a consistent or systematic manner (Swan and Brown 2017: Table 2). Swan and Brown (2017) did not set a restoration criterion for site inclusion in their study, instead including all sites regardless of the combination of applied restoration treatments. I suggest that this oversight leads to unnecessary assumptions about the efficacy of restoration by assuming the effects of all treatments combinations are equivalent, and this issue could have been partially resolved a priori by hypothesizing how each restoration reach would affect headwater and mainstem streams and then setting requirements for site inclusion in the analyses.

I contend that the various restoration treatments differ not only in their overall effects but also if the treatment is applied in headwater or mainstem streams, and, for these reasons, criteria for site selection could be set. I suggest that bank stabilization and in-channel manipulation treatments are more likely to have consistent effects in both headwaters and mainstems, while riparian reforestation would likely have stronger effects in headwater compared to mainstem streams. A similar argument was made by Swan and Brown, though it was not explicitly noted until the erratum (Swan and Brown 2018). Bank stabilization and in-channel manipulation can increase bed stability and substrate availability and diversity in both headwater and mainstem streams; however, the effects of riparian reforestation could act on a gradient from headwaters to mainstems. For example, leaf litter is an important source of habitat and nutrients in headwaters but less so in mainstems (Vannote et al. 1980, Rosi-Marshall and Wallace 2002). Additionally, the utility of riparian reforestation on reducing nutrient inputs notwithstanding (Collins et al. 2013), the effects of riparian reforestation could be stronger in headwater streams because they are isolated systems, whereas mainstem streams receive flows of water, nutrients, and organisms from many tributaries (Vannote et al. 1980). Effectively, mainstems are dependent on other tributaries and any local restoration effects via riparian reforestation could be overwhelmed by incoming flows from unrestored streams (Wahl et al. 2013).

Here, I present a re-analysis of the data provided by Swan and Brown (2017, 2018). I hypothesized that stream-channel manipulations would have a more consistent effect between headwaters and mainstems relative to the effect of riparian reforestation, with stronger effects of restoration in headwaters relative to mainstems; I used this hypothesis to guide and inform site selection in my re-analysis. I required sites in the re-analysis to have received both the bank stabilization and in-channel manipulations treatments (hereafter “revised” sites), although sites

receiving riparian reforestation were also included if they received both the bank stabilization and in-channel manipulations treatments. I also re-analyzed the full data (hereafter “full” sites) to determine if any differences, or lack thereof, between the full and revised sites analyses could be attributed to increased variation in the revised sites due to decreased sample size. Finally, I compare the interpretation and conclusions from my re-analysis to those in Swan and Brown (2017) and the erratum (Swan and Brown 2018).

Methods

Sampling Design

Swan and Brown (2017) conducted their study in 5 headwater and 8 mainstem streams in Baltimore County, Maryland, U.S.A. Each stream had a paired structure, where restored and adjacent, unrestored reaches were sampled; restored and adjacent reaches were separated by < 10 m. The sampling design was explicitly constructed to permit comparisons between paired restored-adjacent reaches in each of the focal streams. Each of the 13 focal streams was sampled quarterly in 2011 (spring, summer, and fall) and 2012 (winter; Swan and Brown 2017).

Statistical Analyses

I generally followed the analyses as written by Swan and Brown (2017), with modifications made when necessary. The three community response variables were local diversity, spatial dissimilarity, and temporal variation. Local diversity was calculated as taxonomic richness (i.e. number of different taxa present) and taxonomic diversity (i.e. Shannon’s diversity) and compared using an analysis of variance (ANOVA). The model was constructed to examine the individual effects of reach (restored or adjacent), order (headwater or mainstem), and season (spring, summer, fall, and winter) and all two- and three-way interactions, with individual ANOVAs for richness and diversity; I also fit the full and reduced taxonomic

richness models proposed in the erratum (Swan and Brown 2018) as a separate set of ANOVAs. Spatial dissimilarity between communities in restored and adjacent reaches for each order-by-season combination was quantified using the modified Gower index (Anderson et al. 2006) with a logarithm with a base of 5 on an untransformed abundance matrix. Values of the modified Gower dissimilarities were then compared using an ANOVA with the individual effects of season and order as well as their interaction. Temporal variability was measured as the multivariate dispersion (i.e. mean distance to the centroid) of repeated samples for each stream-by-reach-by-order combination (Anderson et al. 2006). Distances were calculated in principal coordinates space after Bray-Curtis dissimilarity was performed on the untransformed abundance matrix. Temporal variability values were then compared using an ANOVA with the individual effects of order and reach and their interaction. All ANOVAs were performed for both the full and revised sites, with stream identity fitted as a random blocking factor in each ANOVA; all ANOVAs were fitted by restricted maximum likelihood.

Exploratory data analysis was conducted prior to any model fitting to determine if the data met test assumptions (Zuur et al. 2010). For the full sites analyses, numerical summaries demonstrated an unbalanced design, with equal representation of restored and unrestored reaches but a large disparity in the number of samples between headwaters and mainstems for each of the taxonomic richness and diversity (headwater $n = 38$, mainstem $n = 62$), spatial dissimilarity (headwater $n = 19$, mainstem $n = 31$), and temporal variation (headwater $n = 10$, mainstem $n = 16$) analyses. Additionally, the assumption of homogeneity of variance was violated for the taxonomic richness and diversity and the spatial dissimilarity analyses. The unbalanced design was greatly reduced for the revised sites analyses: taxonomic richness and diversity (headwater $n = 30$, mainstem $n = 40$), spatial dissimilarity (headwater $n = 15$, mainstem $n = 20$), and temporal

variation (headwater $n = 8$, mainstem $n = 10$); however, the assumption of homogeneity of variance was still violated. To better meet the assumption of equal variance, taxonomic richness was \ln -transformed, taxonomic diversity was square root-transformed, and spatial dissimilarity was \ln -transformed for all analyses. Along with using transformations to response variables to better meet model assumptions, I used Type III sums of squares for evaluating main and interactive effects of factors included in the ANOVA. Swan and Brown (2017, 2018) used Type I sums of squares, which are inadequate for unbalanced and multi-factor designs with interactions between or among factors (Shaw and Mitchell-Olds 1993, Quinn and Keough 2002). Model assumptions were inspected graphically, and significance was considered at $P < 0.050$. I removed the spring sample from the restored reach of site 227 from analyses because there was no corresponding sample from the adjacent reach, which would have precluded paired comparisons of restored-adjacent sites; however, I did not remove any sites prior to fitting the full and reduced model ANOVAs set by Swan and Brown (2018). Additionally, the only difference between the full and reduced ANOVAs set by Swan and Brown (2018) was the use of Type III sums of squares instead of Type I sums of squares. Untransformed values of variables are presented in the results and figures. All analyses were conducted using R version 3.5.3 (R Core Team 2019) with the nlme (version 3.1-139, Pinheiro et al. 2019) and vegan (version 2.5-4, Oksanen et al. 2019) packages; data and R code are deposited in the figshare repository ([10.6084/m9.figshare.6448010](https://doi.org/10.6084/m9.figshare.6448010)). Given I made necessary modifications to the analyses written by Swan and Brown (2017, 2018), later comparisons between the re-analysis presented here and the results presented by Swan and Brown (2017, 2018) will only be in terms of statistical and ecological interpretation and not exact values of test statistics. To facilitate discussion among the initial study (Swan and Brown 2017), erratum (Swan and Brown 2018), and this re-analysis,

effect sizes were calculated for each factor and interaction in the ANOVA models. Local diversity and temporal variability effect sizes were calculated for the erratum (Swan and Brown 2018) and full and reduced models in this re-analysis. Spatial dissimilarity effect sizes were calculated for the initial study (Swan and Brown 2017) and the full and reduced models in this re-analysis; effect sizes were not calculated for local diversity and temporal variability of the initial study (Swan and Brown 2017) as results were later corrected in the erratum (Swan and Brown 2018), and it would be illogical to make comparisons to deprecated analyses. All effect sizes were calculated as partial η^2 (Cohen 1973):

$$\eta = \frac{df_{between} \times F}{df_{between} \times F + df_{within}}$$

whereby $df_{between}$ is the degrees of freedom associated with the factor or interaction, F is the F statistic associated with the factor or interaction, and df_{within} is the degrees of freedom associated with the residual error. Effect sizes were classified as small = 0.01, medium = 0.06, and large = 0.14.

Results & Discussion

There were no main or interactive effects of season, order, or reach on taxonomic richness for either the full or revised sites analyses (Table 1, Figure 1). The full model of taxonomic richness proposed in the erratum (Swan and Brown 2018) did not show any main or interactive effects of season, order, or reach (Table 2); however, the reduced model of taxonomic richness demonstrated an interaction between order and season ($F_{3, 80} = 4.105$, $P = 0.009$) and significant main effects of season ($F_{3, 80} = 4.358$, $P = 0.007$) and reach ($F_{1, 80} = 4.844$, $P = 0.031$). In contrast to taxonomic richness, taxonomic diversity varied by season for the full ($F_{3, 80} = 12.267$, $P < 0.001$) and revised ($F_{3, 80} = 10.999$, $P < 0.001$) sites (Table 1, Figure 2). There were no further main or interactive effects of season, order, or reach on taxonomic diversity for either

the full or revised sites (Table 1, Figure 2). Spatial dissimilarity did not vary by any of the main or interactive effects of season and order for both the full and revised sites (Table 1, Figure 3). Additionally, temporal variation did not vary by the main effects of or interaction between reach and order for the full and revised sites (Table 1, Figure 4).

Analyses by Swan and Brown (2017, 2018) overestimated some effect sizes for the local diversity and spatial dissimilarity analyses (Table 3). Large effect sizes were observed for season in the local diversity ($\eta^2 = 0.1405$) and spatial dissimilarity analyses ($\eta^2 = 0.1817$) by Swan and Brown (2017, 2018), but there was a medium effect size of season for the full ($\eta^2 = 0.0756$) and revised ($\eta^2 = 0.0826$) sites analyses of local diversity; a medium effect size of season was observed for the full sites ($\eta^2 = 0.0999$) and a small effect size was observed for the revised sites ($\eta^2 = 0.0599$) for the spatial dissimilarity analyses. Similar effect sizes for the local diversity analyses were observed for order for the Swan and Brown (2017, 2018) analyses ($\eta^2 = 0.1255$) and the full sites ($\eta^2 = 0.1143$) analysis, but the revised sites analysis had a small effect size of order ($\eta^2 = 0.0006$). Noticeably, equivalent and small effect sizes of reach were observed for the local diversity analyses for all analyses (Table 3). In contrast to local diversity and spatial dissimilarity, Swan and Brown (2017, 2018) frequently underestimated effect sizes for the temporal variability analyses (Table 3). A medium effect size of order ($\eta^2 = 0.1203$) and large effect sizes of reach and the order-by-reach interaction were observed in the revised sites analysis (reach $\eta^2 = 0.2779$, order-by-reach $\eta^2 = 0.4026$), with the largest effect size from all analyses and community diversity metrics derived from the order-by-reach interaction in the revised sites analysis ($\eta^2 = 0.4026$). Although this effect was not statistically significant ($P = 0.066$), it suggests that restoration treatments could have an effect that is dependent on network position, but the statistical power was insufficient.

Differences in significant main effects or interactions within the full and revised sites in the re-analysis did not seem to be the result of increased variation in the revised sites. In fact, variance, as measured by 95% confidence intervals, was either similar or even reduced for each of local diversity, spatial dissimilarity, and temporal variability for the revised sites compared to the full sites (Figures 1-4). It is therefore unlikely that that revised sites analysis was unable to detect effects due to increased variation and more likely due to reduced statistical power associated with a smaller sample size or the true lack of an effect of restoration on different facets of biodiversity in this system.

Effectiveness of Local Restoration

I hypothesized that stream-channel manipulations would have a more consistent effect relative to the effect of riparian reforestation between headwaters and mainstems, with stronger effects of restoration in headwaters relative to mainstems. As there were no significant effects of restoration on any of the community metrics between headwaters and mainstems, this hypothesis could be invalid or, at a minimum, revised and re-tested in restoration experiments. I should note that I was unable to directly test this hypothesis because I was re-analyzing data from a previous study and the experimental design precluded any robust test to isolate the effects; however, the hypothesis was intended to guide criteria for site selection and reduce variation in restoration treatments among sites and not to necessarily or strictly compare the effects in-channel manipulations and riparian reforestation treatments on biodiversity in restored streams. Despite these limitations, there is some evidence of a large effect of the revised sites on temporal variation in community composition, with this effect dependent on network position, but the analysis was lacking sufficient statistical power to combine statistical significance with ecological relevance; further evaluation of this interactive effect of restoration and network

position on temporal variability could be an efficacious avenue for bridging metacommunity ecology and restoration efforts in rivers and streams and increasing positive outcomes.

Regarding evidence for stream-channel manipulations and other treatments for effective restoration, previous research suggests local habitat manipulations are ineffective for structuring communities and increasing biodiversity (Palmer et al. 2010). An emerging hypothesis is that local factors, such as habitat complexity and water quality, are overwhelmed by regional factors, such as dispersal and position within the larger network (Heino 2013, Tonkin et al. 2014). Given restoration did not have a statistically-significant effect on any diversity measure of communities in either headwaters or mainstems and that the majority of effect sizes of restoration were small-to-medium (Table 3), this could suggest either restoration was wholly inadequate for both headwaters and mainstems or that the larger network and regional species pool were already degraded (Sundermann et al. 2011), overwhelming any mitigating effects of restoration.

Restoration Ecology & Experimental Design

Restoration of the streams was done in isolation of the study design and prior to data collection, resulting in variation in the types of treatments applied to and the time since restoration of the streams (Swan and Brown 2017). Although Swan and Brown (2017) noted this limitation of their study, they did not acknowledge they could have better controlled for this variation by setting strict criteria for site selection and inclusion, which informed my hypothesis and was the foundation for my re-analysis. This concern was briefly acknowledged in the erratum (Swan and Brown 2018), where the data quality control process removed sites if they only received riparian reforestation treatments without at least one of either the bank stabilization or in-channel manipulation treatments; however, Swan and Brown (2017, 2018) proceeded to analyze data from sites receiving any combination of restoration treatments, despite suggesting

that in-stream modification treatments would have stronger effects on communities relative to riparian reforestation (Swan and Brown 2018). Setting a more stringent criterion for site inclusion, as was done in this re-analysis of the revised sites, would have reduced the variation in the applied restoration treatments and provided a more balanced experimental design.

The inconsistent application of restoration treatments prohibited a robust evaluation that could have been possible with a factorial experiment; therefore, the singular and interactive effects of the restoration treatments in the study system remain untested. This further complicates the indiscriminate usage of “restoration” by Swan and Brown (2017, 2018) and in the full sites analysis presented here, as the underlying mechanism of restoration on the stream invertebrate communities remains an unknown quantity. Identifying how individual and combinations of restoration treatments affect stream communities would provide valuable insight for maximizing the effectiveness of restoration efforts. In the absence of this knowledge, reducing the variation in which restoration treatments were applied to the streams, as done with revised sites analysis, arguably would have been a better avenue.

Statistical Inconsistencies

Restoration was not found to have a significant effect on local diversity, spatial dissimilarity, or temporal variability of stream invertebrate communities between paired restored and unrestored reaches in headwaters and mainstems. These results presented here, not exact values of test statistics but in terms of interpretation, directly contradict the results presented in the original paper (Swan and Brown 2017) and in the erratum (Swan and Brown 2018; Table 3). This is concerning, as any data management and analytical errors in the original paper were supposedly resolved in the erratum (Swan and Brown 2018); however, the discrepancies can be partially explained by the erroneous reporting and implementation of statistical analyses. First, as

was noted above, Swan and Brown (2017, 2018) analyzed an unbalanced design with unequal variance using an ANOVA with Type I sums of squares, when transformations to response variables were necessary to better meet test assumptions and Type III sums of squares were more appropriate for investigating the main and interactive effects (Shaw and Mitchell-Olds 1993, Quinn and Keough 2002). Second, the fitting of the random effects in the ANOVAs was incorrect. Swan and Brown (2018) reported fitting stream identity as a random effect. With the R code provided with the erratum (Swan and Brown 2018: Supporting Information), each site-by-reach combination was fitted as a random effect, despite adjacent and restored reaches being separated by < 10 m (Swan and Brown 2017). Fitting stream identity alone would have been a more appropriate fitting of a random effect that reflected the non-independence of the reaches at such small spatial scales, and, importantly, an accurate implementation of the written methods. Additionally, with the evidence available, only the local diversity analysis had a random effect fitted with the model, despite a random effect being applicable to all analyses based on the experimental design. Third, removal of the restored site without a paired observation is necessary for the fundamental goal of the study: comparing community diversity between paired restored and adjacent reaches across headwater and mainstem streams. Without removing the site, comparisons would be made to an unpaired reach, violating the experimental design and central goal of the study.

Finally, and of greatest concern, is the wholesale disagreement between the reported analytical procedure and what was actually conducted when analyzing temporal variability. Temporal variability was reportedly quantified as the mean distance to the group centroid (Anderson et al. 2006) after applying a Bray-Curtis dissimilarity index on an untransformed abundance matrix (Swan and Brown 2017). Results presented in the erratum were actually

derived from the spatial median after a Jaccard index was applied to a presence-absence matrix (Swan and Brown 2018: Supporting Information); in the initial study, Swan and Brown (2017) stated that a Jaccard index applied to a presence-absence matrix would produce stronger results than a Bray-Curtis index applied to an abundance matrix. No random effect of stream identity was fitted for this ANOVA, although it would have been appropriate given the study design (Quinn and Keough 2002). Most importantly, none of the changes to the analytical procedure were reported in the erratum, and these alterations were only found upon evaluation of the provided R code (Swan and Brown 2018: Supporting Information). Without consulting the supporting information or if no R code was provided, it would have been assumed the results presented in the erratum (Swan and Brown 2018) were derived from the analytical procedure described in the original study (Swan and Brown 2017), just with the corrected dataset; this assumption would have been incorrect.

Ecological Implications & Prospective Suggestions

Based on my re-analysis, I have concluded that, given restoration had no effect on any community diversity metric, local restoration of streams can be ineffective if (1) both dispersal and habitat quality are structuring biodiversity (Heino et al. 2015, Smith et al. 2015, Downes et al. 2017) and (2) the larger regional community is already degraded (Sundermann et al. 2011). Future research projects should experimentally- and factorially-manipulate different environmental factors to evaluate the relative impact and effectiveness of restoration techniques, which could allow for the identification of how in-stream and riparian reforestation treatments affect stream biodiversity. More broadly, restoration of individual reaches of rivers and streams might be insufficient to reach objectives of restoration projects (Bernhardt and Palmer 2011, Sundermann et al. 2011). Streams and rivers will likely have great restoration success if a

regional approach is taken (Bernhardt and Palmer 2011), whereby heterogeneity among rivers and streams in ecological, geographic, and spatial context is incorporated (Palmer et al. 2010, 2014, Booth et al. 2016). In this respect, I agree with Swan and Brown (2017, 2018), along with previous research, that local manipulations are insufficient, and a regional perspective is needed for more effective restoration in rivers and streams.

Acknowledgements

I am grateful for the comments provided by Karl Cottenie, Mariana Perez Rocha, Eric Harvey that improved the manuscript.

Literature Cited

- Altermatt, F. 2013. Diversity in riverine metacommunities: A network perspective. *Aquatic Ecology* 47:365–377.
- Anderson, M. J., K. E. Ellingsen, and B. H. McArdle. 2006. Multivariate dispersion as a measure of beta diversity. *Ecology Letters* 9:683–693.
- Bernhardt, E. S., and M. A. Palmer. 2011. River restoration: the fuzzy logic of repairing reaches to reverse catchment scale degradation. *Ecological Applications* 21:1926–1931.
- Booth, D. B., A. H. Roy, B. Smith, and K. A. Capps. 2016. Global perspectives on the urban stream syndrome. *Freshwater Science* 35:412–420.
- Cohen, J. 1973. Eta-Squared and Partial Eta-Squared in Fixed Factor Anova Designs. *Educational and Psychological Measurement* 33:107–112.
- Collins, K. E., C. Doscher, H. G. Rennie, and J. G. Ross. 2013. The effectiveness of riparian ‘restoration’ on water quality: A case study of lowland streams in Canterbury, New Zealand. *Restoration Ecology* 21:40–48.
- Downes, B. J., J. Lancaster, A. Glaister, and W. D. Bovill. 2017. A fresh approach reveals how

332 dispersal shapes metacommunity structure in a human-altered landscape. *Journal of*
333 *Applied Ecology* 54:588–598.

334 Grant, E. H. C., W. H. Lowe, and W. F. Fagan. 2007. Living in the branches: Population
335 dynamics and ecological processes in dendritic networks. *Ecology Letters* 10:165–175.

336 Heino, J. 2013. Does dispersal ability affect the relative importance of environmental control and
337 spatial structuring of littoral macroinvertebrate communities? *Oecologia* 171:971–980.

338 Heino, J., A. S. Melo, T. Siqueira, J. Soininen, S. Valanko, and L. M. Bini. 2015.
339 Metacommunity organisation, spatial extent and dispersal in aquatic systems: Patterns,
340 processes and prospects. *Freshwater Biology* 60:845–869.

341 Heino, J., T. Muotka, and R. Paavola. 2003. Determinants of macroinvertebrate diversity in
342 headwater streams: Regional and local influences. *Journal of Animal Ecology* 72:425–
343 434.

344 Leibold, M. A., M. Holyoak, N. Mouquet, P. Amarasekare, J. M. Chase, M. F. Hoopes, R. D.
345 Holt, J. B. Shurin, R. Law, D. Tilman, M. Loreau, and A. Gonzalez. 2004. The
346 metacommunity concept: a framework for multi-scale community ecology. *Ecology*
347 *Letters* 7:601–613.

348 Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R.
349 B. O'Hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, and H. Wagner.
350 2019. vegan: Community Ecology Package. R package version 2.5-4. <[https://CRAN.R-](https://CRAN.R-project.org/package=vegan)
351 [project.org/package=vegan](https://CRAN.R-project.org/package=vegan)>

352 Palmer, M. A., H. L. Menninger, and E. Bernhardt. 2010. River restoration, habitat heterogeneity
353 and biodiversity: A failure of theory or practice? *Freshwater Biology* 55:205–222.

354 Palmer, M. A., K. L. Hondula, and B. J. Koch. 2014. Ecological Restoration of Streams and
 355 Rivers: Shifting Strategies and Shifting Goals. *Annual Review of Ecology, Evolution,*
 356 *and Systematics* 45:247–269.

357 Pinheiro J., D. Bates, S. DebRoy, D. Sarkar, and R Core Team. 2019. nlme: Linear and
 358 Nonlinear Mixed Effects Models. R package version 3.1-139. <[https://CRAN.R-](https://CRAN.R-project.org/package=nlme)
 359 [project.org/package=nlme](https://CRAN.R-project.org/package=nlme)>

360 Quinn, G. P., and M. J. Keough. 2002. Experimental design and data analysis for biologists.
 361 Cambridge University Press, Cambridge, UK.

362 R Development Core Team. 2019. R: a language and environment for statistical computing. R
 363 Foundation for Statistical Computing, Vienna, Austria. <<https://cran.r-project.org/>>

364 Rosi-Marshall, E. J., and J. B. Wallace. 2002. Invertebrate food webs along a stream resource
 365 gradient. *Freshwater Biology* 47:129-141.

366 Shaw, R. G., and T. Mitchell-Olds. 1993. ANOVA for unbalanced data: An overview. *Ecology*
 367 74:1638–1645.

368 Smith, R. F., P. D. Venugopal, M. E. Baker, and W. O. Lamp. 2015. Habitat filtering and adult
 369 dispersal determine the taxonomic composition of stream insects in an urbanizing
 370 landscape. *Freshwater Biology* 60:1740–1754.

371 Sundermann, A., S. Stoll, and P. Haase. 2011. River restoration success depends on the species
 372 pool of the immediate surroundings. *Ecological Applications* 21:1962-1971.

373 Swan, C. M., and B. L. Brown. 2017. Metacommunity theory meets restoration: isolation may
 374 mediate how ecological communities respond to stream restoration. *Ecological*
 375 *Applications* 27:2209-2219.

376 Swan, C. M., and B. L. Brown. 2018. Erratum for: Metacommunity theory meets restoration:
377 isolation may mediate how ecological communities respond to stream restoration.
378 Ecological Applications 28:1370–1371.

379 Tonkin, J. D., S. Stoll, A. Sundermann, and P. Haase. 2014. Dispersal distance and the pool of
380 taxa, but not barriers, determine the colonisation of restored river reaches by benthic
381 invertebrates. Freshwater Biology 59:1843-1855.

382 Vannote, R. L., G. W. Minshall, K. W. Cummins, J. R. Sedell, and C. E. Cushing. 1980. The
383 river continuum concept. Canadian Journal of Fisheries and Aquatic Sciences 37:130-
384 137.

385 Wahl, C. M., A. Neils, and D. Hooper. 2013. Impacts of land use at the catchment scale constrain
386 the habitat benefits of stream riparian buffers. Freshwater Biology 58:2310–2324.

387 Zuur, A. F., E. N. Ieno, and C. S. Elphick. 2010. A protocol for data exploration to avoid
388 common statistical problems. Methods in Ecology and Evolution 1:3–14.

389 Tables

390 Table 1: ANOVA results for taxonomic richness, taxonomic diversity, spatial dissimilarity, and
391 temporal variability. Models were fitted using restricted maximum likelihood and with Type III
392 sums of squares for estimating main and interactive effects of factors.

Source of Variation	Full Sites				Revised Sites			
	numDF	denDF	F	P	numDF	denDF	F	P
Taxonomic Richness								
Season	3	73	1.991	0.123	3	47	1.411	0.251
Order	1	11	1.419	0.259	1	7	0.004	0.951
Reach	1	73	0.730	0.396	1	47	0.398	0.531
Season x Order	3	73	1.641	0.187	3	47	0.553	0.649
Season x Reach	3	73	0.045	0.987	3	47	0.143	0.934
Order x Reach	1	73	0.308	0.581	1	47	1.375	0.247
Season x Order x Reach	3	73	0.083	0.969	3	47	0.149	0.930
Taxonomic Diversity								
Season	3	73	12.267	< 0.001	3	47	10.999	< 0.001
Order	1	11	2.073	0.178	1	7	0.253	0.630
Reach	1	73	0.088	0.768	1	47	0.085	0.772
Season x Order	3	73	2.477	0.068	3	47	1.272	0.295
Season x Reach	3	73	0.323	0.809	3	47	0.284	0.837
Order x Reach	1	73	0.139	0.710	1	47	0.917	0.343
Season x Order x Reach	3	73	0.313	0.816	3	47	0.149	0.930
Spatial Dissimilarity								
Season	3	31	1.146	0.346	3	20	0.425	0.737
Order	1	11	0.062	0.808	1	7	0.007	0.934
Season x Order	3	31	1.356	0.275	3	20	0.306	0.821
Temporal Variability								
Order	1	11	0.030	0.866	1	7	0.957	0.361
Reach	1	11	1.906	0.195	1	7	2.693	0.145
Order x Reach	1	11	2.054	0.180	1	7	4.718	0.066

393

Table 2: ANOVA results for the full and reduced models of taxonomic richness proposed by Swan and Brown (2017, 2018). Models were fitted using restricted maximum likelihood and with Type III sums of squares for estimating main and interactive effects of factors. The unpaired sample was not removed from the data prior to fitting the full and reduced model ANOVAs.

Source of Variation	Full Model				Reduced Model			
	numDF	denDF	F	P	numDF	denDF	F	P
Season	3	74	2.478	0.068	3	80	4.360	0.007
Order	1	11	0.925	0.357	1	11	1.579	0.235
Reach	1	74	1.409	0.239	1	80	4.844	0.031
Season x Order	3	74	2.359	0.078	3	80	4.105	0.009
Season x Reach	3	74	0.152	0.928	N/A	N/A	N/A	N/A
Order x Reach	1	74	0.979	0.326	1	80	3.530	0.064
Season x Order x Reach	3	74	0.071	0.976	N/A	N/A	N/A	N/A

Note: N/A indicates a factor or interaction that was removed in the reduced model.

Table 3: Comparison of ANOVA results and effect sizes (η^2) between the initial study (Swan and Brown 2017), the erratum (Swan and Brown 2018), and if results are consistent upon re-analysis in the full (Full) or revised (Revised) sites analyses. Support is denoted as: Yes = consistent with both the initial study and the erratum; No = inconsistent with both the initial study and erratum; NC = no comparisons can be made as the factor or interaction is missing from the previous analysis. Spatial dissimilarity was not re-analyzed in the erratum, so results are only provided for the initial study. Bold values in the Supported column indicate differences in statistical significance between the initial study and/or erratum and the re-analysis.

Source of Variation	Swan and Brown 2017		Swan and Brown 2018		Full Sites		Revised Sites		Supported	
	P	η^2	P	η^2	P	η^2	P	η^2	Full	Revised
Local Diversity										
Season	NULL	NULL	< 0.001	0.1405	0.123	0.0756	0.251	0.0826	No	No
Order	NULL	NULL	0.18	0.1255	0.259	0.1143	0.951	0.0006	Yes	Yes
Reach	NULL	NULL	0.22	0.0571	0.396	0.0099	0.531	0.0084	Yes	Yes
Season x Order	NULL	NULL	0.086	0.1334	0.187	0.0632	0.649	0.0341	No	No
Season x Reach	NULL	NULL	N/A	N/A	0.987	0.0018	0.934	0.0091	NC	NC
Order x Reach	NULL	NULL	0.064	0.0423	0.581	0.0042	0.247	0.0284	Yes	Yes
Season x Order x Reach	NULL	NULL	N/A	N/A	0.969	0.0034	0.930	0.0094	NC	NC
Spatial Dissimilarity										
Season	0.015	0.1817	N/A	N/A	0.346	0.0999	0.361	0.0599	No	No
Order	0.022	0.0972	N/A	N/A	0.808	0.0056	0.145	0.0011	No	No
Season x Order	0.220	0.0796	N/A	N/A	0.275	0.1160	0.066	0.0439	Yes	Yes
Temporal Variability										
Order	NULL	NULL	0.36	0.0385	0.866	0.0027	0.361	0.1203	Yes	Yes
Reach	NULL	NULL	0.095	0.1214	0.195	0.1477	0.145	0.2779	Yes	Yes
Order x Reach	NULL	NULL	0.020	0.2234	0.180	0.1573	0.066	0.4026	No	No

Note: N/A indicates a factor or interaction that was removed in the reduced model or not analyzed by Swan and Brown (2018). NULL indicates a deprecated P-value and an effect size that was not calculated as the model was corrected by Swan and Brown (2018).

Figures




Figure 1: Plots of taxonomic richness in headwaters (A and C) and mainstems (B and D).

Taxonomic richness is reported for restored (triangles) and adjacent (circles) sites; values from the full sites are reported in black (A and B), while those from the revised sites are reported in grey (C and D). Points represent mean \pm 95% CI.




Figure 2: Plots of taxonomic diversity, calculated as Shannon's diversity, in headwaters (A and C) and mainstems (B and D). Taxonomic diversity is reported for restored (triangles) and adjacent (circles) sites; values from the full sites are reported in black (A and B), while those from the revised sites are reported in grey (C and D). Points represent mean \pm 95% CI.




Figure 3: Plots of spatial dissimilarity between paired restored and adjacent reaches. Estimates are reported for headwaters (HW, circles) and mainstems (MS, triangles); values from the full sites are reported in black (A), while those from the revised sites are reported in grey (B). Points represent mean \pm 95% CI.




Figure 4: Plots of temporal variability. Estimates are reported for headwaters (HW, circles) and mainstems (MS, triangles); values from the full sites are reported in black (A), while those from the revised sites are reported in grey (B). Points represent mean \pm 95% CI.