Diversity begets diversity in microbiomes

One Sentence Summary: Making use of the most comprehensive biodiversity catalogue, the Earth Microbiome Project, we show that prokaryote diversification significantly correlates with microbiome diversity, pointing to the importance of species interactions in driving microbial evolution and community structure.

Authors: Naïma Madi[1], Michiel Vos[2], Pierre Legendre[1] and B. Jesse Shapiro[1*]

1. Departement de sciences biologiques, Universite de Montreal, Canada
2. European Centre for Environment and Human Health, University of Exeter, Penryn, UK

*correspondence: jesse.shapiro@umontreal.ca

keywords: microbiome, diversification, evolution, ecology, Earth Microbiome Project, 16S rRNA

Abstract

Microbes are embedded in complex microbiomes where they engage in a wide array of

interspecific interactions. However, how these interactions shape diversification, and

ultimately biodiversity, is not well understood. Two competing hypotheses have been put

forward to explain how species interactions could influence diversification rates.

Ecological Controls (EC) predicts a negative diversity-diversification relationship, where

the evolution of novel types becomes constrained as available niches become filled.

Diversity Begets Diversity (DBD) predicts a positive relationship, with diversity

promoting diversification via niche construction and other species interactions. Using the

Earth Microbiome Project, the largest standardized survey of global biodiversity to date,

we provide support for DBD as the dominant driver of microbiome diversity. Only in the

most diverse microbiomes does DBD reach a plateau, presumably because of

increasingly saturated niche space. Genera that are strongly associated with particular

biomes show a stronger DBD relationship than non-residents, consistent with prolonged

evolutionary interactions driving diversification. Genera with larger genomes also

experience a stronger DBD response, which could be due to a higher potential for

metabolic interactions and niche construction. Our results provide evidence that

microbiome diversity – and its potential for future diversification – is crucially shaped by

species interactions.

Main text

The majority of the Tree of Life consists of microbes (*1–3*) and the functioning of

all Earth's ecosystems is reliant on diverse microbial communities (*4*). High-throughput

16S rRNA gene amplicon sequencing studies continue to yield unprecedented insight into

the taxonomic richness of microbiomes (e.g. (*5*, *6*)), and abiotic drivers of community

composition (e.g. pH (*7*, *8*)) are increasingly uncovered. Microbe-microbe (biotic)

interactions can also be important in determining community composition (*9*), but

comparatively little is known about how such interactions (e.g. cross-feeding (*10*) and

toxin-mediated interference competition (*11*, *12*)) shape microbiome diversity.

The dearth of studies exploring how microbial interactions influence

diversification stands in marked contrast to a long research tradition on biotic controls of

plant and animal diversity (*13*, *14*). In an early study of 49 animal (vertebrate and

invertebrate) community samples, Elton plotted the number of species versus the number

of genera and observed a ~1:1 ratio in each individual sample, but a ~4:1 ratio when all

samples were pooled (*14*). Elton took this observation as evidence for competitive

exclusion, preventing related species, more likely to overlap in niche space, to co-exist.

This concept, more recently referred to as niche filling or "ecological controls" (EC) (*15*)

predicts speciation (or, more generally, diversification) rates to decrease with standing

species diversity because available niches become filled over time (*16*). In contrast, the

Diversity Begets Diversity (DBD) model predicts that diversity favors further

diversification when species interactions create novel niches (*17*, *18*). For example, niche

construction (i.e. the physical, chemical or biological alteration of the environment by

microbes) could influence the evolution of the species constructing the niche, and/or that

of co-occurring species (*19, 20*). Evidence for the action of EC vs. DBD in plant and

animal communities has been mixed (*18, 21-24)*, and models suggest that intraspecific

competition can promote diversification, while interspecific competition and predation

inhibit diversification (*25*). Laboratory evolution experiments tracking the diversification

of a focal bacterial lineage in communities of varying complexity have also yielded

contradicting results, with support for EC, DBD, or intermediate scenarios (*26-30*).

To test whether natural microbial communities are shaped by EC or DBD

dynamics, we used 2,000 microbiome samples each rarefied to 5,000 16S Amplicon

Sequence Variants (ASVs; exact, error-corrected DNA sequences, directly comparable

across samples (*31*)) archived in the Earth Microbiome Project (EMP), the largest

available repository of biodiversity based on standardized sampling and sequencing

protocols (*32*). All samples used were subjected to rarefaction, as diversification rate

estimates are highly sensitive to sampling effort (*33,34*). In the absence of time series or a

fossil record, microbial diversification must be inferred from extant diversity. Rather than

taking a phylogenetic approach requiring complex assumptions (*35, 36*), we use the

equivalent of the Species:Genus (S:G) ratios that Elton used three quarters of a century

ago (*14*) to infer bacterial diversification rates. We use a range of taxonomic ratios

(ASV:Genus, Genus:Family, Family:Order, Order:Class, and Class:Phylum) as proxies

for diversification of a focal lineage of interest, from shallow to deep evolutionary time,

and plot these as a function of the number of non-focal lineages (Genera, Families,

Orders, Classes, and Phyla, respectively) with which the focal lineage could interact. A

negative relationship is consistent with the EC hypothesis, whereas a positive relationship

is consistent with the DBD hypothesis (**Figure 1**).

We used generalized linear mixed models (GLMMs) (*37*) to determine how the

diversification of a focal lineage (*e.g.* its ASV:Genus ratio) is affected by the diversity of

other lineages (*e.g.* non-focal genera) in the community. The effects of environment (as

defined by the EMP Ontology 'level 3 biomes;' Methods) and the identity of the focal

lineage were included by fitting these as random effects on the slope and intercept. The

submitting laboratory (identified by the principal investigator) and the EMP unique

sample identifier (i.e. if two taxa were part of the same sample) were also included as

random effects. The DBD model was supported across all taxonomic ratios, which all had

significantly positive slopes fitting the diversity-diversification relationship (**Table 1,**

Supplementary Data file 1 Section 1). In some cases, the variation due to lineage,

lineage*environment, or environment* laboratory interactions could reverse the sign of

the slope (**Table 1**), but the vast majority of slope estimates are positive, supporting DBD

(**Figure S1**).

As an illustration, **Figure 2** shows diversification within the Proteobacteria

(Classes:Phylum) as a function of all other Phyla in each sample across 17 biomes. All

slopes were significantly positive ($P < 0.001$ after Bonferroni correction), except in

hypersaline and non-saline sediments. The strongest DBD effect was found in animal

proximal gut microbiomes (slope=0.3, P=1.03e−08). For each taxonomic ratio, the three

most prevalent taxa followed positive slopes in most environments (**Figures S2-S6**), with

only a few instances of significantly negative slopes (Alphaproteobacteria, Rhizobiales,

Flavobacteriales, Flavobacteriaceae, Planctomyces in the plant rhizosphere, and

Clostridium in non-saline sediment). Overall, the majority of significant slope estimates

were positive (81.7-99.5% depending on the taxonomic ratio considered), favoring DBD

as the dominant (but not universal) model supported by the data (**Table S1**).

To test for any potential confounding effects of data structure or sampling bias,

we sought to remove any patterns of co-occurrence between ASVs in the same sample

via permutation. We took 2,000 simulated samples by selecting from the overall

distribution of 155,002 unique ASVs across all samples, weighted by their abundance

(total number of sequence counts). This resulted in a slightly negative diversity-

diversification relationship (slope = −0.002; Pearson correlation = −0.61; P<2.2.e−16;

Figure S7), indicating that the observed positive relationships (**Table 1; Figure 2**) are

not the effect of data structure.

The taxonomic approach considered thus far has certain limitations. First, not all

taxonomic ranks have the same phylogenetic depth (*38*) and not all named taxa are

monophyletic (*39*). Second, unclassified ASVs had to be excluded from taxonomic

analysis. Although these limitations are not expected to bias the results toward either EC

or DBD, we sought to validate the results with a taxonomy-independent approach,

clustering ASVs at decreasing levels of nucleotide identity, from 100% identical ASVs

down to 75% identity (roughly equivalent to phyla (*40*)). We estimated diversification as

the mean number of descendants per cluster (e.g. number of 100% clusters per 97%

cluster) and plotted this against the total number of non focal clusters (97% identity in

this example). For each of the six nucleotide divergence ratios tested, the relationship

between diversity and diversification remained positive (**Figure S8**), consistent with

DBD and suggesting that the taxonomic analyses were largely unbiased.

To exclude the possibility that our results were driven by abiotic confounders, we

repeated the taxonomic analysis on a subset of 192 EMP samples for which

measurements of four important abiotic drivers of diversity, temperature, pH, latitude,

and elevation (*7*, *8*, *15*, *41*) were available. We fitted a GLMM with diversification rate

as the dependent variable, and with the number of non-focal lineages, the four abiotic

factors and their interactions as predictors (fixed effects). As in the full dataset (**Table 1**),

diversification was positively associated with diversity at all taxonomic ratios (**Table S2**).

As expected, certain abiotic factors, alone or in combination with diversity, had

significant effects on diversification. However, the effects of abiotic factors were always

weaker than the effect of community diversity (**Table S2**; **Supplementary Data file 1**

Section 2). Although only a small subset of abiotic factors was considered, this analysis

suggests that the DBD trend is unlikely to be mainly driven by variation in the abiotic

environment.

The DBD hypothesis rests on the premise that species interactions drive

diversification (*15*, *18*). We therefore expect that lineages that are more tightly associated

with a specific biome (i.e. long-term residents) are more likely to have had a long history

of interaction and thus are more likely to experience DBD than lineages that are not

tightly associated with that biome (i.e. poorly adapted migrants or broadly adapted

generalists). To test this prediction, we clustered environmental samples by their genus-

level community composition using fuzzy *k*-means clustering (**Figure 3A**). Three clusters

were identified, which we refer to as 'animal-associated', 'saline,' and 'non-saline'

clusters. All three clusters included some unexpected outliers (*e.g.* plant corpus grouping

with animals), but were generally intuitive and consistent with known distinctions

between host-associated vs. free-living (*32*), and saline vs. non-saline communities (*42*).

Resident genera were defined as those with a strong preference for a particular

environment cluster, using indicator species analysis (permutation test, *P*<0.05; **Figure**

3A; **Figure S9**; **Supplementary Data file 2**), and genera without a strong preference

were considered generalists. For each environment cluster, we ran a GLMM with resident

genus-level diversity (number of non-focal genera) as a predictor of diversification

(ASV:Genus ratio) for residents, generalists, or migrants (residents of one cluster found

in a different cluster) (**Supplementary Data file 1 Section 3**). Resident diversity had no

significant effect on the diversification of generalists (z=0.646, *P*=0.518; z=0.279,

P=0.780; z=0.347, *P*=0.729, respectively for animal-associated, saline and non-saline

clusters), but did significantly increase resident diversification (z=7.1, *P*= 1.25e-12;

z=3.316, *P*=0.0009; z=7.109, *P*=1.17e-12, respectively). Resident diversity significantly

decreased migrant diversification in saline (z=-3.194, *P*=0.0014) and non-saline

environment clusters (z=-2.840, *P*=0.0045), but had no significant effect in the animal-

associated cluster (z=-0.566, *P*=0.571) (**Figure 3B**). These results suggest that diversity

begets diversification among lineages sharing the same environment over a long

evolutionary time period, but that this is not the case for lineages that do not consistently

occur in the same microbiome and presumably interact less frequently. The

diversification of migrants in a new environment might even be impeded, presumably

because most niches are already occupied by residents.

The positive effect of diversity on diversification should eventually reach a

plateau as niches, including those constructed by biotic interactions, become saturated. In

the animal distal gut, a relatively low-diversity biome, we observed a strong linear DBD

relationship at most sequence identity ratios; in contrast, the more diverse soil biome

clearly attained a plateau (**Figure S10**). To further test the hypothesis that increasingly

diverse microbiomes experience weaker DBD due to saturated niche space, we used a

GLMM to model Family:Order, Order:Class and Class:Phylum diversification (as DBD

slope variation by environment was statistically significant for these taxonomic ratios;

Table 1). We included the interaction between diversity and environment type as a fixed

effect. Consistent with our hypothesis, DBD slopes were significantly more positive in

less diverse (often host-associated) biomes (**Figure 4A**, **Figure S11, Supplementary**

Data file 1 Section 4). We also found evidence for a DBD plateau in the nucleotide

identity analyses, in which cubic models provided slightly better fits than linear models

across all biomes (**Figure S8**).

The Black Queen hypothesis posits that microbes embedded in complex

communities can exploit the production of extracellular public goods produced by other

species, resulting in selection for loss of genes encoding these goods – as long as the

essential trait is not lost from the community as a whole (*43*). Lineages that interact more

frequently with other lineages through such public good exploitation can be expected to

experience greater genome reduction and also to experience stronger DBD, because their

survival and diversification is dependent on other community members. To test whether

genome size is negatively correlated with DBD, we assigned genome sizes to 576 genera

for which at least one whole-genome sequence was available and added an interaction

term between genome size and diversity as a fixed effect to the GLMM (Methods).

Contrary to the Black Queen prediction, we observed a slight but significant positive

effect of genome size on the slope ($z=2.5$, *P*=0.01; **Figure 4B, Supplementary Data file**

1 Section 5). The positive relationship may even be stronger than estimated, because

genus-level genome size estimates are likely quite noisy. This result supports a model in

which biotic interactions (and resulting diversification) drive genome expansion (*e.g.*

through the accumulation of toxin- and resistance-gene diversity during antagonistic

coevolution (*11*)). Alternatively (or additionally), species with larger biosynthetic gene

repertoires and greater opportunity to engage in niche construction (*19*) could be more

prone to interact with other species, driving DBD.

Using 10 million individual marker sequences, we demonstrate a pervasive

positive relationship between prokaryotic diversity and diversification, which holds

across a broad range of microbiome types. The strength of the DBD relationship

dissipates with increasing microbiome diversity which might be due to niche saturation,

or potentially due to the fact that highly diverse communities prevent species from

reliably interacting with each other. DBD appears to be particularly strong among deeply

diverged lineages (*e.g.* phyla), suggesting the importance of DBD in the ancient

diversification of bacterial lineages and supporting the view that high taxonomic ranks

are ecologically coherent (*44*, *45*). The very early stages of diversification are

inaccessible at the resolution of 16S ASVs, but could be addressed in the future using

genomic and metagenomic methods. Due to the correlational nature of our data, it is not

possible to test whether the positive relationship between diversification and diversity is

primarily due to the creation of novel niches via biotic interactions and niche construction

(*20*), or potentially due to increased competition leading to adaptation to underexploited

resources (*12*, *29*). Regardless of the underlying mechanisms, our results clearly

demonstrate the importance of biotic interactions in shaping microbiome diversity, which

has important implications for their function and stability (*46, 47*). The answer to the

question 'why are microbiomes so diverse?' (*48*) might in a large part be because

microbiomes are so diverse (*23*).

Acknowledgements

We thank Luke Thompson for assistance obtaining EMP data and Zofia Ecaterina

Taranu, Vincent Fugère and Guillaume Larocque for advice on Generalized Linear

Mixed Models. We are also grateful to Steven Kembel and Tom Battin for critical

comments that improved the manuscript. **Funding:** This project was made possible by an

NSERC Discovery Grant and Canada Research Chair to BJS. **Authors contributions:**

Conceptualization: BJS, MV. Data curation: NM. Formal analysis: NM, MV, BJS.

Funding acquisition: BJS. Investigation: NM, MV, PL, BJS. Methodology: NM, MV, PL,

BJS. Resources: BJS, PL. Supervision: PL, BJS. Software: NM. Visualization: NM.

Writing original draft: NM, MV, BJS. Writing - review & editing: NM, MV, PL, BJS.

Competing interests: none to declare. **Data and materials availability:** All data is

available from the Earth Microbiome Project (ftp.microbio.me), as detailed in the

Methods. All computer code used for analysis are available at

https://github.com/Naima16/dbd.git.

Supplementary Materials

Materials and Methods

Tables S1 – S2

Fig S1 – S11

File 1. Full GLMM outputs.

File 2. Indicator species analysis.

Table 1. Diversity has a positive effect on diversification across taxonomic ratios.

The GLMMs showed statistically significant positive effect of diversity on

diversification. Each row reports the effect of diversity on diversification, as well as its

standard deviation, Wald z-statistic for its effect size and the corresponding *P*-value (left

section), or standard deviation on the slope for the significant random effects (right

section). SE=standard error, Env=environment type, Lin=lineage type, Lab=Principal

Investigator ID, Sample=Sample ID. Interactions are denoted as '*'. n.s.=not significant

(likelihood-ratio test).

	Slope (fixed effects)				Standard deviation on the slope (random effects)				
	Diversity	SE	z	*P*	Env	Lin	Lin*Env	Env*Lab	Sample
ASV: Genus	0.091	0.016	5.792	6.95e-09	n.s.	0.074	0.142	0.114	0.067
Genus: Family	0.047	0.008	5.911	3.41e-09	n.s.	0.071	0.07	0.039	n.s.
Family: Order	0.119	0.017	7.001	2.54e-12	0.023	0.094	0.092	0.106	n.s.
Order: Class	0.109	0.020	5.447	5.13e-08	0.05	0.141	0.078	0.051	n.s.
Class: Phylum	0.272	0.043	6.341	2.29e-10	0.119	0.174	0.119	0.114	n.s.

Figure 1. Contrasting the Diversity Begets Diversity (DBD) and Ecological Controls (EC) models of diversification. We consider the diversification of a focal lineage as a function of initial diversity present at the time of diversification. (A) For example, sample 1 contains one non-focal genus, and two ASVs diversify within the focal genus (point at x=1, y=2 in the plot). Sample 2 contains three non-focal genera, and four ASVs diversify within the focal genus (point at x=3, y=4). Tracing a line through these points yields a positive slope, supporting the Diversity Begets Diversification (DBD) model (red). (B) Alternatively, a negative slope would support the Ecological Controls (EC) model (blue line).

Figure 2. Diversification as a function of diversity across biomes in the phylum

Proteobacteria. Linear models are shown for diversification (the number of classes

within Proteobacteria, y-axis) as a function of diversity (the number of non-

proteobacterial phyla, x-axis) in each of the 17 environments (EMPO3 biomes). P-values

are Bonferroni corrected for 17 tests. Significant (*P* <0.05) models are shown with red

trend lines; non-significant (*P* > 0.05) trends are shown in blue.

Figure 3. Diversity begets diversification in resident versus non resident genera. (A)

PCA showing genera clustering into their preferred environment clusters. Circles

indicate genera and triangles indicate environments (EMPO 3 biomes). The three

environment clusters identified by fuzzy *k*-means clustering are: Non-saline (NS, blue),

saline (S, green) and animal-associated (purple). Resident genera were identified by

indicator species analysis. **(B) DBD in resident versus migrant genera across**

environment clusters. Results of GLMMs modeling diversification as a function of

diversity in resident, migrant, or generalist groups. The x-axis shows the standardized

number of non-focal resident genera (diversity); the y-axis shows the number of ASVs

per focal genus (diversification). Resident focal genera are shown in orange, migrant

focal genera in red, and generalist focal genera in black.

Figure 4. Ecological and evolutionary mechanisms to explain variation in the

strength of DBD. (A) DBD slope is higher in low-diversity (often host-associated)

microbiomes. The x-axis shows the mean number of phyla in each biome. On the y-axis,

DBD slope was estimated by the GLMM predicting diversification as a function of the

interaction between diversity and environment type at the Class:Phylum ratio

(**Supplementary Data file 1 Section 4.3**). The line represents a regression line; the

shaded area depicts 95% confidence limits of the fitted values. **(B) Positive correlation**

between genome size and DBD slope. Results are shown from a GLMM predicting

diversification as a function of the interaction between diversity and genome size at the

ASV:Genus ratio (**Supplementary Data file 1 Section 5**). The x-axis is genus-level

genome size in Mbp (min=0.97, max=14.78); the y-axis is DBD slope (the effect of

diversity on diversification). Vertical bars indicate 95% confidence limits of the fitted

values.

References

1. S. Sunagawa *et al.*, Ocean plankton. Structure and function of the global ocean microbiome. *Science*. **348**, 1261359 (2015).

2. P. Lapierre, J. P. Gogarten, Estimating the size of the bacterial pan-genome. *Trends Genet.* **25**, 107–110 (2009).

3. L. A. Hug *et al.*, A new view of the tree of life. *Nat. Microbiol.* **1**, 16048 (2016).

4. P. G. Falkowski, T. Fenchel, E. F. Delong, The microbial engines that drive Earth's biogeochemical cycles. *Science*. **320**, 1034–1039 (2008).

5. M. L. Sogin *et al.*, Microbial diversity in the deep sea and the underexplored "rare biosphere." *Proc. Natl. Acad. Sci. U. S. A*. **103**, 12115–12120 (2006).

6. S. Louca, F. Mazel, M. Doebeli, L. W. Parfrey, A census-based estimate of Earth's bacterial and archaeal diversity. *PLoS Biol.* **17**, e3000106 (2019).

7. C. L. Lauber, M. Hamady, R. Knight, N. Fierer, Soil pH as a predictor of soil bacterial community structure at the continental scale: a pyrosequencing-based assessment. *Appl. Environ. Microbiol.* **75**, 5111-5120 (2009).

8. J. F. Power *et al.*, Microbial biogeography of 925 geothermal springs in New Zealand. *Nat. Commun.* **9**, 2876 (2018).

9. D. M. Needham, J. A. Fuhrman, Pronounced daily succession of phytoplankton, archaea and bacteria following a spring bloom. *Nature Microbiology*, 1, 16005 (2016).

10. E. C. Seth, M. E. Taga, Nutrient cross-feeding in the microbial world. *Front. Microbiol.* **5**, 350 (2014).

11. T. L. Czárán, R. F. Hoekstra, L. Pagie, Chemical warfare between microbes promotes biodiversity. *Proc. Natl. Acad. Sci. U. S. A*. **99**, 786–790 (2002).

12. M. E. Hibbing, C. Fuqua, M. R. Parsek, S. B. Peterson, Bacterial competition: surviving and thriving in the microbial jungle. *Nat. Rev. Microbiol.* **8**, 15–25 (2010).

13. G. F. Gause, *The Struggle for Existence* (Courier Corporation, 2003).

14. C. Elton, Competition and the Structure of Ecological Communities. *J. Anim. Ecol.* **15**, 54–68 (1946).

15. D. Schluter, M. W. Pennell, Speciation gradients and the distribution of biodiversity. *Nature*. **546**, 48–55 (2017).

16. D. L. Rabosky, A. H. Hurlbert, Species richness at continental scales is dominated by ecological limits. *Am. Nat.* **185**, 572–583 (2015).

17. R. H. Whittaker, Evolution and Measurement of Species Diversity. *Taxon*. **21**, 213–251 (1972).

18. V. Calcagno, P. Jarne, M. Loreau, N. Mouquet, P. David, Diversity spurs diversification in
ecological communities. *Nat. Commun.* **8**, 15810 (2017).

19. M. San Roman, A. Wagner, An enormous potential for niche construction through bacterial
cross-feeding in a homogeneous environment. *PLoS Comput. Biol.* **14**, e1006340 (2018).

20. K. N. Laland, F. J. Odling-Smee, M. W. Feldman, Evolutionary consequences of niche
construction and their implications for ecology. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 10242–
10247 (1999).

21. T. D. Price *et al.*, Niche filling slows the diversification of Himalayan songbirds. *Nature*
509, 222-225 (2014).

22. D. L. Rabosky *et al.*, An inverse latitudinal gradient in speciation rate for marine fishes.
Nature. **559**, 392–395 (2018).

23. B. C. Emerson, N. Kolm, Species diversity can drive speciation. *Nature*. **434**, 1015–1017
(2005).

24. M. W. Palmer, T. A. Maurer, Does Diversity Beget Diversity? A Case Study of Crops and
Weeds. *J. Veg. Sci.* **8**, 235–240 (1997).

25. Pontarp Mikael, Petchey Owen L., Ecological opportunity and predator–prey interactions:
linking eco-evolutionary processes and diversification in adaptive radiations. *Proceedings of*
the Royal Society B: Biological Sciences. **285**, 20172550 (2018).

26. M. A. Brockhurst, N. Colegrave, D. J. Hodgson, A. Buckling, Niche occupation limits
adaptive radiation in experimental microcosms. *PLoS One*. **2**, e193 (2007).

27. P. Gómez, A. Buckling, Real-time microbial adaptive diversification in soil. *Ecol. Lett.* **16**,
650–655 (2013).

28. J. R. Meyer, R. Kassen, The effects of competition and predation on diversification in a
model adaptive radiation. *Nature*. **446**, 432–435 (2007).

29. A. Jousset, N. Eisenhauer, M. Merker, N. Mouquet, S. Scheu, High functional diversity
stimulates diversification in experimental microbial communities. *Sci Adv*. **2**, e1600124
(2016).

30. S. F. Bailey, J. R. Dettman, P. B. Rainey, R. Kassen, Competition both drives and impedes
diversification in a model adaptive radiation. *Proc. Biol. Sci.* **280**, 20131253 (2013).

31. B. J. Callahan, P. J. McMurdie, S. P. Holmes, Exact sequence variants should replace
operational taxonomic units in marker-gene data analysis. *ISME J.* **11**, 2639–2643 (2017).

32. L. R. Thompson *et al.*, A communal catalogue reveals Earth's multiscale microbial diversity.
Nature, nature24621 (2017).

33. N. J. Gotelli, R. K. Colwell, Quantifying biodiversity: procedures and pitfalls in the
measurement and comparison of species richness. *Ecol. Lett.* **4**, 379–391 (2001).

34. A. Z. Krug, D. Jablonski, J. W. Valentine, Species-genus ratios reflect a global history of
diversification and range expansion in marine bivalves. *Proc. Biol. Sci.* **275**, 1117–1123

(2008).

35. R. S. Etienne, A. L. Pigot, A. B. Phillimore, How reliably can we infer diversity-dependent
diversification from phylogenies? *Methods Ecol. Evol.* **7**, 1092–1099 (2016).

36. S. Louca *et al.*, Bacterial diversification through geological time. *Nat Ecol Evol.* **2**, 1458–
1467 (2018).

37. B. M. Bolker *et al.*, Generalized linear mixed models: a practical guide for ecology and
evolution. *Trends Ecol. Evol.* **24**, 127–135 (2009).

38. M. Vos, A species concept for bacteria based on adaptive divergence. *Trends Microbiol.* **19**,
1–7 (2011).

39. D. H. Parks *et al.*, A standardized bacterial taxonomy based on genome phylogeny
substantially revises the tree of life. *Nat. Biotechnol.* **36**, 996–1004 (2018).

40. K. T. Konstantinidis, J. M. Tiedje, Towards a genome-based taxonomy for prokaryotes. *J.*
Bacteriol. **187**, 6258–6264 (2005).

41. M. Delgado-Baquerizo *et al.*, A global atlas of the dominant bacteria found in soil. *Science.*
359, 320–325 (2018).

42. C. A. Lozupone, R. Knight, Global patterns in bacterial diversity. *Proc. Natl. Acad. Sci. U.*
S. A. **104**, 11436–11440 (2007).

43. J. J. Morris, R. E. Lenski, The Black Queen Hypothesis: evolution of dependencies through
adaptive gene loss. *MBio* **3**, e00036-12 (2012).

44. L. Philippot *et al.*, The ecological coherence of high bacterial taxonomic ranks. *Nat. Rev.*
Microbiol. **8**, 523–529 (2010).

45. J. B. H. Martiny, S. E. Jones, J. T. Lennon, A. C. Martiny, Microbiomes in light of traits: A
phylogenetic perspective. *Science.* **350**, aac9323 (2015).

46. K. Z. Coyte, J. Schluter, K. R. Foster, The ecology of the microbiome: Networks,
competition, and stability. *Science.* **350**, 663–666 (2015).

47. F. Pennekamp *et al.*, Biodiversity increases and decreases ecosystem stability. *Nature.* **563**,
109–112 (2018).

48. D. E. Dykhuizen, Santa Rosalia revisited: why are there so many species of bacteria?
Antonie Van Leeuwenhoek. **73**, 25–33 (1998).

49. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software, Articles.* **67**, 1–48 (2015).

50. X. A. Harrison *et al.*, A brief introduction to mixed effects modelling and multi-model
inference in ecology. *PeerJ.* **6**, e4794 (2018).

51. W. Forstmeier, H. Schielzeth, Cryptic multiple hypotheses testing in linear models:
overestimated effect sizes and the winner's curse. *Behav. Ecol. Sociobiol.* **65**, 47–55 (2011).

52. M. Dufrene, P. Legendre, Species Assemblages and Indicator Species: The Need for a
Flexible Asymmetrical Approach. *Ecol. Monogr.* **67**, 345–366 (1997).