Does diversity beget diversity in microbiomes?

Authors: Naïma Madi[1], Michiel Vos[2], Carmen Lia Murall[1],

Pierre Legendre[1] and B. Jesse Shapiro[1,3,4*]

1. Département de sciences biologiques, Université de Montréal, Canada
2. European Centre for Environment and Human Health, University of Exeter, Penryn, UK
3. Department of Microbiology and Immunology, McGill University, Canada
4. McGill Genome Centre, McGill University, Canada

*correspondence: jesse.shapiro@mcgill.ca

keywords: microbiome, evolution, ecology, Earth Microbiome Project, 16S rRNA

Abstract

Microbes are embedded in complex communities where they engage in a wide array of

intra- and inter-specific interactions. The extent to which these interactions drive or

impede microbiome diversity is not well understood. Historically, two contrasting

hypotheses have been suggested to explain how species interactions could influence

diversity. 'Ecological Controls' (EC) predicts a negative relationship, where the evolution

or migration of novel types is constrained as niches become filled. In contrast, 'Diversity

Begets Diversity' (DBD) predicts a positive relationship, with existing diversity

promoting the accumulation of further diversity via niche construction and other

interactions. Using high-throughput amplicon sequencing data from the Earth

Microbiome Project, we provide evidence that DBD is strongest in low-diversity biomes,

but weaker in more diverse biomes, consistent with biotic interactions initially favoring

the accumulation of diversity (as predicted by DBD). However, as niches become

increasingly filled, diversity hits a plateau (as predicted by EC).

Impact statement:

Microbiome diversity favors further diversity in a positive feedback that is strongest in

lower-diversity biomes (*e.g.* guts) but which plateaus as niches are increasingly filled in

higher-diversity biomes (*e.g.* soils).

Introduction

The majority of the genetic diversity on Earth is encoded by microbes (Hug et al., 2016; Lapierre & Gogarten, 2009; Sunagawa et al., 2015) and the functioning of all Earth's ecosystems is reliant on diverse microbial communities (Falkowski et al., 2008). High-throughput 16S rRNA gene amplicon sequencing studies continue to yield unprecedented insight into the taxonomic richness of microbiomes (e.g. (Louca et al., 2019; Sogin et al., 2006)), and abiotic drivers of community composition (e.g. pH; Lauber et al., 2009; Power et al., 2018) are increasingly characterized. Although it is known that biotic (microbe-microbe) interactions can also be important in determining community composition (Needham & Fuhrman, 2016), comparatively little is known about how such interactions, either positive (e.g. cross-feeding; Seth & Taga, 2014) or negative (e.g. toxin-mediated interference competition; Czárán et al., 2002; Hibbing et al., 2010), shape microbiome diversity as a whole.

The dearth of studies exploring how microbial interactions could influence diversity stands in marked contrast to a long research tradition on biotic controls of plant and animal diversity (Elton, 1946; Gause, 2003). In an early study of 49 animal (vertebrate and invertebrate) community samples, Elton plotted the number of species versus the number of genera and observed a ~1:1 ratio in each individual sample, but a ~4:1 ratio when all samples were pooled (Elton, 1946). He took this observation as evidence for competitive exclusion preventing related species, more likely to overlap in niche space, to co-exist. This concept, more recently referred to as niche filling or Ecological Controls (EC) (Schluter & Pennell, 2017), predicts speciation (or, more generally, diversification) rates to decrease with increasing standing species diversity

because less niche space is available (Rabosky & Hurlbert, 2015). In contrast, the

Diversity Begets Diversity (DBD) model predicts that when species interactions create

novel niches, standing biodiversity favors further diversification (Calcagno et al., 2017;

Whittaker, 1972). For example, niche construction (i.e. the physical, chemical or

biological alteration of the environment) could influence the evolution of the species

constructing the niche, as well as that of co-occurring species (Laland et al., 1999; San

Roman & Wagner, 2018). An alternative to either EC or DBD is The Neutral Theory of

Biodiversity and Biogeography, in which all species are functionally equivalent and

communities assemble via random sampling (Hubbell, 2001). Neutral Theory serves as a

null hypothesis of community assembly in macrobes (Azaele et al., 2016; N. J. Gotelli &

McGill, 2006), and more recently in microbiome research (Harris et al., 2017; Li & Ma,

2016).

Empirical evidence for the action of EC vs. DBD in natural plant and animal

communities has been mixed (Calcagno et al., 2017; Emerson & Kolm, 2005; Palmer &

Maurer, 1997; Price et al., 2014; Rabosky et al., 2018). Laboratory evolution experiments

tracking the diversification of a focal bacterial lineage in communities of varying

complexity have also yielded contradictory results, with support for EC, DBD, or

intermediate scenarios (Brockhurst et al., 2007; Meyer & Kassen, 2007). For example,

diversification of a focal *Pseudomonas* clone was favored by increasing community

diversity in the range of 0-20 other strains or species within the same genus (Calcagno et

al., 2017; Jousset et al., 2016) but diversification was inhibited in highly diverse

communities (*e.g.* hundreds or thousands of species in compost; (Gómez & Buckling,

2013)). These experiments are consistent with interspecific competition initially driving

(Bailey et al., 2013), but eventually inhibiting diversification as niches are filled.

Most laboratory experiments are restricted to relatively short evolutionary time

scales and include only a small number of taxa; it is therefore unclear if they can be

generalized to natural communities consisting of many more taxa evolving and

assembling over much longer periods, spanning more environmental change, greater

evolutionary diversification, and frequent migration events. Although the absence of a

substantial prokaryotic fossil record hinders deconvoluting speciation and extinction rates

(Louca & Pennell, 2020; Marshall, 2017), Louca et al. (Louca et al., 2018) recently

estimated that bacterial diversity has mostly increased over the past billion years, with

speciation rates slightly exceeding extinction rates. However, because many free-living

microbes have high migration rates ("everything is everywhere, but the environment

selects" (de Wit & Bouvier, 2006)), we expect that the majority of diversity present

within a typical microbiome sample is selected from a pool of migrants rather than

having evolved *in situ*. As such, here we broadly define "diversity begets diversity"

(DBD) to include the combined effects of community assembly from a migrant pool

('ecological species sorting') and *in situ* evolutionary diversification (**Fig. 1**).

Fig. 1. Contrasting the Diversity Begets Diversity (DBD) and Ecological Controls (EC) models. (A) In this hypothetical scenario, microbiome sample 1 contains one non-focal genus, and two amplicon sequence variants (ASVs) within the focal genus (point at x=1, y=2 in the plot). Sample 2 contains three non-focal genera, and four ASVs within the focal genus (point at x=3, y=4). Tracing a line through these points yields a positive diversity slope, supporting the DBD model (red). **(B)** Alternatively, a negative slope would support the Ecological Controls (EC) model (blue line). In the middle panel, we consider a community assembly model to explain the hypothetical data of the top panel, in which standing diversity (black points) in a community selects (for or against) new types (referred to here as ASVs) which arrive via migration (purple points & arrows). In the bottom panel, we consider an evolutionary diversification model of a focal lineage (genus) into ASVs as a function of initial genus-level community diversity present at the time of diversification.

To test whether patterns of diversity in natural communities conform to EC or

DBD dynamics, we used 2,000 microbiome samples from the Earth Microbiome Project

(EMP), the largest available repository of biodiversity based on standardized sampling

and sequencing protocols, with 16S rRNA gene amplicon sequence variants (ASVs) as

the finest-grained taxonomic unit (Thompson et al., 2017). Following Elton (Elton,

1946), we use the equivalent of Species:Genus ratios, calculating a range of taxonomic

diversity ratios (up to the Class:Phylum level) as proxies for diversity within a focal

taxon, from shallow to deep evolutionary time. We then plot each ratio as a function of

the number of non-focal taxa (Genera, Families, Orders, Classes, and Phyla, respectively)

with which the focal taxon could interact. We refer to the slope of these plots as the

"diversity slope", with negative slopes supporting EC and positive slopes supporting

DBD (**Fig. 1**). As a null, we compare these slopes to the expectation under Neutral

Theory. To avoid a trivially positive diversity slope due to variation in sequencing effort,

all samples were rarefied to 5,000 observations (counts of 16S rRNA gene sequences), as

diversity estimates are highly sensitive to sampling effort (Nicholas J. Gotelli & Colwell,

2001). As 16S evolves at a rate of roughly 1-2 substitutions per million years (Kuo &

Ochman, 2009b), evolutionary diversification within individual EMP samples cannot be

uncovered using this marker; rather our data represent mainly a record of community

assembly.

Results

Quantifying the DBD-EC continuum in prokaryote communities compared to neutral null models. We used generalized linear mixed models (GLMMs) to estimate the diversity slope at each taxonomic level in the EMP data, which revealed a tendency toward positive slopes with significant variation explained by the random effects of lineage, environment, and their interaction (**Table 1, Figure 2, Figure 2 supplements 1-6, Supplementary Data file 1 Section 1**). All models reported here provide significantly better fits compared to models without the fixed effect of community diversity, and coefficients of determination (R^2) are higher with the inclusion of random effects, showing their importance (**Supplementary Data file 2**). Examples of how the diversity slope varies across lineages and environments are shown in **Figure 2** and **Figure 2 supplements 2-6**. To assess the significance of these slope estimates in light of potential sampling bias and data structure (Gotelli & Colwell, 2001; Jarvinen, 1982), we considered null models, all of which randomize the associations between ASVs within a sample, thus randomizing any true biotic interactions. Models 1 and 2 are based on draws from the zero-sum multinomial (ZSM) distribution, which arises from the standard Neutral Theory of Biodiversity (**Methods**). Model 1, in which each microbiome sample is drawn from the same ZSM distribution, produces a significantly negative diversity slope (**Figure 2 supplement 7; Table 2**). Model 2, in which each environment draws from a separate distribution, is effectively a composite of Model 1 in which different environments, each with a negative slope, are 'stacked' to yield an overall positive slope (**Figure 2 supplement 7**). However, the Model 2 slope is not significant in a GLMM

accounting for variation across environments (**Table 2, Supplementary Data file 3**

Section 1.2). In the real EMP data, most individual environments tend toward a positive

slope (**Figure 2 supplement 8**). The tendency toward positive diversity slopes in the

EMP is therefore not straightforwardly explained by neutral processes.

To estimate the power to detect either DBD or EC, we specifically added each of

these effects to data simulated under a null model. As expected, adding DBD reversed the

negative slope and rendered it positive (**Table 2; Figure 2 supplement 7,**

Supplementary Data file 3 Section 2.1), suggesting reasonable power to detect DBD

when truly present. In contrast, the addition of EC had little effect on the slope,

suggesting low power to detect EC under some null models. Taken together, these

modelling results suggest that positive diversity slopes observed in the EMP are more

readily explained by DBD than by Neutral Theory, whereas negative slopes could be

explained by EC, Neutral Theory, or some combination of the two.

Because taxonomic labels can be unavailable or inconsistent with phylogenetic

relationships (Parks et al., 2018; Vos, 2011) we repeated the analyses using nucleotide

sequence identity in the 16S rRNA gene instead of taxonomy, and again recovered

generally positive diversity slopes (**Methods**). As a final sensitivity analysis, we repeated

the GLMMs using unrarefied community Shannon diversity instead of richness

(**Methods**) and obtained similar results, with generally positive diversity slopes that

could in some cases be reversed depending on the lineage or environment (**Table 3,**

Supplementary Data file 1 Section 2). The Shannon diversity metric is robust to

sampling effort, suggesting that the results are not biased by undersampling in diverse

biomes. Even if undersampling could bias the diversity slope downward in more diverse

samples, the effect is unlikely to be large at a rarefaction to 5,000 sequences, and only to

occur at the extremes of diversity (*e.g.* very many genera and high ASV:genus ratios) and

not at higher taxonomic levels (*e.g.* Class:Phylum) (**Figure 2 supplement 9**).

Fig. 2. Focal lineage diversity as a function of community diversity in the top two

most prevalent taxa at each taxonomic level. As in **Fig. 1**, the x-axes show community

diversity in units of the number of non-focal taxa (*e.g.* the number of non-Proteobacteria

phyla for the left-most column), and the y-axes show the taxonomic ratio within the focal

taxon (*e.g.* the number of classes within Proteobacteria). Significant positive diversity

slopes are shown in red, negative in blue (linear models, *P* <0.05, Bonferroni corrected

for 17 tests), and non-significant in grey. Note that linear models are distinct from

GLMMs, and are for illustrative purposes only. Four representative environments are

shown (see **Figure 2 supplements 2-6** for plots in all 17 environments).

DBD reaches a plateau at high diversity. It is expected from theory and experimental

studies that a positive DBD relationship should eventually reach a plateau, giving way to

EC as niches become saturated (Brockhurst et al., 2007; Gómez & Buckling, 2013). This

expectation is borne out in our dataset, particularly in the nucleotide sequence-based

analyses which support quadratic or cubic relationships over linear diversity slopes

(**Figure 2 supplement 10**). For example, in the animal distal gut, a relatively low-

diversity biome, we observed a strong linear DBD relationship at most phylogenetic

depths; in contrast, the much more diverse soil biome clearly reaches a plateau (**Figure 2**

supplement 11).

To comprehensively test the hypothesis that more diverse microbiomes

experience weaker DBD due to saturated niche space, we used a GLMM including the

interaction between diversity and environment as a fixed effect. We considered this

model only for taxonomic ratios with significant diversity slope variation by environment

(**Table 1**): Family:Order, Order:Class, and Class:Phylum. Diversity slopes were

significantly higher in less diverse (often host-associated) biomes, suggesting that niche

filling leads to a plateau of DBD in more diverse biomes (**Fig. 3, Supplementary Data**

file 1 Section 3). The interaction observed in the real EMP data between community

diversity and biome type in shaping focal lineage diversity was not observed under a

neutral null (Model 2, in which each environment has its own characteristic level of

diversity) (**Supplementary Data file 3 Section 1.2**). The DBD plateau observed in more

diverse biomes is thus not readily explained by a neutral model, nor is rarefaction

expected to bias the diversity slope estimates, particularly at the Class:Phylum level

(**Figure 2 supplement 9**). This suggests that the plateau of DBD at higher levels of

community diversity is not an artefact of data structure or sampling effort. Finally, we

considered whether variation along the EC-DBD continuum could be explained by

differential cell density across environments, which could affect both the frequency of

cell-cell interactions (a biological effect) or the sampling depth (a technical artefact).

Although precise estimates of cell densities in all EMP biomes are not available, we

extracted plausible ranges for eight biomes from the literature (Kennedy & de Luna,

2005; Lindow & Brandl, 2003; Sender et al., 2016; Whitman et al., 1998) and annotated

these in **Figure 3**. It is clear from this figure that relatively high- and low-density samples

are found along the range of community taxonomic diversities, demonstrating that cell

density is unlikely to drive the trend of decreasing diversity slopes with increasing

community diversity.

Fig. 3. The diversity slope of focal taxa is higher in low-diversity (often host-associated) microbiomes. The x-axis shows the mean number of non-focal taxa: (A) phyla, B) classes, and C) orders in each biome. On the y-axis, the diversity slope was estimated by a GLMM predicting focal lineage diversity as a function of the interaction between community diversity and environment type at the level of A) Class:Phylum, B) Order:Class, and C) Family:Order ratios (**Supplementary Data file 1 Section 3**). The line represents a linear regression; the shaded area depicts 95% confidence limits of the fitted values. Adjusted R^2 and P-values from the linear fits are shown at the top right of each panel. See **Supplementary Data file 2** for model goodness of fit. Slopes not significantly different from zero are shown as empty circles. Estimates of bacterial cell

density from the literature are indicated in grey text, in units of bacteria/mm^3. For animal

(skin) and plant surface, units of bacteria/mm^2 were converted to mm^3 assuming layers of

bacteria 1 micron thick. For rhizosphere samples we assume a density of 1-2g/cm^3

(Kennedy & de Luna, 2005).

Abiotic drivers of diversity. Our results thus far suggest that community diversity is a

major determinant of the EC-DBD continuum, and by extension that biotic interactions

may override abiotic factors in determining where a community lies on the continuum.

To formally test for the additional role abiotic drivers might play in generating the

observed EC-DBD continuum, we analyzed two data sets in more detail.

First, we analyzed a subset of 192 EMP samples with measurements of four key

abiotic factors shown to affect microbial diversity (pH, temperature, latitude, and

elevation; (Delgado-Baquerizo et al., 2018; Lauber et al., 2009; Power et al., 2018;

Schluter & Pennell, 2017)). We fitted a GLMM with focal lineage-specific diversity as

the dependent variable, and with the number of non-focal lineages, the four abiotic

factors and their interactions as predictors (fixed effects). As in the full EMP dataset

(**Table 1**), focal lineage diversity was positively associated with community diversity at

all taxonomic ratios in the EMP subset (**Table 4**). As expected, certain abiotic factors,

alone or in combination with diversity, had significant effects on focal lineage diversity

(**Table 4**). However, the effects of abiotic factors were always weaker than the effect of

community diversity (**Table 4**; **Supplementary Data file 1 Section 4**).

Second, we used a global 16S sequencing dataset of 237 soil samples associated

with more detailed environmental metadata (Delgado-Baquerizo et al., 2018) which we

reprocessed to yield ASVs comparable to those in the EMP (**Methods**). This dataset

revealed weaker evidence for DBD and stronger effects of abiotic variables on diversity.

Community diversity generally had significant positive effects on focal-lineage diversity,

but the effect was weak and not detectable at all taxonomic ratios (**Table 5**). Known

abiotic drivers of soil bacterial diversity such as pH (Lauber et al., 2009) and latitude

(Delgado-Baquerizo et al., 2018) had effects of similar or stronger magnitude compared

to the effect of community diversity (**Table 5, Supplementary Data file 4**). The

relatively weak effect of DBD and strong effect of abiotic drivers on diversity in this soil

dataset can be explained by the fact that soils generally are highly diverse and have

relatively low diversity slopes (**Figure 3**).

We note that it remains possible that unmeasured abiotic effects could explain

some of the DBD effects observed in the EMP. Although only a small subset of abiotic

factors was considered, the generally positive diversity slopes in the EMP are not likely

to be driven by these factors in the abiotic environment (**Table 4**). Specifically, we

consider it unlikely that unmeasured abiotic factors would always act similarly, and in the

same direction across multiple different environments, to drive DBD. However, as

demonstrated in soil (**Table 5**), abiotic factors may become increasingly important in

highly diverse biomes with weak DBD.

DBD is more pronounced in resident taxa than in migrant- or generalist taxa. A

recent meta-analysis of 16S sequence data from a variety of biomes suggests there is an

important distinction between generalist lineages found in many environments, compared

to specialists with a more restricted distribution (Sriswasdi et al., 2017). Generalists were

inferred to have higher speciation rates, suggesting that the DBD-EC balance might differ

between generalists and specialists (Sriswasdi et al., 2017). To further investigate this

difference, we defined 'residents', taxa with a strong preference for a specific biome, in

addition to generalists without a strong biome preference in the EMP dataset. We first

clustered environmental samples by their genus-level community composition using

fuzzy k-means clustering (**Fig. 4a**), which identified three major clusters: 'animal-

associated', 'saline', and 'non-saline'. The clustering included some outliers (*e.g.* plant

corpus grouping with animals), but was generally consistent with known distinctions

between host-associated vs. free-living (Thompson et al., 2017), and saline vs. non-saline

communities (Auguet et al., 2010; Lozupone & Knight, 2007). Resident genera were

defined as those with a strong preference for a particular environment cluster (whether

due to dispersal limitation or narrow niche breadth) using indicator species analysis

(permutation test, $P<0.05$; **Fig. 4a**; **Figure 4 supplement 1**; **Supplementary Data file 5**),

and genera without a strong preference were considered generalists. When residents of

one environmental cluster were (relatively infrequently) observed in a different cluster,

we defined them as "migrants" in that sample. For each environment cluster, we ran a

GLMM with resident genus-level diversity (the number of non-focal genera) as a

predictor of focal-lineage diversity (the ASV:Genus ratio) for residents, generalists, or

migrants to that sample (**Supplementary Data file 1 Section 5**).

      Resident community diversity had no significant effect on the diversity of

generalists in animal-associated, saline and non-saline clusters (GLMM, Wald test,

$P>0.05$), but was positively correlated with lineage-specific resident diversity (GLMM,

Wald test, $z=7.1$, $P=1.25e-12$; $z=3.316$, $P=0.0009$; $z=7.109$, $P=1.17e-12$, respectively).

Resident community diversity significantly decreased migrant diversity in saline

(GLMM, z=-3.194, *P*=0.0014) and non-saline environment clusters (GLMM, z=-2.840,

P=0.0045), but had no significant effect in the animal-associated cluster (GLMM,

P>0.05) (**Fig. 4b**). These results suggest that, although generalist lineages may have

higher speciation rates and colonize more habitats than specialists (Sriswasdi et al.,

2017), they have lower diversity slopes. Migrants to the "wrong" environment experience

even less DBD, and are even subject to EC in two out of three environment types (**Fig.**

4b). The accumulation of diversity via successful establishment of migrants may thus be

limited, presumably because most niches are already occupied by residents.

Fig. 4. The DBD relationship varies between resident and non-resident genera. (A) Ordination showing genera clustering into their preferred environment clusters. The matrix of 1128 genera (rows) by 17 environments (columns), with the matrix entries indicating the percentage of samples from a given environment in which each genus is present, was subjected to principal components analysis (PCA). Circles indicate genera and triangles indicate environments (EMPO 3 biomes). Colored circles are genera inferred by indicator species analysis to be residents of a certain environmental cluster, and grey circles are generalist genera. The three environment clusters identified by fuzzy *k*-means clustering are: Non-saline (NS, blue), saline (S, green) and animal-associated (purple). Triangles of the same color indicate EMPO 3 biomes clustered into the same environmental cluster. **(B) DBD in resident versus non-resident genera across environment clusters.** Results of GLMMs modeling focal lineage diversity as a function of the interaction between community diversity and resident/migrant/generalist status. The x-axis shows the standardized number of non-focal resident genera (community diversity); the y-axis shows the number of ASVs per focal genus. Resident focal genera are shown in orange, migrant focal genera in red, and generalist focal genera in black. Red stars indicate a significantly positive or negative slope (Wald test, *P*<0.005). See **Supplementary Data file 2** for model goodness of fit.

Discussion

Using ~10 million individual marker sequences from the EMP, we demonstrate an overall trend for diversity in focal lineages to be positively associated with overall community diversity, albeit with significant variation across lineages and environments. The strength of the DBD relationship dissipates with increasing microbiome diversity, which we hypothesize is caused by niche saturation. In more diverse biomes such as soil, abiotic factors therefore may become relatively more important in driving focal-lineage diversity. The effect of DBD is strongest among habitat specialists (residents), suggesting that long-term niche adaptation tends to select against the establishment of migrant diversity.

While most of the DBD literature considers a model of evolutionary

diversification (Schluter & Pennell, 2017; Whittaker, 1972), our results pertain mainly to

ecological community assembly dynamics. At the limited resolution of 16S rRNA gene

sequences, we do not expect measurable diversification within an individual microbiome

sample (Kuo & Ochman, 2009b); however, community diversity could still select for (as

in DBD) or against (as in EC) increasing diversity in a focal lineage, even if this lineage

diversified before the sampled community assembled. Future work with higher resolution

genomic or metagenomic data will enable testing if and how DBD arises in microbial

communities via evolutionary diversification, and also how prokaryote diversification is

affected by other community members including phages (Brockhurst et al., 2005),

protists (Meyer & Kassen, 2007), and fungi (Kastman et al., 2016). Predator-prey, cross-

feeding, and other biotic interactions with these non-prokaryotic community members

could explain some of the unaccounted variation we observed in diversity slopes across

environments.

Our dataset also provides an opportunity to explore how DBD relates with

genome size evolution. Bacteria with larger repertoires of accessory genes, and thus

larger genomes, are able to occupy a wider range of niches (Barberán et al., 2014). Taxa

with larger genomes might therefore be hypothesized to better survive and thrive when

they disperse into a new location, exhibiting stronger DBD. Although a comprehensive

test of this hypothesis will require higher resolution genomic or metagenomic data, as a

preliminary exploration we assigned genome sizes to 576 focal genera for which at least

one whole genome sequence was available (using the largest recorded genome size for

each genus) and added an interaction term between genome size and diversity as a fixed

effect in the GLMM (**Methods**). Consistent with our expectation, we observed a

significant positive effect of genome size on the diversity slope (GLMM, Wald test,

$z=2.5$, $P=0.01$; **Fig. 5, Supplementary Data file 1 Section 6**). This effect was not

observed in null models, in which the interaction between community diversity and focal

genus genome size was never significant (**Supplementary Data file 3 Section 1.3 and**

2.2) and so this effect of genome size cannot be trivially explained by data structure. The

positive relationship between genome size and DBD is likely even stronger than

estimated, because assigning genome sizes to entire genera is imprecise (*i.e.* there is

variation in genome size within a genus, or even within species), therefore weakening the

correlation.

The positive correlation between genome size and DBD observed here could be

driven by larger metabolic repertoires encoded by larger genomes (40), potentially

creating more opportunities to benefit from cross-feeding, niche construction (San Roman

& Wagner, 2018), and other interspecies interactions. This tendency appears to be at odds

with the Black Queen hypothesis, which predicts that social conflict between interacting

species leads to the inactivation and loss of genes involved in shareable metabolites

(public goods), eventually resulting in reduced genome size (Morris & Lenski, 2012).

Such a process would produce a negative correlation between the degree of species

interactions (*i.e.* community diversity) and genome size (Morris & Lenski, 2012). The

interaction between genome size, biotic interactions and diversification thus deserves

further study.

Fig. 5. Positive effect of genome size on DBD. Results are shown from a GLMM
predicting focal lineage diversity as a function of the interaction between community
diversity and genome size at the ASV:Genus ratio (**Supplementary Data file 1 Section
6**). The x-axis shows the standardized number of non-focal genera (community diversity);
the y-axis shows the number of ASVs per focal genus. Variable diversity slopes
corresponding to different genome sizes are shown in a blue color gradient; the shaded
area depicts 95% confidence limits of the fitted values. See **Supplementary Data file 2**
for model goodness of fit.

Alongside theory and experimental data, the EMP survey data provide a window
into the biotic drivers of microbial diversity in nature. In particular, our correlational
results support previous experimental and theoretical results showing that DBD is strong

when community diversity is low (Calcagno et al., 2017; Jousset et al., 2016), driving the

accumulation of diversity in a positive feedback loop until niches are filled and EC starts

to predominate (Bailey et al., 2013; Brockhurst et al., 2007; Gómez & Buckling, 2013;

Meyer & Kassen, 2007). However, due to the correlational nature of the EMP data, it is

not possible to test whether DBD is primarily due to the creation of novel niches via

biotic interactions and niche construction (Laland et al., 1999), or due to increased

competition leading to specialization on underexploited resources (Hibbing et al., 2010;

Jousset et al., 2016). We hope future higher resolution genomic studies, and

complementary experiments, will be able to elucidate the types of biotic interactions that

promote microbiome diversity. Regardless of the underlying mechanisms, our results

demonstrate a general scaling between different levels of community diversity, which has

important implications for modeling and predicting community function and stability in

response to perturbations (Coyte et al., 2015; Pennekamp et al., 2018). The answer to the

question 'why are microbiomes so diverse?' might in a large part be because

microbiomes are so diverse (Emerson & Kolm, 2005).

Acknowledgements.

We thank Luke Thompson for assistance obtaining EMP data and Zofia Ecaterina

Taranu, Vincent Fugère and Guillaume Larocque for advice on GLMMs. We are also

grateful to Steven Kembel, Tom Battin, the reviewers Eric Kemen and Benjamin E.

Wolfe, and the editor Detlef Weigel for critical comments that improved the manuscript.

Funding: This project was made possible by an NSERC Discovery Grant and Canada

Research Chair to BJS.

Competing interests: none to declare.

Data and materials availability: All data is available from the Earth Microbiome

Project (ftp.microbio.me), as detailed in the Methods. All computer code used for

analysis are available at https://github.com/Naima16/dbd.git.

 Tables

Table 1. Effects of community diversity on focal lineage diversity across taxonomic
ratios. The GLMMs showed statistically a significant positive effect of community
diversity on focal lineage diversity. Each row reports the effect of community diversity
on focal lineage diversity (Div), as well as its standard error, Wald z-statistic for its effect
size and the corresponding *P*-value (left section), or standard deviation on the slope for
the significant random effects (right section). SE=standard error, Env=environment type,
Lin=lineage type, Lab=Principal Investigator ID, Sample=EMP Sample ID. Interactions
are denoted as '*'. n.s.=not significant (likelihood-ratio test). All models provide a
significantly better fit than null models without fixed effects (ΔAIC > 10 and *P* < 0.05;
Supplementary Data file 2).
447
448

	Slope (fixed effects)				Standard deviation on the slope (random effects)				
	Div	SE	z	*P*	Env	Lin	Lin*Env	Env*Lab	Sample
ASV:Genus	0.091	0.016	5.792	6.95e-09	n.s.	0.074	0.142	0.114	0.067
Genus:Family	0.047	0.008	5.911	3.41e-09	n.s.	0.071	0.07	0.039	n.s.
Family:Order	0.119	0.017	7.001	2.54e-12	0.023	0.094	0.092	0.106	n.s.
Order:Class	0.109	0.020	5.447	5.13e-08	0.05	0.141	0.078	0.051	n.s.
Class:Phylum	0.272	0.043	6.341	2.29e-10	0.119	0.174	0.119	0.114	n.s.

Table 2. GLMMs applied to data simulated under null models. Null models 1 and 2 were generated under the ZSM distribution, with a single distribution for the whole dataset (Model 1) or one distribution per environment (Model 2). Model 3 is similar to Model 1, except with a single Poisson distribution for the whole dataset, and +DBD or +EC refer to adding these effects to 100% of ASVs (see **Methods** and **Figure 2 supplement 7**). Each row reports the effect of community diversity on focal lineage diversity (Div), as well as its standard error, Wald z-statistic for its effect size and the corresponding *P*-value (Wald test) (left section), or standard deviation on the slope for the significant random effects (right section). SE=standard error, Env=environment type, Lin=lineage type, Sample=EMP Sample ID. n.s.=not significant (likelihood-ratio test), n.t.= not tested, because separate environments were not included in Models 1 or 3.

	Slope (fixed effects)				Stand dev on the slope (random effects)			
	Div	SE	z	*P*	Env	Lin	Lin*Env	Sample
Model 1	**-0.005**	0.000	-9.807	**<2e -16**	n.t.	0.639	n.t.	n.s.
Model 2	n.s.							
Model 3	**-0.012**	0.002	-6.552	**5.69e-11**	n.t.	0.021	n.t.	n.s.
Model 3 + DBD	**0.016**	0.001	11.48	**<2e-16**	n.t.	0.008	n.t.	n.s.
Model 3 + EC	**-0.011**	0.002	-6.14	**8.26e-10**	n.t.	ns	n.t.	n.s.

Table 3. GLMMs with community diversity measured using Shannon diversity.
Results are shown from GLMMs with Shannon diversity of non-focal taxa (Div) as a
predictor of ASVs richness of focal taxa. Each row reports the estimate (Div), as well as
its standard error, Wald z-statistic for its effect size and the corresponding *P*-value (Wald
test) (left section), or standard deviation on the slope for the significant random effects
(right section). SE=standard error, Env=environment type, Lin=lineage type,
Lab=Principal Investigator ID, Sample=EMP Sample ID. n.s.=not significant (likelihood-
ratio test).
471

	Fixed effects				Random effects				
	Div	SE	z	*P*	Env	Lin	Env*Lin	Env*Lab	Sample
Genus	0.055	0.013	4.33	1.49e-05	n.s.	0.08	0.15	0.085	0.054
Family	0.148	0227	6.491	8.51e-11	n.s.	0.184	0.268	0.16	0.134
Order	0.378	0.038	9.864	<2e-16	n.s.	0.34	0.417	0.258	0.202
Class	0.398	0.05	7.973	1.54e-15	n.s.	0.369	0.46	0.326	0.262
Phylum	0.319	0.088	3.614	0.0003	0.169	0.316	0.5	0.495	0.378

Table 4. Community diversity has a stronger effect than abiotic factors on focal lineage
diversity (EMP dataset). Results are shown from GLMMs with community diversity, four
abiotic factors (temperature, elevation, pH, and latitude), and their interactions with community
diversity, as predictors of focal lineage diversity. Random effects on the intercept included
environment, lineage, lab ID and sample ID. Each row reports the taxonomic ratio, the predictors
used in the GLMM (fixed effects only), their estimate (Est), standard error (SE) and *P*-value (P)
(Wald test). Interactions are denoted as '*'. Random effects are not shown.
480

	Predictor	Est	SE	P
ASV:Genus	**Div**	0.128	0.013	< 2e-16
	Temperature	0.04	0.014	0.00479
	Div*Temperature	0.043	0.014	0.00175
	Div*Latitude	0.031	0.013	0.02119
	Div*Elevation	-0.031	0.014	0.02829
Genus:Family	**Div**	0.094	0.009	< 2e-16
	Temperature	0.026	0.009	0.00268
	pH	-0.042	0.009	5.88e-06
Family:Order	**Div**	0.131	0.01	< 2e-16
Order:Class	**Div**	0.184	0.01	< 2e-16
	Div*Temperature	0.032	0.009	0.000827
	Div*Latitude	0.023	0.008	0.005403
Class:Phylum	**Div**	0.236	0.011	< 2e-16
	Div*Temperature	0.059	0.014	2.15e-05
	Div*Latitude	0.03	0.011	0.00884

Table 5. GLMMs applied to a soil dataset. Each row reports the taxonomic ratio, the predictors used in the GLMM (fixed effects only), their estimate (Est), standard error (SE) and *P*-value (P) (Wald test). Left columns: GLMM with community diversity (Div) and all abiotic variables considered separately, as predictors of focal lineage diversity. Right columns: GLMM with community diversity (Div) and the three first principle components (PCs) representing abiotic variables, as predictors of focal lineage diversity. n.s., non-significant (LRT test). All models provide a significantly better fit than null models without fixed effects (ΔAIC > 10 and $P < 0.05$; **Supplementary Data file 2**), except for the GLMM with abiotic factors at the Family:Order level, where latitude has a significant effect on focal lineage diversity but its effect is nearly null, with a ΔAIC between full and null model of 4 and a null marginal R^2.

	GLMMs with abiotic variables				GLMMs with the 3 first PCs			
	Predictor	Est	SE	P	Predictor	Est	SE	P
ASV:Genus	Div	n.s.			Div	0.064	0.016	9.47e-05
	Latitude	0.294	0.025	< 2e-16	PC1	-0.065	0.007	< 2e-16
	UV_light	-0.177	0.016	< 2e-16	PC2	-0.03	0.006	1.98e-05
	MDR	0.028	0.006	7.12e-06				
	NPP2003_2015	-0.066	0.005	< 2e-16				
	Latitude^2	-0.3	0.029	< 2e-16				
	Clay_silt^2	-0.012	0.004	0.003				
	Soil_N^2	-0.007	0.001	1.66e-06				
	Soil_C_N_ratio	0.003	0.001	0.004				
	PSEA^2	0.01	0.002	4.84e-06				
	MDR^2	0.017	0.003	2.40e-08				
	NPP2003_2015	-0.016	0.004	0.0001				
Genus:Family	Div	0.032	0.01	0.0011	Div	0.033	0.01	0.001
	Latitude	-0.035	0.006	2.04e-09	PC1	-0.016	0.006	0.02
					PC2	0.02	0.006	0.00089
Family:Order	Div	n.s.			Div	n.s.		
	Latitude	-0.0005	0.0002	0.0105	PC1	-0.026	0.007	0.00032
					Div*PC1	0.04	0.006	2.14e-12
					Div*PC3	0.023	0.005	1.68e-06
Order:Class	Null model with no predictor was significant							
Class:Phylum	Div	0.032	0.01	0.00174	Div	0.032	0.01	0.003
	pH	0.074	0.01	4.37e-13	PC1	-0.051	0.01	3.54e-07
					PC2	-0.028	0.01	0.006

Supplementary Figure Legends

Figure 2 supplement 1. Distributions of diversity slope estimates across different random effects, from the GLMMs predicting focal lineage diversity as a function of community diversity. (A) Class:Phylum, **(B)** Order:Class, **(C)** Family:Order, **(D)** Genus:Family, and **(E)** ASV:Genus. Estimation of random effect coefficients from the GLMMs (Table S1), shows that the effect of diversity on focal lineage diversity (slope estimates) are generally positive but could be negative in some lineages or combinations of environment, lineage (Environment*Lineage), and the laboratory that submitted the dataset (Environment*Lab).

Figure 2 supplement 2. Focal lineage diversity as a function of community diversity across biomes in the three most prevalent phyla. (A) Proteobacteria, (B) Bacteroidetes, (C) Actinobacteria. Linear models are shown for the number of classes per phylum (y-axis) as a function of community diversity (number of non-focal phyla, x-axis) in each of the 17 environments (EMPO3 biomes). Only environments containing the focal lineage are shown. P-values are Bonferroni corrected for 17 tests. Significant ($P <0.05$) models are shown with red trend lines, non-significant ($P > 0.05$) trends are shown in blue.

Figure 2 supplement 3. Focal lineage diversity as a function of community diversity across biomes in the three most prevalent classes. Linear models are shown for the number of orders per class (y-axis) as a function of community diversity (non-focal classes, x-axis) in each of the 17 environments (EMPO3 biomes). Only environments containing the focal lineage are shown. Significant positive diversity slopes are shown in red, negative in blue (linear models, $P <0.05$, Bonferroni corrected for 17 tests), and non-significant in grey.

Figure 2 supplement 4. Focal lineage diversity as a function of community diversity across biomes in the three most prevalent orders. Linear models are shown for the number of families per order (y-axis) as a function of community diversity (non-focal orders, x-axis) in each of the 17 environments (EMPO3 biomes). Only environments containing the focal lineage are shown. Significant positive diversity slopes are shown in red, negative in blue (linear models, $P <0.05$, Bonferroni corrected for 17 tests), and non-significant in grey.

Figure 2 supplement 5. Focal lineage diversity as a function of community diversity across biomes in the three most prevalent families. Linear models are shown for genera per family (y-axis) as a function of community diversity (non-focal families, x-axis) in each of the 17 environments (EMPO3 biomes). Only environments containing the focal lineage are shown. Significant positive diversity slopes are shown in red, negative in blue (linear models, $P <0.05$, Bonferroni corrected for 17 tests), and non-significant in grey.

Figure 2 supplement 6. Focal lineage diversity as a function of community diversity across biomes in the three most prevalent genera. Linear models are shown for ASVs per genus (y-axis) as a function of community diversity (non-focal genera, x-axis) in each

544   of the 17 environments (EMPO3 biomes). Only environments containing the focal
545   lineage are shown. Significant positive diversity slopes are shown in red, negative in blue
546   (linear models, P <0.05, Bonferroni corrected for 17 tests), and non-significant in grey.
547
Figure 2 supplement 7. Null models based on Neutral Theory. Results are shown from
data simulated under (A) neutral Model 1, (B) neutral Model 2, or (C) neutral Model 3.
Model 1 is sampled from the zero-sum multinomial distribution with a single distribution
for the whole dataset, while Model 2 includes a separate distribution for each of the 17
different environments (EMPO 3 biomes). In Model 3 (C), the effect of DBD (top rows)
or EC (bottom rows) are "spiked in" at different levels, ranging from 0 to 100% of ASVs
in a sample. Blue lines show a linear fit, with slopes (m) estimated by GLMM in selected
panels. See Methods for model details, and Table 2 and Supplementary Data file 3,
Section 1.2 for full GLMM results.
557
558   **Figure 2 supplement 8. Lineage diversity (mean ASV:Genus ratio among all**
559   **lineages) as a function of community diversity (number of genera) in the EMP data**.
560   Samples from different environments (EMPO level 3) are shown in different colors, each
561   with their corresponding linear model fit.
562
Figure 2 supplement 9. Taxonomic ratios estimated from simulated rarefied
sequence data. Each panel simulates a set of microbiome samples that differ in their
diversity (number of genera in left panels **A** and **B**, number of phyla in right panels **C** and
D) while maintaining a set true taxonomic ratio (horizontal black line). **(A)** True ratio set
to 2 ASVs/genus, close to the per-sample mean and median in the real EMP data, in a
range of samples between 1 and 1128 named genera, as observed in the real EMP data.
(B) True ratio set to 20 ASVs/genus, equal to the overall mean of 22,014 named ASVs in
1128 named genera, and close to the maximum ratios observed in individual samples
(Fig. 2 supplement 6). Insets show the ranges of 1-50 and 51-150 genera, approximating
observations from lower- or higher-diversity samples such as gut and soil, respectively
(Fig. 2 supplement 6). The insets only show the rarefaction to 5,000 sequences, as used in
the real EMP dataset. **(C)** True ratio set to 3 classes/phylum, close to the per-sample
mean and median in the real EMP data, in a range of samples between 1 and 84 named
phyla, as observed in the real EMP data. **(D)** True ratio set to 10 classes/phylum, close to
the maximum ratios observed in individual samples (Fig. S2). Different rarefaction levels
are shown as different colored lines.
579
Figure 2 supplement 10. Linear, quadratic and cubic models for the relationship
between focal lineage diversity and community diversity for varying levels of %
nucleotide identity. Community diversity was estimated as the number of clusters at a
focal level (d_i) and focal lineage diversity as the mean of the clusters at the rank above
(d_{i+1}/d_i). All P-values are < 0.001. Linear fit (grey); quadratic fit (blue), cubic fit (red);
same colors for the associated adjusted R^2. The x-axis (diversity) shows the number of
clusters at the focal percent-identity level (d_i), and the y-axis (diversification) is the mean
of the clusters at the rank above (d_{i+1}/d_i).
588

Figure 2 supplement 11. Linear, quadratic and cubic models for each environment
type for varying levels of % nucleotide identity. Community diversity was estimated as
the number of clusters at a focal level (d_i) and focal lineage diversity as the mean of the
clusters at the rank above (d_{i+1}/d_i). Linear (grey), quadratic (blue) and cubic (red), with
corresponding adjusted R-squared values in the same colour. *P*-values are Bonferroni
corrected for 17 tests. Significant, $P < 0.05$ (solid lines), non-significant (dashed lines).
The x-axis shows the number of clusters at the focal percent-identity level (d_i), and the y-
axis is the mean of the clusters at the rank above (d_{i+1}/d_i).
597
598 **Figure 4 supplement 1. Resident genera of environment clusters.** Results from
599 indicator species analysis illustrated as a heatmap**.** Only the 25 resident genera with the
600 highest indval indices and *P*<0.05 (permutation test) are shown for every environment
601 cluster (animal-associated, non-saline and saline free). For the full results see
602 **Supplementary Data file 5.**
603
604
605
606 **Supplementary File legends**
607
608 **File 1. Full GLMM outputs for the EMP data.**
609
610 **File 2. Goodness of fit for the GLMMs.**
611
612 **File 3. Full GLMM output for simulated data under Neutral Theory models**
613
614 **File 4. Full GLMM output for soil data (Delgado et al.)**
615
616 **File 5. Indicator species analysis.** The table shows the assignment of each genus to one
617 of three environment types.
618
619 **File 6. Genome size assignment.** The table shows genome sizes assigned to each genus.
620

Materials and Methods

Earth Microbiome Project dataset. We used the EMP '2000 subset' of 16S rRNA gene

sequences, rarefied to 5000 sequences per sample. This subset contains 155,002 ASVs

from 2,000 samples with an even distribution across 17 natural environments (EMP

Ontology level 3). Data were downloaded from the EMP FTP server (ftp.microbio.me),

on February 9, 2018.

Specifically, 16S rRNA-V4 region reads (90 bp, GreenGenes 13.8 taxonomy) along with

environmental data and EMPO3 designations

(http://press.igsb.anl.gov/earthmicrobiome/protocols-and-standards/empo/) were

downloaded from the EMP FTP server (ftp.microbio.me), on February 9, 2018. Sequence

summaries were downloaded from :

ftp://ftp.microbio.me/emp/release1/otu_distributions/otu_summary.emp_deblur_90bp.sub

set_2k.rare_5000.tsv, environmental data from:

ftp://ftp.microbio.me/emp/release1/mapping_files/emp_qiime_mapping_release1.tsv, and

EMPO3 designations from :

ftp://ftp.microbio.me/emp/release1/mapping_files/emp_qiime_mapping_subset_2k.tsv.

The list of the associated 97 studies and 61 corresponding principal investigator identities

were downloaded from https://www.nature.com/articles/nature24621#s1.

Based on the ASV annotations across samples, we estimated the taxonomic ratio for each

focal lineage (ASV:Genus, Genus:Family, Family:Order, Order:Class and Class:Phylum),

along with the number of non-focal lineages (dbd_analys_input.py,

glmm_analys_input.py, Python Version 2.7). Unclassified ASVs were removed from the

analyses.

Generalized linear mixed model (GLMM) analyses. We used GLMMs to determine

how focal lineage diversity (*e.g.* its ASV:Genus ratio) is affected by community diversity

(*e.g.* non-focal genera). The effects of environment (as defined by the EMP Ontology

'level 3 biomes') and the focal lineage identity were included as random effects on the

slope and intercept. We also controlled for the submitting laboratory (identified by the

principal investigator) and the EMP unique sample identifier (i.e. if two taxa were part of

the same sample).

All models were fitted in Rstudio (Version 1.1.442, R Version 3.5.2) using the

glmer function of the lme4 package (Bates et al., 2015). Data standardization

(transformation to a mean of zero and a standard deviation of one) was applied to all

predictors to get comparable estimates. In models with only one predictor, applying

standardization resolved convergence warnings and considerably sped up the

optimization. We first tested the significance of random effects, by using likelihood-ratio

tests (LRTs, implemented in the anova function in the R stats package) on nested models

where each random effect was dropped one at a time. We then assessed the significance

of fixed effects using drop1 function from stats package with the likelihood-ratio test

option (this function drops individual terms from the full model and compares models

based on the AIC). We calculated the Akaike information criterion (AIC) of each

significant model and a null model including all random effects but no fixed effects other

than the intercept. We then report the difference in AIC between the full and null models

(ΔAIC), along with a likelihood ratio test p-value to assess the significance of the full

model relative to the null. Only significant models ($P<0.05$) are reported.

As an additional test of the goodness of fit for the significant models, we

estimated the coefficient of determination (R^2) using the r.squaredGLMM function from

the MuMIn R package. This function implements a method developed by Nakagawa and

Schielzeth and its extension for random slopes (Johnson, 2014; Nakagawa & Schielzeth,

2013). Two values were estimated: the marginal R^2, as a measure of the variance

explained only by fixed effects, and the conditional R^2 as a measure of the variance

explained by the entire model (both fixed effects and random effects). Only results from

R^2 estimation based on lognormal and trigamma methods were reported because they are

specific to the logarithmic link function used in all GLMMs.

Diagnostic plots (plot and qqnorm R functions in base and stats packages) were

checked for each model to ensure that residual homoscedasticity (homogeneity of

variance) was fulfilled: no increase of the variance with fitted values and residuals were

symmetrically distributed tending to cluster around the 0 of the ordinate, but with an

expected pattern due to count data. Normality plots were imperfect, but they generally

showed that the residuals were close to being normally distributed. The assumption of

normality is often difficult to fulfill with high numbers of observations, as is the case in

our models (https://www.statisticshowto.datasciencecentral.com/shapiro-wilk-test/), and

non-normality is less of concern than heteroscedastic for the validity of GLMMs

(https://bbolker.github.io/mixedmodels-misc/ecostats_chap.html#diagnostics).

We tested for overdispersion using the overdisp_fun R function available at

https://bbolker.github.io/mixedmodels-misc/glmmFAQ.html, and found that all the

models were not overdispersed, but rather were underdispersed : the ratio of the sum of

squared Pearson residuals to residual degrees of freedom was < 1 and non-significant

when tested with a chi-squared test. The only exception was Shannon diversity-based

GLMMs. In case of underdispersion and given that underdispersion leads to more

conservative results, we retained the GLMMs with Poisson error distribution, despite the

underdispersion. (GLMM FAQ; Ben Bolker and others; 25 September 2018;

https://bbolker.github.io/mixedmodels-misc/glmmFAQ.html#underdispersion). For

Shannon diversity-based GLMMs, we accounted for overdispersion by adding an

observation-level random effect to the GLMMs (Elston et al., 2001).

Taxonomy-based GLMMs

To test how focal lineage diversity (*e.g.* its ASV:Genus ratio) is affected by community

diversity (*e.g.* non-focal genera richness), for different environment types and lineages

across all taxonomic ratios, we used generalized linear mixed models (GLMMs) fitted on

the EMP dataset. As the dependent variable (focal lineage diversity, defined as taxonomic

ratios, ASV:Genus, Genus:Family, Family:Order, Order:Class, and Class:Phylum) was a

count response, we used a Poisson error distribution with a log link function. Community

diversity (number of non-focal lineages: non-focal Genera, Families, Orders, Classes, and

Phyla), standardized to a mean of zero and a standard deviation of one, was specified as

the predictor (fixed effect). We included the following random effects on the slope and

intercept: lineage (Lin), environment (Env), environment nested within lineage (a lineage

may be present in different environments) and lab (the principal investigator who

conducted the EMP study) nested within environment (different labs sampled and

sequenced a given environment) (as suggested in http://bbolker.github.io/mixedmodels-

misc/glmmFAQ.html). Defining random effects on the slope enabled us to test slope

variation across groups of each categorical variable (*e.g.* slope variation between different

environments or different lineages). We included the EMP unique sample ID as a random

effect to control for dependencies between observations (if two taxa were part of the

same sample) (**Table 1**, **Supplementary file 1 section 1)**.

Shannon diversity-based GLMMs

We also tested whether ASV diversity in a focal taxon is dependent on the diversity of all

other ASVs in that sample (rather than the diversity at only the focal taxonomic level, as

in the taxonomy-based GLMMs above). We used the Shannon diversity index, which is

robust to differences in sampling effort, and generally reaches a plateau at 5,000

sequences or fewer (48, 49). To do so, we fitted a GLMM with the number of ASVs per

focal taxon as the response variable, and the Shannon diversity based on ASVs across all

non-focal taxa (z-standardized) as the predictor (fixed effect), the random effects were

kept as in the taxonomy-based GLMMs, but we added an observation-level random effect

to account for overdispersion (**Table 3**, **Supplementary file 1 section 2**). To avoid

dependence between the response and predictor variables, we used the rarefied ASV

dataset (5,000 ASVs/sample as above) as the response variable, and the Shannon

diversity calculated on unrarefied data from the same samples as the predictor.

Null models. We considered three null models, all of which randomize the associations

between ASVs within a sample, thus breaking any true biotic interactions. These null

models were randomly generated matrices of the same size as the real EMP dataset, but

based on a distribution that arises from the Neutral Theory of Biodiversity. Neutral

Theory postulates that the biodiversity of a metacommunity is governed by independent

random population dynamics across species. The aggregate behaviour is quantified by the

fundamental biodiversity number θ, such that $\theta = 2 J_M \upsilon$, where J_M is the size of the

metacommunity and v is the speciation rate. Parametrized by θ, the metacommunity zero-

sum multinomial distribution (mZSM) was developed to obtain random samples of size J

(Alonso & McKane, 2004). We used this mZSM distribution (implemented with the *sads*

package in R; http://search.r-project.org/library/sads/html/dmzsm.html) to generate the

counts of the ASVs for each dataset in models 1 and 2. Model 1 assumes that the whole

dataset follows the same species abundance distribution (SAD), characterized by a

mZSM with $\theta = 50$. Model 2 assumes that each environment has its own SAD and thus

all the samples of a single environment are assigned the same θ but are distinct across

environments (θ was chosen uniformly between 1 and 100). The number of samples per

environment were the same as the EMP dataset. To obtain similar mean counts as the real

dataset, we set $J = 1000$ for both models 1 and 2, in order to vary θ from 1 to 100. These

values are reasonable based on previous studies that estimated these parameters from

microbiome data (Li & Ma, 2016). We included a down-sampling step to replicate the

zero-inflated nature of the real dataset (on average there were only 96 ASVs per sample

while there was a total of 22,014 ASVs in the entire EMP dataset). To replicate the

sampling effect due to rarefaction, we first created a vector of all individuals from a

single sample. We then selected 5000 individuals at random whose identities determined

which ASVs were found in that sample. These neutrally-derived random matrices, null

models 1 and 2, were plotted using the same plots (ASV:Genus vs number of genera) as

the real EMP dataset and were then analyzed using GLMMs with community diversity as

a predictor of focal lineage diversity (fixed effect), with lineage identity and EMP sample

ID as random effects. For Model 1, the slope was significantly negative (GLMM, Wald

test, $z=-9.807$, $P<2e-16$). For Model 2, the null GLMM (including the intercept only) was

significant, meaning that the community diversity has no significant effect on focal

lineages diversity (Likelihood-ratio test between the model with the predictor and the

intercept-only model, $P=0.9399$).

To generate a null model for a metacommunity assembled by niche processes,

null model 3 was made by sampling from a single Poisson distribution ($\lambda = 0.01$) for each

element of the data matrix. We used the Poisson distribution as a sensitivity analysis

compared to the ZSM, and found the two behave quite similarly (*i.e.* Model 1 and 3

produce qualitatively similar results). The probability of size zero was sufficiently large

that the down-sampling step was not needed for this model. Next, DBD and EC effects

were added to null model 3 according to the following procedure. An element was chosen

at random in a sample and tested if it is empty or full (*i.e.* checks the presence/absence of

a particular ASV). If the element is full then the DBD algorithm fills an empty element

chosen at random in the same sample, while the EC algorithm empties a filled element in

the same sample. This is to mimic the effect of DBD creating a niche for a new ASV, or

EC removing a niche based on the existing diversity. The strength of DBD or EC effects

were determined by the percent of elements tested. These data were analyzed with

GLMMs to test the power of our models to detect DBD or EC (**Table 2**, Supplementary

Data file 3 Section 2.1).

Rarefaction simulation

We constructed a simple simulation in which each microbiome sample may differ in total

diversity (*e.g.* in the observed range of genera) while maintaining a constant taxonomic

ratio (*e.g.* ASV:genus ratio = 2). To mimic rarefaction, we then sampled a set number of

sequencing reads from each synthetic community, assuming ASVs are sampled with

equal probability and plotted the observed taxonomic ratio (**Fig. 2 supplement 9**). This

simple simulation is implemented in permute_ASVs_synthetic.pl.

Nucleotide sequence-based analysis. We clustered ASVs at decreasing levels of

nucleotide identity, from 100% identical ASVs down to 75% identity (roughly equivalent

to phyla (Konstantinidis & Tiedje, 2005)). We estimated focal cluster diversity as the

mean number of descendants per cluster (e.g. number of 100% clusters per 97% cluster)

and plotted this against the total number of clusters (97% identity in this example). This

approach has the advantage of including sequences even if they come from unnamed

taxa. For each of the six nucleotide divergence ratios tested, the relationship between total

number of clusters and focal cluster diversity was positive (**Fig. 2 supplement 10**),

consistent with DBD and suggesting that the taxonomic analyses were qualitatively

unbiased.

Fasta files with all ASVs per sample were produced by a python script

(Construct_fasta_per_sample.py, Python Version 2.7) from the sequences summary file

(otu_summary.emp_deblur_90bp.subset_2k.rare_5000 from EMP ftp server). We

clustered sequences from each sample using USEARCH V9.2 and estimated sample

diversity as the total number of clusters at a given level (*e.g.* 97% identity) and focal

cluster diversity as the mean number of descendent clusters (*e.g.* number of 100%

clusters per 97% cluster). To describe the putative DBD or EC relationships, we tested

three models: linear, quadratic and cubic (lm function in R). Model comparisons were

based on the adjusted R^2 (**Figure 2 supplement 10**).

We note that diversity at level i (d_i) and at level $i+1$ (d_{i+1}/d_i) are not independent

in this analysis because d_{i+1} must be greater than or equal to d_i. To assess the effects of

this non-independence on the results, we conducted permutation tests by randomizing the

associations between d_i and d_{i+1}. Using 999 permutations, *P*-values were calculated based

on how many times we observed a correlation greater than that seen in the real data

(cor.test R function with kendall method). In each permutation, we recalculated the

significance test (Wald z) for the correlation in the randomized data, and then computed

the *P*-value based on how many times we observed a z value greater than that of the

original data. At all six levels of nucleotide identity, the real data always showed a

significantly stronger positive correlation when compared to permuted data ($P = 0.001$),

indicating that the DBD patterns was not an artefact of the dependence structure in the

data.

The effect of community diversity on focal cluster diversity was also tested across

different environments analyzed separately. We modelled this relationship with linear,

quadratic and cubic fits, and compared those models based on the adjusted R^2 (**Figure 2**

supplement 11).

DBD variation across environments

We tested the variation of focal lineage diversity slopes across different environments by

including EMPO 3 biome type as a fixed effect. We fitted a GLMM with the interaction

between community diversity and environment type as a predictor of focal lineage

diversity. All other random effects on intercept and slope were kept as in the previous

GLMMs (**Figure 3**, **Supplementary Data file 1 Section 3**). DBD variation across

environments was tested for Family:Order, Order:Class and Class:Phylum taxonomic

ratios, as diversity slope variation by environment was statistically significant

(likelihood-ratio test, $P<0.05$) for these ratios in the taxonomy based models (**Table 1**).

Abiotic effects

To test for the relative effect of biotic and abiotic environmental variables on focal

lineage diversity across different taxonomic ratios, we used a separate GLMM, with

Poisson error distribution and a log link function, for every ratio. We fitted the GLMM on

a subset (~10%) of the whole dataset, 192 samples (from water: saline (19) and non-

saline (44), surface: saline (42) and non-saline (19), sediment: saline (22) and non-saline

(31), soil (8) and plant rhizosphere (7)), for which measurements of four key abiotic

variables (temperature, pH, latitude and elevation) were available. As predictors of focal

lineage diversity (fixed effects), we included non-focal community diversity and abiotic

variables, as well as their interactions. All predictors were standardized to a mean of zero

and a standard deviation of one to obtain comparable estimates. The GLMM had the

same random effects as in the previous analysis, but only on the intercept for simplicity

(**Table 4**, **Supplementary file 1 section 4**).

Soil dataset analysis

We used the Delgado-Baquerizo et al. 2018 soil microbiome survey (237 samples from

18 countries) to further test the relative impacts of biotic versus abiotic drivers of

diversity. Raw data and abiotic measurements were downloaded from Figshare

(https://figshare.com/s/82a2d3f5d38ace925492; DOI: 10.6084/m9.figshare.5611321).

16S bioinformatic processing was performed using QIIME2 and Deblur with the same

protocol as in Thompson et al. 2017. Raw data 16S rRNA gene (V3-V4 region), were

processed by trimming the primers (341F/805R primer set) with qiime cutadapt trim-

paired, then merged using qiime vsearch join-pairs. Sequences were quality filtered and

denoised using Deblur with a trimming length of 400bp. The resulting 400-bp Deblur

BIOM table was filtered to keep only ASVs with at least 25 reads total over all samples

and rarefied to a depth of 5000. Taxonomy was assigned with a Naive Bayes classifier

trained on the V4-V3 region of 99% OTU Greengenes 13.8 sequences with qiime feature-

classifier. We obtained a final dataset of 186 samples and 24,252 ASVs which was used

as input for all statistical analysis as in the EMP dataset analysis. This data set included

14 environmental factors: aridity index (Aridity_Index), minimum and maximum

temperature (MINT and MAXT), precipitation seasonality (PSEA), mean diurnal

temperature range (MDR), ultra-violet (UV) radiation (UV_Light), net primary

productivity (NPP2003_2015), soil texture (Clay_silt), pH; total C (Soil_C), N (Soil_N)

and P (Soil_P) concentrations, C:N ratio (Soil_C_N_ratio) and Latitude.

We used a separate GLMM with Poisson error distribution and a log link function to test

for the effect of biotic (non-focal community diversity) and abiotic environmental

variables on focal lineage diversity (*e.g.* the ASV:Genus ratio for a focal genus), across

different taxonomic ratios. We defined non-focal taxa diversity and abiotic variables as

predictors (fixed effects) and the lineage identity as a random effect.

We also fitted the same model but with the first three principal components (PCs) from the

principal component analysis (PCA, rda function, vegan R package) of the abiotic variables

(a matrix of 237 samples (rows) by 14 abiotic variables (columns)), as well as the

interactions between diversity and each PC, and the interaction between PCs as predictors

(fixed effects).

Because of possible non-linear relationships between abiotic variables and diversity,

GLMMs were fitted with a linear and a quadratic term for every abiotic variable. The

quadratic terms were not significant, except for the ASV:genus ratio (**Table 5**; likelihood-

ratio test, $P < 2.2e-16$). The interaction terms were not significant except the interaction

between diversity and PCs at Family:Order ratio (likelihood-ratio test, $P= 2.182e-05$;

Table 5, **Supplementary file 4**).

Defining residents, generalists, and migrants. We defined a genus-level community

composition matrix as a matrix of 1128 genera (rows) by 17 environments (columns),

with the matrix entries indicating the percentage of samples from a given environment in

which each genus is present. We clustered the environmental samples based on their

genus-level community composition using fuzzy k-means clustering. The clustering

(cmeans function, package e1071 in R) was done on the 'hellinger' transformed data

(decostand function, vegan R package). To identify resident genera to each cluster, we

used indicator species analysis (Dufrene & Legendre, 1997) as implemented in the indval

function (labdsv R package). We defined residents as genera with indval indices between

0.4 and 0.9, with permutation test $P < 0.05$. Genera not associated with any cluster were

considered generalists. We used principal component analysis (PCA) on the community

composition matrix to visualize the clustering and the indicator genera (rda function,

vegan R package) (**Figure 4**). We then ran a separate GLMM for each environmental

cluster, with resident genus-level diversity (number of non-focal genera) as a predictor of

focal genus diversity (ASV:Genus ratio) for resident, migrant (residents of one cluster

found in a different cluster) and generalist genera. The fixed effect was specified as the

interaction between diversity and a factor defining the genus-cluster association (with

three levels: resident, migrant and generalist). Random effects on intercept and slope

were kept as in the GLMMs described above.

Genome size analysis. We chose a subset of genera represented by one or more

sequenced genomes in the NCBI microbial genomes database

(https://www.ncbi.nlm.nih.gov/genome/browse#!/prokaryotes/). For these genera, a

representative genome size was assigned by selecting the genome with the lowest number

of scaffolds (if no closed genomes were available) (**Supplementary file 6**). If multiple

genomes were available with the same level of completion, the largest genome size was

used, as smaller genomes could be artefacts of incomplete assembly which would bias the

mean and median downward. Moreover, given the deletional bias in bacterial genomes

(Kuo & Ochman, 2009a), the largest genome is likely more reflective of the ancestral

genome size of the genus. Only genera with two or more ASVs in at least one sample

were included in the analysis. Intracellular symbionts were excluded. We fitted a GLMM

on the subset of data with known genome size (576 genera, ranging from ~1 to 15 Mbp)

with the interaction between community diversity and genome size as a predictor of focal

lineage diversity at the ASV:Genus level. All the other random effects on intercept and

slope were kept as in the previous GLMMs (**Supplementary file 1 section 6**).

References

Alonso, D., & McKane, A. J. (2004). Sampling Hubbell's neutral theory of biodiversity:

Sampling neutral theory. *Ecology Letters*, *7*(10), 901–910.

https://doi.org/10.1111/j.1461-0248.2004.00640.x

Auguet, J.-C., Barberan, A., & Casamayor, E. O. (2010). Global ecological patterns in

uncultured Archaea. *The ISME Journal*, *4*(2), 182–190.

https://doi.org/10.1038/ismej.2009.109

Azaele, S., Suweis, S., Grilli, J., Volkov, I., Banavar, J. R., & Maritan, A. (2016).

Statistical mechanics of ecological systems: Neutral theory and beyond. *Reviews of*

Modern Physics, *88*(3), 035003. https://doi.org/10.1103/RevModPhys.88.035003

Bailey, S. F., Dettman, J. R., Rainey, P. B., & Kassen, R. (2013). Competition both drives

and impedes diversification in a model adaptive radiation. *Proceedings. Biological*

Sciences / The Royal Society, *280*(1766), 20131253.

https://doi.org/10.1098/rspb.2013.1253

Barberán, A., Ramirez, K. S., Leff, J. W., Bradford, M. a., Wall, D. H., & Fierer, N.

(2014). Why are some microbes more ubiquitous than others? Predicting the habitat

breadth of soil bacteria. *Ecology Letters*, *17*(7), 794–802.

https://doi.org/10.1111/ele.12282

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects

Models Using lme4. *Journal of Statistical Software*, *67*(1), 1–48.

https://doi.org/10.18637/jss.v067.i01

Brockhurst, M. A., Buckling, A., & Rainey, P. B. (2005). The effect of a bacteriophage

on diversification of the opportunistic bacterial pathogen, *Pseudomonas aeruginosa*.

Proceedings. Biological Sciences / The Royal Society, *272*(1570), 1385–1391.

https://doi.org/10.1098/rspb.2005.3086

Brockhurst, M. A., Colegrave, N., Hodgson, D. J., & Buckling, A. (2007). Niche

occupation limits adaptive radiation in experimental microcosms. *PloS One*, *2*(2),

e193. https://doi.org/10.1371/journal.pone.0000193

Calcagno, V., Jarne, P., Loreau, M., Mouquet, N., & David, P. (2017). Diversity spurs

diversification in ecological communities. *Nature Communications*, *8*, 15810.

https://doi.org/10.1038/ncomms15810

Coyte, K. Z., Schluter, J., & Foster, K. R. (2015). The ecology of the microbiome:

Networks, competition, and stability. *Science*, *350*(6261), 663–666.

https://doi.org/10.1126/science.aad2602

Czárán, T. L., Hoekstra, R. F., & Pagie, L. (2002). Chemical warfare between microbes

promotes biodiversity. *Proceedings of the National Academy of Sciences of the*

United States of America, *99*(2), 786–790. https://doi.org/10.1073/pnas.012399899

Delgado-Baquerizo, M., Oliverio, A. M., Brewer, T. E., Benavent-González, A.,

Eldridge, D. J., Bardgett, R. D., Maestre, F. T., Singh, B. K., & Fierer, N. (2018). A

global atlas of the dominant bacteria found in soil. *Science*, *359*(6373), 320–325.

https://doi.org/10.1126/science.aap9516

de Wit, R., & Bouvier, T. (2006). "Everything is everywhere, but, the environment

selects"; what did Baas Becking and Beijerinck really say? *Environmental*

Microbiology, *8*(4), 755–758. https://doi.org/10.1111/j.1462-2920.2006.01017.x

Dufrene, M., & Legendre, P. (1997). Species Assemblages and Indicator Species: The

Need for a Flexible Asymmetrical Approach. *Ecological Monographs*, *67*(3), 345–

366. https://doi.org/10.2307/2963459

Elston, D. A., Moss, R., Boulinier, T., Arrowsmith, C., & Lambin, X. (2001). Analysis of

aggregation, a worked example: numbers of ticks on red grouse chicks.

Parasitology, *122*(Pt 5), 563–569. https://doi.org/10.1017/s0031182001007740

Elton, C. (1946). Competition and the Structure of Ecological Communities. *The Journal

of Animal Ecology*, *15*(1), 54–68. https://doi.org/10.2307/1625

Emerson, B. C., & Kolm, N. (2005). Species diversity can drive speciation. *Nature*,

434(7036), 1015–1017. https://doi.org/10.1038/nature03450

Falkowski, P. G., Fenchel, T., & Delong, E. F. (2008). The microbial engines that drive

Earth's biogeochemical cycles. *Science*, *320*(5879), 1034–1039.

https://doi.org/10.1126/science.1153213

Gause, G. F. (2003). *The Struggle for Existence* (Williams & Wilkins, Baltimore, 1934).

Gómez, P., & Buckling, A. (2013). Real-time microbial adaptive diversification in soil.

Ecology Letters, *16*(5), 650–655. https://doi.org/10.1111/ele.12093

Gotelli, N. J., & Colwell, R. K. (2001). Quantifying biodiversity: procedures and pitfalls

in the measurement and comparison of species richness. *Ecology Letters*, *4*(4), 379–

391. https://doi.org/10.1046/j.1461-0248.2001.00230.x

Gotelli, N. J., & McGill, B. J. (2006). Null Versus Neutral Models: What's The

Difference? *Ecography*, *29*(5), 793–800. https://doi.org/10.1111/j.2006.0906-

7590.04714.x

Harris, K., Parsons, T. L., Ijaz, U. Z., Lahti, L., Holmes, I., & Quince, C. (2017). Linking

Statistical and Ecological Theory: Hubbell's Unified Neutral Theory of Biodiversity

as a Hierarchical Dirichlet Process. *Proceedings of the IEEE*, *105*(3), 516–529.

https://doi.org/10.1109/JPROC.2015.2428213

Hibbing, M. E., Fuqua, C., Parsek, M. R., & Peterson, S. B. (2010). Bacterial

competition: surviving and thriving in the microbial jungle. *Nature Reviews*

Microbiology, *8*(1), 15–25. https://doi.org/10.1038/nrmicro2259

Hubbell, S. P. (2001). *The Unified Neutral Theory of Biodiversity and Biogeography*.

Princeton University Press.

Hug, L. A., Baker, B. J., Anantharaman, K., Brown, C. T., Probst, A. J., Castelle, C. J.,

Butterfield, C. N., Hernsdorf, A. W., Amano, Y., Kotaro, I., Suzuki, Y., Dudek, N.,

Relman, D. A., Finstad, K. M., Amundson, R., Thomas, B. C., & Banfield, J. F.

(2016). A new view of the tree and life's diversity. *Nature Microbiology*, 1, 16048

https://doi.org/10.1038/nmicrobiol.2016.48

Jarvinen, O. (1982). Species-To-Genus Ratios in Biogeography: A Historical Note.

Journal of Biogeography, *9*(4), 363–370. https://doi.org/10.2307/2844723

Johnson, P. C. (2014). Extension of Nakagawa & Schielzeth's R2GLMM to random

slopes models. *Methods in Ecology and Evolution / British Ecological Society*, *5*(9),

944–946. https://doi.org/10.1111/2041-210X.12225

Jousset, A., Eisenhauer, N., Merker, M., Mouquet, N., & Scheu, S. (2016). High

functional diversity stimulates diversification in experimental microbial

communities. *Science Advances*, *2*(6), e1600124.

https://doi.org/10.1126/sciadv.1600124

Kastman, E. K., Kamelamela, N., Norville, J. W., Cosetta, C. M., Dutton, R. J., & Wolfe,

B. E. (2016). Biotic Interactions Shape the Ecological Distributions of

Staphylococcus Species. *mBio*, *7*(5). https://doi.org/10.1128/mBio.01157-16

1014 Kennedy, A. C., & de Luna, L. Z. (2005). Rhizhosphere. In D. Hillel (Ed.), *Encyclopedia*

1015   *of Soils in the Environment* (pp. 399–406). Elsevier. https://doi.org/10.1016/B0-12-

1016   348530-4/00163-6

1017 Konstantinidis, K. T., & Tiedje, J. M. (2005). Towards a genome-based taxonomy for

1018   prokaryotes. *Journal of Bacteriology*, *187*(18), 6258–6264.

1019 Kuo, C.-H., & Ochman, H. (2009a). Deletional bias across the three domains of life.

1020   *Genome Biology and Evolution*, *1*, 145–152. https://doi.org/10.1093/gbe/evp016

1021 Kuo, C.-H., & Ochman, H. (2009b). Inferring clocks when lacking rocks: the variable

1022   rates of molecular evolution in bacteria. *Biology Direct*, *4*, 35.

1023   https://doi.org/10.1186/1745-6150-4-35

1024 Laland, K. N., Odling-Smee, F. J., & Feldman, M. W. (1999). Evolutionary consequences

1025   of niche construction and their implications for ecology. *Proceedings of the National*

1026   *Academy of Sciences of the United States of America*, *96*(18), 10242–10247.

1027   https://doi.org/10.1073/pnas.96.18.10242

1028 Lapierre, P., & Gogarten, J. P. (2009). Estimating the size of the bacterial pan-genome.

1029   *Trends in Genetics: TIG*, *25*(3), 107–110. https://doi.org/10.1016/j.tig.2008.12.004

1030 Lauber, C. L., Hamady, M., Knight, R., & Fierer, N. (2009). Soil pH as a predictor of soil

1031   bacterial community structure at the continental scale: a pyrosequencing-based

1032   assessment. *Applied and Environmental Microbiology*. 75, 5111-5120.

1033   http://aem.asm.org/content/early/2009/06/05/AEM.00335-09.short

1034 Li, L., & Ma, Z. S. (2016). Testing the Neutral Theory of Biodiversity with Human

1035   Microbiome Datasets. *Scientific Reports*, *6*, 31448.

1036   https://doi.org/10.1038/srep31448

Lindow, S. E., & Brandl, M. T. (2003). Microbiology of the phyllosphere. *Applied and*

Environmental Microbiology, *69*(4), 1875–1883.

https://doi.org/10.1128/aem.69.4.1875-1883.2003

Louca, S., Mazel, F., Doebeli, M., & Parfrey, L. W. (2019). A census-based estimate of

Earth's bacterial and archaeal diversity. *PLoS Biology*, *17*(2), e3000106.

https://doi.org/10.1371/journal.pbio.3000106

Louca, S., & Pennell, M. W. (2020). Extant timetrees are consistent with a myriad of

diversification histories. *Nature*, *580*(7804), 502–505.

https://doi.org/10.1038/s41586-020-2176-1

Louca, S., Shih, P. M., Pennell, M. W., Fischer, W. W., Parfrey, L. W., & Doebeli, M.

(2018). Bacterial diversification through geological time. *Nature Ecology &*

Evolution, *2*(9), 1458–1467. https://doi.org/10.1038/s41559-018-0625-0

Lozupone, C. A., & Knight, R. (2007). Global patterns in bacterial diversity. *Proceedings*

of the National Academy of Sciences of the United States of America, *104*(27),

11436–11440. https://doi.org/10.1073/pnas.0611525104

Marshall, C. R. (2017). Five palaeobiological laws needed to understand the evolution of

the living biota. *Nature Ecology & Evolution*, *1*(6), 165.

https://doi.org/10.1038/s41559-017-0165

Meyer, J. R., & Kassen, R. (2007). The effects of competition and predation on

diversification in a model adaptive radiation. *Nature*, *446*(7134), 432–435.

http://www.nature.com/doifinder/10.1038/nature05599

Morris, J. J., & Lenski, R. E. (2012). The Black Queen Hypothesis: evolution of

dependencies through adaptive gene loss. *mBio*. 3, e00036-12

http://mbio.asm.org/content/3/2/e00036-12.short

Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining R 2

from generalized linear mixed-effects models. *Methods in Ecology and Evolution /*

British Ecological Society, *4*(2), 133–142. https://doi.org/10.1111/j.2041-

210x.2012.00261.x

Needham, D. M., & Fuhrman, J. A. (2016). Pronounced daily succession of

phytoplankton , archaea and bacteria following a spring bloom. *Nature*

Microbiology, 1, 16005. https://doi.org/10.1038/NMICROBIOL.2016.5

Palmer, M. W., & Maurer, T. A. (1997). Does Diversity Beget Diversity? A Case Study

of Crops and Weeds. *Journal of Vegetation Science*, *8*(2), 235–240.

https://doi.org/10.2307/3237352

Parks, D. H., Chuvochina, M., Waite, D. W., Rinke, C., Skarshewski, A., Chaumeil, P.-

A., & Hugenholtz, P. (2018). A standardized bacterial taxonomy based on genome

phylogeny substantially revises the tree of life. *Nature Biotechnology*, *36*(10), 996–

1004. https://doi.org/10.1038/nbt.4229

Pennekamp, F., Pontarp, M., Tabi, A., Altermatt, F., Alther, R., Choffat, Y., Fronhofer, E.

A., Ganesanandamoorthy, P., Garnier, A., Griffiths, J. I., Greene, S., Horgan, K.,

Massie, T. M., Mächler, E., Palamara, G. M., Seymour, M., & Petchey, O. L.

(2018). Biodiversity increases and decreases ecosystem stability. *Nature*, *563*(7729),

109–112. https://doi.org/10.1038/s41586-018-0627-8

Power, J. F., Carere, C. R., Lee, C. K., Wakerley, G. L. J., Evans, D. W., Button, M.,

White, D., Climo, M. D., Hinze, A. M., Morgan, X. C., McDonald, I. R., Cary, S.

C., & Stott, M. B. (2018). Microbial biogeography of 925 geothermal springs in

New Zealand. *Nature Communications*, *9*(1), 2876. https://doi.org/10.1038/s41467-

018-05020-y

Price, T. D., Hooper, D. M., Buchanan, C. D., Johansson, U. S., Tietze, D. T., Alström,

P., Olsson, U., Ghosh-Harihar, M., Ishtiaq, F., Gupta, S. K., Martens, J., Harr, B.,

Singh, P., & Mohan, D. (2014). Niche filling slows the diversification of Himalayan

songbirds. *Nature*. 509, 222-225 https://doi.org/10.1038/nature13272

Rabosky, D. L., Chang, J., Title, P. O., Cowman, P. F., Sallan, L., Friedman, M.,

Kaschner, K., Garilao, C., Near, T. J., Coll, M., & Alfaro, M. E. (2018). An inverse

latitudinal gradient in speciation rate for marine fishes. *Nature*, *559*(7714), 392–395.

https://doi.org/10.1038/s41586-018-0273-1

Rabosky, D. L., & Hurlbert, A. H. (2015). Species richness at continental scales is

dominated by ecological limits. *The American Naturalist*, *185*(5), 572–583.

https://doi.org/10.1086/680850

San Roman, M., & Wagner, A. (2018). An enormous potential for niche construction

through bacterial cross-feeding in a homogeneous environment. *PLoS*

Computational Biology, *14*(7), e1006340.

https://doi.org/10.1371/journal.pcbi.1006340

Schluter, D., & Pennell, M. W. (2017). Speciation gradients and the distribution of

biodiversity. *Nature*, *546*(7656), 48–55. https://doi.org/10.1038/nature22897

Sender, R., Fuchs, S., & Milo, R. (2016). Revised Estimates for the Number of Human

and Bacteria Cells in the Body. *PLoS Biology*, *14*(8), e1002533.

https://doi.org/10.1371/journal.pbio.1002533

Seth, E. C., & Taga, M. E. (2014). Nutrient cross-feeding in the microbial world.

Frontiers in Microbiology, *5*, 350. https://doi.org/10.3389/fmicb.2014.00350

Sogin, M. L., Morrison, H. G., Huber, J. A., Welch, D. M., Huse, S. M., Neal, P. R.,

Arrieta, J. M., & Herndl, G. J. (2006). Microbial diversity in the deep sea and the

underexplored "rare biosphere." *Proceedings of the National Academy of Sciences of*

the United States of America, *103*(32), 12115–12120.

Sriswasdi, S., Yang, C.-C., & Iwasaki, W. (2017). Generalist species drive microbial

dispersion and evolution. *Nature Communications*, *8*(1), 1162.

https://doi.org/10.1038/s41467-017-01265-1

Sunagawa, S., Coelho, L. P., Chaffron, S., Kultima, J. R., Labadie, K., Salazar, G.,

Djahanschiri, B., Zeller, G., Mende, D. R., Alberti, A., Cornejo-Castillo, F. M.,

Costea, P. I., Cruaud, C., d'Ovidio, F., Engelen, S., Ferrera, I., Gasol, J. M., Guidi,

L., Hildebrand, F., … Bork, P. (2015). Ocean plankton. Structure and function of the

global ocean microbiome. *Science*, *348*(6237), 1261359.

https://doi.org/10.1126/science.1261359

Thompson, L. R., Sanders, J. G., McDonald, D., Amir, A., Ladau, J., Locey, K. J., Prill,

R. J., Tripathi, A., Gibbons, S. M., Ackermann, G., Navas-Molina, J. A., Janssen, S.,

Kopylova, E., Vázquez-Baeza, Y., González, A., Morton, J. T., Mirarab, S., Xu, Z.

Z., Jiang, L., … Zhao, H. (2017). A communal catalogue reveals Earth's multiscale

microbial diversity. *Nature*, 551, 457-463. https://doi.org/10.1038/nature24621

Vos, M. (2011). A species concept for bacteria based on adaptive divergence. *Trends in*

Microbiology, *19*(1), 1–7. https://doi.org/10.1016/j.tim.2010.10.003

Whitman, W. B., Coleman, D. C., & Wiebe, W. J. (1998). Prokaryotes: the unseen

majority. *Proceedings of the National Academy of Sciences of the United States of*

1129   *America*, *95*(12), 6578–6583. https://doi.org/10.1073/pnas.95.12.6578

1130   Whittaker, R. H. (1972). Evolution and Measurement of Species Diversity. *Taxon*,

1131   *21*(2/3), 213–251. https://doi.org/10.2307/1218190